

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(3-Phenylisoquinolin-1-yl)hydrazine

 P. Manivel,^a Venkatesha R. Hathwar,^b P. Nithya,^a
 K. Prabakaran^a and F. Nawaz Khan^{a*}
^aChemistry Division, School of Science and Humanities, VIT University, Vellore 632 014, Tamil Nadu, India, and ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India

Correspondence e-mail: nawaz_f@yahoo.co.in


Received 17 November 2008; accepted 10 December 2008

 Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.047; wR factor = 0.116; data-to-parameter ratio = 10.6.

The title compound, $\text{C}_{15}\text{H}_{13}\text{N}_3$, contains two independent molecules in the asymmetric unit. The isoquinoline moiety and phenyl rings form dihedral angles of 4.38 (2) and 10.14 (3)° in the two independent molecules. The crystal packing is stabilized by $\text{N}-\text{H}\cdots\text{N}$ molecular dimers formed across a center of symmetry.

Related literature

For general background to hydrazine compounds, see: Broadhurst *et al.* (2001); Behrens (1999); Broadhurst (1991); Chao *et al.* (1999); Kametani (1968). For related crystal structures, see: Yang *et al.* (2008); Choudhury & Guru Row (2006); Choudhury *et al.* (2002); Hathwar *et al.* (2008). For bond-length data, see: Allen *et al.* (1998). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{13}\text{N}_3$
 $M_r = 235.28$
 Triclinic, $P\bar{1}$
 $a = 6.672$ (2) Å
 $b = 13.825$ (4) Å
 $c = 14.934$ (5) Å
 $\alpha = 63.836$ (5)°
 $\beta = 86.895$ (6)°

 $\gamma = 82.106$ (5)°
 $V = 1224.5$ (7) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 290$ (2) K
 $0.15 \times 0.12 \times 0.05$ mm

Data collection

 Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.953$, $T_{\max} = 0.996$
 12381 measured reflections
 4546 independent reflections
 2926 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.116$
 $S = 1.02$
 4546 reflections
 429 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.14$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2'-\text{H}2'N\cdots\text{N}3^{\text{ii}}$	0.91 (2)	2.15 (2)	2.967 (2)	151 (2)
$\text{N}2-\text{H}2N\cdots\text{N}3^{\text{ii}}$	0.90 (2)	2.20 (2)	3.027 (2)	152 (2)
$\text{N}3'-\text{H}3'B\cdots\text{N}1^{\text{iii}}$	0.89 (2)	2.24 (2)	3.119 (2)	169 (2)
$\text{N}3-\text{H}3A\cdots\text{N}1^{\text{iv}}$	0.92 (2)	2.26 (2)	3.170 (3)	168 (2)

 Symmetry codes: (i) $-x, -y, -z$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y, -z$; (iv) $-x+2, -y, -z+1$.

Data collection: SMART (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1999) and CAMERON (Watkin *et al.*, 1993); software used to prepare material for publication: PLATON (Spek, 2003).

We thank the Department of Science and Technology, India, for use of the CCD facility set up under the IRHPA-DST program at IISc. We thank Prof T. N. Guru Row, IISc, Bangalore, for useful crystallographic discussions. FNK thanks the DST for Fast Track Proposal funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CS2101).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1998). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Behrens, C. H. (1999). US Patent 4 942 163.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Broadhurst, M. D. (1991). US Patent 5 070 097.
- Broadhurst, M. D., Michael, J. J., William, H. W. & Daryl, S. (2001). US Patent 6 235 787.
- Bruker (2004). SMART and SAINT Bruker AXS Inc., Madison, Wisconsin, USA.
- Chao, Q., Deng, L., Shih, H., Leoni, L. M., Genini, D., Carson, D. A. & Cottam, H. B. (1999). *J. Med. Chem.* **2**, 3860–3873.
- Choudhury, A. R. & Guru Row, T. N. (2006). *CrystEngComm* **8**, 265–274.
- Choudhury, A. R., Urs, U. K., Guru Row, T. N. & Nagarajan, K. (2002). *J. Mol. Struct.* **605**, 71–77.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hathwar, V. R., Prabakaran, K., Subashini, R., Manivel, P. & Khan, F. N. (2008). *Acta Cryst. E* **64**, o2295.
- Kametani, T. (1968). *The Chemistry of the Isoquinoline Alkaloids*. Tokyo: Hirokawa and Amsterdam: Elsevier.

Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

Watkin, D. J., Pearce, L. & Prout, C. K. (1993). *CAMERON*. Chemical
Crystallography Laboratory, University of Oxford, England.
Yang, Y., Yang, P., Zhang, C. & Wu, B. (2008). *Anal. Sci.* **24**, x97–x98.

supporting information

Acta Cryst. (2009). E65, o137–o138 [doi:10.1107/S1600536808042062]

1-(3-Phenylisoquinolin-1-yl)hydrazine

P. Manivel, Venkatesha R. Hathwar, P. Nithya, K. Prabakaran and F. Nawaz Khan

S1. Comment

The title compound belongs to the class isoquinolines. Isoquinolines and isoquinolinones are an integral part of many naturally occurring fused heterocycles and find applications in synthetic and pharmaceutical chemistry (Kametani *et al.*, 1968). Isoquinolinones and isoquinoline amines were reported as cancer chemotherapeutic agents (Behrens, 1999) whereas quinolyl and isoquinolyl derivatives have been reported as insecticidal compounds (Broadhurst, 1991). 3-Substituted isoquinolines are of potent use in medicine (Chao, *et al.*, 1999) and in general, hydrazine derivatives can be used as medicaments (Broadhurst *et al.*, 2001). Choudhury, *et al.* (2002, 2006) reported crystal structures of substituted isoquinolines while Hathwar, *et al.* (2008) reports the crystal structure of an isoquinolinyl diselenide.


The asymmetric unit of the crystal structure of the title compound contains two independent molecules (Fig. 1). The isoquinoline moiety and phenyl rings form dihedral angles of 4.38 (2) and 10.14 (3)°, respectively, in the two independent molecules. All bond lengths and angles are normal (Allen *et al.*, 1998). The packing (Fig. 2) is consolidated by four N—H···N hydrogen bonds. All the four N—H···N hydrogen bonds generate dimers across centres of symmetry (Table 1) resulting in tight molecular packing in the crystal. The N2'—H2'N···N3' and the N2—H2N···N3 hydrogen bonds form a $R^2_2(6)$ motif whereas the N3'—H3'B···N1' and the N3—H3A···N1 hydrogen bond dimers form a $R^2_2(10)$ motif (Bernstein *et al.*, 1995) in the crystal (Fig. 2).

S2. Experimental


The solution of 1-chloro-3-phenylisoquinoline in ethanol was treated with hydrazine hydrate and stirred at 323 K for 3hr. The product was filtered. The solid was washed with water and diethyl ether and dried under vacuum. Single crystals of the title compound were obtained via recrystallization from a dichloromethane solution.

S3. Refinement

All the H atoms in the title compound were located from difference electron density maps and refined isotropically resulting in C—H and N—H bond lengths of 0.91 (4) - 1.02 (2) Å and 0.89 (2) - 0.97 (3) Å, respectively.

**Figure 1**

ORTEP diagram of the asymmetric unit of (I) with 50% probability displacement ellipsoids.

**Figure 2**

A packing excerpt from the crystal with dotted lines indicating intermolecular N—H...N hydrogen bonds. H atoms not involved in the interactions are omitted for clarity.

1-(3-Phenylisoquinolin-1-yl)hydrazine

Crystal data

$C_{15}H_{13}N_3$
 $M_r = 235.28$
 Triclinic, $P\bar{1}$
 Hall symbol: $-P\ 1$
 $a = 6.672\ (2)\ \text{\AA}$
 $b = 13.825\ (4)\ \text{\AA}$
 $c = 14.934\ (5)\ \text{\AA}$
 $\alpha = 63.836\ (5)^\circ$
 $\beta = 86.895\ (6)^\circ$
 $\gamma = 82.106\ (5)^\circ$
 $V = 1224.5\ (7)\ \text{\AA}^3$

$Z = 4$
 $F(000) = 496$
 $D_x = 1.276\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 832 reflections
 $\theta = 1.7\text{--}25.3^\circ$
 $\mu = 0.08\ \text{mm}^{-1}$
 $T = 290\ \text{K}$
 Needle, colourless
 $0.15 \times 0.12 \times 0.05\ \text{mm}$

Data collection

Bruker SMART CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans

Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.953$, $T_{\max} = 0.996$
 12381 measured reflections
 4546 independent reflections

2926 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 1.5^\circ$

$h = -8 \rightarrow 8$
 $k = -16 \rightarrow 16$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.116$
 $S = 1.02$
 4546 reflections
 429 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0567P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.14 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.91992 (19)	0.17183 (10)	0.48294 (10)	0.0386 (3)
N2	0.6714 (2)	0.05976 (11)	0.52799 (11)	0.0433 (4)
N3	0.7300 (2)	0.02645 (15)	0.45200 (14)	0.0471 (4)
N1'	0.4205 (2)	0.13567 (10)	0.03997 (10)	0.0391 (3)
N2'	0.1689 (2)	0.07962 (11)	-0.01722 (11)	0.0466 (4)
N3'	0.2325 (2)	-0.03027 (12)	0.04958 (15)	0.0495 (4)
C1	0.7552 (2)	0.14107 (12)	0.53468 (12)	0.0366 (4)
C2	1.0076 (2)	0.25388 (13)	0.48758 (12)	0.0406 (4)
C3	0.9290 (3)	0.30370 (15)	0.54472 (14)	0.0529 (5)
C4	0.6656 (4)	0.32080 (18)	0.66201 (16)	0.0718 (7)
C5	0.4979 (4)	0.28801 (19)	0.71627 (17)	0.0768 (7)
C6	0.4084 (4)	0.20448 (17)	0.71345 (16)	0.0666 (6)
C7	0.4885 (3)	0.15463 (16)	0.65585 (14)	0.0515 (5)
C8	0.6624 (2)	0.18705 (12)	0.59912 (12)	0.0395 (4)
C9	0.7543 (3)	0.27114 (14)	0.60246 (13)	0.0482 (5)
C10	1.1910 (2)	0.28398 (13)	0.42535 (12)	0.0431 (4)
C11	1.2553 (3)	0.23496 (16)	0.36341 (13)	0.0523 (5)
C12	1.4255 (3)	0.26016 (18)	0.30579 (15)	0.0639 (6)
C13	1.5362 (4)	0.33508 (19)	0.30874 (17)	0.0679 (6)
C14	1.4761 (3)	0.38509 (18)	0.36893 (17)	0.0655 (6)
C15	1.3048 (3)	0.36044 (16)	0.42676 (15)	0.0561 (5)
C1'	0.2527 (2)	0.16171 (13)	-0.01268 (12)	0.0385 (4)
C2'	0.5085 (2)	0.21556 (13)	0.04790 (12)	0.0406 (4)
C3'	0.4252 (3)	0.32116 (15)	0.00313 (14)	0.0531 (5)
C4'	0.1531 (4)	0.45988 (17)	-0.10239 (18)	0.0780 (7)
C5'	-0.0202 (4)	0.48553 (19)	-0.15637 (19)	0.0876 (8)
C6'	-0.1088 (4)	0.40594 (18)	-0.16610 (16)	0.0735 (7)
C7'	-0.0217 (3)	0.30071 (16)	-0.12234 (14)	0.0547 (5)
C8'	0.1569 (3)	0.27129 (13)	-0.06543 (12)	0.0411 (4)
C9'	0.2471 (3)	0.35185 (14)	-0.05511 (13)	0.0502 (5)
C10'	0.6945 (2)	0.17691 (14)	0.11012 (12)	0.0412 (4)

C11'	0.7519 (3)	0.06689 (16)	0.16619 (14)	0.0525 (5)
C12'	0.9251 (3)	0.02867 (18)	0.22499 (15)	0.0599 (5)
C13'	1.0450 (3)	0.09997 (19)	0.22820 (15)	0.0587 (5)
C14'	0.9911 (3)	0.2093 (2)	0.17264 (15)	0.0608 (6)
C15'	0.8188 (3)	0.24744 (17)	0.11461 (15)	0.0532 (5)
H2N	0.542 (3)	0.0483 (14)	0.5454 (13)	0.060 (6)*
H3	0.992 (3)	0.3609 (14)	0.5471 (12)	0.056 (5)*
H3A	0.843 (3)	-0.0248 (16)	0.4726 (14)	0.075 (7)*
H3B	0.761 (3)	0.0880 (17)	0.3956 (15)	0.078 (7)*
H4	0.733 (3)	0.3797 (16)	0.6607 (14)	0.082 (7)*
H5	0.438 (3)	0.3227 (17)	0.7544 (16)	0.094 (7)*
H2'N	0.038 (3)	0.0910 (13)	-0.0365 (12)	0.053 (5)*
H6	0.286 (3)	0.1827 (16)	0.7500 (15)	0.084 (7)*
H3'A	0.269 (3)	-0.0320 (14)	0.1126 (14)	0.065 (6)*
H3'B	0.341 (3)	-0.0536 (16)	0.0238 (15)	0.076 (7)*
H7	0.423 (3)	0.1003 (14)	0.6516 (12)	0.055 (5)*
H11	1.172 (3)	0.1806 (14)	0.3645 (13)	0.062 (5)*
H12	1.470 (3)	0.2194 (16)	0.2639 (15)	0.090 (7)*
H13	1.658 (3)	0.3473 (17)	0.2734 (16)	0.095 (8)*
H14	1.549 (3)	0.4356 (15)	0.3753 (13)	0.069 (6)*
H15	1.263 (3)	0.3971 (14)	0.4688 (13)	0.059 (6)*
H3'	0.492 (3)	0.3731 (15)	0.0110 (13)	0.064 (6)*
H4'	0.219 (3)	0.5136 (17)	-0.0959 (14)	0.082 (7)*
H5'	-0.082 (4)	0.5558 (19)	-0.1851 (17)	0.102 (8)*
H6'	-0.234 (3)	0.4247 (16)	-0.2025 (15)	0.080 (7)*
H7'	-0.081 (3)	0.2459 (15)	-0.1303 (13)	0.065 (6)*
H11'	0.661 (3)	0.0177 (14)	0.1642 (12)	0.057 (5)*
H12'	0.964 (3)	-0.0520 (18)	0.2615 (15)	0.092 (7)*
H13'	1.164 (3)	0.0723 (15)	0.2692 (14)	0.070 (6)*
H14'	1.073 (3)	0.2590 (16)	0.1730 (14)	0.078 (6)*
H15'	0.786 (3)	0.3249 (15)	0.0727 (14)	0.069 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0333 (8)	0.0399 (8)	0.0440 (8)	-0.0050 (6)	-0.0015 (6)	-0.0194 (7)
N2	0.0385 (9)	0.0477 (9)	0.0538 (9)	-0.0105 (7)	0.0057 (7)	-0.0306 (8)
N3	0.0416 (9)	0.0510 (10)	0.0609 (11)	-0.0049 (8)	0.0013 (8)	-0.0358 (9)
N1'	0.0349 (8)	0.0393 (8)	0.0494 (8)	-0.0077 (6)	0.0027 (7)	-0.0245 (7)
N2'	0.0365 (9)	0.0386 (9)	0.0688 (11)	-0.0028 (7)	-0.0065 (8)	-0.0271 (8)
N3'	0.0420 (10)	0.0382 (9)	0.0720 (12)	-0.0014 (7)	-0.0050 (9)	-0.0281 (9)
C1	0.0341 (9)	0.0347 (9)	0.0398 (9)	-0.0008 (7)	-0.0057 (8)	-0.0155 (8)
C2	0.0388 (10)	0.0401 (10)	0.0416 (10)	-0.0052 (8)	-0.0054 (8)	-0.0161 (8)
C3	0.0572 (12)	0.0539 (12)	0.0606 (12)	-0.0214 (10)	0.0080 (10)	-0.0337 (10)
C4	0.0929 (18)	0.0715 (15)	0.0772 (15)	-0.0321 (13)	0.0289 (13)	-0.0532 (13)
C5	0.0979 (19)	0.0770 (16)	0.0769 (16)	-0.0248 (14)	0.0364 (14)	-0.0532 (14)
C6	0.0735 (15)	0.0659 (14)	0.0659 (14)	-0.0203 (12)	0.0312 (12)	-0.0341 (12)
C7	0.0539 (12)	0.0495 (11)	0.0541 (12)	-0.0111 (10)	0.0095 (9)	-0.0250 (10)

C8	0.0416 (10)	0.0374 (9)	0.0368 (9)	-0.0013 (8)	-0.0019 (8)	-0.0145 (8)
C9	0.0549 (12)	0.0462 (11)	0.0496 (11)	-0.0100 (9)	0.0041 (9)	-0.0259 (9)
C10	0.0393 (10)	0.0408 (10)	0.0415 (10)	-0.0048 (8)	-0.0055 (8)	-0.0106 (8)
C11	0.0525 (12)	0.0569 (12)	0.0458 (11)	-0.0127 (10)	0.0048 (9)	-0.0198 (10)
C12	0.0592 (14)	0.0723 (15)	0.0525 (13)	-0.0113 (11)	0.0097 (10)	-0.0205 (12)
C13	0.0505 (14)	0.0723 (15)	0.0578 (14)	-0.0092 (12)	0.0081 (11)	-0.0081 (12)
C14	0.0508 (13)	0.0575 (13)	0.0733 (15)	-0.0191 (11)	-0.0029 (12)	-0.0115 (12)
C15	0.0510 (12)	0.0508 (12)	0.0652 (14)	-0.0133 (10)	0.0005 (10)	-0.0223 (11)
C1'	0.0354 (10)	0.0413 (10)	0.0454 (10)	-0.0068 (8)	0.0065 (8)	-0.0252 (8)
C2'	0.0408 (10)	0.0411 (10)	0.0447 (10)	-0.0115 (8)	0.0077 (8)	-0.0223 (8)
C3'	0.0611 (13)	0.0420 (11)	0.0600 (12)	-0.0146 (10)	-0.0037 (10)	-0.0230 (10)
C4'	0.1021 (19)	0.0388 (12)	0.0861 (17)	-0.0059 (12)	-0.0263 (15)	-0.0189 (12)
C5'	0.115 (2)	0.0430 (14)	0.0886 (18)	0.0131 (14)	-0.0384 (16)	-0.0160 (13)
C6'	0.0877 (18)	0.0566 (14)	0.0677 (15)	0.0104 (13)	-0.0303 (13)	-0.0215 (12)
C7'	0.0602 (13)	0.0500 (12)	0.0537 (12)	-0.0013 (10)	-0.0095 (10)	-0.0231 (10)
C8'	0.0450 (10)	0.0400 (10)	0.0381 (10)	-0.0051 (8)	0.0030 (8)	-0.0174 (8)
C9'	0.0620 (13)	0.0359 (10)	0.0505 (11)	-0.0080 (9)	-0.0019 (10)	-0.0161 (9)
C10'	0.0406 (10)	0.0492 (11)	0.0424 (10)	-0.0116 (8)	0.0061 (8)	-0.0269 (9)
C11'	0.0573 (13)	0.0514 (12)	0.0513 (12)	-0.0153 (10)	-0.0051 (10)	-0.0217 (10)
C12'	0.0654 (14)	0.0609 (14)	0.0523 (12)	-0.0064 (11)	-0.0107 (10)	-0.0232 (11)
C13'	0.0533 (13)	0.0777 (16)	0.0502 (12)	-0.0078 (12)	-0.0060 (10)	-0.0322 (12)
C14'	0.0571 (13)	0.0779 (16)	0.0612 (13)	-0.0257 (12)	-0.0016 (11)	-0.0380 (13)
C15'	0.0556 (12)	0.0543 (13)	0.0575 (13)	-0.0162 (10)	-0.0002 (10)	-0.0288 (11)

Geometric parameters (Å, °)

N1—C1	1.3163 (19)	C12—C13	1.369 (3)
N1—C2	1.3748 (19)	C12—H12	1.02 (2)
N2—C1	1.3652 (19)	C13—C14	1.371 (3)
N2—N3	1.420 (2)	C13—H13	0.94 (2)
N2—H2N	0.904 (18)	C14—C15	1.381 (3)
N3—H3A	0.920 (19)	C14—H14	0.94 (2)
N3—H3B	0.93 (2)	C15—H15	0.976 (17)
N1'—C1'	1.317 (2)	C1'—C8'	1.435 (2)
N1'—C2'	1.3712 (19)	C2'—C3'	1.358 (2)
N2'—C1'	1.360 (2)	C2'—C10'	1.481 (2)
N2'—N3'	1.420 (2)	C3'—C9'	1.413 (3)
N2'—H2'N	0.903 (18)	C3'—H3'	0.946 (18)
N3'—H3'A	0.976 (19)	C4'—C5'	1.360 (3)
N3'—H3'B	0.89 (2)	C4'—C9'	1.408 (3)
C1—C8	1.442 (2)	C4'—H4'	0.96 (2)
C2—C3	1.359 (2)	C5'—C6'	1.382 (3)
C2—C10	1.487 (2)	C5'—H5'	0.92 (2)
C3—C9	1.414 (2)	C6'—C7'	1.361 (3)
C3—H3	0.959 (17)	C6'—H6'	0.96 (2)
C4—C5	1.353 (3)	C7'—C8'	1.407 (2)
C4—C9	1.408 (2)	C7'—H7'	0.955 (18)
C4—H4	0.98 (2)	C8'—C9'	1.404 (2)

C5—C6	1.388 (3)	C10'—C11'	1.383 (2)
C5—H5	0.94 (2)	C10'—C15'	1.389 (2)
C6—C7	1.368 (3)	C11'—C12'	1.386 (3)
C6—H6	0.96 (2)	C11'—H11'	0.983 (17)
C7—C8	1.402 (2)	C12'—C13'	1.370 (3)
C7—H7	0.945 (17)	C12'—H12'	1.00 (2)
C8—C9	1.407 (2)	C13'—C14'	1.370 (3)
C10—C11	1.389 (2)	C13'—H13'	0.95 (2)
C10—C15	1.391 (2)	C14'—C15'	1.374 (3)
C11—C12	1.375 (3)	C14'—H14'	0.94 (2)
C11—H11	0.988 (18)	C15'—H15'	0.972 (18)
C1—N1—C2	119.22 (14)	C14—C13—H13	120.8 (13)
C1—N2—N3	121.14 (14)	C13—C14—C15	120.6 (2)
C1—N2—H2N	122.1 (11)	C13—C14—H14	123.3 (12)
N3—N2—H2N	109.5 (11)	C15—C14—H14	116.1 (12)
N2—N3—H3A	109.6 (12)	C14—C15—C10	120.6 (2)
N2—N3—H3B	107.2 (12)	C14—C15—H15	119.7 (11)
H3A—N3—H3B	109.9 (18)	C10—C15—H15	119.8 (11)
C1'—N1'—C2'	119.52 (14)	N1'—C1'—N2'	117.49 (15)
C1'—N2'—N3'	120.66 (15)	N1'—C1'—C8'	123.38 (14)
C1'—N2'—H2'N	119.4 (11)	N2'—C1'—C8'	119.13 (15)
N3'—N2'—H2'N	112.8 (11)	C3'—C2'—N1'	121.27 (16)
N2'—N3'—H3'A	107.7 (11)	C3'—C2'—C10'	123.79 (16)
N2'—N3'—H3'B	108.0 (13)	N1'—C2'—C10'	114.93 (15)
H3'A—N3'—H3'B	109.2 (17)	C2'—C3'—C9'	120.53 (17)
N1—C1—N2	117.71 (15)	C2'—C3'—H3'	117.9 (11)
N1—C1—C8	123.47 (14)	C9'—C3'—H3'	121.5 (11)
N2—C1—C8	118.81 (15)	C5'—C4'—C9'	121.0 (2)
C3—C2—N1	121.64 (16)	C5'—C4'—H4'	122.4 (12)
C3—C2—C10	123.14 (16)	C9'—C4'—H4'	116.6 (12)
N1—C2—C10	115.22 (15)	C4'—C5'—C6'	120.7 (2)
C2—C3—C9	120.43 (17)	C4'—C5'—H5'	120.1 (15)
C2—C3—H3	120.7 (10)	C6'—C5'—H5'	119.2 (15)
C9—C3—H3	118.9 (10)	C7'—C6'—C5'	120.1 (2)
C5—C4—C9	121.3 (2)	C7'—C6'—H6'	119.9 (12)
C5—C4—H4	123.4 (12)	C5'—C6'—H6'	120.0 (12)
C9—C4—H4	115.3 (12)	C6'—C7'—C8'	120.6 (2)
C4—C5—C6	120.2 (2)	C6'—C7'—H7'	120.2 (11)
C4—C5—H5	120.7 (14)	C8'—C7'—H7'	119.2 (11)
C6—C5—H5	119.0 (14)	C9'—C8'—C7'	119.44 (17)
C7—C6—C5	120.5 (2)	C9'—C8'—C1'	116.33 (15)
C7—C6—H6	119.1 (12)	C7'—C8'—C1'	124.19 (16)
C5—C6—H6	120.4 (12)	C8'—C9'—C4'	118.12 (19)
C6—C7—C8	120.42 (19)	C8'—C9'—C3'	118.93 (16)
C6—C7—H7	120.1 (10)	C4'—C9'—C3'	122.94 (18)
C8—C7—H7	119.4 (10)	C11'—C10'—C15'	117.18 (17)
C7—C8—C9	119.24 (16)	C11'—C10'—C2'	120.30 (16)

C7—C8—C1	124.44 (16)	C15'—C10'—C2'	122.51 (17)
C9—C8—C1	116.31 (15)	C10'—C11'—C12'	121.42 (18)
C8—C9—C4	118.37 (18)	C10'—C11'—H11'	116.7 (10)
C8—C9—C3	118.91 (16)	C12'—C11'—H11'	121.9 (10)
C4—C9—C3	122.71 (18)	C13'—C12'—C11'	120.2 (2)
C11—C10—C15	117.55 (18)	C13'—C12'—H12'	121.5 (12)
C11—C10—C2	120.17 (16)	C11'—C12'—H12'	118.2 (12)
C15—C10—C2	122.27 (17)	C12'—C13'—C14'	119.1 (2)
C12—C11—C10	121.6 (2)	C12'—C13'—H13'	119.1 (12)
C12—C11—H11	122.8 (10)	C14'—C13'—H13'	121.8 (12)
C10—C11—H11	115.6 (10)	C13'—C14'—C15'	120.8 (2)
C13—C12—C11	120.0 (2)	C13'—C14'—H14'	120.1 (12)
C13—C12—H12	121.3 (12)	C15'—C14'—H14'	119.1 (12)
C11—C12—H12	118.6 (12)	C14'—C15'—C10'	121.3 (2)
C12—C13—C14	119.8 (2)	C14'—C15'—H15'	119.9 (11)
C12—C13—H13	119.3 (13)	C10'—C15'—H15'	118.8 (11)
C2—N1—C1—N2	179.67 (14)	C2'—N1'—C1'—N2'	-179.00 (14)
C2—N1—C1—C8	-1.7 (2)	C2'—N1'—C1'—C8'	1.6 (2)
N3—N2—C1—N1	-14.0 (2)	N3'—N2'—C1'—N1'	14.0 (2)
N3—N2—C1—C8	167.31 (15)	N3'—N2'—C1'—C8'	-166.61 (15)
C1—N1—C2—C3	0.4 (2)	C1'—N1'—C2'—C3'	0.5 (2)
C1—N1—C2—C10	-179.08 (13)	C1'—N1'—C2'—C10'	179.06 (14)
N1—C2—C3—C9	0.7 (3)	N1'—C2'—C3'—C9'	-1.5 (3)
C10—C2—C3—C9	-179.86 (15)	C10'—C2'—C3'—C9'	-179.91 (15)
C9—C4—C5—C6	-0.6 (4)	C9'—C4'—C5'—C6'	0.1 (4)
C4—C5—C6—C7	-0.1 (4)	C4'—C5'—C6'—C7'	-0.7 (4)
C5—C6—C7—C8	0.2 (3)	C5'—C6'—C7'—C8'	1.1 (3)
C6—C7—C8—C9	0.3 (3)	C6'—C7'—C8'—C9'	-0.9 (3)
C6—C7—C8—C1	-178.62 (17)	C6'—C7'—C8'—C1'	176.68 (18)
N1—C1—C8—C7	-179.17 (15)	N1'—C1'—C8'—C9'	-2.6 (2)
N2—C1—C8—C7	-0.6 (2)	N2'—C1'—C8'—C9'	177.99 (15)
N1—C1—C8—C9	1.9 (2)	N1'—C1'—C8'—C7'	179.72 (16)
N2—C1—C8—C9	-179.52 (14)	N2'—C1'—C8'—C7'	0.3 (2)
C7—C8—C9—C4	-0.9 (3)	C7'—C8'—C9'—C4'	0.3 (3)
C1—C8—C9—C4	178.12 (17)	C1'—C8'—C9'—C4'	-177.49 (17)
C7—C8—C9—C3	-179.72 (16)	C7'—C8'—C9'—C3'	179.33 (17)
C1—C8—C9—C3	-0.7 (2)	C1'—C8'—C9'—C3'	1.6 (2)
C5—C4—C9—C8	1.0 (3)	C5'—C4'—C9'—C8'	0.1 (3)
C5—C4—C9—C3	179.8 (2)	C5'—C4'—C9'—C3'	-178.9 (2)
C2—C3—C9—C8	-0.5 (3)	C2'—C3'—C9'—C8'	0.4 (3)
C2—C3—C9—C4	-179.28 (18)	C2'—C3'—C9'—C4'	179.37 (19)
C3—C2—C10—C11	-175.55 (17)	C3'—C2'—C10'—C11'	169.50 (17)
N1—C2—C10—C11	3.9 (2)	N1'—C2'—C10'—C11'	-9.0 (2)
C3—C2—C10—C15	4.8 (3)	C3'—C2'—C10'—C15'	-11.0 (3)
N1—C2—C10—C15	-175.73 (15)	N1'—C2'—C10'—C15'	170.47 (15)
C15—C10—C11—C12	0.4 (3)	C15'—C10'—C11'—C12'	0.5 (3)
C2—C10—C11—C12	-179.27 (16)	C2'—C10'—C11'—C12'	180.00 (16)

C10—C11—C12—C13	0.1 (3)	C10'—C11'—C12'—C13'	-0.4 (3)
C11—C12—C13—C14	-0.3 (3)	C11'—C12'—C13'—C14'	0.1 (3)
C12—C13—C14—C15	0.1 (3)	C12'—C13'—C14'—C15'	0.3 (3)
C13—C14—C15—C10	0.4 (3)	C13'—C14'—C15'—C10'	-0.2 (3)
C11—C10—C15—C14	-0.6 (3)	C11'—C10'—C15'—C14'	-0.2 (3)
C2—C10—C15—C14	179.05 (16)	C2'—C10'—C15'—C14'	-179.68 (16)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2'—H2' <i>N</i> ···N3' ⁱ	0.91 (2)	2.15 (2)	2.967 (2)	151 (2)
N2—H2 <i>N</i> ···N3 ⁱⁱ	0.90 (2)	2.20 (2)	3.027 (2)	152 (2)
N3'—H3' <i>B</i> ···N1' ⁱⁱⁱ	0.89 (2)	2.24 (2)	3.119 (2)	169 (2)
N3—H3 <i>A</i> ···N1 ^{iv}	0.92 (2)	2.26 (2)	3.170 (3)	168 (2)

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y, -z$; (iv) $-x+2, -y, -z+1$.