

RESEARCH ARTICLE

17 β -Hydroxysteroid Dehydrogenase Type 2 Inhibition: Discovery of Selective and Metabolically Stable Compounds Inhibiting Both the Human Enzyme and Its Murine Ortholog

Emanuele M. Gargano¹, Giuseppe Allegretta², Enrico Perspicace¹, Angelo Carotti³, Chris Van Koppen⁴, Martin Frotscher¹, Sandrine Marchais-Oberwinkler^{1,2*}, Rolf W. Hartmann^{1,2*}

1 Pharmaceutical and Medicinal Chemistry, Saarland University, Saarbrücken, Germany, **2** Helmholtz Institute for Pharmaceutical Research Saarland, Saarbrücken, Germany, **3** Dipartimento Farmaco-Chimico, Università degli Studi di Bari, Bari, Italy, **4** ElexoPharm GmbH, Saarbrücken, Germany

* rolf.hartmann@helmholtz-hzi.de (RWH); s.marchais@mx.uni-saarland.de (SMO)



CrossMark
click for updates

OPEN ACCESS

Citation: Gargano EM, Allegretta G, Perspicace E, Carotti A, Van Koppen C, Frotscher M, et al. (2015) 17 β -Hydroxysteroid Dehydrogenase Type 2 Inhibition: Discovery of Selective and Metabolically Stable Compounds Inhibiting Both the Human Enzyme and Its Murine Ortholog. PLoS ONE 10(7): e0134754. doi:10.1371/journal.pone.0134754

Editor: Alessio Lodola, University of Parma, ITALY

Received: June 10, 2015

Accepted: July 13, 2015

Published: July 31, 2015

Copyright: © 2015 Gargano et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The authors acknowledge the Deutsche Forschungsgemeinschaft (DFG) for financial support (Grants HA1315/12-1). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Elexopharm GmbH had no financial interest in this collaboration.

Competing Interests: The authors have declared that no competing interests exist. Chris Van Koppen is an employee of Elexopharm GmbH. He was in

Abstract

Design and synthesis of a new class of inhibitors for the treatment of osteoporosis and its comparative *h*17 β -HSD2 and *m*17 β -HSD2 SAR study are described. **17a** is the first compound to show strong inhibition of both *h*17 β -HSD2 and *m*17 β -HSD2, intracellular activity, metabolic stability, selectivity toward *h*17 β -HSD1, *m*17 β -HSD1 and estrogen receptors α and β as well as appropriate physicochemical properties for oral bioavailability. These properties make it eligible for pre-clinical animal studies, prior to human studies.

Introduction

Osteoporosis is a common, age-related disease, characterized by a systemic impairment of bone mass and microarchitecture, increasing bone fragility and risk of fractures [1]. It has been shown that the drop in 17 β -estradiol (E2) and testosterone (T) levels, occurring with ageing, is the main factor driving the onset and progression of this disease [2]. 17 β -Hydroxysteroid dehydrogenase type 2 (17 β -HSD2) catalyzes the conversion of the highly active E2 and T into the weakly potent 17-ketosteroids estrone (E1) and Δ 4-androstene-3,17-dione (Δ 4-AD), respectively [3]. It is expressed in osteoblastic cells [4], therefore its inhibition can lead to the desired increase of E2 and T levels in the bone tissue and may thus be a novel approach for the treatment of osteoporosis.

Some steroidal [5–7] and non steroidal [8, 9] 17 β -HSD2 inhibitors are already described. In our group we also developed and reported about several classes of non-steroidal 17 β -HSD2 inhibitors [10–14], with a strong inhibition of human 17 β -hydroxysteroid dehydrogenase type 2 (*h*17 β -HSD2) and a good selectivity toward *h*17 β -HSD1. Since *h*17 β -HSD1 is the biological

charge of the biological testing of the compounds. This work was based on a scientific collaboration between Elexopharm GmbH, the Saarland University and the Helmholtz Institute for Pharmaceutical Research Saarland. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials related to this study.

counterpart of *h17 β -HSD2*, catalyzing the opposite conversion, selectivity toward this enzyme is an important feature to take into consideration. Potent and selective *h17 β -HSD1* inhibitors have also been described for the treatment of estrogen-dependent diseases [15, 16].

Given that the most commonly used animal model for osteoporosis studies are established in rodents [17, 18], we aimed at the development of new inhibitors displaying a good inhibition of mouse 17 β -hydroxysteroid dehydrogenase type 2 (*m17 β -HSD2*) and a reasonable selectivity toward *m17 β -HSD1*. In order to have a compound suitable for animal testing and following human studies, the designed inhibitors should also display *h17 β -HSD2* inhibitory activity and selectivity toward *h17 β -HSD1*. Other characteristics to be implemented are preferably low affinity to the estrogen receptors (ERs) α and β in order to maximize the E2 local effect and to minimize systemic side effects as well as metabolic stability.

Since the 3D-structure for both human and mouse 17 β -HSD2 is up to date not available a ligand based approach was chosen for the design of new inhibitors. A set of *h17 β -HSD2* inhibitors was selected and tested for *m17 β -HSD2* inhibitory activity in order to get insight in the SAR and tracking the lead for the rational design of new inhibitors.

Results and Discussion

h17 β -HSD2, *h17 β -HSD1*, *m17 β -HSD2* and *m17 β -HSD1* cell-free assays were performed similarly, by incubating enzyme, tritiated substrate, cofactor and inhibitor, according to described procedures [19–22].

As starting point 25 previously described *h17 β -HSD2* inhibitors (Fig 1), belonging to the 2,5-thiophene amide, 1,3-phenyl amide and 1,4-phenyl amide class [12–14], were tested for *m17 β -HSD2* inhibition, in order to elaborate a comparative SAR and to develop an optimization strategy.

IC₅₀ or percent of inhibition values for both *h17 β -HSD2* and *m17 β -HSD2* are given (Table 1, compounds 1a–2g, 3a, 4a–4f) to facilitate comparison.

In the 2,5-thiophene amide class (compounds 1a–2g), *h17 β -HSD2* inhibitors [13, 14], a broad range of inhibitory activities was detected, depending on the substitution pattern; the most active compounds show IC₅₀ values around 60 nM (Table 1). Conversely, the inhibitory activity towards *m17 β -HSD2* was only marginally affected by these changes (inhibitory activity around 30% at 1 μ M). Only compound 2a shows a more pronounced *m17 β -HSD2* inhibition combined with good inhibition of the human enzyme (65% at 1 μ M and IC₅₀ = 61 nM). Unfortunately, compound 2a turned out to be metabolically very unstable, with a half-life of only 4 minutes in the human liver S9 fraction [13]. In a further screening, all the tested 2,5-thiophene amide displayed a high metabolic instability [13].

It is striking that neither the nature of substituents on ring A and C or their substitution pattern does appear to exert an effect on *m17 β -HSD2* inhibition, whereas it is decisive for the *h17 β -HSD2* one. This result suggests that the inhibitors in this class are likely to have different binding modes in the two enzyme isoforms.




Fig 1. Previously described *h17 β -HSD2* inhibitors, tested for *m17 β -HSD2* inhibition.

doi:10.1371/journal.pone.0134754.g001

Table 1. Inhibition of Human and Mouse 17 β -HSD2 by 2,5-Thiophene Amide, 1,3-Phenyl Amide and 1,4-Phenyl Amide Derivatives in a Cell-Free Assay.

Cmpd	R ₁	R ₂	IC ₅₀ (nM) ^a or % inh. at 1 μ M ^{a, d}	
			<i>h</i> 17 β -HSD2 ^b	<i>m</i> 17 β -HSD2 ^c
1a	3-OMe	3-OMe	68	29%
1b	3-Me	3-Me	52	42%
1c	3-Me	3-OMe	58	30%
1d	2-OMe	3-OMe	490	19%
1e	3-OH	3-OH	33%	n.i.
1f	2-OH	3-OH	410	18%
1g	3-N(Me) ₂	3-OMe	170	40%
1h	3-F	3-OMe	510	n.i.
1i	4-CN	3-OMe	48%	n.i.
1j	2-F, 3-OMe	3-OMe	62	26%
1k	2-F, 3-OMe	3-Me	62	45%
2a	2-F, 3-OMe	3-OH	61	65%
2b	3-OMe	3-OMe	370	16%
2c	3-OH	3-OH	390	26%
2d	4-OH	3-OH	330	35%
2e	3-Me	3-OH	160	45%
2f	3-F	3-OH	330	37%
2g	4-CN	3-OH	n.i.	n.i.
3a	3-Me	3-Me	1100	50%
4a	3-OMe	3-OMe	520	25%
4b	4-OMe	3-OMe	1200	11%
4c	3-OH	3-OH	35%	n.i.
4d	3-OMe	3-OMe	11%	n.i.
4e	3-OH	3-OH	640	29%
4f	4-OH	3-OH	480	22%

^aMean value of at least two determinations, standard deviation less than 20%.

^bHuman placental microsomal fraction, substrate E2 [500 nM], cofactor NAD⁺ [1500 μ M].

^cMouse liver microsomal fraction, substrate E2 [500 nM], cofactor NAD⁺ [1500 μ M].

^dn.i.: no inhibition, i.e., inhibition \leq 10%.

doi:10.1371/journal.pone.0134754.t001

Exchange of the central thiophene by a 1,3-disubstituted phenyl led to compounds **4a-4c** for $n = 0$ and **4d-4f** for $n = 1$, with weak inhibitory activity towards both the human and the mouse enzyme (Table 1).

In contrast, compound **3a** (Table 1), bearing a 1,4-disubstituted phenyl moiety as central ring, shows moderate inhibition of both *h*17 β -HSD2 and *m*17 β -HSD2 and also revealed exceptional metabolic stability in the human liver S9 fraction, with a half-life time >120 minutes [13]. It was therefore taken as starting point for the design of a small library of inhibitors where the substitution pattern and the physicochemical nature of substituents on the A and C rings was varied (Fig 2). A larger number of derivatives, bearing substituents with different physicochemical properties on the A ring were prepared, according to their chemical accessibilities. Compounds **25a** and **25** were also synthesized to investigate the effect of the methylene linker between the amide function and the C ring.

The synthesis of the 1,4-phenyl derivatives **5a-23a**, **25a**, **7** and **25**, depicted in Fig 3, was accomplished following a two- or three-step reaction pathway.




Fig 2. Chemical Structures of the Designed Compounds.

doi:10.1371/journal.pone.0134754.g002

First, amidation was carried out by reaction of the commercially available 4-bromobenzoyl chloride **5d** with substituted anilines **5c** or with the 1-(3-methoxyphenyl)-*N*-methylmethanamine **25c** using standard conditions (method A: triethylamine, dichloromethane, from 0°C to room temperature, overnight) affording the brominated intermediates **5b**, **6b** and **11b** in almost quantitative yields. Subsequently, Suzuki coupling (Method B) using tetrakis(triphenylphosphine)palladium and cesium carbonate in a mixture of DME/EtOH/H₂O (1:1:1, 3 mL) as solvent and microwave irradiation (150°C, 150 W for 20 minutes), provided the biphenyl derivatives **5a-21a** and **25a**, and the 1,3-benzodioxole derivatives **22a** and **23a** in good yields. Compounds **7a** and **25a** were submitted to ether cleavage using boron trifluoride-dimethyl sulfide complex, yielding the hydroxylated molecules **7** and **25**.

The synthesis of the retroamide **24a**, displayed in Fig 4, follows a two-step procedure.

First the commercially available 3-methoxybenzoyl chloride **24c** was reacted with 4-bromo-*N*-methyl aniline **24d** according to method A and afforded the brominated intermediate **24b** with 70% yield. Subsequently, Suzuki coupling following method B afforded compound **24a** in 68% yield.

Compounds **5a-25a**, **7** and **25** were tested for *h*17 β -HSD2, *m*17 β -HSD2 and *h*17 β -HSD1 inhibition (Table 2, results expressed as percentage of inhibition or IC₅₀ value).

Compounds **18a** and **24a** were not tested for inhibition of *m*17 β -HSD2, due to their low inhibitory activity towards the human enzyme. Among the different synthesized 1,4-biphenyl




Fig 3. Synthesis of 1,4-Phenyl Derivatives 5a-23a, 25a, 7, 25. Reagents and conditions: (i) Et₃N, CH₂Cl₂, room temperature, overnight; (ii) DME/EtOH/H₂O (1:1:1), Cs₂CO₃, Pd(PPh₃)₄, microwave irradiation (150°C, 150W, 20 min); (iii) BF₃·SMe₂, CH₂Cl₂, room temperature, overnight.

doi:10.1371/journal.pone.0134754.g003




Fig 4. Synthesis of 1,4-Phenyl Retroamide Derivative 24a. Reagents and conditions: (i) Et₃N, CH₂Cl₂, room temperature, overnight; (ii) DME/EtOH/H₂O (1:1:1), Cs₂CO₃, Pd(PPh₃)₄, microwave irradiation (150°C, 150W, 20 min).

doi:10.1371/journal.pone.0134754.g004

amides without methylene linker ($n = 0$, Table 2, compounds 5a-21a), the best *h*17β-HSD2 inhibitory activity and selectivity toward *h*17β-HSD1 was achieved for compounds 6a, 7a and

Table 2. Inhibition of *h*17β-HSD2, *m*17β-HSD2 and *h*17β-HSD1 by Biphenyl Amide Derivatives with Different Substitution Patterns on the A and C Rings in Cell-Free System.

Cmpd	R ₁	R ₂	IC ₅₀ (nM) ^a or % inh. at 1 μM ^{a, c}			s. f. ^{d, f}	IC ₅₀ (nM) ^a or % inh. at 1 μM ^{a, c}
			<i>h</i> 17β-HSD2 ^b	<i>h</i> 17β-HSD1 ^c	<i>m</i> 17β-HSD2 ^{f, g}		
3a	3-Me	3-Me	1100	11500	10	50%	
5a	3-OMe	3-Me	44%	n.i.	n.d.	56%	
6a	3-Me	3-OMe	260	6400	25	260	
7a	3-OMe	3-OMe	330	6400	20	290	
8a	-H	3-OMe	51%	22%	n.d.	48%	
9a	2-OMe	3-OMe	50%	n.i.	n.d.	33%	
10a	4-OMe	3-OMe	710	27000	38	66%	
11a	3-OMe	4-OMe	27%	n.i.	n.d.	58%	
12a	4-OMe	4-OMe	19%	n.i.	n.d.	62%	
7	3-OH	3-OH	36%	15%	n.d.	42%	
13a	3-F	3-OMe	1000	8000	8	57%	
14a	3-Cl	3-OMe	950	12600	13	73%	
15a	3-N(Me) ₂	3-OMe	650	5000	8	67%	
16a	3-OMe,4-OMe	3-OMe	520	85500	164	52%	
17a	3-Me,4-Me	3-OMe	300	13300	44	140	
18a	3-F,4-F	3-OMe	31%	n.i.	n.d.	n.d.	
19a	2-F, 3-OMe	3-OMe	64%	35%	n.d.	67%	
20a	2-F, 3-Me	3-OMe	460	11300	24	48%	
21a	2-F, 3-Me	3-Me	330	3710	11	56%	
22a	-	3-OMe	51%	n.i.	n.d.	56%	
23a	-	3-Me	560	10900	20	70%	
24a	-	-	11%	n.i.	n.d.	n.d.	
25a	2-F, 3Me	3-OMe	310	87000	283	43%	
25	2-F, 3-Me	3-OH	260	31000	118	190	

^aMean value of at least three determinations, standard deviation less than 20% except for 11a (*h*HSD2): 26%, 7(*h*HSD1): 25%,.

^bHuman placental, microsomal fraction, substrate E2 [500 nM], cofactor NAD⁺ [1500 μM].

^cHuman placental, cytosolic fraction, substrate E1 [500 nM], cofactor NADH [500 μM].

^ds.f. (selectivity factor) = IC₅₀(17β-HSD1)/IC₅₀(17β-HSD2).

^en.i.: no inhibition (inhibition of <10%).

^fn.d.: not determined.

^gMouse liver microsomal fraction, substrate E2 [500 nM], cofactor NAD⁺ [1500 μM].

doi:10.1371/journal.pone.0134754.t002

17a (Table 2, IC₅₀ values between 260 and 330nM, s.f. between 20 and 44), showing that a 3-OMe-group on ring C in combination with either a 3-OMe- or a 3-Me-group on ring A leads to a maximum in potency and selectivity in this series of compounds.

The presence of a methyl group in 4-position of ring A is tolerated by the *h17 β -HSD2* (compound **17a**, IC₅₀ = 310nM) and increases the selectivity toward *h17 β -HSD1* (**17a**: s.f. 44; **6a**: s.f. 25). In contrast, compound **16a** bearing a 3,4-dimethoxy substituted A ring displays a slight decrease in *h17 β -HSD2* inhibitory activity if compared to the corresponding compound **7a** with only one methoxy group on that ring. The rigidification of the two methoxy substituents by the synthesis of a 1,3-benzodioxole ring (compounds **22a** and **23a**) could not overcome the drop in potency. Compounds **6a**, **7a** and **17a** displayed the strongest *m17 β -HSD2* inhibitory activity (IC₅₀ = 260, 290 and 140 nM, respectively).

In general, the introduction of methyl- or methoxy-groups, especially in the 3-positions of rings A and B, had a positive impact on inhibitory activity, which is similar for both 17 β -HSD2 isoforms (Table 2). Therefore, inhibitors belonging to the 1,4-phenyl class are likely to bind in a conserved area common to both enzyme, in contrast to the 2,5-thiophene amide class. As 17 β -HSD2 belongs to the SDR superfamily, characterized by the conserved Rossmann fold and catalytic triad [23], it is possible that these inhibitors bind in or very close to these regions. Furthermore, since compounds **6a** and **17a** lack of the two oxygen functions to mimic the E2 interactions with the enzyme, they are likely to bind to the active site in an alternative mode, significantly influenced by the methyl substitution patterns.

Compounds with a methylene linker **25a** and **25** (Table 2), showed *h17 β -HSD2* inhibitory activity in the same order of magnitude as the corresponding derivative **20a** lacking the methylene group, but displayed improved selectivities over *h17 β -HSD1*. In contrast to what was observed for human 17 β -HSD2, compounds **25a** and **25** showed a significant difference in terms of *m17 β -HSD2* inhibition, indicating that the hydroxy group of compound **25** might function as H-bond donor in the interaction with the enzyme. A similar behavior was observed in the 2,5-thiophene amide class with regard to the *h17 β -HSD2* inhibitory activity[13]: all the tested 2,5-thiophene amides with a methylene linker achieved the highest *h17 β -HSD2* inhibition when substituted with an hydroxy group on the C ring. It might be therefore speculated, that the addition of the methylene linker can influence the binding mode of the inhibitors in both classes.

The 1,4-phenyl amides are likely to be competitive inhibitors, as derivatives with a similar structure were found to inhibit the enzyme following this mode of action.

The most potent *m17 β -HSD2* inhibitors **6a-8a**, **10a**, **14a**, **15a**, **17a**, **19a**, **21a**, **23a** and **25** were also tested for their *m17 β -HSD1* inhibitory activity. None of the tested compounds showed any inhibition of *m17 β -HSD1* at a concentration of 1 μ M, indicating a significant selectivity toward this enzyme.

The inhibitory activity of compounds **6a**, **17a** and **25** was also evaluated in the human mammary cell line MDA-MB-231 containing endogenous 17 β -HSD2 (Table 3). The compounds

Table 3. Human 17 β -HSD2 Cellular Inhibition of Compounds 6a, 17a and 25.

Cmpd	% inh ^b HSD2 ^a at 250 nM	% inh ^b HSD2 ^a at 1250 nM
6a	60%	87%
17a	66%	90%
25	67%	88%

^aMDA-MB-231 cell line, substrate E2 [200 nM].

^bMean value of two determinations, standard deviation less than 15%.

doi:10.1371/journal.pone.0134754.t003

were tested at 250 nM and 1250 nM, representing approximately the IC₅₀ observed in the cell-free assay, and its 5-fold value. As displayed in Table 3, all three compounds showed an inhibition between 60% and 67% at the lower concentration and approximately 90% inhibition at the higher concentration, indicating that the inhibitors can permeate the membrane and are able to inhibit the enzyme in a concentration dependent manner.

Compounds 6a, 7a, 17a, 21a, 25a and 25 were tested for their affinity toward the ERs α and β according to described methods [24] (assay details are available in the Materials and Methods). Even when applied in a 1000-fold excess relative to E2, no inhibitor was able to displace more than 20% of the steroid from the corresponding receptor, indicating a very low binding affinity to the ERs.

The metabolic stabilities of the most active compounds 6a, 17a and 25 were evaluated using human liver microsomes (S9 fraction). In addition, compounds 5a, 14a, 20a and 25a were also tested in order to investigate whether structure modifications might exert an effect on metabolic stability (Table 4, assay details are available in Materials and Methods). All compounds, except 25, revealed a very high stability, which was not influenced by the nature of the substituents. Compound 25, exhibiting a short half-life, bears a hydroxy group, potentially susceptible to phase II metabolism. Interestingly, all the tested inhibitors from the 1,4-phenyl amide class, for $n = 0$, demonstrated high metabolic stability, which seemingly constitutes a positive feature of the whole class. As species differences for the metabolism of drugs that are not or only partly metabolized are usually small [25], sufficient metabolic stability in species other than human can be anticipated.

Conclusion

The aim of this work was the design of a compound, which should be suitable for application in both an animal model of osteoporosis and in humans. We report here the discovery of compound 17a, which is the first to show an appropriate profile for this purpose, with strong inhibition of both human and mouse 17 β -HSD2 and selectivity toward the respective type 1 enzymes and the ERs. It also displayed good cellular inhibitory activity, high metabolic stability and good physicochemical parameters (MW = 345 and cLogP = 4.75) predictor for good oral bio-availability [26]. A comparative SAR study for *h*17 β -HSD2 and *m*17 β -HSD2 is also described for the 1,4-phenyl and the 2,5-thiophene classes of inhibitors, revealing that only

Table 4. Half-life in Human Liver Microsomes S9 Fraction of Representative Compounds 3a, 5a, 6a, 14a, 17a, 20a, 25a and 25.

Cmpd	R ₁	R ₂	Inhibitor ^c t _{1/2} (min) ^{a,b}
3a	3-Me	3-Me	>120
5a	3-OMe	3-Me	116
6a	3-Me	3-OMe	106
14a	3-Cl	3-OMe	82
17a	3-Me,4-Me	3-OMe	107
20a	2-F, 3-Me	3-OMe	104
25a	2-F, 3-Me	3-OMe	103
25	2-F, 3-Me	3-OH	6

^aMean of at least two determinations, standard deviation less than 25%.

^b1 mg/ml pooled mammalian liver S9 fraction (BD Gentest), 2 mM NADPH regenerating system, 1 mM UDPGA and 0.1 mM PAPS at 37°C for 0, 5, 15 and 60 minutes.

^cInhibitor tested at a final concentration of 1 μ M.

doi:10.1371/journal.pone.0134754.t004

compounds belonging to the first series similarly inhibit the two enzymes, probably through a similar binding mode. The species specific characterization of the thiophene and the phenyl derivatives pointed out the superiority of the latter class of inhibitors, which is able to equally inhibit the two isoenzymes and additionally displays a high metabolic stability. *In vivo* assays in a mouse osteoporosis model will be carried out soon and the results reported in due course in a specialized journal dealing with bone diseases.

Materials and Methods

Chemical Methods

Chemical names follow IUPAC nomenclature. Starting materials were purchased from Aldrich, Acros, Combi-Blocks or Fluorochem and were used without purification. Column chromatography was performed on silica gel (70–200 μ m) and reaction progress was monitored by TLC on Alugram SIL G/UV254 (Macherey-Nagel). Visualization was accomplished with UV light. ^1H NMR and ^{13}C NMR spectra were measured on a Bruker AM500 spectrometer (at 500 MHz and 125 MHz, respectively) at 300 K and on Bruker Fourier 300 (at 300 MHz and 75 MHz, respectively) at 300K. Chemical shifts are reported in δ (parts per million: ppm), by reference to the hydrogenated residues of deuteriated solvent as internal standard: 2.05 ppm (^1H NMR) and 29.8 and 206.3 ppm (^{13}C NMR) for CD_3COCD_3 , 7.26 ppm (^1H NMR) and 77.0 ppm (^{13}C NMR) for CDCl_3 . Signals are described as br (broad), s (singlet), d (doublet), t (triplet), dd (doublet of doublets), ddd (doublet of doublet of doublets), dt (doublet of triplets) and m (multiplet). All coupling constants (J) are given in Hertz (Hz).

Melting points (mp) were measured in open capillaries on a Stuart Scientific SMP3 apparatus and are uncorrected.

Mass spectrometry was performed on a TSQ Quantum (ThermoFisher, Dreieich, Germany). The triple quadrupole mass spectrometer was equipped with an electrospray interface (ESI). The Surveyor-LC-system consisted of a pump, an auto sampler, and a PDA detector. The system was operated by the standard software Xcalibur. A RP C18 NUCLEODUR 100–5 (3 mm) column (Macherey-Nagel GmbH, Dühren, Germany) was used as stationary phase. All solvents were HPLC grade. In a gradient run (acetonitrile/water) the percentage of acetonitrile (containing 0.1% trifluoroacetic acid) was increased from an initial concentration of 0% at 0 min to 100% at 15 min and kept at 100% for 5 min. The injection volume was 15 μ L and flow rate was set to 800 μ L/min. MS analysis was carried out at a needle voltage of 3000 V and a capillary temperature of 350°C. Mass spectra were acquired in positive mode from 100 to 1000 m/z and UV spectra were recorded at the wave length of 254 nm and in some cases at 360 nm.

All microwave irradiation experiments were carried out in a 507 CEM-Discover microwave apparatus.

All tested compounds exhibited $\geq 95\%$ chemical purity as measured by LC/MS.

The following compounds were prepared according to previously described procedures: 4-bromo-*N*-methyl-*N*-(*m*-tolyl)benzamide **7a** [13].

Method A, general procedure for amide formation. To a solution of bromobenzoylchloride (2 mmol) was added the corresponding *N*-methylaniline (2 mmol) followed by Et_3N (2 mmol) in CH_2Cl_2 (10 mL) at 0°C. After a few minutes, the ice bath was removed and the reaction mixture was warmed up to room temperature and stirred at room temperature overnight. The reaction mixture was extracted twice with CH_2Cl_2 (2×15 mL). The organic layer was dried over MgSO_4 , filtered and the solution was concentrated under reduced pressure. The residue was purified by silica gel column chromatography using hexanes and EtOAc as eluent or by trituration in a mixture of diethyl ether / petroleum ether to afford the desired compound.

Method B, general procedure for Suzuki-Miyaura coupling. In a sealed tube the previously prepared bromo-*N*-heteroarylcarboxamide derivative (1 eq.) was introduced, followed by the corresponding boronic acid (1.5 eq.), cesium carbonate (3 eq.), tetrakis(triphenylphosphine)palladium (0.02 eq.) and a mixture of DME/EtOH/H₂O (1:1:1, v:v:v, 3 mL) as solvent. The reactor was flushed with N₂ and submitted to microwave irradiation (150°C, 150 W) for 20 minutes. After cooling to room temperature, a mixture of EtOAc/H₂O (1:1, v:v, 2 mL) was added to stop the reaction. The aqueous layer was extracted with EtOAc (3 × 10 mL). The organic layer was washed once with brine and once with water, dried over MgSO₄, filtered and the solution was concentrated under reduced pressure. The residue was purified by column chromatography using hexanes and EtOAc as eluent to afford the desired compound.

Detailed synthesis procedures of the compounds

4-Bromo-*N*-(4-methoxyphenyl)-*N*-methylbenzamide (11b). The title compound was prepared by reaction of 4-bromobenzoyl chloride (**5c**) (1272 mg, 5.8mmol) and 4-methoxy-*N*-methylaniline (**11d**) (400 mg, 2.9mmol) according to method A. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as yellow solid (932 mg, 99%). C₁₅H₁₄BrNO₂; MW 320; mp: 61–62°C; MS (ESI) 320, 322 [M]⁺; IR (cm⁻¹) 1635, 2840, 2850, 2917; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.36 (s, 3H), 3.74 (s, 3H), 6.82 (d, *J* = 9.0 Hz, 2H), 7.09 (d, *J* = 9.0 Hz, 2H), 7.23 (d, *J* = 8.0 Hz, 2H), 7.38 (d, *J* = 8.0 Hz, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.5, 55.7, 115.2, 123.7, 129.3, 131.4, 131.6, 137.1, 138.6, 159.1, 169.5.

4-Bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (6b). The title compound was prepared by reaction of 4-bromobenzoyl chloride (**5c**) (1272 mg, 5.8mmol) and 3-methoxy-*N*-methylaniline (**6d**) (400 mg, 2.9mmol) according to method A. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as yellow solid (935 mg, quantitative). C₁₅H₁₄BrNO₂; MW 320; mp: 98–100°C; MS (ESI) 320, 322 [M]⁺; IR (cm⁻¹) 1638, 2840, 2945, 3012; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.40 (s, 3H), 3.71 (s, 3H), 6.70–6.72 (m, 1H), 6.74–6.78 (m, 2), 7.17 (t, *J* = 8.0 Hz, 1H), 7.26–7.28 (m, 2H), 7.39–7.41 (m, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.2, 55.7, 113.2, 113.8, 120.2, 123.9, 130.7, 131.3, 131.6, 137.0, 147.0, 161.2, 169.4.

4-Bromo-*N*-methyl-*N*-(*m*-tolyl)benzamide (5b). The title compound was prepared by reaction of 4-bromobenzoyl chloride (**5c**) (724 mg, 3.3mmol) and *N*-methyl-*m*-toluidine (**5d**) (200 mg, 1.6mmol) according to method A. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as white solid (500 mg, 99%). C₁₅H₁₄BrNO; MW 304; mp: 90–92°C; MS (ESI) 304, 306 [M]⁺; IR (cm⁻¹) 1586, 1638, 2853, 2923, 3058; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 2.24 (s, 3H), 3.39 (s, 3H), 6.90–6.93 (d, *J* = 8.0Hz, 1H), 6.99–7.05 (m, 2H), 7.14 (t, *J* = 8.0 Hz, 1H), 7.23–7.26 (m, 2H), 7.37–7.40 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 21.2, 38.4, 123.9, 125.2, 128.1, 128.5, 129.8, 131.4, 131.6, 137.0, 140.0, 145.8, 169.4.

4-Bromo-*N*-(3-methoxybenzyl)-*N*-methylbenzamide (25b). The title compound was prepared by reaction of 4-bromobenzoyl chloride (**5c**) (579 mg, 2.6mmol) and 3-methoxy-*N*-methylbenzylamine (**25d**) (200 mg, 1.3mmol) according to method A. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as yellow oil (430 mg, 97%). C₁₆H₁₆BrNO₂; MW 334; MS (ESI) 334, 336 [M]⁺; IR (cm⁻¹) 1630, 2834, 2923; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.91 (s, 3H), 3.79 (s, 3H), 4.52–4.70 (m, 2H), 6.76–6.80 (m, 1H), 6.86 (dd, *J* = 8.8 Hz, 5Hz, 1H), 6.95 (s, 1H), 7.28 (t, *J* = 8.0 Hz, 1H), 7.44 (d, *J* = 7.6 Hz, 2H), 7.62 (s, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 37.2, 51.0, 55.5, 113.3, 113.6, 114.4, 119.7, 121.0, 123.9, 129.9, 130.5, 132.3, 137.0, 161.1.

***N*-(4-bromophenyl)-3-methoxy-*N*-methylbenzamide (24b).** The title compound was prepared by reaction of 3-methoxybenzoyl chloride (**24c**) (336 mg, 2.0mmol) and 4-bromo-*N*-methylaniline (**24d**) (367 mg, 2.0mmol) according to method A. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as white solid (440 mg, 70%). C₁₅H₁₄BrNO₂; MW 320; mp: 97–100°C; MS (ESI) 320, 322 [M]⁺; IR (cm⁻¹) 1634, 2834, 2929, 3009; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.41 (s, 3H), 3.69 (s, 3H), 6.84–6.88 (m, 3H), 7.12–7.17 (m, 3H), 7.44–7.47 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.3, 55.6, 114.8, 116.3, 120.1, 121.7, 129.8, 130.0, 133.1, 138.8, 145.7, 160.1, 170.3.

***N*-(3-methoxyphenyl)-*N*-methyl-[1,1'-biphenyl]-4-carboxamide (8a).** The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (80 mg, 0.25mmol), phenylboronic acid (40 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) in DME/EtOH/H₂O (1/1/1, 3 mL) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow solid (77 mg, 97%). C₂₁H₁₉NO₂; MW 317; mp: 102–104°C; MS (ESI) 318 [M+H]⁺; IR (cm⁻¹) 1588, 1636, 2831, 2917, 2960, 3040; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.44 (s, 3H), 3.70 (s, 3H), 6.72–6.77 (m, 2H), 6.80 (t, *J* = 2 Hz, 1H), 7.17 (t, *J* = 8 Hz, 1H), 7.33–3.37 (s, 1H), 7.41–7.46 (m, 4H), 7.51–7.54 (m, 2H), 7.60–7.63 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.4, 55.7, 113.0, 113.8, 120.1, 126.9, 127.7, 128.6, 129.8, 130.1, 130.6, 136.7, 140.8, 142.6, 147.4, 161.2, 170.2.

2'-Methoxy-*N*-(3-methoxyphenyl)-*N*-methyl-[1,1'-biphenyl]-4-carboxamide (9a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (112 mg, 0.35mmol), 2-methoxyphenylboronic acid (70 mg, 0.46mmol), cesium carbonate (342 mg, 1.05mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 8 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as colorless oil (113 mg, 93%). C₂₂H₂₁NO₃; MW 347; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1597, 1640, 2837, 2939; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.44 (s, 3H), 3.69 (s, 3H), 3.77 (s, 3H), 6.73–6.78 (m, 3H), 6.99 (dt, *J* = 1.0, 7.3 Hz, 1H), 7.07 (dd, *J* = 1.0, 8.3 Hz, 1H), 7.18 (dt, *J* = 1.0, 7.5 Hz, 1H), 7.24 (dd, *J* = 2.0, 7.5 Hz, 1H), 7.30–7.34 (m, 1H), 7.35–7.39 (m, 4H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.4, 55.6, 55.9, 112.5, 113.0, 113.7, 120.0, 121.7, 129.1, 129.5, 130.0, 130.4, 130.6, 131.3, 136.0, 140.7, 147.4, 157.5, 161.1, 170.4.

3'-Methoxy-*N*-(3-methoxyphenyl)-*N*-methyl-[1,1'-biphenyl]-4-carboxamide (7a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (81 mg, 0.25mmol), 3-methoxyphenylboronic acid (46 mg, 0.3mmol), cesium carbonate (247 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as white solid (74 mg, 85%). C₂₂H₂₁NO₃; MW 347; mp: 131–134°C; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1582, 1635, 2837, 2941, 2963; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.43 (s, 3H), 3.70 (s, 3H), 3.84 (s, 3H), 6.73–6.76 (m, 2H), 6.80–6.81 (m, 1H), 6.92(ddd, *J* = 1.0, 2.0, 8.0 Hz, 1H), 7.14–7.19 (m, 3H), 7.32–7.35 (m, 1H), 7.41–7.42 (m, 2H), 7.51–7.53 (m, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.4, 55.6, 55.7, 113.0, 113.1, 113.8, 114.3, 120.0, 120.1, 126.9, 130.0, 130.6, 130.8, 136.8, 142.2, 142.5, 147.37, 147.38, 161.2, 170.2.

3'-Hydroxy-*N*-(3-hydroxyphenyl)-*N*-methyl-[1,1'-biphenyl]-4-carboxamide (7). To a solution of 3'-methoxy-*N*-(3-methoxyphenyl)-*N*-methyl-[1,1'-biphenyl]-4-carboxamide (**7a**) (100 mg, 0.29mmol) in CH₂Cl₂ (10 mL) was added trifluorobromine dimethyl sulphide complex BF₃·SMe₂ (0.36 mL, 3.46 mM) at room temperature. The reaction mixture was stirred overnight. The reaction was quenched with MeOH (10 mL). The solvent was removed under reduced pressure at 25°C. Water was added to dissolve inorganic salts and the precipitate was

filtered off and triturated with diethyl ether to afford the desired product as white solid (80 mg, 87%). C₂₀H₁₇NO₃; MW 319; mp: 168–171°C; MS (ESI) 320[M+H]⁺; IR (cm⁻¹) 1589, 3191; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.42 (s, 3H), 6.64–6.68 (m, 3H), 6.82–6.86 (m, 1H), 7.07–7.12 (m, 3H), 7.26 (t, *J* = 8.0 Hz, 1H), 7.39–7.44 (m, 2H), 7.47–7.49 (m, 2H), 8.42 (s, 1H), 8.48 (s, 1H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.4, 114.4, 114.6, 115.0, 115.6, 119.0, 119.1, 126.8, 130.1, 130.7, 130.8, 136.6, 142.3, 142.7, 147.4, 158.8, 158.9, 170.1.

4'-Methoxy-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (10a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (123 mg, 0.38mmol), 4-methoxyphenylboronic acid (75 mg, 0.49mmol), cesium carbonate (371 mg, 1.14mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 9 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc80:20) to afford the desired product as white solid (70 mg, 53%). C₂₂H₂₁NO₃; MW 347; mp: 128–130°C; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1601, 1625, 2840, 2935, 2963; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.43 (s, 3H), 3.70 (s, 3H), 3.82(s, 3H), 6.72–6.75 (m, 2H), 6.78–6.79 (m, 1H), 6.98 (d, *J* = 2.3 Hz, 1H), 7.00 (d, *J* = 2.3 Hz, 1H), 7.15–7.18 (m, 1H), 7.37–7.40 (m, 2H), 7.46–7.48 (m, 2H), 7.55–7.58 (m,2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.4, 55.6, 55.7, 112.9, 113.8, 115.2, 120.1, 126.3, 128.8, 130.1, 130.6, 130.1, 135.9, 142.4, 147.5, 160.7, 161.2, 170.3.

3'-Fluoro-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (13a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (100 mg,0.31mmol), 3-fluorophenylboronic acid (57 mg, 0.41mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow solid (91 mg, 89%). C₂₁H₁₈FNO₂; MW 335; mp: 97–100°C; MS (ESI) 336 [M+H]⁺; IR (cm⁻¹) 1587, 1637, 2834, 2923, 2960; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.44 (s, 3H), 3.71 (s, 3H), 6.73–6.77 (m, 2H), 6.80–6.81 (m, 1H), 7.09–7.21 (m, 2H), 7.36–7.50 (m, 5H), 7.55–7.57 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.4, 55.8, 113.1, 113.9, 114.3, 114.6, 115.2, 115.4, 120.2, 123.7, 127.1, 130.2, 130.8, 131.6, 131.8, 137.4, 141.2, 143.3, 143.4, 147.4, 161.3, 162.6, 165.8, 170.1.

3'-Chloro-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (14a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (80 mg, 0.25mmol), 3-chlorophenylboronic acid (52 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as colorless oil (47 mg, 53%). C₂₁H₁₈ClNO₂; MW 352; MS (ESI) 352, 354 [M]⁺; IR (cm⁻¹) 1594, 1639, 2837, 2938; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.43 (s, 3H), 3.70 (s, 3H), 6.73–6.76 (m, 2H), 6.80–6.81(m, 1H), 7.17 (t,*J* = 8 Hz, 1H), 7.38–7.40 (m, 1H), 7.43–7.47 (m, 3H), 7.54–7.59 (m, 3H), 7.63–7.64 (m, 1H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.3, 55.7, 113.1, 113.8, 120.2, 126.3, 127.0, 127.6, 128.5, 130.1, 130.7, 131.4, 135.3, 137.4, 141.0, 142.9, 147.3, 161.2, 170.0.

N-(3-methoxyphenyl)-N,3'-dimethyl-[1,1'-biphenyl]-4-carboxamide (6a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (77 mg, 0.24mmol), 3-methylphenylboronic acid (42 mg, 0.31mmol), cesium carbonate (235 mg, 0.72mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 70:30) to afford the desired product as colorless oil (56 mg, 71%). C₂₂H₂₁NO₂; MW 331; MS (ESI) 332 [M+H]⁺; IR (cm⁻¹) 1586, 1640, 2840, 2929; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.38 (s, 3H), 3.44 (s, 3H), 3.70 (s, 3H), 6.74–6.76 (m, 2H), 6.79–6.80 (m, 1H), 7.15–7.20 (m, 2H), 7.31 (t,*J* = 8 Hz, 1H), 7.39–7.43 (m, 4H), 7.49–7.52 (m, 2H); ¹³C NMR (C₂D₆CO, 75

MHz) δ (ppm) 21.5, 38.4, 55.7, 113.0, 113.9, 120.2, 124.9, 126.9, 128.5, 129.4, 129.8, 130.1, 130.7, 136.6, 137.4, 140.9, 142.9, 147.5, 161.3, 170.3.

3'-(Dimethylamino)-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (15a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (80 mg, 0.25mmol), 3-(dimethylamino)phenylboronic acid (54 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow oil (49 mg, 54%). C₂₃H₂₄N₂O₂; MW 360; MS (ESI) 361 [M+H]⁺; IR (cm⁻¹) 1597, 1639, 2797, 2834, 2939; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 2.97 (s, 6H), 3.43 (s, 3H), 3.70 (s, 3H), 6.72–6.76 (m, 3H), 6.79–6.80 (m, 1H), 6.86–6.93 (m, 2H), 7.17 (t, *J* = 8.0 Hz, 1H), 7.23 (t, *J* = 8.0 Hz, 1H), 7.38–7.41 (m, 2H), 7.49–7.52 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.5, 40.8, 55.7, 111.8, 113.0, 113.9, 116.0, 120.2, 127.0, 130.0, 130.4, 130.7, 136.5, 141.6, 143.9, 147.5, 152.2, 161.3, 170.4.

N-(3-methoxyphenyl)-N,3',4'-trimethyl-[1,1'-biphenyl]-4-carboxamide (17a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (100 mg, 0.31mmol), 3,4-dimethylphenylboronic acid (62 mg, 0.41mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 70:30) to afford the desired product as yellow oil (106 mg, 99%). C₂₃H₂₃NO₂; MW 345; MS (ESI) 346[M+H]⁺; IR (cm⁻¹) 1599, 1640, 2840, 2914, 2942; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.26 (s, 3H), 2.29 (s, 3H), 3.43 (s, 3H), 3.70 (s, 3H), 6.73–6.75 (m, 2H), 6.79–6.80 (m, 1H), 7.15–7.19 (m, 2H), 7.32 (dd, *J* = 2.0Hz, 8.0Hz, 1H), 7.38–7.40 (m, 3H), 7.47–7.49 (m, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 19.4, 19.8, 38.4, 55.6, 112.9, 113.8, 120.1, 125.0, 126.5, 128.8, 130.0, 130.6, 131.0, 136.2, 137.0, 137.8, 138.3, 142.8, 147.4, 161.2, 170.3.

3',4'-Difluoro-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (18a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (100 mg, 0.31mmol), 3,4-difluorophenylboronic acid (65 mg, 0.41mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow oil (97 mg, 88%). C₂₁H₁₇F₂NO₂; MW 353; MS (ESI) 354[M+H]⁺; IR (cm⁻¹) 1599, 1639, 2837, 2948; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 3.43 (s, 3H), 3.70 (s, 3H), 6.73–6.76 (m, 2H), 6.80–6.81 (m, 1H), 7.17 (t, *J* = 8.0 Hz, 1H), 7.35–7.41 (m, 1H), 7.42–7.44 (m, 2H), 7.45–7.48 (m, 1H), 7.52–7.54 (m, 2H), 7.56 (dd, *J* = 2.0 Hz, 8.0 Hz, 1H), 7.60 (dd, *J* = 2.0 Hz, 8.0 Hz, 1H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 38.3, 55.7, 113.0, 113.8, 116.6, 116.7, 118.5, 118.7, 120.2, 124.3, 124.4, 126.9, 130.1, 130.7, 137.3, 140.3, 147.3, 161.2, 170.0.

3',4'-Dimethoxy-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (16a). The title compound was prepared by reaction of 4-bromo-*N*-(3-methoxyphenyl)-*N*-methylbenzamide (**6b**) (100 mg, 0.31mmol), 3,4-dimethoxyphenylboronic acid (75 mg, 0.41mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as white solid (83 mg, 71%). C₂₃H₂₃NO₄; MW 377; mp: 153–156°C; MS (ESI) 378[M+H]⁺; IR (cm⁻¹) 1604, 1635, 2843, 2938, 2969, 3003, 3068; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.51 (s, 3H), 3.69 (s, 3H), 3.90 (s, 3H), 3.92 (s, 3H), 6.62–6.72 (m, 3H), 6.89–6.91 (d, *J* = 8.0 Hz, 1H), 7.04–7.17 (m, 3H), 7.38 (s, 4H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.6, 55.6, 56.1, 110.4, 111.6, 112.2, 113.0, 119.4, 119.6, 126.1, 129.4, 130.0, 133.2, 134.4, 142.2, 146.3, 149.2, 149.3, 160.2, 170.5.

2'-Fluoro-3'-methoxy-N-(3-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (19a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (80 mg, 0.25mmol), 2-fluoro-3-methoxyphenylboronic acid (56 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 90:10) to afford the desired product as brown solid (43 mg, 47%). $C_{22}H_{20}FNO_3$; MW 365; mp: 125–127°C; MS (ESI) 366[M+H]⁺; IR (cm⁻¹) 1592, 1644, 2837, 2948, 2975, 3009; ¹H NMR (C_2D_6CO , 300 MHz) δ (ppm) 3.44 (s, 3H), 3.70 (s, 3H), 3.90 (s, 3H), 6.72–6.81 (m, 3H), 6.95–7.00 (m, 1H), 7.10–7.20 (m, 3H), 7.39–7.45 (m, 4H); ¹³C NMR (C_2D_6CO , 75 MHz) δ (ppm) 38.3, 55.7, 56.6, 113.1, 113.8, 113.9, 120.1, 122.4, 125.2, 125.3, 129.0, 129.1, 129.6, 130.6, 137.1, 137.5, 147.3, 161.2, 170.1.

2'-Fluoro-N-(3-methoxyphenyl)-N,3'-dimethyl-[1,1'-biphenyl]-4-carboxamide (20a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (72 mg, 0.22mmol), 2-fluoro-3-methylphenylboronic acid (45 mg, 0.29mmol), cesium carbonate (215 mg, 0.66mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 5 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow solid (70 mg, 91%). $C_{22}H_{20}FNO_2$; MW 349; mp: 91–94°C; MS (ESI) 350 [M+H]⁺; IR (cm⁻¹) 1599, 1635, 2840, 2920, 2963, 3055; ¹H NMR (C_2D_6CO , 500 MHz) δ (ppm) 2.29 (d, $J = 2.5$ Hz, 3H), 3.44 (s, 3H), 3.70 (s, 3H), 6.73–6.77 (m, 2H), 6.79–6.80 (m, 1H), 7.13 (t, $J = 8.0$ Hz, 1H), 7.18 (t, $J = 8.0$ Hz, 1H), 7.25 (t, $J = 8.0$ Hz, 1H), 7.39–7.44 (m, 4H); ¹³C NMR (C_2D_6CO , 125 MHz) δ (ppm) 14.6, 14.7, 38.3, 55.7, 113.1, 113.8, 120.1, 125.0, 125.1, 126.2, 126.3, 128.6, 128.7, 129.0, 129.07, 129.09, 129.6, 130.6, 131.9, 132.0, 137.0, 137.9, 147.3, 157.9, 159.9, 170.1.

4-(Benzo[d][1,3]dioxol-5-yl)-N-(3-methoxyphenyl)-N-methylbenzamide (22a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxyphenyl)-N-methylbenzamide (**6b**) (80 mg, 0.25mmol), benzo[d][1,3]dioxol-5-ylboronic acid (55 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow solid (68 mg, 76%). $C_{22}H_{19}NO_4$; MW 361; mp: 139–141°C; MS (ESI) 362 [M+H]⁺; IR (cm⁻¹) 1592, 1644, 2837, 2948, 2975, 3009; ¹H NMR (C_2D_6CO , 500 MHz) δ (ppm) 3.43 (s, 3H), 3.70 (s, 3H), 6.02 (s, 2H), 6.72–6.73 (m, 1H), 6.74–6.75 (m, 1H), 6.79–6.80 (m, 1H), 6.89–6.90 (m, 1H), 7.10–7.12 (m, 2H), 7.17 (t, $J = 8.0$ Hz, 1H), 7.37–7.39 (m, 2H), 7.44–7.46 (m, 2H); ¹³C NMR (C_2D_6CO , 125 MHz) δ (ppm) 38.4, 55.7, 102.3, 108.0, 109.4, 112.9, 113.8, 120.1, 121.4, 126.5, 130.0, 130.6, 135.0, 136.2, 142.4, 147.4, 148.5, 149.3, 161.2, 170.2.

3'-Methoxy-N-(4-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (11a). The title compound was prepared by reaction of 4-bromo-N-(4-methoxyphenyl)-N-methylbenzamide (**11b**) (100 mg, 0.31mmol), 3-methoxyphenylboronic acid (61 mg, 0.40mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as colorless oil (106 mg, 98%). $C_{22}H_{21}NO_3$; MW 347; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1635, 2840, 2935; ¹H NMR (C_2D_6CO , 300 MHz) δ (ppm) 3.40 (s, 3H), 3.75 (s, 3H), 3.85 (s, 3H), 6.82–6.86 (m, 2H), 6.92 (ddd, $J = 1.0$ Hz, 2.6 Hz, 8.3 Hz, 1H), 7.12–7.19 (m, 4H), 7.32–7.40 (m, 3H), 7.50–7.53 (m, 2H); ¹³C NMR (C_2D_6CO , 75 MHz) δ (ppm) 38.7, 55.6, 55.7, 113.1, 114.3, 115.2, 120.0, 126.9, 129.2, 130.1, 130.8, 136.9, 139.0, 142.2, 142.3, 159.0, 161.2, 170.2.

4'-Methoxy-N-(4-methoxyphenyl)-N-methyl-[1,1'-biphenyl]-4-carboxamide (12a). The title compound was prepared by reaction of 4-bromo-N-(4-methoxyphenyl)-N-methylbenzamide (**11b**) (78 mg, 0.24mmol), 4-methoxyphenylboronic acid (48 mg, 0.32mmol), cesium

carbonate (235 mg, 0.72mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 90:10) to afford the desired product as yellow solid (73 mg, 88%). C₂₂H₂₁NO₃; MW 347; mp: 129–132°C; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1635, 2840, 2932, 2957; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.39 (s, 3H), 3.74 (s, 3H), 3.83 (s, 3H), 6.82–6.82 (m, 2H), 6.98–7.01 (m, 2H), 7.11–7.14 (m, 2H), 7.34–7.37 (m, 2H), 7.44–7.47 (m, 2H), 7.54–7.57 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.7, 55.6, 55.7, 115.2, 126.2, 128.8, 129.2, 130.2, 133.0, 135.9, 139.1, 142.1, 159.0, 160.6, 170.3.

3'-Methoxy-N-methyl-N-(*m*-tolyl)-[1,1'-biphenyl]-4-carboxamide (5a). The title compound was prepared by reaction of 4-bromo-N-(3-methylphenyl)-N-methylbenzamide (**5b**) (123 mg, 0.40mmol), 3-methoxyphenylboronic acid (79 mg, 0.52mmol), cesium carbonate (393 mg, 1.20mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 9 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 70:30) to afford the desired product as yellow oil (101 mg, 77%). C₂₂H₂₁NO₃; MW 331; MS (ESI) 332 [M+H]⁺; IR (cm⁻¹) 1582, 1601, 1639, 2834, 2935, 3034; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 2.24 (s, 3H), 3.42 (s, 3H), 3.84 (s, 3H), 6.90–7.00 (m, 3H), 7.08 (s, 1H), 7.11–7.18 (m, 3H), 7.33 (t, *J* = 8.0 Hz, 1H), 7.38–7.41 (m, 2H), 7.49–7.52 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 21.2, 38.5, 55.6, 113.1, 114.3, 120.0, 125.1, 126.9, 127.9, 128.4, 129.8, 130.1, 130.8, 136.7, 139.9, 142.3, 142.5, 146.2, 161.2, 170.2.

2'-Fluoro-N,3'-dimethyl-N-(*m*-tolyl)-[1,1'-biphenyl]-4-carboxamide (21a). The title compound was prepared by reaction of 4-bromo-N-(3-methylphenyl)-N-methylbenzamide (**5b**) (93 mg, 0.31mmol), 2-fluoro-3-methylphenylboronic acid (62 mg, 0.40mmol), cesium carbonate (303 mg, 0.93mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 7 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as white solid (103 mg, quantitative). C₂₂H₂₀FNO; MW 333; mp: 73–76°C; MS (ESI) 334 [M+H]⁺; IR (cm⁻¹) 1604, 1634, 2853, 2920; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.24 (s, 3H), 2.28 (d, *J* = 2.4 Hz, 3H), 3.43 (s, 3H), 6.94–6.97 (m, 1H), 6.98–7.01 (m, 1H), 7.07 (s, 1H), 7.11–7.16 (m, 2H), 7.23 –7.26 (m, 2H), 7.38–7.41 (m, 4H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 14.6, 14.7, 21.2, 38.5, 125.05, 125.09, 125.12, 126.2, 126.3, 128.5, 128.6, 128.7, 128.99, 129.01, 129.07, 129.09, 129.6, 129.8, 131.9, 132.0, 136.9, 137.8, 139.9, 146.1, 157.9, 159.9, 170.1.

4-(Benzo[d][1,3]dioxol-5-yl)-N-methyl-N-(*m*-tolyl)benzamide (22a). The title compound was prepared by reaction of 4-bromo-N-(3-methylphenyl)-N-methylbenzamide (**5b**) (79 mg, 0.26mmol), benzo[d][1,3]dioxol-5-ylboronic acid (56 mg, 0.34mmol), cesium carbonate (254 mg, 0.78mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 70:30) to afford the desired product as yellow solid (44 mg, 49%). C₂₂H₁₉NO₃; MW 345; mp: 111–113°C; MS (ESI) 346[M+H]⁺; IR (cm⁻¹) 1602, 1634, 2788, 2911, 2966; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.24 (s, 3H), 3.42 (s, 3H), 6.02 (s, 2H), 6.89 (d, *J* = 8.0 Hz, 1H), 6.94 (d, *J* = 8.0 Hz, 1H), 6.99 (d, *J* = 8.0 Hz, 1H), 7.07–7.11 (m, 3H), 7.14 (t, *J* = 8.0 Hz, 1H), 7.35–7.37 (m, 2H), 7.42–7.44 (m, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 21.2, 38.5, 102.3, 107.9, 109.4, 121.4, 125.1, 126.5, 127.9, 128.4, 129.8, 130.2, 135.1, 136.2, 139.9, 142.3, 146.3, 148.5, 149.3, 170.2.

2'-Fluoro-N-(3-methoxybenzyl)-N,3'-dimethyl-[1,1'-biphenyl]-4-carboxamide (25a). The title compound was prepared by reaction of 4-bromo-N-(3-methoxybenzyl)-N-methylbenzamide (**25b**) (200 mg, 0.60mmol), 2-fluoro-3-methylphenylboronic acid (120 mg, 0.78mmol), cesium carbonate (587 mg, 1.80mmol) and tetrakis(triphenylphosphine) palladium (0.02 eq., 14 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc70:30) to afford the desired product as yellow oil (200 mg, 87%).

C₂₃H₂₂FNO₂; MW 363; MS (ESI) 364 [M+H]⁺; IR (cm⁻¹) 1630, 2834, 2923, 3034; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.32 (s, 3H), 2.97 (s, 3H), 3.81 (s, 3H), 4.59–4.73 (m, 2H), 6.80–7.00 (m, 3H), 7.17 (t, *J* = 8 Hz, 1H), 7.27–7.32 (m, 2H), 7.33–7.36 (m, 1H), 7.57–7.59 (m, 2H), 7.62 (s, 2H); ¹³C NMR (C₂D₆CO, 125 MHz) δ (ppm) 14.6, 14.7, 37.3, 51.0, 55.5, 113.6, 114.4, 121.0, 125.1, 125.2, 126.2, 126.3, 128.1, 128.7, 128.8, 129.1, 129.2, 129.7, 129.8, 130.5, 132.0, 132.1, 137.1, 138.0, 158.0, 159.9, 161.1.

2'-Fluoro-*N*-(3-hydroxybenzyl)-*N*,3'-dimethyl-[1,1'-biphenyl]-4-carboxamide (25). To a solution of 2'-fluoro-*N*-(3-methoxybenzyl)-*N*,3'-dimethyl-[1,1'-biphenyl]-4-carboxamide (**25a**) (100 mg, 0.28mmol) in CH₂Cl₂ (10 mL) was added trifluorobromine dimethyl sulphide complex BF₃.SMe₂ (0.18 mL, 1.68mM) at room temperature. The reaction mixture was stirred overnight. The reaction was quenched with MeOH (10 mL). The solvent was removed under reduced pressure at 25°C. Water was added to dissolve inorganic salts and the precipitated was filtered off and triturated with diethyl ether to afford the desired product as yellow solid (66 mg, 67%). C₂₂H₂₀FNO₂; MW 349; mp: 123–126°C; MS (ESI) 350 [M+H]⁺; IR (cm⁻¹) 1600, 2917, 3181; ¹H NMR (C₂D₆CO, 500 MHz) δ (ppm) 2.32 (s, 3H), 2.96 (s, 3H), 4.55–4.69 (m, 2H), 6.72–6.90 (m, 3H), 7.16–7.22 (m, 2H), 7.29 (t, *J* = 8.0 Hz, 1H), 7.33 –7.36 (m, 1H), 7.57–7.58 (m, 2H), 7.62 (s, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 14.6, 14.7, 115.2, 125.1, 125.2, 126.2, 126.4, 128.0, 128.7, 128.9, 129.17, 129.21, 129.7, 129.8, 130.6, 132.0, 132.1, 137.0, 138.0, 157.3, 158.6, 160.6

3-Methoxy-*N*-(3'-methoxy-[1,1'-biphenyl]-4-yl)-*N*-methylbenzamide (24a). The title compound was prepared by reaction of *N*-(4-bromophenyl)-3-methoxy-*N*-methylbenzamide (**24b**) (80 mg, 0.25mmol), 3-methoxyphenylboronic acid (50 mg, 0.33mmol), cesium carbonate (244 mg, 0.75mmol) and tetrakis(triphenyl)phosphine palladium (0.02 eq., 6 mg) according to method B. The residue was purified by silica gel column chromatography (hexanes/EtOAc 70:30) to afford the desired product as yellow oil (59 mg, 68%). C₂₂H₂₁NO₃; MW 347; MS (ESI) 348 [M+H]⁺; IR (cm⁻¹) 1640, 2840, 2936; ¹H NMR (C₂D₆CO, 300 MHz) δ (ppm) 3.45 (s, 3H), 3.66 (s, 3H), 3.85 (s, 3H), 6.80 –6.84 (m, 1H), 6.90–6.94 (m, 3H), 7.09–7.19 (m, 3H), 7.24–7.27 (m, 2H), 7.34 (t, *J* = 8.0 Hz, 1H), 7.57–7.60 (m, 2H); ¹³C NMR (C₂D₆CO, 75 MHz) δ (ppm) 38.4, 55.6, 55.7, 113.2, 114.0, 114.9, 116.4, 120.0, 121.9, 128.3, 128.4, 129.7, 130.9, 139.0, 139.6, 142.3, 145.7, 160.1, 161.3, 170.3.

logP Determination

The logP values were calculated from ACD/Labs Percepta 2012 Release program. The logarithm of partition constant P (log P) was calculated using the “GALAS” method (Global Adjusted Locally According to Similarity). The program predicts clogP by comparing the molecule with structurally similar molecules where experimental data are known.

Biological methods

[2,4,6,7-³H]-E2 and [2,4,6,7-³H]-E1 were purchased from Perkin-Elmer, Boston. Quickszint Flow 302 scintillator fluid was bought from Zinsser Analytic, Frankfurt. Other chemicals were purchased from Sigma, Roth or Merck.

Ethics statement. Anonymized placental samples were obtained from Saarbrücken-Dudweiler Hospital's Department of Gynecology. No author involved in this study has received information about the patients. The microsomal fraction of the mouse enzyme (m17 β -HSD2) was obtained from mouse livers, which were bought from Pharmacelsus GmbH (Saarbrücken, Germany).

***h17 β -HSD1* and *h17 β -HSD2* enzyme preparation.** Cytosolic (*h17 β -HSD1*) and microsomal (*h17 β -HSD2*) fractions were obtained from human placenta according to previously

described procedures [20, 27]. Fresh tissue was homogenized and the enzymes were separated from the mitochondria, cell membrane. Nucleus and other rests by fractional centrifugation at 1000 g, 10.000 g and 150.000 g. The pellet fraction containing the microsomal *h17 β -HSD2* was used for the determination of *h17 β -HSD2* inhibition, while *h17 β -HSD1* was obtained after precipitation with ammonium sulfate from the cytosolic fraction for use of testing of *h17 β -HSD1* inhibition. Aliquots containing *h17 β -HSD1* or *h17 β -HSD2* were stored frozen.

***m17 β -HSD2* enzyme preparation and inhibition.** The microsomal fraction (*m17 β -HSD2*) was obtained from mouse liver as described for *h17 β -HSD2*. Inhibitory activities were evaluated by a method identical to the one described for the human enzyme.

***m17 β -HSD1* enzyme preparation.** Recombinant *m17 β -HSD1* enzyme was produced by transfection of HEK 293 cells with a *m17 β -HSD1* expression plasmid (coding sequence of NM_010475 in pCMV6Entry vector, OriGene Technologies, Inc.) according to a described procedure [28]. 48 hours after transfection cells were homogenized by sonication (3 x 10 s) in a buffer containing saccharose (40 mM Tris, 250 mM saccharose, 5 mM EDTA, 7 mM DTT, 1 mM PMSF, pH 7.5). Cell lysate was centrifuged (1000 g, 15 min, 4°C) and 20% glycerol was added to the supernatant before aliquots were frozen and stored at -70°C.

Inhibition of *h17 β -HSD2* and *m17 β -HSD2* in cell-free assay. Inhibitory activities were evaluated following an established method with minor modifications [29–31]. Briefly, the enzyme preparation was incubated with NAD⁺ [1500 μ M] in the presence of potential inhibitors at 37°C in a phosphate buffer (50 mM) supplemented with 20% of glycerol and EDTA 1 mM. Inhibitor stock solutions were prepared in DMSO. Final concentration of DMSO was adjusted to 1% in all samples. The enzymatic reaction was started by addition of a mixture of unlabelled- and [³H]-E2 (final concentration: 500 nM, 0.11 μ Ci). After 20 min, the incubation was stopped with HgCl₂ and the mixture was extracted with ether. After evaporation, the steroids were dissolved in acetonitrile/water (45:55). E1 and E2 were separated using acetonitrile/water (45:55) as mobile phase in a C18 RP chromatography column (Nucleodur C18, 3 μ m, Macherey-Nagel, Düren) connected to a HPLC-system (Agilent 1100 Series, Agilent Technologies, Waldbronn). Detection and quantification of the steroids were performed using a radio-flow detector (Berthold Technologies, Bad Wildbad). The conversion rate was calculated according to the following equation: %conversion = (%E1/(%E1+%E2)) \times 100. Each value was calculated from at least two independent experiments.

Inhibition of *h17 β -HSD1* and *m17 β -HSD1* in cell-free assay. The 17 β -HSD1 inhibition assay was performed similarly to the *h17 β -HSD2* test. The human cytosolic enzyme was incubated with NADH [500 μ M] while the mouse recombinant enzyme was reacted with NADPH [500 μ M]. Test compound and a mixture of unlabelled- and [³H]-E1 (final concentration: 500 nM, 0.15 μ Ci) were added and mixed for 10 min at 37°C. Further treatment of the samples and HPLC separation was carried out as mentioned above for *h17 β -HSD2*.

Inhibition of *h17 β -HSD2* in a cellular assay. Cellular *h17 β -HSD2* inhibitory activity is measured using the breast cancer cell-line MDA-MB-231 [32] (17 β -HSD1 activity negligible). [³H]-E2 (200 nM) is taken as substrate and is incubated with the inhibitor for 6 h at 37°C. After ether extraction, substrate and product are separated by HPLC and detected with a radio-flow detector. Potency is evaluated as percentage of inhibition (inhibitor concentrations used: 1250 nM and 250 nM).

Estrogen receptor affinity in a cell-free assay. The binding affinity of selected compounds to ER α and ER β was determined according to the recommendations of the US Environmental Protection Agency (EPA) by their Endocrine Disruptor Screening Program (EDSP) [24] using recombinant human proteins. Briefly, 1 nM of ER α and 4 nM of ER β , respectively, were incubated with [³H]-E2 (3 nM for ER α and 10 nM for ER β) and test compound for 16–20 h at 4°C.

The potential inhibitors were dissolved in DMSO (5% final concentration). Evaluation of non-specific-binding was performed with unlabeled E2 at concentrations 100-fold of [³H]-E2 (300 nM for ER α and 1000 nM for ER β). After incubation, ligand-receptor complexes were selectively bound to hydroxyapatite (83.5 g/LinTE-buffer). The bound complex was washed three times and resuspended in ethanol. For radiodetection, scintillator cocktail (Quickszint 212, Zinsser Analytic, Frankfurt) was added and samples were measured in a liquid scintillation counter (1450 LSC & Luminescence Counter, Perkin Elmer).

From these results the percentage of [³H]-E2 displacement by the compounds was calculated. The plot of % displacement versus compound concentration resulted in sigmoidal binding curves. The compound concentrations necessary to displace 50% of the receptor bound [³H]-E2 were determined. Unlabeled E2 IC₅₀ values were determined in each experiment and used as reference. The E2 IC₅₀ determined were 3 \pm 20% nM for ER α and 10 \pm 20% nM for ER β .

Relative Binding Affinity was determined by applying the following equation: RBA[%] = (IC₅₀(E2)/IC₅₀(compound)) · 100[24]. This results in a RBA value of 100% for E2.

After the assay was established and validated, a modification was made to increase throughput. Compounds were tested at concentrations of 1000 times the IC₅₀(E2). Compounds with less than 50% displacement of [³H]-E2 at a concentration of 1000 times IC₅₀(E2) were classified as RBA <0.1%.

Metabolic Stability in a cell-free assay. Compounds **7a**, **8a**, **16a**, **19a**, **22a**, **27a** and **27** were tested according to established method[33–35] For evaluation of phase I and II metabolic stability 1 μ M compound was incubated with 1 mg/ml pooled mammalian liver S9 fraction (BD Gentest), 2 mM NADPH regenerating system, 1 mM UDPGA and 0.1 mM PAPS at 37°C for 0, 5, 15 and 60 minutes at a final volume of 100 μ L. The incubation was stopped by precipitation of S9 enzymes with 2 volumes of cold acetonitrile containing internal standard. Concentration of the remaining test compound at the different time points was analyzed by LC-MS/MS and used to determine half-life (t_{1/2}).

Acknowledgments

We thank Josef Zapp for the NMR measurements, Larissa Müller, Isabella Mang and Martina Jankowski for the biological tests.

Author Contributions

Conceived and designed the experiments: EMG SMO. Performed the experiments: EMG GA EP CVK. Analyzed the data: EMG SMO. Contributed reagents/materials/analysis tools: EMG EP. Wrote the paper: EMG AC SMO MF RWH.

References

1. NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis and Therapy. Osteoporosis prevention, diagnosis, and therapy. *J Am Med Assoc.* 2001; 285: 785–795.
2. Vanderschueren D, Gaytant J, Boonen S, Venken K. Androgens and bone. *Curr Opin Endocrinol.* 2008; 15: 250–254.
3. Wu L, Einstein M, Geissler WM, Chan HK, Elliston KO, Andersson S. Expression cloning and characterization of human 17 beta-hydroxysteroid dehydrogenase type 2, a microsomal enzyme possessing 20 alpha-hydroxysteroid dehydrogenase activity. *J Biol Chem.* 1993; 268: 12964–12969. PMID: [8099587](#)
4. Dong Y, Qiu QQ, Debear J, Lathrop WF, Bertolini DR, Tamburini PP. 17Beta-hydroxysteroid dehydrogenases in human bone cells. *J Bone Miner Res.* 1998; 13: 1539–1546. PMID: [9783542](#)
5. Bydal P, Auger S, Poirier D. Inhibition of type 2 17beta-hydroxysteroid dehydrogenase by estradiol derivatives bearing a lactone on the D-ring: structure-activity relationships. 2004; 69: 325–342. PMID: [15219411](#)

6. Poirier D, Bydal P, Tremblay MR, Sam KM, Luu-The V. Inhibitors of type II 17 β -hydroxysteroid dehydrogenase. *Mol Cell Endocrinol*. 2001; 171: 119–128. PMID: [11165020](#)
7. Sam KM, Auger S, Luu-The V, Poirier D. Steroidal spiro-gamma-lactones that inhibit 17 β -hydroxysteroid dehydrogenase activity in human placental microsomes. *J Med Chem*. 1995; 38: 4518–4528. PMID: [7473580](#)
8. Gunn D, Akuche C, Baryza J, Blue ML, Brennan C, Campbell AM et al. 4,5-Disubstituted cis-pyrrolidones as inhibitors of type II 17 β -hydroxysteroid dehydrogenase. Part 2. SAR. *Bioorg Med Chem Lett*. 2005; 15: 3053–3057. PMID: [15890515](#)
9. Wood J, Bagi CM, Akuche C, Bacchiocchi A, Baryza J, Blue ML et al. 4,5-Disubstituted cis-pyrrolidones as inhibitors of type II 17 β -hydroxysteroid dehydrogenase. Part 3. Identification of lead candidate. *Bioorg Med Chem Lett*. 2006; 16: 4965–4968. PMID: [16806919](#)
10. Wetzel M, Marchais-Oberwinkler S, Perspicace E, Möller G, Adamski J, Hartmann RW. Introduction of an electron withdrawing group on the hydroxyphenyl naphthol scaffold improves the potency of 17 β -hydroxysteroid dehydrogenase type 2 (17 β -HSD2) inhibitors. *J Med Chem*. 2011; 54: 7547–7557. doi: [10.1021/jm2008453](#) PMID: [21972996](#)
11. Wetzel M, Gargano EM, Hinsberger S, Marchais-Oberwinkler S, Hartmann RW. Discovery of a new class of bicyclic substituted hydroxyphenylmethanones as 17 β -hydroxysteroid dehydrogenase type 2 (17 β -HSD2) inhibitors for the treatment of osteoporosis. *Eur J Med Chem*. 2012; 47: 1–17. doi: [10.1016/j.ejmech.2011.09.004](#) PMID: [21945251](#)
12. Marchais-Oberwinkler S, Xu K, Wetzel M, Perspicace E, Negri M, Meyer A et al. Structural optimization of 2,5-thiophene amides as highly potent and selective 17 β -hydroxysteroid dehydrogenase type 2 inhibitors for the treatment of osteoporosis. *J Med Chem*. 2013; 56: 167–181. doi: [10.1021/jm3014053](#) PMID: [23145773](#)
13. Gargano EM, Perspicace E, Hanke N, Carotti A, Marchais-Oberwinkler S, Hartmann RW. Metabolic stability optimization and metabolite identification of 2,5-thiophene amide 17 β -hydroxysteroid dehydrogenase type 2 inhibitors. *Eur J Med Chem*. 2014; 87: 203–219. doi: [10.1016/j.ejmech.2014.09.061](#) PMID: [25259513](#)
14. Xu K, Al-Soud YA, Wetzel M, Hartmann RW, Marchais-Oberwinkler S. Triazole ring-opening leads to the discovery of potent nonsteroidal 17 β -hydroxysteroid dehydrogenase type 2 inhibitors. *Eur J Med Chem*. 2011; 46: 5978–5990. doi: [10.1016/j.ejmech.2011.10.010](#) PMID: [22037253](#)
15. Marchais-Oberwinkler S, Kruchten P, Frotscher M, Ziegler E, Neugebauer A, Bhoga U et al. Substituted 6-phenyl-2-naphthols. Potent and selective nonsteroidal inhibitors of 17 β -hydroxysteroid dehydrogenase type 1 (17 β -HSD1): design, synthesis, biological evaluation, and pharmacokinetics. *J Med Chem*. 2008; 51: 4685–4698. doi: [10.1021/jm800367k](#) PMID: [18630892](#)
16. Bey E, Marchais-Oberwinkler S, Kruchten P, Frotscher M, Werth R, Oster A et al. Design, synthesis and biological evaluation of bis(hydroxyphenyl) azoles as potent and selective non-steroidal inhibitors of 17 β -hydroxysteroid dehydrogenase type 1 (17 β -HSD1) for the treatment of estrogen-dependent diseases. *Bioorg Med Chem*. 2008; 16: 6423–6435. doi: [10.1016/j.bmc.2008.04.073](#) PMID: [18514529](#)
17. Turner AS. Animal models of osteoporosis-necessity and limitations. *Eur Cell Mater*. 2001; 1: 66–81. PMID: [14562261](#)
18. Reinwald S, Burr D. Review of nonprimate, large animal models for osteoporosis research. *J Bone Miner Res*. 2008; 9: 1353–1368.
19. Kruchten P, Werth R, Marchais-Oberwinkler S, Frotscher M, Hartmann RW. Development of a biological screening system for the evaluation of highly active and selective 17 β -HSD1-inhibitors as potential therapeutic agents. *Mol Cell Endocrinol*. 2009; 301: 154–157. doi: [10.1016/j.mce.2008.09.035](#) PMID: [18984028](#)
20. Marchais-Oberwinkler S, Wetzel M, Ziegler E, Kruchten P, Werth R, Henn C et al. New drug-like hydroxyphenyl naphthol steroidomimetics as potent and selective 17 β -hydroxysteroid dehydrogenase type 1 inhibitors for the treatment of estrogen-dependent diseases. *J Med Chem*. 2011; 54: 534–547. doi: [10.1021/jm1009082](#) PMID: [21189020](#)
21. Tremblay MR, Lin SX, Poirier D. Chemical synthesis of 16 β -propylaminoacyl derivatives of estradiol and their inhibitory potency on type 1 17 β -hydroxysteroid dehydrogenase and binding affinity on steroid receptors. *Steroids*. 2001; 66: 821–831. PMID: [11576622](#)
22. Perspicace E, Giorgio A, Carotti A, Marchais-Oberwinkler S, Hartmann RW. Novel N-methylsulfonamide and retro-N-methylsulfonamide derivatives as 17 β -hydroxysteroid dehydrogenase type 2 (17 β -HSD2) inhibitors with good ADME-related physicochemical parameters. *Eur J Med Chem*. 2013; 69: 201–215. doi: [10.1016/j.ejmech.2013.08.026](#) PMID: [24036043](#)
23. Kavanagh KL, Jornvall H, Persson B, Oppermann U. Medium- and short-chain dehydrogenase/reductase gene and protein families: the SDR superfamily: functional and structural diversity within a family

- of metabolic and regulatory enzymes. *Cell Mol Life Sci.* 2008; 65: 3895–3906. doi: [10.1007/s00018-008-8588-y](https://doi.org/10.1007/s00018-008-8588-y) PMID: [19011750](https://pubmed.ncbi.nlm.nih.gov/19011750/)
24. U.S. Environmental Protection Agency. Protocol for the In Vitro Estrogen Receptor Saturation Binding and Competitive Binding Assays Using Rat Uterine Cytosol 2015; Available: http://www.epa.gov/endo/pubs/assayvalidation/appendix1_er_ruc.pdf
 25. Martignoni M, Groothuis GM, de Kanter R. Species differences between mouse, rat, dog, monkey and human CYP-mediated drug metabolism, inhibition and induction. *Expert Opin Drug Metab Toxicol.* 2006; 2: 875–894. PMID: [17125407](https://pubmed.ncbi.nlm.nih.gov/17125407/)
 26. Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliver Rev.* 2001; 46: 3–26.
 27. Zhu DW, Lee X, Breton R, Ghosh D, Pangborn W, Duax WL et al. Crystallization and preliminary X-ray diffraction analysis of the complex of human placental 17 beta-hydroxysteroid dehydrogenase with NADP+. *J Mol Biol.* 1993; 234: 242–244. PMID: [8230203](https://pubmed.ncbi.nlm.nih.gov/8230203/)
 28. Sam KM, Boivin RP, Tremblay MR, Auger S, Poirier D. C16 and C17 derivatives of estradiol as inhibitors of 17 beta-hydroxysteroid dehydrogenase type 1: chemical synthesis and structure-activity relationships. *Drug Des Discov. Drug* 1998; 15: 157–180.
 29. Xu K, Wetzel M, Hartmann RW, Marchais-Oberwinkler S. Synthesis and biological evaluation of spiro- δ -lactones as inhibitors of 17 β -hydroxysteroid dehydrogenase type 2 (17 β -HSD2). *Lett Drug Des Discov.* 2011; 8: 406–421.
 30. Lin SX, Yang F, Jin JZ, Breton R, Zhu DW, Luu-The V et al. Subunit identity of the dimeric 17 beta-hydroxysteroid dehydrogenase from human placenta. *J Biol Chem.* 1992; 267: 16182–16187. PMID: [1322895](https://pubmed.ncbi.nlm.nih.gov/1322895/)
 31. Puranen TJ, Poutanen MH, Peltoketo HE, Vihko PT, Vihko RK. Site-directed mutagenesis of the putative active site of human 17 beta-hydroxysteroid dehydrogenase type 1. *Biochem J.* 1994; 304: 289–293. PMID: [7998947](https://pubmed.ncbi.nlm.nih.gov/7998947/)
 32. Day JM, Tutill HJ, Newman SP, Purohit A, Lawrence HR, Nicker N et al. 17Beta-hydroxysteroid dehydrogenase Type 1 and Type 2: association between mRNA expression and activity in cell lines. *Mol Cell Endocrinol.* 2006; 248: 246–249. PMID: [16490301](https://pubmed.ncbi.nlm.nih.gov/16490301/)
 33. Di L, Kerns EH, Hong Y, Kleintop TA, McConnell OJ, Huryn DM. Optimization of a higher throughput microsomal stability screening assay for profiling drug discovery candidates. *J Biomol Screen* 2003; 4: 453–462.
 34. Moreno-Farre JW, P.; Raynud F. Analysis of Potential Drug-Drug Interactions for Anticancer Agents in Human Liver Microsomes by High Throughput Liquid Chromatography/Mass Spectrometry Assay. *Aust-Asian J Cancer* 2007; 6: 55–69.
 35. Hui JP, Stuart Grossert J, Cutler MJ, Melanson JE. Strategic identification of in vitro metabolites of 13-desmethyl spirolicide C using liquid chromatography/high-resolution mass spectrometry. *Rapid Commun Mass Spectrom.* 2012; 26: 345–354. doi: [10.1002/rcm.5336](https://doi.org/10.1002/rcm.5336) PMID: [22223322](https://pubmed.ncbi.nlm.nih.gov/22223322/)