

18 β -Glycyrrhetic acid exerts protective effects against cyclophosphamide-induced hepatotoxicity: potential role of PPAR γ and Nrf2 upregulation

Ayman M. Mahmoud¹ · Hussein S. Al Dera²

Received: 2 July 2015 / Accepted: 8 September 2015 / Published online: 19 September 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract 18 β -Glycyrrhetic acid (18 β -GA) has been proposed as a promising hepatoprotective agent. The current study aimed to investigate the protective action and the possible mechanisms of 18 β -GA against cyclophosphamide (CP)-induced liver injury in rats, focusing on the role of peroxisome proliferator-activated receptor gamma (PPAR γ) and NF-E2-related factor-2 (Nrf2). Rats were administered 18 β -GA at doses 25 and 50 mg/kg 2 weeks prior to CP injection. Five days after CP administration, animals were sacrificed and samples were collected. CP induced hepatic damage evidenced by the histopathological changes and significant increase in serum pro-inflammatory cytokines, liver marker enzymes, and liver lipid peroxidation and nitric oxide (NO) levels. 18 β -GA counteracted CP-induced oxidative stress and inflammation as assessed by restoration of the antioxidant defenses and diminishing of pro-inflammatory cytokines, lipid peroxidation, and NO production. These hepatoprotective effects appear to depend on activation of Nrf2 and PPAR γ , and subsequent suppression of nuclear factor-kappa B. In conclusion, the present study provides evidence that 18 β -GA exerts hepatoprotective effects against CP through induction of antioxidant defenses and suppression of inflammatory response. This report also confers new information that 18 β -GA protects liver against the toxic effect of

chemotherapeutic alkylating agents via activation of Nrf2 and PPAR γ .

Keywords Chemotherapy · Hepatotoxicity · Inflammation · PPARs · Nrf2

Abbreviations

18 β -GA	18 β -Glycyrrhetic acid
CP	Cyclophosphamide
PPAR γ	Peroxisome proliferator-activated receptor gamma
Nrf2	NF-E2-related factor-2
NO	Nitric oxide
NF- κ B	Nuclear factor-kappa B
iNOS	Inducible nitric oxide synthase
ROS	Reactive oxygen species
ARE	Antioxidant response element
bZIP	Basic leucine zipper
RXR	Retinoid X receptor
GL	Glycyrrhizin
CCl ₄	Carbon tetrachloride
GSH	Reduced glutathione
TBA	Thiobarbituric acid
DTNB	5,5'-Dithiobis-(2-nitrobenzoic acid)
BSA	Bovine serum albumin
CMC	Carboxymethylcellulose
ip	Intraperitoneal
PBS	Phosphate-buffered saline
ALT	Alanine aminotransferase
AST	Aspartate aminotransferase
ALP	Alkaline phosphatase
LDH	Lactate dehydrogenase
γ GT	Gamma-glutamyl transferase
TNF- α	Tumor necrosis factor alpha
IL-6	Interleukin 6

✉ Ayman M. Mahmoud
ayman.mahmoud@science.bsu.edu.eg;
aymano911@yahoo.com

¹ Physiology Division, Zoology Department, Faculty of Science, Beni-Suef University, Beni Suef 62514, Egypt

² Basic Medical Sciences Department, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

IL-1 β	Interleukin 1beta
ELISA	Enzyme-linked immunosorbent assay
H&E	Hematoxylin and eosin
MDA	Malondialdehyde
SOD	Superoxide dismutase
CAT	Catalase
GPx	Glutathione peroxidase
qPCR	Quantitative polymerase chain reaction
TBS	Tris-buffered saline
ANOVA	Analysis of variance
SD	Standard deviation
Keap 1	Kelch-like ECH2-associated protein
PPREs	Peroxisome proliferator response elements
STATs	Signal transducer and activator of transcription
AP-1	Activator protein 1

Introduction

Hepatotoxicity represents the most important cause of the non-approval and withdrawal of drugs by the Food and Drug Administration (Mahmoud 2014). Cyclophosphamide (CP) is an alkylating agent used for treating a variety of cancers and as an immunosuppressive agent for organ transplantation (Shanafelt et al. 2007; Uber et al. 2007). The therapeutic uses of CP are often restricted due to its toxicity and wide adverse side effects (Papaldo et al. 2005). As a metabolically active organ, liver is particularly susceptible to reactive oxygen species (ROS) produced as by-products of normal metabolism and detoxification processes (Klaassen et al. 2008). CP-induced ROS generation and oxidative stress have been implicated in its hepatotoxic effects (Motawi et al. 2010). In addition, the cytochrome p450-mediated biotransformation of CP leads to formation of phosphoramidate mustard and acrolein which are highly toxic (Kern and Kehrer 2002) and have the potential to generate superfluous ROS (Ahmadi et al. 2008).

Elevated ROS and electrophiles activate the antioxidant response element (ARE) leading to induction of antioxidant genes to protect cells against oxidative stress (Mathers et al. 2004). ARE-driven antioxidant gene expression is primarily regulated by NF-E2-related factor-2 (Nrf2; Cople et al. 2010; Wang et al. 2010). Nrf2 is a member of the cap 'n' collar family of basic leucine zipper (bZIP) transcription factors (Kobayashi and Yamamoto 2005). Nrf2 plays complex and multicellular roles in hepatic fibrosis, inflammation, hepatocarcinogenesis, and regeneration via its target gene induction (Shin et al. 2013). Upon cell stimulation, activated Nrf2 is translocated into the nucleus where it binds to the ARE and leads to expression of target genes (Farombi et al. 2008; Hong et al. 2010). Therefore, Nrf2 plays a role as multiorgan protector against

oxidative stress through target gene induction (Lee et al. 2005).

Peroxisome proliferator-activated receptors (PPARs) are proteins that belong to the nuclear receptor family of ligand-activated transcription factors. Upon ligand binding, they form heterodimers with retinoid X receptor (RXR) and result in modulation of gene transcription (Barish et al. 2006). There are three major isoforms of PPARs, including PPAR α , PPAR β/δ , and PPAR γ (Michalik and Wahli 2008). In the liver, PPARs regulate inflammatory responses, cholesterol and bile acid homeostasis, carbohydrate/lipid metabolism, regenerative mechanisms, and cell differentiation/proliferation (Peyrou et al. 2012). Dysregulations of specific PPAR isoforms contribute to the development of a wide range of hepatic diseases (Peyrou et al. 2012). Also, several studies reported that PPAR γ deficiency in hepatic stellate cells is associated with excessive formation of fibrotic tissue in the liver (Zhang et al. 2012) and activation of PPAR γ signaling protects the liver against fibrosis (Nan et al. 2009) and drug-induced inflammation (Mahmoud 2014; Mahmoud et al. 2014). Thus, modulation of PPAR γ might represent an important strategy for the treatment of liver diseases.

Glycyrrhiza glabra L. (Liquorice) root and its ingredients are widely used as a conditioning and flavoring agent and in herbal medicines for the treatment of various inflammatory diseases (Eisenbrand 2006). The major active ingredients of liquorice root are glycyrrhizin (GL), 18 β -glycyrrhetic acid (18 β -GA), and 18 α -GA (Montoro et al. 2011). Studies have demonstrated several pharmacological effects of 18 β -GA including anti-inflammatory, anti-ulcer, anti-viral, antioxidant, and hepatoprotective properties (Pezzuto 1997; Maitraie et al. 2009; Wang et al. 2011). Recently, 18 β -GA was reported to attenuate 2-acetylaminofluorene-induced hepatotoxicity in Wistar rats (Hasan et al. 2015). Also, Chen et al. (2013) stated that upregulation of Nrf2 by glycyrrhetic acid protected mice against carbon tetrachloride (CCl₄)-induced chronic liver fibrosis. However, the issues of an involvement of PPAR γ and Nrf2 activation in the protective effects of 18 β -GA against CP-induced hepatotoxicity have not been previously investigated. In this study, we have investigated whether 18 β -GA, an active ingredient of *G. glabra*, might upregulate PPAR γ and Nrf2, leading to protection against CP-induced hepatotoxicity in rats.

Materials and methods

Chemicals

Cyclophosphamide (Endoxan) was supplied as vials from Baxter Oncology (Dusseldorf, Germany). 18 β -GA, reduced glutathione (GSH), thiobarbituric acid (TBA), 5,5'-

dithiobis-(2-nitrobenzoic acid) (DTNB), pyrogallol, Tween 20, and bovine serum albumin (BSA) were purchased from Sigma (USA). All other chemicals were of analytical grade and obtained from standard commercial supplies.

Animals and experimental design

Male Wistar rats weighing 130–150 g, obtained from the animal house of the National Research Centre (El-Giza, Egypt) were included in the present investigation. The animals were housed in plastic well-aerated cages (4 rats/cage) at normal atmospheric temperature (25 ± 2 °C) and normal 12-h light/dark cycle. Rats had free access to water and were supplied daily with laboratory standard diet of known composition. All animal procedures were undertaken with the approval of Institutional Animal Ethics Committee of Beni-Suef University (Egypt).

Rats were kept under observation for 1 week before the onset of the experiment for acclimatization and to exclude any intercurrent infection. To study the protective effects of 18 β -GA against CP-induced hepatotoxicity, twenty-four male Wistar rats were randomly allocated into four groups having six in each as follows:

Group 1 (control): received the vehicle 0.5 % carboxymethylcellulose (CMC) by oral gavage for 15 days and a single intraperitoneal (ip) injection of saline at day 16.

Group 2 (CP): received 0.5 % CMC by oral gavage for 15 days and a single ip dose of 150 mg/kg CP at day 16.

Group 3 (CP + 25 mg GA): received 25 mg/kg GA suspended in 0.5 % CMC by oral gavage for 15 days and a single ip dose of 150 mg/kg CP at day 16.

Group 4 (CP + 50 mg GA): received 50 mg/kg GA suspended in 0.5 % CMC by oral gavage for 15 days and a single ip dose of 150 mg/kg CP at day 16.

The doses of 18 β -GA were balanced consistently as indicated by any change in body weight to keep up comparable dosage for every kg body weight over the entire period of study.

Samples preparation

At the end of the experiment (5 days after CP administration), animals were sacrificed under ether anesthesia and blood samples were collected, left to coagulate, and centrifuged at $1000 \times g$ for 15 min to separate serum. Sera were then collected and kept at -20 °C as aliquots for subsequent biochemical assays. Liver samples were immediately excised and perfused with ice-cold phosphate-buffered saline (PBS). Frozen samples (10 % w/v) were homogenized in chilled PBS, and the homogenates were cen-

trifuged at $1500 \times g$ for 10 min. The clear homogenates were collected and used for subsequent assays. Samples from the liver were taken at the same time on RIPA buffer supplemented with proteinase inhibitors for western blotting analysis. Other liver samples were immediately excised, perfused with ice-cold PBS, and fixed in 10 % formalin for histological processing or kept frozen at -80 °C for gene expression analysis.

Biochemical assays

Assay of liver function markers

Serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were determined according to the method of Schumann and Klauke (2003) using reagent kit purchased from Biosystems (Spain). Serum alkaline phosphatase (ALP) and lactate dehydrogenase (LDH) activities were measured using Spinreact (Spain) reagent kit according to the method of Wenger et al. (1984) and Teitz (1986), respectively. Serum gamma-glutamyl transferase (γ GT) assay was performed according to Persijn and van der Slik (1976) using kits from Reactivos GPL (Spain). Concentration of albumin in serum was assayed following the methods of Webster (1974) using reagent kits supplied by Spinreact (Spain).

Assay of serum cytokines

Serum levels of tumor necrosis factor alpha (TNF- α), interleukin 6 (IL-6), and IL-1 β were determined using specific ELISA kits (R&D Systems, Inc., Minneapolis, MN, USA) following the manufacturer's instructions. The concentrations of assayed cytokines were measured spectrophotometrically at 450 nm. Standard curves were constructed by using standard cytokines, and concentrations of the unknown samples were determined from the standard plots.

Assay of oxidative stress and antioxidant defense system

Lipid peroxidation content was assayed in liver homogenates by measurement of malondialdehyde (MDA) formation according to the method of Preuss et al. (1998). Nitric oxide (NO) was determined according to the method of Montgomery and Dymock (1961) using reagent kit purchased from Biodiagnostics (Egypt). Reduced glutathione (GSH) content and activities of the antioxidant enzymes, glutathione peroxidase (GPx), superoxide dismutase (SOD), and catalase (CAT) were measured according to the methods of Beutler et al. (1963), Matkovic et al. (1998), Marklund and Marklund (1974), and Cohen et al. (1970), respectively.

Histopathological study

The liver samples were flushed with PBS and then fixed in 10 % buffered formalin for 24 h. After fixation, the specimens were dehydrated in ascending series of ethanol, cleared in xylene, and embedded in paraffin wax. Blocks were made, and 4- μ m-thick sections were cut by a sledge microtome. The paraffin-embedded liver sections were deparaffinized, washed with PBS, and stained with hematoxylin and eosin (H&E). The stained slides were examined under light microscope.

RNA isolation and quantitative reverse transcription polymerase chain reaction (qRT-PCR)

Gene expression analysis was performed as we previously described (Mahmoud 2014). In brief, total RNA was isolated from frozen liver samples using Fermentas RNA isolation kit and concentrations were quantified at 260 nm. RNA samples with A260/A280 ratios ≥ 1.7 were selected. Reverse transcription of RNA to cDNA was performed with 1 μ g RNA using reverse transcription kit (Fermentas). Synthesized cDNA was amplified by SYBR Green master mix (Fermentas) in a total volume of 20 μ L using the primer set described in Table 1. Reactions were seeded in a 96-well plate, and the PCR cycles included initial denaturation at 95 °C for 10 min and 35 cycles of denaturation at 95 °C for 30 s, annealing at T_m-5 for 60 s and extension at 72 °C for 30 s. The obtained amplification data were analyzed by the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen 2001), and the values were normalized to GAPDH.

Western blotting analysis

Western blotting for liver Nrf2, PPAR γ , NF- κ B, and iNOS was performed by using the standard method. Equal amounts of proteins were separated by 10 % SDS polyacrylamide gel electrophoresis and electro-transferred to PVDF membrane. The membranes were blocked in 5 % w/v skimmed milk powder in Tris-buffered saline (TBS)/Tween 20 (TBST) for 1 h at room temperature. The membranes were then incubated with rabbit primary antibodies for Nrf2, PPAR γ , NF- κ B p65, iNOS, and GAPDH (Santa Cruz Biotechnology, USA) diluted in blocking buffer overnight at 4 °C. After washing with TBST, the membranes were incubated with peroxidase-conjugated anti-rabbit secondary antibody for 1 h at room temperature and then washed and developed using an enhanced chemiluminescence kit (Thermo Scientific). The band intensity was quantified using ImageJ, normalized to GAPDH, and presented as % of control.

Statistical analysis

Data were analyzed using GraphPad Prism 5 software (GraphPad Software, San Diego, CA, USA), and all statistical comparisons were made by means of the one-way ANOVA test followed by Tukey's post hoc analysis. Results were articulated as mean \pm standard deviation (SD), and a P value <0.05 was considered significant.

Table 1 Primer pairs used for qPCR

Gene	GenBank accession number	Sequence 5'-3'
SOD3	NM_012880.1	F: ACACCTATGCACTCCACAGAC R: ACATTCGACCTCTGGGGGTA
GPX2	NM_183403.2	F: GCATGGCTTACATCGCCAAG R: AGTCCCGGTAGTTGTTCT
CAT	M25670.1	F: GCGGGAACCCAATAGGAGAT R: CAGGTTAGGTGTGAGGGACA
PPAR γ	NM_001145367.1	F: GGACGCTGAAGAAGAGACCTG R: CCGGGTCCTGTCTGAGTATG
Nrf2	NM_031789.2	F: TTGTAGATGACCATGAGTCGC R: TGTCTGCTGTATGCTGCTT
iNOS	U03699.1	F: ATCCCAGCCCAACAACACA R: GCAGCTTGCCAGGGATTCT
NF- κ B	AF079314.1	F: TCTCAGCTGCGACCCCG R: TGGGCTGCTCAATGATCTCC
β -Actin	NM_031144.3	F: TACAACCTTCTTGCAGCTCCT R: CCTTCTGACCCATACCCACC

Results

Effect of 18 β -GA on liver function markers

The effects of CP-induced hepatotoxicity and the preventive effects of 18 β -GA on serum markers of liver function are shown in Table 2. Serum activity of the liver function marker enzymes (AST, ALT, ALP, LDH, and γ GT) was significantly ($P < 0.001$) increased in CP-administered rats in comparison with the control rats. Conversely, serum levels of albumin were significantly decreased in CP-administered rats compared with control rats. Oral administration of 18 β -GA to CP-intoxicated rats significantly ameliorated the altered liver markers in a dose-dependent manner. The 25-mg dose of 18 β -GA significantly ameliorated serum activities of all assayed enzymes, while its effect on serum albumin levels was nonsignificant ($P > 0.05$) compared to the CP-administered rats. On the other hand, the 50-mg dose of 18 β -GA significantly ameliorated serum ALT, AST, LDH, and albumin when compared with the lower dose.

Effect of 18 β -GA on serum cytokines

Serum levels of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 were significantly ($P < 0.001$) elevated in CP-administered rats when compared with the control group (Table 3). Both the 25- and 50-mg doses of 18 β -GA significantly ($P < 0.001$) decreased the elevated cytokines level when supplemented prior to CP administration. The 50-mg dose of 18 β -GA seemed to be more effective ($P < 0.05$) in reducing TNF- α when compared with the lower dose.

Histopathological changes

Microscopic investigation of H&E-stained liver section from control rats demonstrated normal hepatic strands,

hepatocytes, and sinusoids (Fig. 1a). Liver sections from CP-induced rats showed hepatocellular focal degeneration and necrosis, hyperchromatic nuclei, hepatic cell karyolysis (Fig. 1b), fatty degeneration, cytoplasmic vacuolations, hydropic degeneration of hepatocytes (Fig. 1c), leukocytes infiltration and hyperproliferation around the deformed bile ducts, dysplasia, karyomegaly, nuclear atypia, and some fibrotic changes (Fig. 1d). On the contrary, 18 β -GA-pretreated rats exhibited marked alleviation of the liver architecture with mild hepatocyte degenerations (Fig. 1e, f).

Effect of 18 β -GA on hepatic oxidative stress and antioxidant markers

CP administration produced a significant ($P < 0.001$) increase in hepatic levels of the lipid peroxidation marker MDA and NO when compared with the corresponding control rats as depicted in Fig. 2a, b, respectively. Supplementation of CP-administered rats with either dose of 18 β -GA potentially ($P < 0.001$) ameliorated the altered levels of MDA and NO. Although nonsignificant, the higher dose of 18 β -GA was more effective in decreasing MDA and NO levels in the liver of CP-intoxicated rats when compared with the lower supplemented dose.

Conversely, GSH levels in the liver of CP-administered rats showed a significant ($P < 0.05$) decrease when compared with the control group (Fig. 2c). Oral supplementation of 25 mg 18 β -GA significantly ($P < 0.05$) prevented the decline in GSH levels in liver tissue of the CP-administered rats. Administration of the higher 18 β -GA dose produced a significant ($P < 0.01$) increase in hepatic GSH levels.

Similarly, activity of the antioxidant enzymes GPx, SOD, and CAT showed a significant ($P < 0.01$) decrease in liver of the CP-administered rats when compared with the control group (Fig. 2d–f). Oral supplementation of 25 mg

Table 2 Serum markers of liver function in control, CP, and CP rats pretreated with 18 β -GA

Parameter	Control	CP	CP + 25 mg GA	CP + 50 mg GA
ALT (U/L)	23.58 \pm 4.51	85.44 \pm 9.77***	49.22 \pm 5.78####	33.26 \pm 5.02#### ^{SS}
AST (U/L)	39.48 \pm 8.03	106.51 \pm 10.72***	65.18 \pm 11.31###	45.31 \pm 8.69### ^S
ALP (U/L)	60.11 \pm 6.29	149.43 \pm 13.42***	81.55 \pm 10.58###	72.19 \pm 15.70###
γ GT (U/L)	11.68 \pm 4.07	31.42 \pm 5.16***	17.44 \pm 3.89##	14.73 \pm 4.26###
LDH (U/L)	241.19 \pm 18.81	467.22 \pm 29.71***	317.55 \pm 18.93####	246.02 \pm 21.82#### ^{SS}
Albumin (g/dL)	3.09 \pm 0.63	1.97 \pm 0.31*	2.80 \pm 0.42	3.11 \pm 0.53 [#]

Data are M \pm SD ($N = 6$)

ALT alanine aminotransferase, AST aspartate aminotransferase, ALP alkaline phosphatase, γ GT gamma-glutamyl transferase, LDH lactate dehydrogenase, CP cyclophosphamide, GA glycyrrhethinic acid, SD standard deviation, vs versus

* $P < 0.05$ and *** $P < 0.001$ versus control, # $P < 0.05$, ## $P < 0.01$, and #### $P < 0.001$ versus CP, and ^{SS} $P < 0.01$ versus CP + 25 mg GA group

Table 3 Serum levels of TNF- α , IL-1 β , and IL-6 in control, CP, and CP rats pretreated with 18 β -GA

Parameter	Control	CP	CP + 25 mg GA	CP + 50 mg GA
TNF- α (pg/mL)	34.17 \pm 5.02	124.33 \pm 14.32***	62.08 \pm 8.79 ^{###}	40.19 \pm 3.99 ^{###S}
IL-1 β (pg/mL)	26.75 \pm 4.36	89.43 \pm 13.76***	39.34 \pm 7.76 ^{###}	32.11 \pm 6.27 ^{###}
IL-6 (pg/mL)	39.29 \pm 6.58	93.69 \pm 7.81***	65.92 \pm 8.94 ^{###}	47.01 \pm 7.07 ^{###}

Data are M \pm SD (N = 6)

TNF- α tumor necrosis factor alpha, IL interleukin, CP cyclophosphamide, GA glycyrrhetic acid, SD standard deviation, vs versus

*** P < 0.001 versus control, ^{###} P < 0.001 versus CP, and ^S P < 0.05 versus CP + 25 mg GA group




Fig. 1 Photomicrographs of H&E-stained liver sections of control, CP, and CP rats pretreated with 18 β -GA. **a** Liver section of control rat showing normal histological structure, **b–d** liver sections of CP-induced rats showing hepatocellular focal degeneration and necrosis (n), hyperchromatic nuclei (blue arrows), hepatic cell karyolysis (asterisk), fatty degeneration (white arrow), cytoplasmic vacuolations (v), hydropic degeneration of hepatocytes, leukocytes infiltration, and


hyperproliferation around the deformed bile ducts, dysplasia, karyomegaly, nuclear atypia (brown arrows), and some fibrotic changes, **e** liver section of rats pretreated with 25 mg 18- β GA showing normal hepatocytes and mild central vein and sinusoid congestion, and **f** liver section of rats pretreated with 50 mg 18- β GA showing normal hepatocytes with mild degeneration (\times 400)

18 β -GA increased the activity of GPx ($P < 0.01$) and SOD ($P < 0.05$); however, its effect on CAT activity was non-significant ($P > 0.05$) as compared with the CP-administered group. On the other hand, the 50-mg dose of 18 β -GA significantly alleviated the activity of GPx ($P < 0.01$), SOD ($P < 0.01$), and CAT ($P < 0.05$) when compared with the CP control group of rats.

Effect of 18 β -GA on SOD, GPx, and CAT gene expression

Gene expression analysis showed significant downregulation of hepatic SOD ($P < 0.01$), GPx ($P < 0.001$), and CAT ($P < 0.01$) in CP-intoxicated rats when compared with the corresponding controls (Fig. 3a–c). Pretreatment

Fig. 2 Effect of 18 β -GA on liver oxidative stress and antioxidant defense system parameters. Data are M \pm SD ($N = 6$). * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ versus control, and # $P < 0.05$, ## $P < 0.01$, and ### $P < 0.001$ versus CP group. *MDA* malondialdehyde, *NO* nitric oxide, *GSH* glutathione, *GPx* glutathione peroxidase, *SOD* superoxide dismutase, *CAT* catalase, *CP* cyclophosphamide, *GA* glycyrrhethinic acid, *SD* standard deviation, *vs* versus



of the CP-induced rats with both doses of 18 β -GA produced significant ($P < 0.001$) upregulation of the antioxidant enzymes mRNA. The 50 mg 18 β -GA dosage seemed to be more effective in ameliorating the altered expression levels of SOD and GPx.

Effect of 18 β -GA on Nrf2, PPAR γ , NF- κ B, and iNOS gene expression

qPCR analysis of gene expression showed a significant downregulation of both Nrf2 ($P < 0.05$) and PPAR γ ($P < 0.01$) in the liver of CP-intoxicated rats when compared with the corresponding controls (Fig. 4a, b). Oral supplementation of 25 mg 18 β -GA produced nonsignificant ($P < 0.05$) upregulation of PPAR γ mRNA when compared with the CP-administered rats, while Nrf2 was significantly ($P < 0.05$) upregulated. The 50 mg 18 β -GA dose was more effective where it produced significant

($P < 0.01$) upregulation of both Nrf2 and PPAR γ in liver of CP-administered rats.

On the other hand, CP-intoxicated rats exhibited significant ($P < 0.001$) upregulation of NF- κ B and iNOS mRNA expression when compared with the control rats, as represented in Fig. 4c, d. Supplementation of either dose of 18 β -GA potentially downregulated NF- κ B and iNOS mRNA expression in liver of the CP-intoxicated rats.

Effect of 18 β -GA on Nrf2, PPAR γ , NF- κ B, and iNOS protein expression

CP-administered rats exhibited significant ($P < 0.001$) decrease in hepatic Nrf2 and PPAR γ protein expression when compared with the corresponding control rats as shown by western blotting (Fig. 5a, b). While supplementation of CP-administered rats with the lower dose of 18 β -GA produced a nonsignificant ($P > 0.05$) effect on hepatic




Fig. 3 Liver SOD, GPx, and CAT mRNA expression in control, CP, and CP rats pretreated with 18 β -GA. Data are M \pm SD ($N = 6$). ** $P < 0.01$ and *** $P < 0.001$ versus control, ### $P < 0.001$ versus CP, and \$\$\$ $P < 0.001$ versus CP + 25 mg GA group. SOD superoxide dismutase, GPx glutathione peroxidase, CAT catalase, CP cyclophosphamide, GA glycyrrhetic acid, SD standard deviation, vs versus

PPAR γ protein expression, it produced a significant ($P < 0.001$) increase in Nrf2 protein levels. On the other hand, oral supplementation of 50-mg dose of 18 β -GA to CP-administered rats significantly ($P < 0.001$) increased the protein levels of both Nrf2 and PPAR γ when compared with the CP control group. In addition, the higher 18 β -GA dose significantly increased PPAR γ ($P < 0.01$) and Nrf2

($P < 0.001$) protein expression when compared with its lower dose.

On the contrary, CP-intoxicated rats showed significant upregulation of NF- κ B ($P < 0.001$) and iNOS ($P < 0.01$) protein expression when compared with the control rats, as represented in Fig. 5c, d. Pretreatment of the CP-induced rats with either dose of 18 β -GA markedly downregulated NF- κ B and iNOS expression. The higher 18 β -GA produced a significant ($P < 0.05$) downregulation of iNOS when compared with the lower dose.

Discussion

The therapeutic uses of CP against various cancers are often restricted because of its toxicity and wide adverse side effects. CP-induced hepatotoxicity occurs at high chemotherapeutic dosage (de Jonge et al. 2006) and at lower doses attained during treating patients with autoimmune diseases (Akay et al. 2006; Martínez-Gabarrón et al. 2011). In the present study, CP-induced acute hepatotoxicity is evident by increased serum activities of ALT, AST, LDH, ALP, and γ GT, along with the declined albumin levels. These findings are in consistent with our recent studies demonstrating increased liver marker enzymes in serum of CP-intoxicated rats (Mahmoud et al. 2013; Germoush and Mahmoud 2014; Mahmoud 2014). The elevated serum enzymes might be attributed to cell damage caused by CP-induced oxidative stress and inflammation (Dang et al. 2008). The assayed serum transaminases and other enzymes are sensitive markers of liver injury as they are found in the cytoplasm of liver cells and leak into blood circulation following cell damage (Ramaiah 2007). The induced hepatotoxicity due to CP administration was further confirmed by the histological alterations including fatty degeneration of hepatocytes, necrosis, karyolysis, inflammatory cell infiltrations, and other manifestations. Pretreatment of the CP-induced rats with either dose of 18 β -GA significantly ameliorated serum levels of liver marker enzymes in a dose-dependent manner, proving the hepatoprotective and membrane stabilizing efficacies of 18 β -GA. Accordingly, studies have demonstrated that 18 β -GA decreased serum transaminases in 2-acetylaminofluorene-induced hepatotoxicity in rats (Hasan et al. 2015) and in CCl $_4$ -induced chronic liver fibrosis in mice (Chen et al. 2013).

The current findings showed that CP administration induced marked increase in the hepatic lipid peroxidation marker MDA. The studies of Fraiser et al. (1991) and Lameire et al. (2011) reported that CP produces highly reactive electrophiles with catastrophic effects on cell membranes. Also, CP-induced hepatotoxicity has been reported to involve lipid peroxidation due to excessive

Fig. 4 Liver Nrf2, PPAR γ , NF- κ B, and iNOS mRNA expression in control, CP, and CP rats pretreated with 18 β -GA. Data are M \pm SD (N = 6). *P < 0.05, **P < 0.01, and ***P < 0.001 versus control, and #P < 0.05, ##P < 0.01, and ###P < 0.001 versus CP group. Nrf2 NF-E2-related factor-2, PPAR γ peroxisome proliferator-activated receptor gamma, NF- κ B nuclear factor-kappa B, iNOS inducible nitric oxide synthase, CP cyclophosphamide, GA glycyrrhethinic acid, SD standard deviation, vs versus





Fig. 5 Western blotting analysis of liver Nrf2, PPAR γ , NF- κ B, and iNOS protein expression in control, CP, and CP rats pretreated with 18 β -GA. (Top) Gel photograph depicting representative proteins and (Bottom) corresponding densitometric analysis. Data are M \pm SD (N = 6). **P < 0.01 and ***P < 0.001 versus control, #P < 0.05, ##P < 0.01, and ###P < 0.001 versus CP, and \$\$\$P < 0.001 versus CP + 25 mg GA group. Nrf2 NF-E2-related factor-2, PPAR γ peroxisome proliferator-activated receptor gamma, NF- κ B nuclear factor-kappa B, iNOS inducible nitric oxide synthase, CP cyclophosphamide, GA glycyrrhethinic acid, SD standard deviation, vs versus



formation of ROS (Bhatia et al. 2008). Acrolein, a metabolite of CP, and MDA belong to the carbonyl compounds which can cause structural and functional changes

in the enzymes (Tripathi and Jena 2009). CP-induced lipid peroxidation and oxidative stress has been previously reported in our studies (Mahmoud et al. 2013; Germoush

and Mahmoud 2014; Mahmoud 2014). Interestingly, pretreatment with 18 β -GA markedly protected rats against CP-induced lipid peroxidation, indicating its radical scavenging activity.

Similarly, CP-intoxicated rats exhibited significant increase in NO levels which could be explained by the induced upregulation of iNOS. The crucial role of NO in pathogenesis of the adverse effects of CP is quite well confirmed in several studies (Al-Yahya et al. 2009). In this context, we recently reported increased NO levels and upregulation of iNOS in liver of CP-induced rats (Mahmoud 2014). The produced NO could react with superoxide anions to produce the potent and versatile oxidant peroxynitrite (McKim et al. 2003), and stimulate the production of pro-inflammatory cytokines through the activation of NF- κ B in Kupffer cells (Matata and Galinanes 2002). Supplementation of 18 β -GA potentially decreased NO levels in the liver of CP-administered rats. These findings could be attributed to the ability of 18 β -GA to downregulate iNOS expression.

The observed depletion of GSH in liver of CP-administered rats could be resulted by the direct conjugation of CP metabolites with GSH (Yousefipour et al. 2005). GSH plays a significant role against oxidative stress by neutralizing the hydroxyl radicals (Circu and Aw 2011) or as a substrate for GPx (Franco et al. 2007). GSH depletion leads to declined cellular defense against free radical-induced injury resulting in necrotic cell death (Srivastava and Shivanandappa 2010). Thus, prevention of GSH depletion could be a part of the hepatoprotective mechanism of 18 β -GA. Accordingly, CP administration produced marked decrease in the hepatic activity as well as gene expression of SOD, CAT, and GPx. We have demonstrated decreased activity of the enzymatic and non-enzymatic antioxidant defenses in the liver of CP-intoxicated rats which was associated with CP-induced oxidative stress (Mahmoud 2014). SOD, CAT, and GPx play a crucial role in protecting the body against the deleterious effects of ROS and free radicals (Wei et al. 2011). Therefore, 18 β -GA seems to exert its protective effects against CP by enhancing the hepatic antioxidant defenses.

The increase in expression and activity of the antioxidant defense enzymes by 18 β -GA in the current study could be directly linked to upregulation of the transcription factor Nrf2. Nrf2 is essential in tissue protection from oxidants through binding to cis-acting ARE and subsequent induction of antioxidant and defense gene expression (Kensler et al. 2007). Under normal physiological conditions, Nrf2 is located in the cytoplasm inactivated by forming a complex with its repressor Kelch-like ECH2-associated protein (Keap 1) (Kang et al. 2004). Upon stimulation, Nrf2 dissociates from Keap 1, translocates into the nucleus, binds to ARE, and induces expression of the

antioxidant enzymes SOD, CAT, and GPx (Patel and Maru 2008; Bardag-Gorce et al. 2011). Therefore, upregulation of Nrf2 can result in a reduction in the level of reactive oxidants by increasing expression of the antioxidant defense enzymes and correspondingly, less cell injury. Our findings showed a dramatic downregulation of Nrf2 mRNA and protein expression in liver tissue of CP-administered rats which was reversed by 18 β -GA pretreatment. The expression level of Nrf2 is strongly correlated with the expression levels of SOD, CAT, and GPx; however, the exact mechanism of 18 β -GA-induced upregulation of Nrf2 needs further investigation.

During drug-induced toxicity, tissue damage leads to generation of inflammatory mediators by the immune cells as well as by injured cells (Akcaay et al. 2009). Various inflammatory mediators produced during drug-induced hepatic injury have been reported to promote tissue damage (Ishida et al. 2002). The generated inflammatory mediators induce migration and infiltration of leukocytes into the site of injury and provoke the primary injury induced by the toxicant (Akcaay et al. 2009). Also, hepatotoxicity-associated inflammatory cytokines were reported to mediate monocyte/Kupffer cell activation and increase vascular permeability and apoptosis of hepatocytes (Mohammed et al. 2004). In the present study, CP administration induced significant increase in serum levels of TNF- α , IL-1 β , and IL-6. These findings are similar to those observed in our recent studies (Germoush and Mahmoud 2014; Mahmoud 2014) where we reported marked increase in circulatory levels of pro-inflammatory cytokines in CP-treated rats. Through binding to their receptors, IL-1 β and TNF- α elicit potent pro-inflammatory actions (Ishida et al. 2002) and activate the pro-apoptotic caspase cascade (Tacke et al. 2009), respectively. The elevated levels of circulatory inflammatory cytokines are in consistent with the recorded significant upregulation of NF- κ B, induced by ROS, which is well known to regulate the expression of various genes including inflammatory cytokines and iNOS (Pikarsky et al. 2004). CP promoted upregulation of NF- κ B and iNOS with subsequent production of NO, and pro-inflammatory cytokines reflect the degree of inflammation in the induced rats. Pretreatment of rats with both doses of 18 β -GA potentially protected against CP-induced inflammation through downregulation of NF- κ B and iNOS expression, and attenuation of pro-inflammatory cytokines production.

These interesting findings provide strong evidence on the anti-inflammatory efficacy of 18 β -GA, which is one of the significant mechanisms for prevention of drug-induced hepatotoxicity. The anti-inflammatory and hepatoprotective effects of 18 β -GA may be explained, at least in part, via its ability to upregulate PPAR γ expression in the liver. After interaction with ligands, PPAR γ translocates into the

nucleus and heterodimerizes with RXR. This complex binds to peroxisome proliferator response elements (PPREs), found in the promoters of PPAR responsive genes, and thereby controls their expression (Berger and Moller 2002). Activated PPAR γ has been reported to decrease the activity of signal transducer and activator of transcription (STATs), NF- κ B, and activator protein 1 (AP-1). This subsequently suppresses synthesis of iNOS (Pascual et al. 2005) and pro-inflammatory cytokines (De Bosscher et al. 2006). In this context, we recently reported that PPAR γ activation protected against CP and isoniazid induced oxidative stress and inflammation in rats (Mahmoud 2014; Mahmoud et al. 2014). Also, studies have demonstrated that PPAR γ activation reduces generation of ROS from leukocytes of obese subjects (Garg et al. 2000), decreases production of superoxide radicals from monocytes in diabetic patients (Marfella et al. 2006), and eliminates oxidative stress in rodents (Dobrian et al. 2004). Furthermore, PPAR γ upregulated the expression of CAT (Okuno et al. 2008) and SOD (Gong et al. 2012) in rodents through binding to its elements in the promoters of their genes.

The hepatoprotective effects of 18 β -GA in the present investigation could be attributed to co-activation and possible interaction between PPAR γ and Nrf2. This interaction occurs through multiple mechanisms. First, ARE and PPRE coexist in the same genes such as CAT (Kwak et al. 2001; Girnun et al. 2002); second, a reciprocal transcriptional regulation exists between PPAR γ and Nrf2 genes, PPAR γ gene contains ARE (Cho et al. 2010), and conversely, Nrf2 gene appears to contain PPREs (Shih et al. 2005); third, the effect of Nrf2 in ameliorating oxidative stress was proposed to be through NF- κ B inhibition (Bowie and O'Neill 2000). Hence, an interaction between Nrf2 and PPAR γ may be through inhibition of NF- κ B. However, a limitation of this study was to determine Nrf2 and PPAR γ in nuclear extracts for achieving better results.

In conclusion, the present study confers new information on the protective mechanism of 18 β -GA against CP-induced liver injury. 18 β -GA has shown strong modulatory potential against CP-induced inflammation and oxidative stress via induction of antioxidant defenses and suppression of ROS production. These hepatoprotective effects appear to depend on the co-activation of Nrf2 and PPAR γ , and subsequent suppression of NF- κ B. It seems that the simultaneous activation of both PPAR γ and Nrf2 may have a synergistic effect in protecting against CP-induced hepatotoxicity.

Funding This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors' contribution AMM designed and conceived the study, performed experiments (animal treatments, and biochemical, gene

expression and histological analysis), analyzed data, and drafted the manuscript. HSA participated in western blotting. Both authors read and approved the manuscript.

Compliance with ethical standards

Conflict of interest Ayman Mahmoud and Hussein Al-Dera declare that they have no conflict of interest.

Ethical standard All institutional and national guidelines for the care and use of laboratory animals were followed.

References

- Ahmadi A, Hosseinimehr S, Naghshvar F, Hajir E, Ghahremani M (2008) Chemoprotective effects of hesperidin against genotoxicity induced by cyclophosphamide in mice bone marrow cells. *Arch Pharm Res* 31:794–797
- Akay H, Akay T, Secilmis S, Kocak Z, Donderici O (2006) Hepatotoxicity after low-dose cyclophosphamide therapy. *South Med J* 99:1399–1400
- Akay A, Nguyen Q, Edelstein CL (2009) Mediators of inflammation in acute kidney injury. *Mediators Inflamm* 2009:137072
- Al-Yahya AA, Al-Majed AA, Gado AM, Daba MH, Al-Shabanah OA, Abd-Allah AR (2009) Acacia senegal gum exudate offers protection against cyclophosphamide-induced urinary bladder cytotoxicity. *Oxid Med Cell Longev* 2:207–213
- Bardag-Gorce F, Oliva J, Lin A, Li J, French BA, French SW (2011) Proteasome inhibitor up regulates liver antioxidant enzymes in rat model of alcoholic liver disease. *Exp Mol Pathol* 90:123–130
- Barish GD, Narkar VA, Evans RM (2006) PPAR δ : a dagger in the heart of the metabolic syndrome. *J Clin Invest* 116:590–597
- Berger J, Moller DE (2002) The mechanisms of action of PPARs. *Annu Rev Med* 53:409–435
- Beutler E, Duron O, Kelly BM (1963) Improved method for the determination of blood glutathione. *J Lab Clin Med* 61:882–888
- Bhatia K, Ahmad F, Rashid H, Raisuddin S (2008) Protective effect of S-allylcysteine against cyclophosphamide-induced bladder hemorrhagic cystitis in mice. *Food Chem Toxicol* 46:3368–3374
- Bowie A, O'Neill LA (2000) Oxidative stress and nuclear factor-kappa B activation: a reassessment of the evidence in the light of recent discoveries. *Biochem Pharmacol* 59:13–23
- Chen S, Zou L, Li L, Wu T, Uversky VN (2013) The protective effect of glycyrrhetic acid on carbon tetrachloride-induced chronic liver fibrosis in mice via upregulation of Nrf2. *PLoS ONE* 8:e53662
- Cho HY, Gladwell W, Wang X et al (2010) Nrf2-regulated PPAR γ expression is critical to protection against acute lung injury in mice. *Am J Respir Crit Care Med* 182:170–182
- Circu ML, Aw TY (2011) Redox biology of the intestine. *Free Radic Res* 45:1245–1266
- Cohen G, Dembiec D, Marcus J (1970) Measurement of catalase activity in tissue extracts. *Anal Biochem* 34:30–38
- Copple IM, Goldring CE, Kitteringham NR, Park BK (2010) The Keap1-Nrf2 cellular defense pathway: mechanisms of regulation and role in protection against drug-induced toxicity. *Handb Exp Pharmacol* 196:233–266
- Dang K, Lamb K, Cohen M, Bielefeldt K, Gebhart GF (2008) Cyclophosphamide-induced bladder inflammation sensitizes and enhances P2X receptor function in rat bladder sensory neurons. *J Neurophysiol* 99:49–59
- De Bosscher K, Vanden Berghe W, Haegeman G (2006) Cross-talk between nuclear receptors and nuclear factor kappa B. *Oncogene* 25:6868–6886

- de Jonge ME, Huitema AD, Beijnen JH, Rodenhuis S (2006) High exposures to bioactivated cyclophosphamide are related to the occurrence of veno-occlusive disease of the liver following high-dose chemotherapy. *Br J Cancer* 94:1226–1230
- Dobrian AD, Schriver SD, Khraibi AA, Prewitt RL (2004) Pioglitazone prevents hypertension and reduces oxidative stress in diet-induced obesity. *Hypertension* 43:48–56
- Eisenbrand G (2006) Glycyrrhizin. *Mol Nutr Food Res* 50:1087–1088
- Farombi EO, Shrotriya S, Na H-K, Kim S-H, Surh Y-J (2008) Curcumin attenuates dimethylnitrosamine-induced liver injury in rats through Nrf2-mediated induction of heme oxygenase-1. *Food Chem Toxicol* 46:1279–1287
- Fraiser LH, Kanekal S, Kehrer JP (1991) Cyclophosphamide toxicity: characterising and avoiding the problem. *Drugs* 42:781–795
- Franco R, Schoneveld OJ, Pappa A, Panayiotidis MI (2007) The central role of glutathione in the pathophysiology of human diseases. *Arch Physiol Biochem* 113:234–258
- Garg R, Kumbkarni Y, Aljada A et al (2000) Troglitazone reduces reactive oxygen species generation by leukocytes and lipid peroxidation and improves flow-mediated vasodilatation in obese subjects. *Hypertension* 36:430–435
- Germoush MO, Mahmoud AM (2014) Berberine mitigates cyclophosphamide-induced hepatotoxicity by modulating antioxidant status and inflammatory cytokines. *J Cancer Res Clin Oncol* 140:1103–1109
- Girnun GD, Domann FE, Moore SA, Robbins ME (2002) Identification of a functional peroxisome proliferator-activated receptor response element in the rat catalase promoter. *Mol Endocrinol* 16:2793–2801
- Gong P, Xu H, Zhang J, Wang Z (2012) PPAR expression and its association with SOD and NF- κ B in rats with obstructive jaundice. *Biomed Res India* 23:551–560
- Hasan S, Khan R, Ali N et al (2015) 18- β glycyrrhetic acid alleviates 2-acetylaminofluorene-induced hepatotoxicity in Wistar rats: role in hyperproliferation, inflammation and oxidative stress. *Hum Exp Toxicol* 34:628–641
- Hong Y, Yan W, Chen S, Cr Sun, Zhang JM (2010) The role of Nrf2 signaling in the regulation of antioxidants and detoxifying enzymes after traumatic brain injury in rats and mice. *Acta Pharmacol Sin* 31(11):1421–1430
- Ishida Y, Kondo T, Ohshima T, Fujiwara H, Iwakura Y, Mukaida N (2002) A pivotal involvement of IFN- γ in the pathogenesis of acetaminophen-induced acute liver injury. *FASEB J* 16:1227–1236
- Kang MI, Kobayashi A, Wakabayashi N, Kim SG, Yamamoto M (2004) Scaffolding of Keap1 to the actin cytoskeleton controls the function of Nrf2 as key regulator of cytoprotective phase 2 genes. *Proc Natl Acad Sci USA* 101:2046–2051
- Kensler TW, Wakabayashi N, Biswal S (2007) Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annu Rev Pharmacol* 47:89–116
- Kern JC, Kehrer JP (2002) Acrolein-induced cell death: a caspase-influenced decision between apoptosis and oncosis/necrosis. *Chem Biol Interact* 139:79–95
- Klaassen CD, Casarett LJ, Doull J (2008) Casarett and Doull's toxicology: the basic science of poisons, 7th edn. McGraw-Hill, New York
- Kobayashi M, Yamamoto M (2005) Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation. *Antioxid Redox Signal* 7:385–394
- Kwak MK, Itoh K, Yamamoto M, Sutter TR, Kensler TW (2001) Role of transcription factor Nrf2 in the induction of hepatic phase 2 and antioxidative enzymes in vivo by the cancer chemoprotective agent, 3H-1, 2-dimethiole-3-thione. *Mol Med* 7:135–145
- Lameire N, Kruse V, Rottey S (2011) Nephrotoxicity of anticancer drugs—an underestimated problem? *Acta Clin Belg* 66:337–345
- Lee JM, Li J, Johnson DA, Stein TD, Kraft AD, Calkins MJ, Jakel RJ, Johnson JA (2005) Nrf2, a multi-organ protector? *FASEB J* 19:1061–1066
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25:402–408
- Mahmoud AM (2014) Hesperidin protects against cyclophosphamide-induced hepatotoxicity by upregulation of PPAR γ and abrogation of oxidative stress and inflammation. *Can J Physiol Pharmacol* 92:717–724
- Mahmoud AM, Hussein OE, Ramadan SA (2013) Amelioration of cyclophosphamide-induced hepatotoxicity by the brown seaweed *Turbinaria ornata*. *Int J Clin Toxicol* 1:9–17
- Mahmoud AM, Germoush MO, Soliman AS (2014) Berberine attenuates isoniazid-induced hepatotoxicity by modulating peroxisome proliferator-activated receptor gamma, oxidative stress and inflammation. *Int J Pharmacol* 10:451–460
- Maitraie D, Hung CF, Tu HY et al (2009) Synthesis, anti-inflammatory, and antioxidant activities of 18 β -glycyrrhetic acid derivatives as chemical mediators and xanthine oxidase inhibitors. *Bioorg Med Chem* 17:2785–2792
- Marfella R, D'Amico M, Esposito K et al (2006) The ubiquitin-proteasome system and inflammatory activity in diabetic atherosclerotic plaques: effects of rosiglitazone treatment. *Diabetes* 55:622–632
- Marklund S, Marklund G (1974) Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 47:469–474
- Martínez-Gabarrón M, Enríquez R, Sirvent AE, García-Sepulcre M, Millán I, Amorós F (2011) Hepatotoxicity following cyclophosphamide treatment in a patient with MPO-ANCA vasculitis. *Nefrologia* 31:496–498
- Matata BM, Galiñanes M (2002) Peroxynitrite is an essential component of cytokines production mechanism in human monocytes through modulation of nuclear factor-kappa B DNA binding activity. *J Biol Chem* 277:2330–2335
- Mathers J, Fraser JA, McMahon M, Saunders RD, Hayes JD, McLellan LI (2004) Antioxidant and cytoprotective responses to redox stress. *Biochem Soc Symp* 71:157–176
- Matkovics B, Szabo L, Varga IS (1998) Determination of enzyme activities in lipid peroxidation and glutathione pathways (in Hungarian). *Lab Diagn* 15:248–249
- McKim SE, Gäbele E, Isayama F et al (2003) Inducible nitric oxide synthase is required in alcohol-induced liver injury: studies with knockout mice. *Gastroenterology* 125:1834–1844
- Michalik L, Wahli W (2008) PPARs mediate lipid signaling in inflammation and cancer. *PPAR Res* 2008:1–15
- Mohammed NA, Abd El-Aleem SA, El-Hafiz HA, McMahon RF (2004) Distribution of constitutive (COX-1) and inducible (COX-2) cyclooxygenase in postviral human liver cirrhosis: a possible role for COX-2 in the pathogenesis of liver cirrhosis. *J Clin Pathol* 57:350–354
- Montgomery HAC, Dymock JF (1961) The determination of nitrite in water. *Analyst* 86:414–416
- Montoro P, Maldini M, Russo M, Postorino S, Piacente S, Pizza C (2011) Metabolic profiling of roots of liquorice (*Glycyrrhiza glabra*) from different geographical areas by ESI/MS/MS and determination of major metabolites by LC-ESI/MS and LC-ESI/MS/MS. *J Pharm Biomed Anal* 54:535–544
- Motawi TMK, Sadik NAH, Refaat A (2010) Cytoprotective effects of DL-alpha-lipoic acid or squalene on cyclophosphamide-induced oxidative injury: an experimental study on rat myocardium, testicles and urinary bladder. *Food Chem Toxicol* 48:2326–2336

- Nan YM, Fu N, Wu WJ et al (2009) Rosiglitazone prevents nutritional fibrosis and steatohepatitis in mice. *Scand J Gastroenterol* 44:358–365
- Okuno Y, Matsuda M, Kobayashi H et al (2008) Adipose expression of catalase is regulated via a novel remote PPAR γ -responsive region. *Biochem Biophys Res Commun* 366:698–704
- Papaldo P, Lopez M, Marolla P et al (2005) Impact of five prophylactic filgrastim schedules on hematologic toxicity in early breast cancer patients treated with epirubicin and cyclophosphamide. *J Clin Oncol* 23:6908–6918
- Pascual G, Fong AL, Ogawa S et al (2005) A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR- γ . *Nature* 437:759–763
- Patel R, Maru G (2008) Polymeric black tea polyphenols induce phase II enzymes via Nrf2 in mouse liver and lungs. *Free Radic Biol Med* 44:1897–1911
- Persijn JP, van der Slik W (1976) A new method for the determination of γ -glutamyltransferase in serum. *Z Klin Chem Klin Biochem* 14:421–427
- Peyrou M, Ramadori P, Bourgoin L, Foti M (2012) PPARs in liver diseases and cancer: epigenetic regulation by microRNAs. *PPAR Res* 2012:757803
- Pezzuto JM (1997) Plant-derived anticancer agents. *Biochem Pharmacol* 53:121–133
- Pikarsky E, Porat RM, Stein I et al (2004) NF- κ B functions as a tumour promoter in inflammation-associated cancer. *Nature* 431:461–466
- Preuss HG, Jarrell ST, Scheckenbach R, Lieberman S, Anderson RA (1998) Comparative effects of chromium, vanadium and gymnema sylvestre on sugar-induced blood pressure elevations in SHR. *J Am Coll Nutr* 17:116–123
- Ramaiah SK (2007) A toxicologist guide to the diagnostic interpretation of hepatic biochemical parameters. *Food Chem Toxicol* 45:1551–1557
- Schumann G, Klauke R (2003) New IFCC reference procedures for the determination of catalytic activity concentrations of five enzymes in serum: preliminary upper reference limits obtained in hospitalized subjects. *Clin Chim Acta* 327:69–79
- Shanafelt TD, Lin T, Geyer SM et al (2007) Pentostatin, cyclophosphamide, and rituximab regimen in older patients with chronic lymphocytic leukemia. *CNCR Cancer* 109:2291–2298
- Shih AY, Imbeault S, Barakauskas V et al (2005) Induction of the Nrf2-driven antioxidant response confers neuroprotection during mitochondrial stress in vivo. *J Biol Chem* 280:22925–22936
- Shin SM, Yang JH, Ki SH (2013) Role of the Nrf2-ARE pathway in liver diseases. *Oxid Med Cell Longev* 2013:763257
- Srivastava A, Shivanandappa T (2010) Hepatoprotective effect of the root extract of *Decalepis hamiltonii* against carbon tetrachloride-induced oxidative stress in rats. *Food Chem* 118:411–417
- Tacke F, Luedde T, Trautwein C (2009) Inflammatory pathways in liver homeostasis and liver injury. *Clin Rev Allergy Immunol* 36:4–12
- Teitz N (1986) *Fundamentals of clinical chemistry*. W. B. Saunders Co., Philadelphia
- Tripathi DN, Jena GB (2009) Intervention of astaxanthin against cyclophosphamide-induced oxidative stress and DNA damage: a study in mice. *Chem Biol Interact* 180:398–406
- Uber WE, Self SE, Van Bakel AB, Pereira NL (2007) Acute antibody-mediated rejection following heart transplantation. *Am J Transplant* 7:2064–2074
- Wang YP, Cheng ML, Zhang BF, Mu M, Wu J (2010) Effects of blueberry on hepatic fibrosis and transcription factor Nrf2 in rats. *World J Gastroenterol* 16:2657–2663
- Wang CY, Kao TC, Lo WH, Yen GC (2011) Glycyrrhizic acid and 18 β -glycyrrhetic acid modulate lipopolysaccharide-induced inflammatory response by suppression of NF- κ B through PI3 K p110 δ and p110 γ inhibitions. *J Agric Food Chem* 59:7726–7733
- Webster D (1974) A study of the interaction of bromocresol green with isolated serum globulin fractions. *Clin Chim Acta* 53:109–115
- Wei XJ, Hu TJ, Chen JR, Wei YY (2011) Inhibitory effect of carboxymethylpachymaran on cyclophosphamide-induced oxidative stress in mice. *Int J Biol Macromol* 49:801–805
- Wenger C, Kaplan A, Rubaltelli FF, Hammerman C (1984) *Alkaline phosphatase*. Clin Chem. The C.V. Mosby Co. St Louis, Toronto. Princeton. 1094–1098
- Yousefipour Z, Ranganna K, Newaz MA, Milton SG (2005) Mechanism of acrolein-induced vascular toxicity. *J Physiol Pharmacol* 56:337–353
- Zhang F, Lu Y, Zheng S (2012) Peroxisome proliferator-activated receptor- γ cross-regulation of signaling events implicated in liver fibrogenesis. *Cell Signal* 24(3):596–605