

Supplementary Material for “A balance of winners and losers in the Anthropocene”

Maria Dornelas, Nicholas J. Gotelli, Hideyasu Shimadzu, Faye Moyes, Anne E. Magurran, Brian McGill

**Data sources**

Metadata for the data sources is available at Figshare: [10.6084/m9.figshare.7687778](https://doi.org/10.6084/m9.figshare.7687778)

	Reference
18	(Zachmann <i>et al.</i> 2010)
33	(Widdicombe <i>et al.</i> 2010a; Widdicombe <i>et al.</i> 2010b)
39	(Holmes & Sturges 1975; Holmes <i>et al.</i> 1986; Holmes & Sherry 1988, 2001)
42	(Beven 1976; Williamson 1987; Gibbons <i>et al.</i> 1993; Standley <i>et al.</i> 1996; Stone <i>et al.</i> 1997; Gaston & Blackburn 2000; Lack 2010)
44	(Halpern & Dyrness 2010; Halpern & Lutz 2013a, b)
46	(Williamson 1983)
47	(Vickery & Nudds 1984)
51	(NERC ; Lindén & Rajala 1981)
52	(NERC ; Pulliainen 1981)
54	(Willig 2016)
56	(Friggens 2008)
57	(LTER)
58	(Waide 2010b, a)
59	(Ernest <i>et al.</i> 2009)
63	(Moore 1991b; Moore 1991a)
67	(CWAC)
68	(Management)
70	(Vanholder 1997)
78	(Zettler 2005)
81	(Robinson <i>et al.</i> 2007; Robinson 2010)
84	(Milchakova 2011)
85	(Addinck & de Kluijver 2003)
92	(Naumov)
97	(Markhaseva <i>et al.</i> 1985)
100	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
101	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
108	(Woehler)
110	(Ostler)
112	(NIWA)
113	(Pugh)
117	(NODC 2011)
119	(Clark & Branton 2007)
121	(CRED 2011)
125	(Reichert 2009)
147	(SWP)
152	(CMarZ)
162	(EPA)
163	(NPRB)
166	(PIROP ; Brown 1975; Diamond <i>et al.</i> 1986; Huettmann 1998; Halpin <i>et al.</i> 2009; Read <i>et al.</i> 2009)

169	(Yen <i>et al.</i> 2004; Jahncke 2006; Rintoul <i>et al.</i> 2006; Yen <i>et al.</i> 2006)
171	(BMMROOS)
172	(POPA ; Machete & Santos 2007; Amorim <i>et al.</i> 2008; Morato <i>et al.</i> 2008)
173	(JAMSTEC)
176	(Kennedy & Spry 2011)
178	(Boutillier 2007)
180	(ECNA)
182	(Wade 2011)
183	(Tremblay & Branton 2007)
187	(NCCOS 2003)
191	(NEFSC 2010)
192	(AADC)
193	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
195	(USGS)
196	(Moore & Howson)
197	(DATRAS 2013b)
198	(DATRAS 2013a)
200	(NEFSC 2010)
204	(Degraer <i>et al.</i> 2006)
206	(DATRAS 2010d)
208	(DATRAS 2010b)
209	(DATRAS 2010a)
210	(DATRAS 2010c)
213	(NMFS 2005)
214	(Harmon 2012)
215	(HMANA)
217	(USFS)
219	(NatureCounts 2012b)
220	(NatureCounts 2012b)
221	(Viereck <i>et al.</i> 2005)
225	(Shochat <i>et al.</i> 2004)
229	(McLarney <i>et al.</i> 2010)
232	(Williams)
236	(Gido)
237	(Lathrop 2000)
238	(LTER 2011)
240	(Muldavin)
243	(Day <i>et al.</i> 2004; Day 2010)
244	(NatureCounts 2012a)
246	(Chen <i>et al.</i> 2015)
247	(Rudstam 2008)
248	(Shi <i>et al.</i> 2015)
249	(Thomsen <i>et al.</i> 2016)
252	(Reichert 2010)
253	(LTER)
254	(LTER)
256	(DATRAS 2010a)
271	(Reed 2014a)
272	(Reed 2014b)
273	(Reed 2014c)
278	(NIWA)
285	(REVIZEE)

288	(Clark & Branton 2007)
290	(CMAR)
291	(CMAR)
292	(CMAR)
297	(Carpenter 2015)
300	(Landis & Gage 2014)
301	(Jonas & Joern 2007; Joern 2016)
305	(Wiley)
306	(Brook <i>et al.</i> 2006; NERC 2010)
308	(Merritt 1999)
310	(Sal <i>et al.</i> 2013)
311	(Kaufman)
313	(Knops & Tilman)
316	(Lightfoot)
318	(SANParks 2011)
319	(Wilgers <i>et al.</i> 2006)
321	(Lightfoot 2007)
327	(Kelt <i>et al.</i> 2013)
329	(Scott <i>et al.</i> )
330	(Davies <i>et al.</i> 2014)
332	(Grossman 2007)
333	(Sandercock)
335	(Grossman <i>et al.</i> 1982)
336	(Ernest <i>et al.</i> 2009)
337	(Vermont Center for Ecostudies <i>et al.</i> 2015)
339	(Svensson <i>et al.</i> 2010)
340	(Lightfoot 2011)
350	(Billett <i>et al.</i> 2001; Billett <i>et al.</i> 2010)
351	(Ruhl & Smith 2004; Kuhn <i>et al.</i> 2014)
354	(Davies <i>et al.</i> 2016)
356	(Bradford <i>et al.</i> 2014)
357	(Stapp 2013)
358	(Dickson <i>et al.</i> 1993)
359	(Reed 2014)
361	(Hall 1984)
363	(Enemar <i>et al.</i> 2004)
366	(Lightfoot 2013)
374	(Monitoring Site 1000 Project <i>et al.</i> 2013)
375	(Monitoring Site 1000 Project <i>et al.</i> 2014)
378	(Benedetti-Cecchi)
379	(Benedetti-Cecchi)
382	(Krefting & Ahlgren 1974; NERC 2010)
413	(Kendeigh 1982)
414	(Kendeigh 1982)
416	(Kendeigh 1982)
418	(Sweatman <i>et al.</i> 2001)
419	(Fraser 2014)
420	(Svensson 2006)
427	(Pomati <i>et al.</i> 2012; Pomati <i>et al.</i> 2015)
428	(Stenseth <i>et al.</i> 1999; Olsen <i>et al.</i> 2009; Rogers <i>et al.</i> 2011; Barceló <i>et al.</i> 2016)
430	(NIWA)
431	(NIWA)

432	(NIWA)
433	(Budy <i>et al.</i> ; Budy <i>et al.</i> )
439	(Zakharov 1998)
440	(Zakharov 1998)
441	(Zakharov 1998)
466	(Neat & Campbell 2011)
468	(Kenner <i>et al.</i> 2013)
469	(Kushner <i>et al.</i> 2013)
473	(Anderson <i>et al.</i> 2011)
475	(Hogstad 1993)
477	(Douglass <i>et al.</i> 2010; Lefcheck 2015)
478	(Wagner <i>et al.</i> 2011)
499	(Institute of Agricultural and Fisheries research (ILVO) 2016a)
500	(Institute of Agricultural and Fisheries research (ILVO) 2016b)
505	(Edelist <i>et al.</i> 2013)

## References

1.

AADC. "Whale Catches in Southern Ocean". OBIS - Australian Antarctic Data Centre. Available at: <http://www.iobis.org>, accessed 2013.

2.

Addinck, W. & de Kluijver, M. (2003). North Sea observations of Crustacea, Polychaeta, Echinodermata, Mollusca and some other groups between 1986 and 2003. Expert Centre for Taxonomic Identification (ETI), the Netherlands. Available at: <http://www.emodnet-biology.eu/data-catalog?module=dataset&david=1037>, accessed 2012.

3.

Amorim, P., Figueiredo, M., Machete, M., Morato, T., Martins, A. & Serrão Santos, R. (2008). Spatial variability of seabird distribution associated with environmental factors: a case study of marine Important Bird Areas in the Azores. *ICES Journal of Marine Science*, 66, 29-40.

4.

Anderson, J., Vermeire, L. & Adler, P.B. (2011). Fourteen years of mapped, permanent quadrats in a northern mixed prairie, USA. *Ecology*, 92, 1703-1703.

5.

Barceló, C., Ciannelli, L., Olsen, E.M., Johannessen, T. & Knutsen, H. (2016). Eight decades of sampling reveal a contemporary novel fish assemblage in coastal nursery habitats. *Global Change Biology*, 22, 1155-1167.

6.

Benedetti-Cecchi, L. "Calafuria Mid-shore Intertidal Dataset (1991-2014)". Department of Biology, University of Pisa. Accessed 2016.

7.

- Beven, G. (1976). Changes in breeding bird populations of an oak-wood on Bookham Common, Surrey, over twenty-seven years. *London Naturalist*, 55, 23-42.
- 8.
- Billett, D., Bett, B., Reid, W., Boorman, B. & Priede, I. (2010). Long-term change in the abyssal NE Atlantic: The 'Amperima Event' revisited. *Deep Sea Research Part II: Topical Studies in Oceanography*, 57, 1406-1417.
- 9.
- Billett, D., Bett, B., Rice, A., Thurston, M., Galéron, J., Sibuet, M. *et al.* (2001). Long-term change in the megabenthos of the Porcupine Abyssal Plain (NE Atlantic). *Progress in Oceanography*, 50, 325-348.
- 10.
- BMMROOS. "Bahamas Marine Mammal Research Organisation Opportunistic Sightings - OBIS SEAMAP". Available at: <http://www.iobis.org>, accessed 2012.
- 11.
- Boutillier, J.A. (2007). "Pacific Shrimp Trawl Survey". In: ShrimpTrawl Bio database, Fisheries and Oceans Canada PBS Shellfish Data Unit. OBIS Canada Digital Collections, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada. Available at: <http://www.iobis.org>, accessed 2012.
- 12.
- Bradford, M.G., Murphy, H.T., Ford, A.J., Hogan, D.L. & Metcalfe, D.J. (2014). Long-term stem inventory data from tropical rain forest plots in Australia. *Ecology*, 95, 2362-2362.
- 13.
- Brook, B.W., Traill, L.W. & Bradshaw, C.J. (2006). Minimum viable population sizes and global extinction risk are unrelated. *Ecology letters*, 9, 375-382.
- 14.
- Brown, R.G. (1975). Atlas of eastern Canadian seabirds.
- 15.
- Budy, P., Luecke, C. & O'Brien, J. "Zooplankton density for lake samples collected near Toolik Lake Arctic LTER in the summer from 1983 to 1992". Arctic Long-Term Ecological Research. Available at: <http://arc-lter.ecosystems.mbl.edu/1983-1992arclterzoops>, accessed 2016.
- 16.
- Budy, P., Luecke, C. & O'Brien, J. "Zooplankton density for lake samples collected near Toolik Lake Arctic LTER in the summers between 1993-2002". Arctic Long-Term Ecological Research. Available at: <http://arc-lter.ecosystems.mbl.edu/1993-2002arcticlterzoops>, accessed 2016.
- 17.
- Carpenter, R. (2015). "MCR LTER: Coral Reef: Long-term Population and Community Dynamics: Other Benthic Invertebrates, ongoing since 2005". Moorea Coral Reef LTER, knb-lter-mcr.7.28. Available at: doi:10.6073/pasta/8e7b3a0c7a8bf315739921861cc79d10, accessed 2016.
- 18.
- Chen, H., Liao, Y.-C., Chen, C.-Y., Tsai, J.-I., Chen, L.-S. & Shao, K.-T. (2015). Long-term monitoring dataset of fish assemblages impinged at nuclear power plants in northern Taiwan. *Scientific data*, 2, 150071.
- 19.

- Clark, D. & Branton, B. (2007). DFO Maritimes Research Vessel Trawl Surveys, OBIS Canada Digital Collections. *Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada, OBIS Canada*.
- 20.
- CMAR. "CSIRO Marine Data Warehouse - OBIS Australia", CSIRO Division of Marine and Atmospheric Research (CMAR), Australia. Available at <http://www.iobis.org>, accessed 2012.
- 21.
- CMarZ. "CMarZ (Census of Marine Zooplankton)-Asia Database". Accessed through OBIS- SCAR-MarBIN., Available at: <http://www.iobis.org/mapper/?dataset=1500>, accessed 2012.
- 22.
- CRED (2011). "Towed-Diver Fish Biomass Surveys in the Pacific Ocean 2000-2010". Coral Reef Ecosystem Division (CRED), Pacific Island Fisheries Sciences Center, National Marine Fisheries Service. Available at: <http://www.iobis.org/mapper/?dataset=1581>, accessed 2012.
- 23.
- CWAC. "Coordinated Waterbird Counts (CWAC) - AfrOBIS". . Available at: <http://www.iobis.org/mapper/?dataset=603>, accessed 2012.
- 24.
- DATRAS (2010a). "Fish trawl survey: ICES Beam Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2761>, accessed 2013.
- 25.
- DATRAS (2010b). "Fish trawl survey: ICES French Southern Atlantic Bottom Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2759>, accessed 2013.
- 26.
- DATRAS (2010c). "Fish trawl survey: ICES North Sea International Bottom Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2763>, accessed 2013.
- 27.
- DATRAS (2010d). "Fish trawl survey: Northern Irish Ground Fish Trawl Survey. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2764>, accessed 2013.
- 28.
- DATRAS (2013a). ICES Baltic International Trawl Survey For Commercial Fish Species (1991 - 2013). Available at <https://datras.ices.dk>, accessed 2013.
- 29.
- DATRAS (2013b). ICES Scottish West Coast Survey For Commercial Fish Species 1985-2013. Available at <https://datras.ices.dk>, accessed 2014.
- 30.

- Davies, C.H., Armstrong, A.J., Baird, M., Coman, F., Edgar, S., Gaughan, D. *et al.* (2014). Over 75 years of zooplankton data from Australia. *Ecology*, 95, 3229-3229.
- 31.
- Davies, C.H., Coughlan, A., Hallegraeff, G., Ajani, P., Armbrrecht, L., Atkins, N. *et al.* (2016). A database of marine phytoplankton abundance, biomass and species composition in Australian waters. *Scientific data*, 3.
- 32.
- Day, F. (2010). "Long-term N-fertilized vegetation plots on Hog Island, Virginia Coastal Barrier Islands, 1992-2014." Virginia Coast Reserve Long-Term Ecological Research Project. Available at: <http://www.vcrlter.virginia.edu/cgi-bin/showDataset.cgi?docid=knb-lter-vcr.106>, accessed 2013.
- 33.
- Day, F.P., Conn, C., Crawford, E. & Stevenson, M. (2004). Long-term effects of nitrogen fertilization on plant community structure on a coastal barrier island dune chronosequence. *Journal of Coastal Research*, 722-730.
- 34.
- Degraer, S., Wittoeck, J., Appeltans, W., Cooreman, K., Deprez, T., Hillewaert, H. *et al.* (2006). "Macrobelt: Long term trends in the macrobenthos of the Belgian Continental Shelf." Oostende, Belgium. Available at: <http://www.emodnet-biology.eu/data-catalog?module=dataset&dasid=145>, accessed 2013.
- 35.
- Diamond, A., Gaston, A. & Brown, R. (1986). Converting PIROP Counts of Seabirds at Sea to Absolute Densities. Progress Notes No 164. *Canadian Wildlife Service, Ottawa*.
- 36.
- Dickson, J.G., Conner, R.N. & Williamson, J.H. (1993). Neotropical migratory bird communities in a developing pine plantation. *Proceedings on the Annual Conference. SEAFWA*, 47, 439-446.
- 37.
- Douglass, J.G., France, K.E., Richardson, J.P. & Duffy, J.E. (2010). Seasonal and interannual change in a Chesapeake Bay eelgrass community: Insights into biotic and abiotic control of community structure. *Limnology and Oceanography*, 55, 1499-1520.
- 38.
- ECNA. "East Coast North America Strategic Assessment Project, Groundfish Atlas for the East Coast of North America". Available at: <http://www.iobis.org>, accessed 2012.
- 39.
- Edelist, D., Rilov, G., Golani, D., Carlton, J.T. & Spanier, E. (2013). Restructuring the Sea: profound shifts in the world's most invaded marine ecosystem. *Diversity and Distributions*, 19, 69-77.
- 40.
- Enemar, A., Sjöstrand, B., Andersson, G. & von Proschwitz, T. (2004). The 37-year dynamics of a subalpine passerine bird community, with special emphasis on the influence of environmental temperature and Epirrita autumnata cycles. *Ornis Svecica*, 14, 63-106.
- 41.
- EPA. "EPA'S EMAP Database". U.S. Environmental Protection Agency through its Environmental Monitoring and Assessment Program (EMAP). Available at: <http://www.iobis.org>, accessed

2012. Also available at <https://archive.epa.gov/emap/archive-emap/web/html/index-2021.html>

42.

Ernest, S., Valone, T.J. & Brown, J.H. (2009). Long-term monitoring and experimental manipulation of a Chihuahuan Desert ecosystem near Portal, Arizona, USA. *Ecology*, 90, 1708-1708.

43.

Fraser, W. (2014). "At-sea seabird censuses. Data on the species encountered (including marine mammals), their abundance, distribution and behavior. Data collected aboard cruises off the coast of the Western Antarctic Peninsula, 1993 - present". Palmer Station Antarctica LTER. Available at: <http://dx.doi.org/10.6073/pasta/e3871e749fa737dd94d5a269ac90e8ce>, accessed 2016.

44.

Friggens, M. (2008). "Sevilleta LTER Small Mammal Population Data", Albuquerque, NM: Sevilleta Long Term Ecological Research Site Database: SEV008. Available at: <http://sev.lternet.edu/data/sev-8>, accessed 2012.

45.

Gaston, K.J. & Blackburn, T.M. (2000). Pattern and process in macroecology. Wiley-Blackwell Oxford, England.

46.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993). *The new atlas of breeding birds in Britain and Ireland: 1988-1991*. T & AD Poyser London.

47.

Gido, K.B. "Fish population on selected watersheds at Konza Prairie - CFP01." Konza Prairie LTER Program. Available at: <http://www.konza.ksu.edu/KNZ/pages/data/Knzdsdetail.aspx?datasetCode=CFP01>, accessed 2012.

48.

Grossman, G.D. (2007). "Stream fish assemblage stability in a southern Appalachian stream at the Coweeta Hydrologic Laboratory from 1984 to 1995". Coweeta Long Term Ecological Research Program. Available at: <http://dx.doi.org/10.6073/pasta/f0baf5f59c89f670e04f537f5cc05290>, accessed 2016.

49.

Grossman, G.D., Moyle, P.B. & Whitaker Jr, J.O. (1982). Stochasticity in structural and functional characteristics of an Indiana stream fish assemblage: a test of community theory. *American Naturalist*, 423-454.

50.

Hall, G.A. (1984). A long-term bird population study in an Appalachian spruce forest. *The Wilson Bulletin*, 228-240.

51.

Halpern, C.B. & Dyrness, C. (2010). "Plant succession and biomass dynamics following logging and burning in the Andrews Experimental Forest Watersheds 1 and 3, 1962-Present". Long-Term Ecological Research. Forest Science Data Bank, Corvallis. Available at: <http://andrewsforest.oregonstate.edu/data/abstract.cfm?dbcode=TP073>, accessed 2012.

52.



- Halpern, C.B. & Lutz, J.A. (2013a). Canopy closure exerts weak controls on understory dynamics: a 30-year study of overstory–understory interactions. *Ecological Monographs*, 83, 221-237.  
53.
- Halpern, C.B. & Lutz, J.A. (2013b). "Canopy closure exerts weak controls on understory dynamics: a 30-year study of overstory–understory interactions". Available at: Dryad DigitalRepository, doi:10.5061/dryad.1q88j, accessed 2013.  
54.
- Halpin, P.N., Read, A.J., Fujioka, E., Best, B.D., Donnelly, B., Hazen, L.J. *et al.* (2009). OBIS-SEAMAP: The world data center for marine mammal, sea bird, and sea turtle distributions. *Oceanography*, 22, 104-115.  
55.
- Harmon, M.F., J. (2012). "Long-term growth, mortality and regeneration of trees in permanent vegetation plots in the Pacific Northwest, 1910 to present." Long-Term Ecological Research. Forest Science Data Bank, Corvallis. Available at: <http://andrewsforest.oregonstate.edu/data/abstract.cfm?dbcode=TV010>, accessed 2012.  
56.
- Henderson, P.A. The long-term study of the fish and crustacean community of the Bristol Channel. Available at <http://www.pisces-conservation.com/>, accessed 2013.  
57.
- Henderson, P.A. & Magurran, A.E. (2014a). Data from: Direct evidence that density-dependent regulation underpins the temporal stability of abundant species in a diverse animal community. Dryad Data Repository.  
58.
- Henderson, P.A. & Magurran, A.E. (2014b). Direct evidence that density-dependent regulation underpins the temporal stability of abundant species in a diverse animal community. *Proceedings of the Royal Society B: Biological Sciences*, 281.  
59.
- HMANA. "Hawk Migration Association of North America (HMANA)". Available at: <http://www.hmana.org/>, accessed 2012.  
60.
- Hogstad, O. (1993). Structure and dynamics of a passerine bird community in a spruce-dominated boreal forest. A 12-year study. In: *Annales Zoologici Fennici*. JSTOR, pp. 43-54.  
61.
- Holmes, R. & Sturges, F. (1975). Bird community dynamics and energetics in a northern hardwoods ecosystem. *The Journal of Animal Ecology*, 175-200.  
62.
- Holmes, R.T. & Sherry, T.W. (1988). Assessing population trends of New Hampshire forest birds: local vs. regional patterns. *The Auk*, 756-768.  
63.
- Holmes, R.T. & Sherry, T.W. (2001). Thirty-year bird population trends in an unfragmented temperate deciduous forest: importance of habitat change. *The Auk*, 118, 589-609.  
64.

- Holmes, R.T., Sherry, T.W. & Sturges, F.W. (1986). Bird Community Dynamics in a Temperate Deciduous Forest: Long-Term Trends at Hubbard Brook. *Ecological Monographs*, 56, 201-220.
- 65.
- Huettmann, F. (1998). An ecological GIS research application for the northern Atlantic-The PIROP database software, environmental data sets and the role of the internet. *Riekert/Tochtermann*.
- 66.
- Institute of Agricultural and Fisheries research (ILVO), B. (2016a). Macrobenthos monitoring at long-term monitoring stations in the Belgian part of the North Sea between 1979 and 1999. Available at: <http://dx.doi.org/10.14284/201>, accessed 2016.
- 67.
- Institute of Agricultural and Fisheries research (ILVO), B. (2016b). Macrobenthos monitoring at long-term monitoring stations in the Belgian part of the North Sea from 2001 on. Available at: <http://dx.doi.org/10.14284/202>, accessed 2016.
- 68.
- Jahncke, J.R., C. (2006). "CalCOFI and NMFS Seabird and Marine Mammal Observation Data, 1987-2006". California Cooperative Oceanic Fisheries Investigations (CalCOFI) and National Marine Fisheries Service (NMFS) cruises, 1987-2006 - OBIS SEAMAP. Available at: <http://www.iobis.org>, accessed 2012.
- 69.
- JAMSTEC. "Marine Biological Sample Database, JAMSTEC" (OBIS\_JAPAN). Available at: <http://www.iobis.org/mapper/?dataset=2289>, accessed 2012.
- 70.
- Joern, A. (2016). CGR02 Sweep Sampling of Grasshoppers on Konza Prairie LTER watersheds (1982-present). Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/7060b2c244229a37e3bfc8c18f14ad02>, accessed 2016.
- 71.
- Jonas, J.L. & Joern, A. (2007). Grasshopper (Orthoptera: Acrididae) communities respond to fire, bison grazing and weather in North American tallgrass prairie: a long-term study. *Oecologia*, 153, 699-711.
- 72.
- Kaufman, D.W. Seasonal summary of numbers of small mammals on 14 LTER traplines in prairie habitats at Konza Prairie. Konza Prairie Long-Term Ecological Research. . Available at: <http://lter.konza.ksu.edu/content/csm01-seasonal-summary-numbers-small-mammals-14-lter-traplines-prairie-habitats-konza>, accessed 2016.
- 73.
- Kelt, D., Meserve, P., Gutiérrez, J., Milstead, W.B. & Previtali, M. (2013). Long-term monitoring of mammals in the face of biotic and abiotic influences at a semiarid site in north-central Chile. *Ecology*, 94, 977-977.
- 74.
- Kendeigh, S.C. (1982). Bird populations in east central Illinois: Fluctuations, variations, and development over a half-century. *University of Illinois Press*.
- 75.

- Kennedy, M. & Spry, J. (2011). Atlantic Zone Monitoring Program Maritimes Region plankton datasets. *Fisheries and Oceans Canada-BioChem archive. OBIS Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada.*
- 76.
- Kenner, M.C., Estes, J.A., Tinker, M.T., Bodkin, J.L., Cowen, R.K., Harrold, C. *et al.* (2013). A multi-decade time series of kelp forest community structure at San Nicolas Island, California (USA). *Ecology*, 94, 2654-2654.
- 77.
- Knops, J. & Tilman, D. Successional Dynamics on a Resampled Chronosequence - Experiment 014. Cedar Creek Ecosystem Science Reserve. . Available at <http://www.cedarcreek.umn.edu/research/data/dataset?ghe014>, accessed 2016.
- 78.
- Krefting, L.W. & Ahlgren, C.E. (1974). Small Mammals and Vegetation Changes After Fire in a Mixed Conifer-Hardwood Forest. *Ecology*, 55, 1391-1398.
- 79.
- Kuhnz, L.A., Ruhl, H.A., Huffard, C.L. & Smith, K.L. (2014). Rapid changes and long-term cycles in the benthic megafaunal community observed over 24years in the abyssal northeast Pacific. *Progress in Oceanography*, 124, 1-11.
- 80.
- Kushner, D.J., Rassweiler, A., McLaughlin, J.P. & Lafferty, K.D. (2013). A multi-decade time series of kelp forest community structure at the California Channel Islands. *Ecology*, 94, 2655-2655.
- 81.
- Lack, P. (2010). *The atlas of wintering birds in Britain and Ireland.* A&C Black.
- 82.
- Landis, D. & Gage, S. (2014). Insect Populations via Sticky Traps at KBS-LTER. Available at: <http://lter.kbs.msu.edu/datatables/67>, accessed 2016.
- 83.
- Lathrop, R. (2000). "Madison Wisconsin Lakes Zooplankton 1976 - 1994." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <http://lter.limnology.wisc.edu/dataset/madison-wisconsin-lakes-zooplankton-1976-1994>, accessed 2013.
- 84.
- Lefcheck, J.S. (2015). The use of functional traits to elucidate the causes and consequences of biological diversity. The College of William & Mary, PhD thesis. Available at: <http://gradworks.umi.com/36/62/3662989.html>, accessed 2016.
- 85.
- Lightfoot, D. "Small Mammal Exclosure Study (SMES)". Sevilleta Long Term Ecological Research Program. Available at: <http://sev.lternet.edu/content/small-mammal-exclosure-study-smes-0>, accessed 2016.
- 86.
- Lightfoot, D. (2007). "Jornada Grasshopper Data". Jornada Basin LTER. Available at: <http://jornada.nmsu.edu/lter/dataset/49712/view>, accessed 2016.
- 87.

- Lightfoot, D. (2011). "Small Mammal Exclosure Study (SMES) Vegetation Data from the Chihuahuan Desert Grassland and Shrubland at the Sevilleta National Wildlife Refuge, New Mexico (2006-2009)". Long Term Ecological Research Network. Available at: <http://dx.doi.org/10.6073/pasta/d80d5e2196cd11ef79df23ebe5a77c19>, accessed 2016.
- 88.
- Lightfoot, D. (2013). "Lizard pitfall trap data (LTER-II, LTER-III)". Jornada Basin LTER. Available at: <http://jornada.nmsu.edu/lter/dataset/49821/view>, accessed 2016.
- 89.
- Lindén, H. & Rajala, P. (1981). Fluctuations and long-term trends in the relative densities of tetraonid populations in Finland, 1964-77. *Finnish Game Research*, 39, 13-34.
- 90.
- LTER, N. "North Temperate Lakes LTER: Phytoplankton - Madison Lakes Area 1995 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-phytoplankton-madison-lakes-area-1995-current>, accessed 2013.
- 91.
- LTER, N. "North Temperate Lakes LTER: Zooplankton - Trout Lake Area 1982 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-zooplankton-trout-lake-area-1982-current>, accessed 2013.
- 92.
- LTER, N. "NTLFI02 North Temperate Lakes LTER: Fish Abundance 1981 - current". North Temperate Lakes Long Term Ecological Research program, NSF. Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-fish-abundance-1981-current>, accessed 2012.
- 93.
- LTER, N. (2011). "North Temperate Lakes LTER: Zooplankton - Madison Lakes Area 1997 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <http://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-zooplankton-madison-lakes-area-1997-current>, accessed 2013.
- 94.
- Machete, M. & Santos, R. (2007). Azores Fisheries Observer Program (POPA): a case study of the multidisciplinary use of observer data. In: *Proceedings of the 5th International Fisheries Observer Conference*, pp. 15-18.
- 95.
- Management, M.a.C. "Marine and Coastal Management - Copepod Surveys - AfrOBIS". Available at: <http://www.iobis.org/>, accessed 2012.
- 96.
- Markhaseva, E.L., Golikov, A.A., Agapova, T.A. & Beig, A.A. (1985). Archives of the Arctic Seas Zooplankton., Available at: <http://www.iobis.org/mapper/?dataset=4470>, accessed 2012.
- 97.

- McLarney, W.O., Meador, J. & Chamblee, J. (2010). "Upper Little Tennessee River Biomonitoring Program Database." Coweeta Long Term Ecological Research Program. Available at: [https://coweeta.uga.edu/dbpublic/dataset\\_details.asp?accession=4045](https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=4045), accessed 2012.
- 98.
- Merritt, J. (1999). Long Term Mammal Data from Powdermill Biological Station 1979-1999. Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/83c888854e239a79597999895bb61cfe>, accessed 2016.
- 99.
- Milchakova, N.A., Ryabogina, V. G. & Chernyshova, E. B. (2011). "Macroalgae of the Crimean coastal zone (the Black Sea, 1967-2007)". Sevastopol, IBSS. Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&david=2690>, accessed 2012.
- 100.
- Monitoring Site 1000 Project, Biodiversity Center & Japan, M.o.E.o. (2013). "Monitoring site 1000 Shorebird Survey" (ShorebirdsDatapackage2012.zip, downloaded from <http://www.biodic.go.jp/moni1000/findings/data/index.html>), Accessed 2016.
- 101.
- Monitoring Site 1000 Project, Biodiversity Center & Japan, M.o.E.o. (2014). "Monitoring site 1000 Forest and grassland research - Surface wandering beetles survey data". (GBDataPackage2014ver1.zip, downloaded from <http://www.biodic.go.jp/moni1000/findings/data/index.html>), Accessed 2016.
- 102.
- Moore, J.J. & Howson, C.M. "Survey of the rocky shores in the region of Sullom Voe, Shetland, A report to SOTEAG from Aquatic Survey & Monitoring Ltd", Cosheton, Pembrokeshire. 29 p. Available at: <http://www.soteag.org.uk>, accessed 2013.
- 103.
- Moore, N.W. (1991a). "The development of dragonfly communities and the consequences of territorial behaviour: A 27-year study on small ponds at Woodwalton Fen, Cambridgeshire, United Kingdom". NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database Version 2.0. Available at: <http://www3.imperial.ac.uk/cpb/databases/gpdd>, accessed 2012.
- 104.
- Moore, N.W. (1991b). The development of dragonfly communities and the consequences of territorial behaviour: a 27 year study on small ponds at Woodwalton Fen, Cambridgeshire, United Kingdom. *Odonatologica*, 20, 203-231.
- 105.
- Morato, T., Varkey, D.A., Damaso, C., Machete, M., Santos, M., Prieto, R. *et al.* (2008). Evidence of a seamount effect on aggregating visitors. *Marine Ecology Progress Series*, 357, 23-32.
- 106.
- Muldavin, E. "Pinon-Juniper (Core Site) Quadrat Data for the Net Primary Production Study at the Sevilleta National Wildlife Refuge, New Mexico (2003-Present)." Sevilleta Long Term Ecological Research Program., Available at: <http://sev.lternet.edu/node/1718>, accessed 2013.
- 107.

- NatureCounts (2012a). Bird Studies Canada "BC Coastal Waterbird Survey (2004)." NatureCounts, a node of the Avian Knowledge Network. Available at: <http://www.birdscanada.org/birdmon/>, accessed 2012.
- 108.
- NatureCounts (2012b). Bird Studies Canada "Marsh Monitoring Program." NatureCounts, a node of the Avian Knowledge Network. Available at: <http://www.birdscanada.org/birdmon/>, accessed 2012.
- 109.
- Naumov, A. "Benthos of the White Sea. A database". White Sea Biological Station, Zoological Institute RAS. Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&david=2769>, accessed 2012.
- 110.
- NCCOS (2003). "National Benthic Infaunal Database (NBID)". NOAA/NOS/NCCOS/CCEHBR/Coastal Ecology Program. NOAA's Ocean Service, National Centers for Coastal Ocean Science (NCCOS). Available at: <https://data.noaa.gov/dataset/national-benthic-infaunal-database-nbid>, accessed 2012.
- 111.
- Neat, F. & Campbell, N. (2011). Demersal fish diversity of the isolated Rockall plateau compared with the adjacent west coast shelf of Scotland. *Biological Journal of the Linnean Society*, 104, 138-147.
- 112.
- NEFSC (2010). "Benthic Database (OBIS-USA)", Northeast Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Department of Commerce., Available at: <http://www.iobis.org/mapper/?dataset=1694>, accessed 2012.
- 113.
- NERC. "Fluctuations and long-term trends in the relative densities of tetraonid populations in Finland, 1964-77." NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database v2.0. . Available at: <https://www.imperial.ac.uk/cpb/gpdd2/secure/register.aspx>, accessed 2012.
- 114.
- NERC. "A transect survey of small land carnivore and red fox populations on a subarctic fell in Finnish Forest Lapland over 13 winters". NERC Centre for Population Biology, Imperial College, The Global Population Dynamics Database v2.0., Available at: <http://www3.imperial.ac.uk/cpb/databases/gpdd>, accessed 2012.
- 115.
- NERC (2010). "The Global Population Dynamics Database Version 2". Centre for Population Biology, Imperial College. Available at: <http://www.sw.ic.ac.uk/cpb/cpb/gpdd.html>, accessed 2016.
- 116.
- NIWA. "The New Zealand Freshwater Fish Database". Available at: <https://www.niwa.co.nz/our-services/online-services/freshwater-fish-database>, accessed 2016.
- 117.
- NIWA. "South Western Pacific Regional OBIS Data Asteroid Subset", NIWA (National Institute of Water and Atmospheric Research - New Zealand) MBIS (Marine Biodata Information System)

- accessed through South Western Pacific OBIS., Available at:  
<http://www.iobis.org/mapper/?dataset=219>, accessed 2012.
- 118.
- NIWA. "South Western Pacific Regional OBIS Data provider for the NIWA Marine Biodata Information System". Ocean Biogeographic Information System. Occurrence Dataset., Available at:  
<https://doi.org/10.15468/zuuiyu>, accessed 2012.
- 119.
- NMFS (2005). "Northeast Fisheries Science Center Bottom Trawl Survey Data (OBIS-USA)." NOAA's National Marine Fisheries Service (NMFS) Northeast Fisheries Science Center. Woods Hole, Massachusetts, USA., Available at: <http://www.iobis.org/mapper/?dataset=1435>, accessed 2013.
- 120.
- NODC (2011). "South TX Outer Continental Shelf and MI, AL, and FL Outer Continental Shelf benthic organism sampling 1974-1978". US National Oceanographic Data Center, Silver Spring, Maryland, USA. Available at [http://www.usgs.gov/obis-usa/data\\_search\\_and\\_access/participants.html](http://www.usgs.gov/obis-usa/data_search_and_access/participants.html), accessed 2012.
- 121.
- NPRB. "The Observer Program database", accessed through the OBIS-USA North Pacific Groundfish Observer (North Pacific Research Board). Available at: <http://www.iobis.org>, accessed 2012.
- 122.
- Olsen, E.M., Carlson, S.M., Gjøsæter, J. & Stenseth, N.C. (2009). Nine decades of decreasing phenotypic variability in Atlantic cod. *Ecology Letters*, 12, 622-631.
- 123.
- Ostler, R. Marine Nature Conservation Review (MNCR) and associated benthic marine data held and managed by JNCC-EurOBIS. *Joint Nature Conservation Committee, Centre for Ecology and hydrology, Aberdeenshire, UK.* (Available at <http://www.emodnet-biology.eu/datacatalog>).
- 124.
- PIROP. "PIROP Northwest Atlantic 1965-1992 - OBIS SEAMAP". Available at:  
<http://www.iobis.org/mapper/?dataset=2245>, accessed 2012.
- 125.
- Pomati, F., Matthews, B., Jokela, J., Schildknecht, A. & Ibelings, B.W. (2012). Effects of re-oligotrophication and climate warming on plankton richness and community stability in a deep mesotrophic lake. *Oikos*, 121, 1317-1327.
- 126.
- Pomati, F., Tellenbach, C., Matthews, B., Venail, P., Ibelings, B.W. & Ptacnik, R. (2015). Challenges and prospects for interpreting long-term phytoplankton diversity changes in Lake Zurich (Switzerland). *Freshwater Biology*, 60, 1052-1059.
- 127.
- POPA. "cetacean, seabird, and sea turtle sightings in the Azores area 1998-2009 - OBIS SEAMAP". . Available at: <http://www.iobis.org/mapper/?dataset=4257>, accessed 2012.
- 128.

- Pugh, P. Discovery Collections Midwater Database. National Oceanography Centre, Southampton, UK., Available at: <http://www.iobis.org/mapper/?dataset=12>, accessed 2012.
- 129.
- Pulliainen, E. (1981). A transect survey of small land carnivore and red fox populations on a subarctic fell in Finnish Forest Lapland over 13 winters. *Annales Zoologici Fennici*, 270-278.
- 130.
- Read, A., Halpin, P., Crowder, L., Best, B. & Fujioka, E. (2009). OBIS-SEAMAP: mapping marine mammals, birds and turtles. *World Wide Web electronic publication*. <http://seamap.duke.edu>. Accessed, 15.
- 131.
- Reed, D.C. (2014). "SBC LTER: Reef: Kelp Forest Community Dynamics: Fish abundance". Santa Barbara Coastal LTER. Available at: doi:10.6073/pasta/e37ed29111b2fddffc08355252b8b8c7, accessed 2016.
- 132.
- Reed, D.C. (2014a). "SBC LTER: Reef: Kelp forest community dynamics: Abundance and size of giant kelp (*Macrocystis pyrifera*), ongoing since 2000". Santa Barbara Coastal LTER. Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.18> doi:10.6073/pasta/d90872297e30026b263a119d4f5bca9f, accessed 2016.
- 133.
- Reed, D.C. (2014b). "SBC LTER: Reef: Kelp forest community dynamics: Fish abundance". Santa Barbara Coastal LTER. Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.17> doi:10.6073/pasta/e37ed29111b2fddffc08355252b8b8c7, accessed 2016.
- 134.
- Reed, D.C. (2014c). "SBC LTER: Reef: Kelp forest community dynamics: Invertebrate and algal density". Santa Barbara Coastal LTER. . Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.19> doi:10.6073/pasta/cd4cf864efecd69891dfe1d73b9ac9c3, accessed 2016.
- 135.
- Reichert, M. (2009). "MARMAP Chevron Trap Survey 1990-2009". SCDNR/NOAA MARMAP Program, SCDNR MARMAP Aggregate Data Surveys, The Marine Resources Monitoring, Assessment, and Prediction (MARMAP) Program, Marine Resources Research Institute, South Carolina Department of Natural Resources U.S.A., Available at: [http://www.usgs.gov/obis-usa/data\\_search\\_and\\_access/participants.html](http://www.usgs.gov/obis-usa/data_search_and_access/participants.html), accessed 2012.
- 136.
- Reichert, M. (2010). "MARMAP Blackfish Trap Survey 1990-2009". SCDNR/NOAA MARMAP Program. SCDNR MARMAP Aggregate Data Surveys. The Marine Resources Monitoring, Assessment, and Prediction (MARMAP) Program. Marine Resources Research Institute. South Carolina Department of Natural Resources USA. Available at: <http://www.usgs.gov/obis-usa/>, accessed 2013.
- 137.
- REVIZEE. "Previous\_fisheries\_REVIZEE\_Program", accessed through Tropical and Subtropical Western South Pacific OBIS., Available at: <http://www.iobis.org/mapper/?dataset=2411>, accessed 2012.
- 138.



- Rintoul, C., Langabeer-Schlagenhauf, B., Hyrenbach, K., Morgan, K. & Sydeman, W. (2006). Atlas of California Current marine birds and mammals: Version 1. *Unpublished Report, PRBO Conservation Science, Petaluma, CA.*  
139.
- Robinson, K.P. (2010). "CRRU (Cetacean Research and Rescue Unit) Cetacean sightings in Scotland waters". Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&david=2819>, accessed 2012.  
140.
- Robinson, K.P., Baumgartner, N., Eisfeld, S.M., Clark, N.M., Culloch, R.M., Haskins, G.N. *et al.* (2007). The summer distribution and occurrence of cetaceans in the coastal waters of the outer southern Moray Firth in northeast Scotland (UK). *Lutra*, 50, 19.  
141.
- Rogers, L.A., Stige, L.C., Olsen, E.M., Knutsen, H., Chan, K.-S. & Stenseth, N.C. (2011). Climate and population density drive changes in cod body size throughout a century on the Norwegian coast. *Proceedings of the National Academy of Sciences*, 108, 1961-1966.  
142.
- Rudstam, L. (2008). "Zooplankton survey of Oneida Lake, New York, 1964 – 2012", KNB Data Repository. Available at: <https://knb.ecoinformatics.org/#view/kgordon.17.56>, accessed 2016.  
143.
- Ruhl, H.A. & Smith, K.L. (2004). Shifts in deep-sea community structure linked to climate and food supply. *Science*, 305, 513-515.  
144.
- Sal, S., López-Urrutia, Á., Irigoien, X., Harbour, D.S. & Harris, R.P. (2013). Marine microplankton diversity database. *Ecology*, 94, 1658-1658.  
145.
- SandercocK, B.K. "Variable distance line-transect sampling of bird population numbers in different habitats on Konza Prairie (1981 - 2009)". Konza Prairie Long Term Ecological Research Program. Available at: <http://www.konza.ksu.edu/knz/pages/data/KnzEntity.aspx?id=CBP011>, accessed 2016.  
146.
- SANParks (2011). "Karoo National Park Census Data. 1994 - 2009". Available at: <http://datadryad.org/handle/10255/dryad.13079?show=full>, accessed 2016.  
147.
- Scott, D., Metts, B. & Lance, S. "The Rainbow Bay Long-term Study". Available at: <http://srelherp.uga.edu/projects/rbay.htm>, accessed 2016.  
148.
- Shi, Z., Sherry, R., Xu, X., Hararuk, O., Souza, L., Jiang, L. *et al.* (2015). Evidence for long-term shift in plant community composition under decadal experimental warming. *Journal of Ecology*, 103, 1131-1140.  
149.
- Shochat, E., Katti, M. & Warren, P. (2004). "Point count bird censusing: long-term monitoring of bird distribution and diversity in central Arizona-Phoenix: period 2000 to 2011". Central Arizona-

- Phoenix Long-Term Ecological Research. Global Institute for Sustainability, Arizona State University. Available at: <https://caplter.asu.edu/data/data-catalog/?id=46>, accessed 2012.
- 150.
- Standley, P., Bucknell, N., Swash, A. & Collins, I. (1996). *The Birds of Berkshire*. Berkshire Atlas Group. Reading, UK.
- 151.
- Stapp, P. (2013). SGS-LTER Long-Term Monitoring Project: Small Mammals on Trapping Webs on the Central Plains Experimental Range, Nunn, Colorado, USA 1994 -2006, ARS Study Number 118. Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/2e311b4e40fea38e573890f473807ba9>, accessed 2017.
- 152.
- Stenseth, N.C., Bjørnstad, O.N., Falck, W., Fromentin, J.M., Gjøsæter, J. & Gray, J.S. (1999). Dynamics of coastal cod populations: intra- and intercohort density dependence and stochastic processes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266, 1645-1654.
- 153.
- Stone, B., Sears, J., Cranswick, P., Gregory, R., Gibbons, D., Rehfisch, M. *et al.* (1997). Population estimates of birds in Britain and in the United Kingdom. *British Birds*, 90, 1-22.
- 154.
- Svensson, S. (2006). Species composition and population fluctuations of alpine bird communities during 38 years in the Scandinavian mountain range. *Ornis Svecica*, 16, 183-210.
- 155.
- Svensson, S., Thorner, A. & Nyholm, N. (2010). Species trends, turnover and composition of a woodland bird community in southern Sweden during a period of fifty-seven years. *Ornis Svecica*, 20, 31-44.
- 156.
- Sweatman, H.P.A., Cheal, A.J., Coleman, G.J., Delean, S., Fitzpatrick, B.M., Miller, I.R. *et al.* (2001). Long-Term Monitoring of the Great Barrier Reef. Status Report Number 5. Available at: <http://www.iobis.org/mapper/?dataset=233>, accessed 2017.
- 157.
- SWP. "South Western Pacific Regional OBIS Data Bryozoan Subset". Accessed through South Western Pacific OBIS., Available at: <http://www.iobis.org/mapper/?dataset=221>, accessed 2012.
- 158.
- Thomsen, P.F., Jørgensen, P.S., Bruun, H.H., Pedersen, J., Riis-Nielsen, T., Jonko, K. *et al.* (2016). Resource specialists lead local insect community turnover associated with temperature – analysis of an 18-year full-seasonal record of moths and beetles. *Journal of Animal Ecology*, 85, 251-261.
- 159.
- Tremblay, J.M. & Branton, B. (2007). DFO Maritimes Research Vessel Trawl Surveys, OBIS Canada Digital Collections. *Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada, OBIS Canada*.
- 160.

- USFS. "Landbird Monitoring Program (UMT-LBMP)." US Forest Service. Available at: <http://www.avianknowledge.net/>, accessed 2012.
- 161.
- USGS. Patuxent Wildlife Research Center "North American Breeding Bird Survey" ftp data set, version 2014.0. Available at: <ftp://ftpext.usgs.gov/pub/er/md/laurel/BBS/DataFiles/>, accessed 2013.
- 162.
- Vanholder, B. (1997). "Belgian Migrating Lepidoptera". NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database v2.0. . Available at: <https://www.imperial.ac.uk/cpb/gpdd2/secure/register.aspx>, accessed 2012.
- 163.
- Vermont Center for Ecostudies, Lambert, J.D. & Hart, J. (2015). "Mountain Birdwatch 1.0". KNB Data Repository, 10.5063/F1DN430G. accessed 2016.
- 164.
- Vickery, W.L. & Nudds, T.D. (1984). Detection of Density-Dependent Effects in Annual Duck Censuses. *Ecology*, 65, 96-104.
- 165.
- Viereck, L.A., Van Cleve, K., Chapin, F.S., Ruess, R.W. & Hollingsworth, T.N. (2005). Vegetation Plots of the Bonanza Creek LTER Control Plots: Species Count (1975 - 2004). Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/8dd0e1ac48e2f82b51adabfbd3c62ae2>, accessed 2012.
- 166.
- Wade, E. (2011). Snow crab research trawl survey database (Southern Gulf of St. Lawrence, Gulf region, Canada) from 1988 to 2010. *OBIS Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada*.
- 167.
- Wagner, R., Marxsen, J., Zwick, P. & Cox, E.J. (2011). *Central European Stream Ecosystems: The Long Term Study of the Breitenbach*. John Wiley & Sons.
- 168.
- Waide, R.B. (2010a). "Bird abundance - point counts. El Verde Field Station, Puerto Rico: Luquillo Long Term Ecological Research Site Database: Data Set 23". Available at: <http://luq.lter.net.edu/data/luqmetadata23>, accessed 2012.
- 169.
- Waide, R.B. (2010b). Bird abundance - point counts. Long Term Ecological Research Network. Available at: <http://dx.doi.org/10.6073/pasta/0d96957379936a038ebbbcc6135b2fab>, accessed 2012.
- 170.
- Widdicombe, C., Eloire, D., Harbour, D., Harris, R. & Somerfield, P. (2010a). Long-term phytoplankton community dynamics in the Western English Channel. *Journal of Plankton Research*, fbp127.
- 171.
- Widdicombe, C.E., Harbour, D. & Harris, R. (2010b). Time series of phytoplankton abundance and composition at station L4 in the English Channel from 1988 to 2009. *PANGAEA*, 754335, 0-0.
- 172.

- Wiley, R.H. "Population estimates of Appalachian salamanders". Coweeta LTER. Available at: <http://coweeta.uga.edu/eml/1044.xml>, accessed 2016.
- 173.
- Wilgers, D.J., Horne, E.A., Sandercock, B.K. & Volkmann, A.W. (2006). Effects of rangeland management on community dynamics of the herpetofauna of the tallgrass prairie. *Herpetologica*, 62, 378-388.
- 174.
- Williams, D. "Pelagic Fish Observations 1968-1999." Australian Antarctic Data Centre. Available at: <http://www.gbif.org/dataset/85b0a82a-f762-11e1-a439-00145eb45e9a>, accessed 2012.
- 175.
- Williamson, M. (1983). The land-bird community of Skokholm: ordination and turnover. *Oikos*, 378-384.
- 176.
- Williamson, M. (1987). Are communities ever stable? In: *Symposium of the British Ecological Society*. 177.
- Willig, M.R.B., C. P. (2016). "El Verde Grid long-term invertebrate data: Luquillo Long Term Ecological Research Site Database: Data Set 107". Available at: <http://luq.lternet.edu/data/luqmetadata107/7427> accessed 2016.
- 178.
- Woehler, E. "Seabirds of the Southern and South Indian Ocean - Australian Antarctic Data Centre". Available at: <http://www.iobis.org>, accessed 2012.
- 179.
- Yen, P., Sydeman, W., Bograd, S. & Hyrenbach, K. (2006). Spring-time distributions of migratory marine birds in the southern California Current: Oceanic eddy associations and coastal habitat hotspots over 17 years. *Deep Sea Research Part II: Topical Studies in Oceanography*, 53, 399-418.
- 180.
- Yen, P.P., Sydeman, W.J. & Hyrenbach, K.D. (2004). Marine bird and cetacean associations with bathymetric habitats and shallow-water topographies: implications for trophic transfer and conservation. *Journal of Marine Systems*, 50, 79-99.
- 181.
- Zachmann, L., Moffet, C. & Adler, P. (2010). Mapped quadrats in sagebrush steppe: long-term data for analyzing demographic rates and plant-plant interactions. *Ecology*, 91, 3427-3427.
- 182.
- Zakharov, V.D. (1998). Biodiversity of bird population of terrestrial habitats in Southern Ural. Miass: IGZ. *Ural Branch of Russian Academy of Sciences*, 158 p.
- 183.
- Zettler, M.L. (2005). Macrozoobenthos baltic sea (1980-2005) as part of the IOW-Monitoring. Institut für Ostseeforschung Warnemünde, Germany., Available at: <http://www.iobis.org/mapper/?dataset=2289>, accessed 2012.