

TR/94

February 1980

A Class of C^2 Piecewise Quintic Polynomials

By

N. PAPAMICHAEL & A. J. WORSEY

W9260300

A B S T R A C T

A new class of C^2 piecewise quintic interpolatory polynomials is defined. It is shown that this new class contains a number of interpolatory functions which present practical advantages, when compared with the conventional cubic spline.

1. Introduction

Given the points

$$a = x_0 < x_1 < \dots < x_k = b, \tag{1.1}$$

and the corresponding values $y_i = y(x_i); i = 0, 1, \dots, k$, let $H_3(x)$ be the piecewise cubic Hermite polynomial, with knots (1.1), which is such that

$$H_3(x_i) = y_i \text{ and } H_3^{(1)}(x_i) = y_i^{(1)}; i = 0, 1, \dots, k.$$

Denote by s the piecewise polynomial obtained from H_3 by replacing the derivatives $y_i^{(1)}; i = 0, 1, \dots, k$, respectively by suitable approximations $m_i; i = 0, 1, \dots, k$. Let p_i be the cubic polynomial interpolating the function y at the points $x_i, x_{i+1}, x_{i+2}, x_{i+3}$, define the quadratic polynomials $q_i; i = 0, 1, \dots, k-2$, by

$$\left. \begin{aligned} q_i &= p_i^{(1)}; i = 0, 1, \dots, k-3, \\ q_{k-2} &= q_{k-3} = p_{k-3}^{(1)}, \end{aligned} \right\} \tag{1.2}$$

and let the approximations m_i satisfy the relations

$$\left. \begin{aligned} m_0 &= y_0^{(1)} \\ \alpha_i m_{i-1} + m_i + \beta_i m_{i+1} &= \alpha_i q_{i-1}(x_{i-1}) + q_{i-1}(x_i) + \beta_i q_{i-1}(x_{i-1}); \\ & \qquad \qquad \qquad i = 1, 2, \dots, k-1, \\ m_k &= y_k^{(1)} \end{aligned} \right\} \tag{1.3}$$

where the α_i and β_i are real numbers. Then, by the definition of Behforooz, Papamichael and Worsey [3], s is a cubic x -spline with parameters $\alpha_i, \beta_i; i = 1, 2, \dots, k-1$. This definition of x -splines is a generalization of an earlier definition due to Clenshaw and Negus [5], and contains the conventional cubic spline s_1 as the special case

$$2\alpha_i = h_{i+1} / (h_i + h_{i+1}), \quad 2\beta_i = 1 - 2\alpha_i; \quad i = 1, 2, \dots, k-1,$$

where $h_i = x_i - x_{i-1}$.

Clearly, a cubic X-spline s is continuous and possesses a continuous first derivative. In general, however, $s^{(2)}$ has a jump discontinuity at each interior knot, and s_I is the only cubic X-spline with C^2 continuity on (a,b) .

Regarding the quality of approximation, it is shown in [3] that for any cubic X-spline s

$$\|s - y\| = O(h^4) \tag{1.4}$$

where $\|\cdot\|$ denotes the uniform norm on $[a,b]$ and $h = \max_i h_i$. Since

(1.4) gives the best order of uniform convergence that can be obtained by an interpolatory piecewise cubic polynomial, it follows that no cubic X-spline can achieve substantially higher accuracy than s_I . However, as it is shown in [33], there are cubic X-splines which produce results of comparable accuracy to those obtained by s_I , with much less computational effort.

In the present paper we generalize the results of [3] to the case of piecewise quintic polynomial interpolation. For this we consider the piecewise quintic Hermite polynomial H_5 with knots (1.1) and, by analogy with the definition of cubic X-splines, we define a quintic X-spline as a C^2 interpolant derived from H_5 by replacing the derivatives $y_i^{(1)}$ and $y_i^{(2)}$; $i = 0, 1, \dots, k$, by approximations which are determined by solving a certain pair of tri-diagonal linear systems. The motivation for this generalization emerges from considering the

problem of constructing C^2 interpolants which lead to $O(h^n)$, $n \geq 5$, convergence and whose construction does not involve excessive computational effort, by comparison with the construction of the conventional cubic spline s_I . The requirements concerning the order of convergence and computational labour are imposed so that the new interpolants may compete, in terms of computational efficiency, with s_I . We show that the class of quintic X-splines, defined in this paper, contains several interpolatory functions which satisfy the above requirements.

2. Interpolatory Piecewise Quintic Polynomials

Given the set of values $y_i = y(x_i)$; $i = 0, 1, \dots, k$, where x_i are the points (1.1), let H_5 be the piecewise quintic Hermite polynomial which is such that

$$H_5(x_i) = y_i, H_5^{(1)}(x_i) = y_i^{(1)} \quad \text{and} \quad H_5^{(2)}(x_i) = y_i^{(2)}; \quad i = 0, 1, \dots, k.$$

Then if $y \in C^6[a, b]$, the following optimal error bound holds

$$\|H_5 - y\| \leq \frac{1}{46,080} h^6 \|y^{(6)}\|, \quad (2.1)$$

where, as before, $\|\cdot\|$ denotes the uniform norm on $[a, b]$,

$h_i = x_i - x_{i-1}$; $i = 1, 2, \dots, k$, and $h = \max_{1 \leq i \leq k} h_i$; see e.g. Birkhoff and Priver [4].

Definition 1. Let Q be the piecewise quintic polynomial derived from H_5 by replacing the derivatives $y_i^{(1)}$ and $y_i^{(2)}$; $i = 0, 1, \dots, k$, respectively by suitable approximations m_i and M_i ; $i = 0, 1, \dots, k$. Then Q will be called a piecewise quintic polynomial (p.q.p.) with derivatives

m_i and M_i ; $i = 0, 1, \dots, k$.

It follows at once from the definition that Q can be written as

$$\begin{aligned}
 Q(x) = & s(x) + Q[x_{i-1}, x_{i-1}, x_{i-1}, x_i, x_i] (x-x_{i-1})^2 (x-x_i)^2 \\
 & + Q[x_{i-1}, x_{i-1}, x_{i-1}, x_i, x_i, x_i] (x-x_{i-1})^3 (x-x_i)^2, \\
 & x \in [x_{i-1}, x_i]; \quad i=1, 2, \dots, k,
 \end{aligned} \tag{2.2}$$

where s is the piecewise cubic polynomial satisfying $s(x_i) = y_i$, $s^{(1)}(x_i) = m_i$; $i = 0, 1, \dots, k$, and, with the usual notation for divided differences ,

$$\left. \begin{aligned}
 Q[x_{i-1}, x_{i-1}, x_i, x_i] &= \frac{1}{2h_i^4} [-6(y_i - y_{i-1}) + 2h_i(m_i - 2m_{i-1}) + h_i^2 m_{i-1}] \\
 \text{and} \\
 Q[x_{i-1}, x_{i-1}, x_{i-1}, x_i, x_i, x_i] &= \frac{1}{2h_i^5} [12(y_i - y_{i-1}) - 6h_i(m_i + m_{i-1}) + h_i^2(M_i - M_{i-1})]; \\
 & i = 1, 2, \dots, k.
 \end{aligned} \right\} \tag{2.3}$$

The following theorem is a trivial generalization of a result due to Hall [6]. It can be established easily by using (2.1) and the cardinal representations of H_5 and Q .

Theorem 1. Let Q be a p. q.p. with derivatives m_i and M_i ; $i = 0, 1, \dots, k$. If $y \in C^6[a, b]$ then, for $x \in [x_{i-1}, x_i]$; $i = 1, 2, \dots, k$,

$$\begin{aligned}
 |Q(x) - y(x)| \leq & \frac{1}{46,080} h^6 \|y^{(6)}\| + \frac{5}{16} h \max\left\{ \left| m_{i-1} - y_{i-1}^{(1)} \right|, \left| m_i - y_i^{(1)} \right| \right\} \\
 & + \frac{1}{32} h^2 \max\left\{ \left| M_{i-1} - y_{i-1}^{(2)} \right|, \left| M_i - y_i^{(2)} \right| \right\}. \tag{2.4}
 \end{aligned}$$

The theorem shows that the best order of approximation that can be achieved by an interpolatory p.q.p. Q is

$$\|Q - y\| = O(h^6).$$

More specifically, the theorem shows that if the approximations m_i and M_i are such that

$$\text{and } \left. \begin{array}{l} m_i - y_i^{(1)} = O(h^r) \\ M_i - y_i^{(2)} = O(h^s) \end{array} \right\} \quad i = 0, 1, \dots, k, \quad (2.5)$$

then,

$$\text{where } \left. \begin{array}{l} \|Q - y\| = O(h^n), \\ n = \min(r + 1, s + 2, 6). \end{array} \right\} \quad (2.6)$$

Since one of our requirements is that the interpolants Q satisfy (2.6) with $n \geq 5$ it follows that, for the purposes of the present paper, the approximations m_i and M_i must satisfy (2.5) with $r \geq 4$ and $s \geq 3$ respectively.

Clearly a p.q.p. Q is continuous and possesses continuous first and second derivatives. In general, however, $s^{(3)}$ has a jump discontinuity $d_i^{(3)}$ at each interior knot x_i . Using (2.2) and (2.3) it can be shown that

$$\begin{aligned} d_i^{(3)} &= Q^{(3)}(x_{i+}) - Q^{(3)}(x_{i-}) \\ &= \frac{3}{h_{i+1}^3 h_i^3} \{ 20 [h_i^3 y_{i+1} - (h_{i+1}^3 h_i^3) y_i + h_{i+1}^3 y_{i-1}] \\ &\quad - 2h_{i+1} h_i [4h_i^2 m_{i+1} + 6(h_i^2 - h_{i+1}^2) m_i - 4h_{i+1}^2 m_{i-1}] \\ &\quad + h_{i+1}^2 h_i^2 [h_i M_{i+1} - 3(h_i + h_{i+1}) M_i + h_{i+1} M_{i-1}] \}; \\ &\quad i = 1, 2, \dots, k-1. \end{aligned} \quad (2.7)$$

Hence, if $y \in C^7[a, b]$,

$$\begin{aligned} d_i^{(3)} &= \frac{3}{h_{i+1}^2 h_i^2} \{ h_{i+1} h_i [h_i (M_{i+1} - y_{i+1}^{(2)}) - 3(h_i + h_{i+1})(M_i - y_i^{(2)}) + h_{i+1} (M_{i-1} - y_{i-1}^{(2)})] \\ &\quad - 2[4h_i^2 (m_{i+1} - y_{i+1}^{(1)}) + 6(h_i^2 - h_{i+1}^2) m_i - y_i^{(1)}] - 4h_{i+1}^2 (m_{i-1} - y_{i-1}^{(1)}) \} \\ &\quad + \frac{1}{5!} (h_i^3 + h_{i+1}^3) y_i^{(6)} + O(h^4); \quad i = 1, 2, \dots, k-1. \end{aligned} \quad (2.8)$$

Equation (2.8) follows from (2.7) by using the result

$$20[h_i^3 y_{i+1} - (h_i^3 + h_{i+1}^3) y_i + h_{i+1}^3 y_{i-1}] - 2h_{i+1} h_i [4h_i^2 y_{i+1}^{(1)} + 6(h_i^2 - h_{i-1}^2) y_i^{(1)} - 4h_{i+1}^2 y_{i-1}^{(1)}] + h_{i+1}^2 h_i^2 [h_i y_{i+1}^{(2)} - 3(h_i h_{i+1}) y_i^{(2)} + h_{i+1} y_{i-1}^{(2)}] = \frac{2}{6!} (h_i^3 + h_{i+1}^3) y_i^{(6)} + O(h^{10});$$

$$i = 1, 2, \dots, k-1,$$

which is established by Taylor series expansions about the point x_i . Thus the magnitude of $d_i^{(3)}$, like the order of convergence of Q , depends only on the quality of the approximations m_i and M_i . More specifically, if the derivatives m_i and M_i of Q satisfy (2.5) then

$$\left. \begin{array}{l} \text{where} \\ d_i^{(3)} = O(h^n), \\ n = \min\{r-2, s-1, 3\}. \end{array} \right\} \quad (2.9)$$

3. Quintic X-splines

Let $q_i; i = 1, 2, \dots, k-2$, be the quadratic polynomials defined by (1.2). Then, by analogy with the definition of cubic X-splines of Behforooz et al [3], we define the class of quintic X-splines as follows.

Definition 2. Let α_i, β_i and $\gamma_i, \delta_i; i = 1, 2, \dots, k-1$, be $4k-4$ real numbers. Then, a p.q.p. Q whose derivatives m_i and $M_i; i = 0, 1, \dots, k$ satisfy respectively the relations

$$\left. \begin{array}{l} m_0 = y_0^{(1)}, \\ \alpha_i m_{i-1} + m_i + \beta_i m_{i+1} = \alpha_i q_{i-1}(x_{i-1}) + q_{i-1}(x_i) + \beta_i q_{i-1}(x_{i+1}); \\ m_k = y_k^{(1)}, \end{array} \right\} \quad \begin{array}{l} \\ \\ i = 1, 2, \dots, k-1, \end{array} \quad (3.1)$$

and

$$\left. \begin{aligned} M_0 &= y_0^{(2)}, \\ \gamma_i M_{i-1} + M_i + \delta_i M_{i+1} &= \gamma_i q_{i-1}^{(1)}(x_{i-1}) + q_{i-1}^{(1)}(x_i) + \delta_i q_{i-1}^{(1)}(x_{i-1}); \\ M_k &= y_k^{(2)}, \end{aligned} \right\} \quad i = 1, 2, \dots, k-1, \quad (3.2)$$

will be called a quintic X-spline with parameters α_i, β_i and γ_i, δ_i ; $i = 1, 2, \dots, k-1$.

By Definition 2 the derivatives m_i and M_i ; $i = 1, 2, \dots, k-1$, of a quintic X-spline Q are determined by solving the two $(k-1) \times (k-1)$ tri-diagonal linear systems defined by (3.1) and (3.2). Thus, a sufficient condition for the unique existence of Q is that its parameters α_i, β_i and γ_i, δ_i satisfy respectively the inequalities

$$|\alpha_i| + |\beta_i| < 1; \quad i = 1, 2, \dots, k-1, \quad (3.3)$$

and

$$|\gamma_i| + |\delta_i| < 1; \quad i = 1, 2, \dots, k-1. \quad (3.4)$$

It follows at once from the definition that, in the representation (2.2) of a quintic X-spline Q , s is a cubic X-spline with parameters α_i, β_i , $i = 1, 2, \dots, k-1$. The convergence properties of such an s are discussed fully in [3]. In particular, it is shown that if $y \in C^6[a, b]$ and (3.3) holds then the derivatives m_i of s satisfy

$$m_i - y_i^{(1)} = O(h^r); \quad i = 1, 2, \dots, k-1,$$

where in general $r = 3$. However, there are several choices of α_i, β_i , for which $r = 4$ and one choice for which $r = 5$. Keeping in mind our requirements concerning the order of $\|Q - y\|$ and the amount of labour involved in computing Q , we conclude from [3: Section 4] that there

are two choices of α_i, β_i , which are of particular interest. These are the values,

$$\left. \begin{aligned} \alpha_i^{(1)} &= \frac{h_{i+1}(h_{i+1} + h_{i+2})}{(h_i + h_{i+1})(h_i + h_{i+1} + h_{i+2})}, \quad \beta_i^{(1)} = 0, i = 1, 2, \dots, k-2, \\ \alpha_{k-1}^{(1)} &= 0, \beta_{k-1}^{(1)} = \frac{h_{k-1}(h_{k-1} + h_{k-2})}{(h_{k-1} + h_k)(h_{k-2} + h_{k-1} + h_k)} \end{aligned} \right\} \quad (3.5)$$

and

$$\left. \begin{aligned} \alpha_i^{(2)} &= \frac{h_{i+1}^2(h_{i+1} + h_{i+2})}{(h_i + h_{i+1} + h_{i+2})(h_i + h_{i+1})^2}, \quad \beta_i^{(2)} = \frac{h_i^2(h_{i+1} + h_{i+2})}{h_{i+2}(h_i + h_{i+1})^2}; \\ & \quad i = 1, 2, \dots, k-1, \end{aligned} \right\} \quad (3.6)$$

$$h_{k+1} = -(h_{k-2} + h_{k-1} + h_k).$$

The values $\alpha_i^{(1)}, \beta_i^{(1)}$ reduce the three-term recurrence relation in (3.1) to a two-term relation. Thus, in this case, the derivatives m_i are determined from (3.1) by forward substitution. It is shown in [3: Section 4.5] that these m_i satisfy

$$m_i - y_i^{(1)} = O(h^4); \quad i = 1, 2, \dots, k-1, \quad (3.7)$$

In particular, when the knots are equally spaced then

$$\left. \begin{aligned} \alpha_i^{(1)} &= 1/3, \quad \beta_i^{(1)} = 0; \quad i = 1, 2, \dots, k-2, \\ \alpha_{k-1}^{(1)} &= 0, \quad \beta_{k-1}^{(1)} = 1/3, \end{aligned} \right\}$$

and, if $y \in C^7[a, b]$,

$$\left| m_i - y_i^{(1)} \right| \leq \frac{1}{40} h^4 \|y^{(5)}\| + \frac{1}{240} h^5 \|y^{(6)}\| + O(h^6); \quad i = 1, 2, \dots, k-1. \quad (3.8)$$

The values (3.6) are the only choice of parameters α_i, β_i for which the derivatives m_i satisfy

$$m_i - y_i^{(1)} = 0(h^5); \quad i = 1, 2, \dots, k-1; \quad (3.9)$$

see [3: Section 4.6]. It should be observed that, in this case, the conditions (3*3) which ensure that the tri-diagonal linear system (3.1) has a unique solution are satisfied only if

$$(h_i + h_{i+1}) (h_i - h_{i+2}) < 2h_{i+2}(h_{i+1} + h_{i+2}); \\ i = 1, 2, \dots, k-1 .$$

When the knots are equally spaced then

$$\alpha_i^{(2)} = 1/6, \quad \beta_i^{(2)} = 1/2; \quad i = 1, 2, \dots, k-2$$

$$\alpha_{k-1}^{(2)} = 1/2, \quad \beta_{k-1}^{(2)} = 1/6,$$

and, if $y \in C^7[a, b]$,

$$\left| m_i - y_i^{(1)} \right| \leq \frac{1}{120} h^5 \|y^{(6)}\| + 0(h^6); \quad i = 1, 2, \dots, k-1 \quad (3.10)$$

We consider now the effect that the parameters γ_i, δ_i have on the quality of the second derivatives of a quintic X-spline Q. For this we assume that the parameters satisfy (3.4), let

$$\varepsilon_i = \gamma_i \{q_{i-1}^{(1)}(x_{i-1}) - y_{i-1}^{(2)}\} + \{q_{i-1}^{(1)}(x_i) - y_{i-1}^{(2)}\} + \delta_i \{q_{i-1}^{(1)}(x_{i-1}) - y_{i+1}^{(2)}\}; \\ i = 1, 2, \dots, k-1, \quad (3.11)$$

and denote by A the matrix of the $(k-1) \times (k-1)$ linear system defined by (3.2). Then, using a result of Lucas [7, p.5763,

$$\|A^{-1}\|_{\infty} \leq v, \quad (3.12)$$

where $v \geq 1$ is such that

$$\left| \gamma_i \right| + \left| \delta_i \right| + 1/v \leq 1; \quad i = 1, 2, \dots, k-1.$$

Hence, from (3.2),

$$\left| M_i - y_i^{(2)} \right| \leq v \max_i \left| \varepsilon_i \right|; \quad i = 1, 2, \dots, k-1 \quad (3.13)$$

Also, by Taylor series expansion about the point x_i we find that if $y \in C^6[a,b]$ then,

$$\varepsilon_i = \frac{1}{12} F_i y_i^{(4)} + \frac{1}{60} G_i y_i^{(5)} + O(h^4) \quad i = 1, 2, \dots, k-1, \quad (3.14)$$

where

$$\left. \begin{aligned} F_i &= \gamma_i \{-h_{i+1}(h_{i+1} + 4h_i + h_{i+2}) - h_i(2h_{i+2} + 3h_i)\} \\ &\quad + \{h_{i+1}(2h_i - h_{i+2} - h_{i+1}) + h_i h_{i+2}\} \\ &\quad + \delta_i \{h_{i+1}(2h_{i+2} - h_{i+1} - h_i) h_i h_{i+2}\}, \\ G_i &= \gamma_i \{-(h_{i+1} + 2h_i)(h_{i+1} + h_{i+2} - h_i)(2h_{i+1} + h_{i+2}) \\ &\quad + h_i(7h_i^2 - 3h_{i+1}^2)\} \\ &\quad - (h_{i+1} - h_i)(h_{i+1} + h_{i+2} - h_i)(2h_{i+1} + h_{i+2}) \\ &\quad + \delta_i \{(2h_{i+1} + h_i)(h_{i+1} + h_{i+2} - h_i)(2h_{i+1} + h_{i+2}) \\ &\quad + h_{i+1}(3h_i^2 - 7h_{i+1}^2)\}; \\ &\quad i = 1, 2, \dots, k-1, \end{aligned} \right\} \quad (3.15)$$

and, as in (3.6),

$$h_{k+1} = - (h_{k-2} + h_{k-1} + h_k) \quad (3.16)$$

When the knots are equally spaced then (3.14) simplifies considerably and, if $y \in C^7[a,b]$, it gives

$$\begin{aligned} \varepsilon_i = & \frac{1}{12} \{-\tilde{\gamma}_i + 1 + \tilde{\delta}_i\} h^2 y_i^{(4)} + \frac{1}{12} \{\delta_i - \gamma_i\} h^3 y_i^{(5)} \\ & + \frac{1}{360} \{16 \tilde{\delta}_i + 1 - 44 \tilde{\gamma}_i\} h^4 y_i^{(6)} + O(h^5); \end{aligned} \quad i = 1, 2, \dots, k-1, \quad (3.17)$$

where

$$\text{and } \left. \begin{aligned} \tilde{\gamma}_i = \gamma_i, \quad \tilde{\delta}_i = \delta_i; \quad i = 1, 2, \dots, k-2 \\ \tilde{\gamma}_{k-1} = \delta_{k-1}, \quad \tilde{\delta}_{k-1} = \gamma_{k-1}. \end{aligned} \right\} \quad (3.18)$$

The results (3.13) and (3.14)- (3.15) show that if the parameters $\gamma_i, \delta_i; i=1,2, \dots, k-1$ of a quintic X-spline satisfy (3.4) then

$$M_i - y_i^{(2)} = O(h^s); \quad i = 1, 2, \dots, k-1,$$

where, in general $s=2$. However, if the γ_i, δ_i , are such that $F_i = 0; i=1,2, \dots, k-1$ then $s=3$, and if $F_i = G_i = 0; i=1,2, \dots, k-1$ then $s=4$.

Corresponding to the two choices (3.5) and (3.6) of the parameters α_i, β_i , there are two choices of the γ_i, δ_i which are of particular interest. These are the values

$$\left. \begin{aligned} \gamma_i^{(1)} = \frac{h_{i+1}(2h_i - h_{i+1} - h_{i+2}) + h_i h_{i+2}}{h_{i+1}(4h_i + h_{i+1} + h_{i+2}) + h_i(3h_i + 2h_{i+2})}, \quad \delta_i^{(1)} = 0; \\ i = 1, 2, \dots, k-2, \\ \gamma_{k-1}^{(1)} = 0, \quad \delta_{k-1}^{(1)} = \frac{h_{k-1}(2h_k - h_{k-1} - h_{k-2}) + h_k h_{k-2}}{h_{k-1}(4h_k - h_{k-1} + h_{k-2}) + h_k(3h_k + 2h_{k-2})}, \end{aligned} \right\} \quad (3.19)$$

and

$$\gamma_i^{(2)} = \frac{h_{i+1}A_i}{D_i(h_i + h_{i+1})}, \quad \delta_i^{(2)} = \frac{h_i B_i}{D_i(h_i + h_{i+1})};$$

$$i = 1, 2, \dots, k-1, \quad (3.20)$$

where

$$\left. \begin{aligned} A_i &= (h_{i+1} + h_{i+2}) (h_{i+1} - h_i) \{-3h_{i+2} (h_i + h_{i+1}) + h_i h_{i+1}^2\} \\ &\quad + h_i h_{i+1} \{3(h_i + 1 + h_{i+2})^2 - h_{i+1}(3h_i + 4h_{i+1})\} . \\ B_i &= (h_{i+1} + h_{i+2}) (h_{i+1} - h_i) \{h_i + h_{i+1} (h_i + h_{i+1} + h_{i+2}) + h_i^2\} \\ &\quad + h_i h_{i+1} \{3(h_{i+1} + h_{i+2})^2 - h_i(4h_i + 3h_{i+1})\}, \\ D_i &= (h_{i+1} + h_{i+2})^2 \{(h_i + 2 - h_{i+1})(4h_i + h_{i+1}) + h_{i+2}(3h_{i+2} + h_{i+1})\} \\ &\quad + 3h_i h_{i+1} h_{i+2} (h_i + h_{i+1} + h_{i+2}), \end{aligned} \right\} \quad (3.21)$$

and h_{k+1} is given by (3.16).

The parameters $\gamma_i^{(1)}, \delta_i^{(1)}$ are such that, in (3.14), $F_i = 0, i = 1, 2, \dots, k-1$.

Therefore, in this case,

$$M_i - y_i^{(2)} = 0(h^3); \quad i = 1, 2, \dots, k-1. \quad (3.22)$$

Also, since the values (3.19) reduce the three-term recurrence relation in (3.2) to a two-term relation, the M_i 's are determined from (3.2) by forward substitution.

When the knots are equally spaced then

$$\gamma_i^{(1)} = 1/11, \quad \delta_i^{(1)} = 0; \quad i = 1, 2, \dots, k-2,$$

$$\gamma_{k-1}^{(1)} = 0, \quad \delta_{k-1}^{(1)} = 1/11,$$

and, since $V = 1/10$, (3.13) and (3.17) give

$$|M_i - y_i^{(2)}| \leq \frac{1}{120} h^3 \|y^{(5)}\| + \frac{11}{1200} h^4 \|y^{(6)}\| + O(h^5);$$

$$i = 1, 2, \dots, k-1. \quad (3.23)$$

The values $\gamma_i^{(2)}, \delta_i^{(2)}$ are the only values of γ_i, δ_i for which $F_i = G_i = 0; i = 1, 2, \dots, k-1$. This implies that (3.20) is the only choice of parameters γ_i, d_i , for which

$$M_i - y_i^{(2)} = O(h^4); i = 1, 2, \dots, k-1. \quad (3.24)$$

Clearly, $\gamma_i^{(2)}, \delta_i^{(2)}$ are defined only if, in (3.20), $D_i \neq 0; i = 1, 2, \dots, k-1$.

A sufficient condition for this to hold is that

$$h_{i+2} (3h_{i+2} + h_{i+1}) > (h_{i+1} - h_{i+2}) (4h_i + h_{i+1});$$

$$i = 1, 2, \dots, k-1. \quad (3.25)$$

It should be observed however that (3.25) does not imply the conditions (3.4) which ensure that the linear system (3.2) has a unique solution.

When the knots are equally spaced then

$$\gamma_i^{(2)} = \delta_i^{(2)} = 1/10; i = 1, 2, \dots, k-1,$$

and, since $v = 5/4$, (3.13) and (3.17) give

$$|M_i - y_i^{(2)}| \leq \frac{1}{160} h^4 \|y^{(6)}\| + O(h^5); i = 1, 2, \dots, k-1. \quad (3.26)$$

The remainder of this paper is concerned with examining the quality of the four quintic X-splines with parameters taken from the four

Possible combinations of the values $\alpha_i^{(r)}, \beta_i^{(r)}$ and $\gamma_i^{(s)} = \delta_i^{(s)}; r, s = 1, 2$.

4. Quintic X-splines of special interest

We let

$$E = \|Q-y\| \quad \text{and} \quad (4.1) \quad D^{(3)} = \max_i |d_i^{(3)}|, \quad (4.1)$$

Where, as in Section 2, $d_i^{(3)}$ denotes the jump discontinuity of $Q^{(3)}$ at an interior knot x_i . We also let $Q_{r,s}$ denote the quintic X-spline with parameters $\alpha_i^{(r)}, \beta_i^{(r)}$ and $\gamma_i^{(s)} = \delta_i^{(s)}$; $r, s = 1, 2$. Then, with this notation, the derivatives m_i and M_i of each of the four $Q_{r,s}$; $r, s = 1, 2$, are determined as follows:

- (i) The m_i of $Q_{1,1}$ and $Q_{1,2}$, by forward substitution, from the lower triangular system defined by (3.1) with $\alpha_i = \alpha_i^{(1)}$, $\beta_i = \beta_i^{(1)}$, where $\alpha_i^{(1)}, \beta_i^{(1)}$ are the values (3.5).
- (ii) The m_i of $Q_{2,1}$ and $Q_{2,2}$, by solving the tri-diagonal system defined by (3.1) with $\alpha_i = \alpha_i^{(2)}$, $\beta_i = \beta_i^{(2)}$, where $\alpha_i^{(2)}, \beta_i^{(2)}$ are the values (3.6).
- (iii) The M_i of $Q_{1,1}$ and $Q_{2,1}$, by forward substitution, from the lower triangular system defined by (3.2) with $\gamma_i = \gamma_i^{(1)}$, $\delta_i = \delta_i^{(1)}$, where $\gamma_i^{(1)}, \delta_i^{(1)}$ are the values (3.19).
- (iv) The M_i of $Q_{1,2}$ and $Q_{2,2}$, by solving the tri-diagonal system defined by (3.2) with $\gamma_i = \gamma_i^{(2)}$, $\delta_i = \delta_i^{(2)}$, where $\gamma_i^{(2)}, \delta_i^{(2)}$ are the values (3.20).

The results of the previous section in conjunction with (2.5) - (2.6) and (2.9) show that for each of the X-splines $Q_{1,1}$, $Q_{1,2}$ and $Q_{2,1}$,

$$E = O(h^5) \quad \text{and} \quad D^{(3)} = O(h^2). \quad (4.2)$$

These results also show that $Q_{2,2}$ is the only quintic X-spline for which

$$E = O(h^6) \quad \text{and} \quad D^{(3)} = O(h^3) . \quad (4.3)$$

we consider now the case of equally spaced knots and, for each of the four $Q_{r,s}$, we list bounds on E and on $D^{(3)}$. These bounds are derived easily from (2.4), (2.8) and (3.8), (3.10), (3.23) and (3.26).

(i) Quintic X-spline $Q_{1,1}$

$$E \leq \frac{31}{3,840} h^5 \|y^{(5)}\| + \frac{371}{230,400} h^6 \|y^{(6)}\| + O(h^7) , \quad (4.4)$$

$$D^{(3)} \leq \frac{7}{5} h^2 \|y^{(5)}\| + \frac{131}{300} h^3 \|y^{(6)}\| + O(h^4) . \quad (4.5)$$

(ii) Quintic X-spline $Q_{1,2}$

$$E \leq \frac{1}{128} h^5 \|y^{(5)}\| + \frac{350}{230,400} h^6 \|y^{(6)}\| + O(h^7) , \quad (4.6)$$

$$D^{(3)} \leq \frac{6}{5} h^2 \|y^{(5)}\| + \frac{110}{300} h^3 \|y^{(6)}\| + O(h^4) . \quad (4.7)$$

(iii) Quintic X-spline $Q_{2,1}$

$$E \leq \frac{1}{840} h^5 \|y^{(5)}\| + \frac{671}{230,400} h^6 \|y^{(6)}\| + O(h^7) , \quad (4.8)$$

$$D^{(3)} \leq \frac{1}{5} h^2 \|y^{(5)}\| + \frac{191}{300} h^3 \|y^{(6)}\| + O(h^4) . \quad (4.9)$$

(iv) Quintic X-spline $Q_{2,2}$

$$E \leq \frac{650}{230,400} h^6 \|y^{(6)}\| + O(h^7) , \quad (4.10)$$

$$D^{(3)} \leq \frac{170}{300} h^3 \|y^{(6)}\| + O(h^4) . \quad (4.11)$$

5. Numerical results and discussion

In Tables 1 and 2 we present numerical results obtained by taking $y(x) = \exp(x)$,

$$x_i = i/20 ; \quad i = 0,1, \dots, 20 , \tag{5.1}$$

and constructing each of the four quintic X-splines considered in section 4. The results listed are values of the absolute error $[Q(x) - y(x)]$, computed at various points between the knots, and the maximum values $D^{(3)}$ of the jump discontinuities in the third derivative at interior knots. The results of Tables 3 and 4 are obtained, in a similar manner, by using the same y and the unequally spaced knots

$$x_i = i^2/8^2 ; \quad i = 0,1, \dots, 8 . \tag{5.2}$$

For comparison purposes, we also include in the first column of each table the corresponding results obtained in [3] by using the conventional cubic spline s_1 ,

The theoretical results of the previous sections indicate that $Q_{2,2}$ is the most 'accurate' X-spline. These results also show that in approximating a smooth function y by a quintic X-spline, the quality of the first derivatives m_i is more critical than that of the second derivatives M_i . This follows from the observation that in (2.4), the magnitudes of the coefficients associated with the terms $(m_j - y_i^{(1)})$ are larger than those associated with $(M_j - y_i^{(2)})$. For this reason we expect $Q_{2,1}$ to produce more accurate approximations than $Q_{1,2}$.

The numerical results in Tables 1 and 3 show that the X-splines $Q_{2,2}$ and $Q_{2,1}$ produce the most accurate results. They also show that there

is no significant overall difference in accuracy between the approximations due to $Q_{2,2}$ and $Q_{2,1}$ and between those of $Q_{1,2}$ and $Q_{1,1}$. This is in accordance with the theory, since when $y(x) = \exp(x)$ and the equally spaced knots (5.1) are used, then the error bounds (4.4), (4.6), (4.8) and (4.10), with the $O(h^7)$ term ignored, give $E \leq n \times 10^{-9}$ where n takes the values 7.0, 6.8, .35 and .13 respectively for each of the X-splines $Q_{1,1}$, $Q_{1,2}$, $Q_{2,1}$ and $Q_{2,2}$. A similar argument would of course explain the results corresponding to the unequally spaced knots (5.2), for which $h = \max_i h_i = 15/64$. Naturally, as h decreases the difference in accuracy between the results due to $Q_{2,2}$ and $Q_{2,1}$ becomes more pronounced. However, $Q_{2,2}$ leads to a marked improvement in accuracy only if h is very small.

Of the four X-splines considered here the construction of $Q_{1,1}$ involves the least computational effort. The derivatives of this X-spline are determined by forward substitution from two lower triangular systems and this involves less computational effort than the determination of the parameters of the conventional cubic spline s_1 . Also, $Q_{1,1}$ is the only X-spline in Section 4 whose unique existence is guaranteed for any distribution of the knots. For this reason, we consider $Q_{1,1}$ to be of greater practical interest than the other three X-splines considered in Section 4.

By Definition 2, the construction of a quintic X-spline requires knowledge of $y^{(1)}$ and $y^{(2)}$ at the two endpoints x_0 , x_k and, in an interpolation problem, this information is not usually available. However, by using techniques similar to those of Behforooz and Papamichael [1 and 2], the end conditions

$$m_0 = y_0^{(1)}, m_k = y_k^{(1)} ; \tag{5.3}$$

$$M_0 = y_0^{(2)}, \quad M_k = y_k^{(2)}, \quad (5.4)$$

can be replaced by conditions which use only the available function values of y at the knots whilst retaining the order of the X-spline approximation. For example, if $\pi_i(x)$ is the quartic polynomial interpolating y at the points $x_i, x_{i+1}, x_{i+2}, x_{i+3}$ and x_{i+4} ; $j = 0, k-4$, then the following end conditions can be used, instead of (5.3), for the construction of $Q_{1,1}$ and $Q_{1,2}$

$$m_0 = \pi_0^{(1)}(x_0), \quad m_k = \pi_{k-4}^{(1)}(x_k). \quad (5.5)$$

Similarly, the end conditions

$$\left. \begin{aligned} m_0 + \alpha_0 m_1 &= \pi_0^{(1)}(x_0) + \alpha_0 \pi_0^{(1)}(x_1), \\ \alpha_k m_{k-1} + m_k &= \alpha_k \pi_{k-4}^{(1)}(x_{k-1}) + \pi_{k-4}^{(1)}(x_k), \end{aligned} \right\} \quad (5.6)$$

can be used for the construction of $Q_{2,1}$ and $Q_{2,2}$, where, in (5.6)

$$\alpha_j = (1 + u_j + v_j + w_j + u_j v_j + v_j w_j + w_j u_j + u_j v_j w_j) / u_j v_j w_j; \quad j = 0,$$

with

$$u_0 = h_2 / h_1, \quad v_0 = (h_1 u_0 + h_3) / h_1, \quad w_0 = (h_1 v_0 + h_4) / h_1,$$

and

$$u_k = h_{k-1}, \quad v_k = (h_k u_k + h_{k-2}) / h_k, \quad w_k = (h_k v_k + h_{k-3}) / h_k.$$

By analogy, the second derivative end conditions (5.4) can be replaced by

$$M_0 = \pi_0^{(2)}(x_0), \quad M_k = \pi_{k-4}^{(2)}(x_k), \quad (5.7)$$

for the construction of $Q_{1,1}$ and $Q_{2,1}$ and

$$\left. \begin{aligned} M_0 + \gamma_0 M_1 &= \pi_0^{(2)}(x_0) + \gamma_0 \pi_0^{(2)}(x_1), \\ \gamma_k M_{k-1} + M_k &= \gamma_k \pi_{k-4}^{(2)}(x_{k-1}) + \pi_{k-4}^{(2)}(x_k), \end{aligned} \right\} \quad (5.8)$$

for $Q_{1,2}$ for $Q_{2,2}$, where in (5.8)

$$\gamma_0 = \frac{[h_1(7h_1^2 - 3h_2^2) + (h_2 + 2h_1)(h_3 + h_2 - h_1)(h_4 + h_3 + h_2 - h_1) + h_1(h_4 + 2h_3 + 3h_2 - h_1)(5h_1 + 3h_2)]}{[(h_1 - h_2)(h_2 + h_3)(h_4 + h_3 + h_2) + h_1h_2(h_4 + 2h_3 + 2h_2)]}$$

and γ_k is obtained from γ_0 by replacing h_j by h_{k+1-j} ; $j = 1, 2, 3, 4$,
throughout.

Table 1

Values of $|Q(x)-y(x)|$. (Knots as in 5.1)

X	S_I	$Q_{1,1}$	$Q_{1,2}$	$Q_{2,1}$	$Q_{2,2}$
0.01	$.674 \times 10^{-8}$	$.114 \times 10^{-9}$	$.120 \times 10^{-9}$	$.733 \times 10^{-11}$	$.803 \times 10^{-12}$
0.02	$.151 \times 10^{-7}$	$.564 \times 10^{-9}$	$.593 \times 10^{-9}$	$.334 \times 10^{-10}$	$.402 \times 10^{-11}$
0.09	$.705 \times 10^{-8}$	$.497 \times 10^{-9}$	$.529 \times 10^{-9}$	$.364 \times 10^{-10}$	$.472 \times 10^{-11}$
0.22	$.189 \times 10^{-7}$	$.446 \times 10^{-9}$	$.366 \times 10^{-9}$	$.797 \times 10^{-10}$	$.519 \times 10^{-12}$
0.36	$.990 \times 10^{-8}$	$.840 \times 10^{-9}$	$.799 \times 10^{-9}$	$.369 \times 10^{-10}$	$.412 \times 10^{-11}$
0.62	$.281 \times 10^{-7}$	$.683 \times 10^{-9}$	$.563 \times 10^{-9}$	$.117 \times 10^{-9}$	$.245 \times 10^{-11}$
0.93	$.374 \times 10^{-7}$	$.152 \times 10^{-8}$	$.148 \times 10^{-8}$	$.102 \times 10^{-9}$	$.621 \times 10^{-10}$
0.96	$.184 \times 10^{-7}$	$.213 \times 10^{-8}$	$.219 \times 10^{-8}$	$.230 \times 10^{-10}$	$.381 \times 10^{-10}$
0.99	$.179 \times 10^{-7}$	$.276 \times 10^{-9}$	$.291 \times 10^{-9}$	$.102 \times 10^{-10}$	$.510 \times 10^{-11}$

Table 2

Values of $D^{(3)}$. (Knots as in 5.1)

	S_I	$Q_{1,1}$	$Q_{1,2}$	$Q_{2,1}$	$Q_{2,2}$
$D^{(3)}$.130	$.285 \times 10^{-2}$	$.186 \times 10^{-2}$	$.921 \times 10^{-3}$	$.714 \times 10^{-4}$

Table 3

Values of $|Q(x)-y(x)|$. (Knots as in 5.2)

X	S_I	$Q_{1,1}$	$Q_{1,2}$	$Q_{2,1}$	$Q_{2,2}$
0.01	$.512 \times 10^{-9}$	$.252 \times 10^{-10}$	$.380 \times 10^{-10}$	$.227 \times 10^{-11}$	$.105 \times 10^{-10}$
0.05	$.287 \times 10^{-8}$	$.200 \times 10^{-8}$	$.253 \times 10^{-8}$	$.842 \times 10^{-9}$	$.315 \times 10^{-9}$
0.1	$.804 \times 10^{-7}$	$.858 \times 10^{-8}$	$.139 \times 10^{-7}$	$.341 \times 10^{-8}$	$.194 \times 10^{-8}$
0.17	$.297 \times 10^{-6}$	$.182 \times 10^{-7}$	$.577 \times 10^{-8}$	$.172 \times 10^{-7}$	$.484 \times 10^{-8}$
0.35	$.589 \times 10^{-6}$	$.293 \times 10^{-6}$	$.352 \times 10^{-6}$	$.314 \times 10^{-7}$	$.277 \times 10^{-7}$
0.5	$.272 \times 10^{-5}$	$.758 \times 10^{-6}$	$.960 \times 10^{-6}$	$.325 \times 10^{-6}$	$.122 \times 10^{-6}$
0.6	$.325 \times 10^{-5}$	$.964 \times 10^{-6}$	$.836 \times 10^{-6}$	$.413 \times 10^{-8}$	$.123 \times 10^{-6}$
0.8	$.721 \times 10^{-5}$	$.233 \times 10^{-5}$	$.229 \times 10^{-5}$	$.194 \times 10^{-6}$	$.154 \times 10^{-6}$
0.9	$.207 \times 10^{-4}$	$.220 \times 10^{-5}$	$.212 \times 10^{-5}$	$.227 \times 10^{-6}$	$.150 \times 10^{-6}$

Table 4

Values of $D^{(3)}$. (Knots as in 5.2)

	S_I	$Q_{1,1}$	$Q_{1,2}$	$Q_{2,1}$	$Q_{2,2}$
$D^{(3)}$.484	$.433 \times 10^{-1}$	$.324 \times 10^{-1}$	$.272 \times 10^{-1}$	$.423 \times 10^{-2}$

R E F E R E N C E S

1. BEHFOROOZ, G.H. and PAPAMICHAEL, N. 1979 End conditions for cubic spline interpolation. J. Inst. Maths Applies . 23, 355 -366.
2. BEHFOROOZ, G.H. and PAPAMICHAEL, N, 1980 End conditions for interpolatory cubic splines with unequally spaced knots. J. Comp. Appl. Maths (to appear).
3. BEHFOROOZ, G.H., PAPAMICHAEL, N. and WORSEY, A.J. 1979 A class of piecewise cubic interpolatory polynomials. J. Inst. Maths Applies, (to appear).
4. BIRKHOFF, G. and PRIVER, A. 1967 Hermite interpolation errors for derivatives. J. Math. Phys. 46, 440 – 447 .
5. CLENSHAW, C.W. and NEGUS, B. 1978 The cubic X-spline and its application to interpolation. J. Inst. Maths Applies. 22 109 - 119.
6. HALL, C.A. 1968 On error bounds for spline interpolation. J. Approx. Theory 1, 209-218.
7. LUCAS, T.R. 1974 Error bounds for interpolating cubic splines under various end conditions. SIAM J. Numer. Anal. 11, 569-584.

2 WEEK LOAN

**NOT TO BE
REMOVED**
FROM THE LIBRARY

XB 2356403 2

