



## A Class of Non-Holonomic Projective Connections on Sub-Riemannian Manifolds

Yanling Han<sup>a,b</sup>, Fengyun Fu<sup>c</sup>, Peibiao Zhao<sup>a</sup>

<sup>a</sup>Department of Applied Mathematics, Nanjing University of Science and Technology, Nanjing 210094, P. R. China

<sup>b</sup>School of Science, Qilu University of Technology, Jinan 250353, P. R. China

<sup>c</sup>School of Mathematics and Statistics, Guangdong University of Finance and Economics, Guangzhou 510320, P. R. China

**Abstract.** The authors define a semi-symmetric non-holonomic (SSNH)-projective connection on sub-Riemannian manifolds and find an invariant of the SSNH-projective transformation. The authors further derive that a sub-Riemannian manifold is of projective flat if and only if the Schouten curvature tensor of a special SSNH-connection is zero.

### 1. Introduction

Since A. Friedmann and J. A. Schouten [8], in the early days of 1924, firstly introduced the concept of semi-symmetric linear connections, the research related to the semi-symmetric connection was unusually brilliant, and made a series of fruitful research results.

K. Yano [21] introduced and studied the semi-symmetric metric connection of Riemannian manifolds. N. S. Agashe and M. R. Chafle [1] introduced a semi-symmetric non-metric connection on a Riemannian manifold and this was further studied by U. C. De etc. [4–7], T. Imai [15], P. B. Zhao and H. Z. Song [27], and so on.

D. K. Sen and J. R. Vastane [19] studied the Weyl manifold by using the idea of semi-symmetric connections. Later the Weyl structure was extended in the semi-Riemannian distribution framework by O. Constantinescu and M. Crasmareanu, see [3]. I. Hinterleitner and J. Mikeš [13] studied the geodesic mappings onto Weyl manifolds. M. Tripathi and N. Nakkar [20] studied the semi-symmetric non-metric connection in a Kenmotsu manifold. B. Barua and A. K. Ray [2] studied the curvature properties of semi-symmetric metric connections and derived a sufficient and necessary condition for the Ricci tensor being of symmetric.

H. B. Yilmaz, F. O. Zengin and S. Aynur Uysal [22] considered a manifold equipped with a semi-symmetric metric connection whose torsion tensor satisfied a special condition and proved that if a manifold

---

2010 *Mathematics Subject Classification.* Primary 53C20; Secondary 53D11

*Keywords.* Sub-Riemannian manifolds, Non-holonomic connection, SSNH-projective connection.

Received: 05 February 2015; Accepted: 07 May 2015

Communicated by Mića Stanković

The first author is supported by a Grant-in-Aid for Scientific Research from Nanjing University of Science and Technology (KN11008).

The second author is supported by NSFC(No. 11526055).

The third author is supported by a Grant-in-Aid for Science Research from Nanjing University of Science and Technology (30920140132035), by NUST Research Funding No.CXZZ11-0258, AD20370 and by the NNSF of China(No.11371194; No.11571172).

*Email addresses:* hanyanling1979@163.com (Yanling Han), ffy1984229@163.com (Fengyun Fu), pbzhao@njjust.edu.cn (Peibiao Zhao)

mentioned as above was conformally flat, then it was a mixed quasi-Einstein manifold. F. Y. Fu, X. P. Yang and P. B. Zhao [9] considered the geometric and physical properties of conformal mappings for the semi-Riemannian manifolds.

J. Mikeš [17, 18] studied the projective and geodesic mappings of special Riemannian spaces. F. Y. Fu and P. B. Zhao [10] studied the geodesic mapping of pseudo-symmetric Riemannian Manifolds. In particular, the second author [23] recently studied the conformal and projective characteristics of sub-Riemannian manifolds by using the so-called non-holonomic sub-Riemannian connection. I. Hinterleitner [11] studied geodesic mappings on compact Riemannian manifolds. I. Hinterleitner and J. Mikeš [12, 13, 16] studied projective and affine connections. M. Zlatanović and etc. [24–26] studied geodesic mappings and similar problems.

However, to the author’s knowledge, the study of geometric and analysis in sub-Riemannian manifolds on view of the semi-symmetric metric connection in sub-Riemannian manifolds is still a gap.

In this paper, we will, based on the setting of [14], investigate a class of semi-symmetric non-holonomic connections, find the sub-Weyl projective invariant, and study the projective flatness of sub-Riemannian manifolds.

## 2. The SSNH-Projective Transformation

Let  $(M, \Delta, g_\Delta)$  be a  $n$ -dimensional sub-Riemannian manifold, where  $\Delta$  is a  $\ell$ -dimensional sub-bundle of tangent bundles, and is called a horizontal bundle, and  $g_\Delta$  is a Riemannian metric defined on  $\Delta$ . In particular, when  $\Delta = TM$ ,  $(M, \Delta, g_\Delta)$  will be degenerated into a Riemannian manifold. Without loss of generality, we assume  $\Delta \neq TM$ . In this subsection, we will define a semi-symmetric non-holonomic(SSNH) metric connection and discuss the SSNH-projective transformation following the work in [14].

We use unless otherwise noted the following ranges for indices:  $i, j, k, h, \dots \in \{1, \dots, \ell\}$ . The repeated indices with one upper index and one lower index indicates the summation over their range. The projection of  $X$  on the horizontal bundle is denoted by  $X_h$ .

**Definition 2.1.** A non-holonomic connection on sub-bundle  $Q \subset TM$  is a mapping  $\nabla : \Gamma(Q) \times \Gamma(Q) \rightarrow \Gamma(Q)$  satisfying the following

$$\nabla_{X_h}(Y_h + Z_h) = \nabla_{X_h}Y_h + \nabla_{X_h}Z_h, \nabla_{X_h}(fY_h) = X_h(f)Y_h + f\nabla_{X_h}Y_h, \nabla_{fX_h+gY_h}Z_h = f\nabla_{X_h}Z_h + g\nabla_{Y_h}Z_h,$$

where  $X, Y, Z \in \Gamma(TM)$ ,  $f, g \in C^\infty(M)$ .

**Definition 2.2.** A non-holonomic connection is said to be metric and symmetric if it satisfies respectively,

$$\begin{aligned} (\nabla_{Z_h}g_\Delta)(X_h, Y_h) &= Z_h(g_\Delta(X_h, Y_h)) - g_\Delta(\nabla_{Z_h}X_h, Y_h) - g_\Delta(X_h, \nabla_{Z_h}Y_h) = 0, \\ T(X_h, Y_h) &= \nabla_{X_h}Y_h - \nabla_{Y_h}X_h - [X_h, Y_h]_h = 0. \end{aligned}$$

**Definition 2.3.** A non-holonomic connection is said to be a sub-Riemannian connection if it is both metric and symmetric.

**Definition 2.4.** A horizontal curve  $\gamma(t) : [0, 1] \rightarrow M$  (i.e.  $\dot{\gamma}(t) \in \Delta_{\gamma(t)}$ ) is said to be a sub-Riemannian parallel (in briefly, SR-parallel) curve if it satisfies

$$\nabla_{\dot{\gamma}}\dot{\gamma} = 0, \tag{1}$$

where  $\nabla$  is the sub-Riemannian connection.

Let  $\gamma : x^a = x^a(t)$ , the corresponding equation (1) is

$$\frac{d^2x^k}{dt^2} + \binom{k}{ij} \frac{dx^i}{dt} \frac{dx^j}{dt} = 0, \tag{2}$$

where  $t$  is an affine parameter.

**Definition 2.5.** Let  $D_1, D_2$  be two classes of non-holonomic connections. If a SR-parallel curve corresponding to  $D_1$  coincides always with one corresponding to  $D_2$ , then we say that  $D_1$  is a projective correspondence to  $D_2$ .

The second author [23] proved that a non-holonomic symmetric connection  $D$  is a projective correspondence if and only if there exists a smooth horizontal 1-form  $\varphi$  (i.e. a 1-form defined on  $\Delta$ ), such that, for any two horizontal vector fields  $X_h, Y_h$ , there holds

$$D_{X_h} Y_h = \nabla_{X_h} Y_h + \varphi(X_h)Y_h + \varphi(Y_h)X_h. \tag{3}$$

If  $D$  is a non-holonomic connection with torsion, then we have the following

**Proposition 2.6.** A non-holonomic connection  $D$  with torsion is a projective correspondence to  $\nabla$  if and only if there exists 1-form  $\lambda$  such that the symmetric part of tensor  $A(X_h, Y_h) = D_{X_h} Y_h - \nabla_{X_h} Y_h$  is of the form

$$(A(X_h, Y_h) + A(Y_h, X_h))/2 = \lambda(X_h)Y_h + \lambda(Y_h)X_h, \text{ for } X, Y \in \Gamma(TM). \tag{4}$$

*Proof.* The necessity is obvious. We only prove the sufficiency. If (4) holds, we denote by  $(D_{X_h} Y_h + D_{Y_h} X_h)/2 = \tilde{D}_{X_h} Y_h, (\nabla_{X_h} Y_h + \nabla_{Y_h} X_h)/2 = \tilde{\nabla}_{X_h} Y_h$ , then (4) is equivalent to  $\tilde{D}_{X_h} Y_h - \tilde{\nabla}_{X_h} Y_h = \lambda(X_h)Y_h + \lambda(Y_h)X_h$ . Hence  $\tilde{D}$  and  $\tilde{\nabla}$  have the same SR-parallel curves by (3).

On the other hand, if  $\gamma(t)$  is a SR-parallel curve of  $D$ , then  $\gamma(t)$  is also a SR-parallel curve of  $\tilde{D}$  by a simple computation. Hence  $\tilde{D}$  and  $D$  also have the same SR-parallel curves, so do  $\tilde{\nabla}$  and  $\nabla$ . Therefore,  $D$  and  $\nabla$  have the same SR-parallel curves, namely,  $D$  is a projective correspondence of  $\nabla$ .  $\square$

**Definition 2.7.** If  $\bar{\nabla}$  is a projective correspondence to  $\nabla$  with torsion,

$$\bar{T}(X_h, Y_h) = \pi(Y_h)X_h - \pi(X_h)Y_h, \tag{5}$$

where  $\pi$  is a given 1-form, then we say that  $\bar{\nabla}$  is a semi-symmetric non-holonomic projective connection, in briefly, a SSNH-projective connection.

**Theorem 2.8.**  $\bar{\nabla}$  is a SSNH-projective connection if and only if there exist two 1-form  $p, q$  such that

$$\bar{\nabla}_{X_h} Y_h = \nabla_{X_h} Y_h + p(X_h)Y_h + q(Y_h)X_h, \tag{6}$$

for any  $X, Y \in \Gamma(TM)$ .

*Proof.* Let  $A(X_h, Y_h) = \bar{\nabla}_{X_h} Y_h - \nabla_{X_h} Y_h$ . Since  $\bar{\nabla}$  is a SSNH-projective connection, from Proposition 2.6, there exists a smooth 1-form  $\varphi$  such that

$$(A(X_h, Y_h) + A(Y_h, X_h))/2 = \varphi(X_h)Y_h + \varphi(Y_h)X_h, \text{ for } X, Y \in TM \tag{7}$$

and 1-form  $\pi$  such that the torsion of  $\bar{\nabla}$  is of the form  $\bar{T}(X_h, Y_h) = \pi(Y_h)X_h - \pi(X_h)Y_h$ , we can deduce from the above equation

$$A(X_h, Y_h) - A(Y_h, X_h) = \pi(Y_h)X_h - \pi(X_h)Y_h. \tag{8}$$

By (6) and (8), we arrive at  $A(X_h, Y_h) = (\varphi - \pi/2)(X_h)Y_h + (\varphi + \pi/2)(Y_h)X_h$  for  $p = \varphi - \pi/2, q = \varphi + \pi/2$ .

Conversely, we assume  $\bar{\nabla}_{X_h} Y_h = \nabla_{X_h} Y_h + p(Y_h)X_h + q(X_h)Y_h$ , then

$$(A(X_h, Y_h) + A(Y_h, X_h))/2 = \frac{p+q}{2}(Y_h)X_h + \frac{p+q}{2}(X_h)Y_h,$$

$$A(X_h, Y_h) - A(Y_h, X_h) = (p - q)(Y_h)X_h - (p - q)(X_h)Y_h.$$

By virtue of Proposition 2.6 again, we know  $\bar{\nabla}$  is a projective correspondence to  $\nabla$ , and we get

$$\bar{T}(X_h, Y_h) = \bar{\nabla}_{X_h} Y_h - \bar{\nabla}_{Y_h} X_h - [X_h, Y_h]_h = (p - q)(Y_h)X_h - (p - q)(X_h)Y_h.$$

Let  $\pi = p - q$ , then  $\bar{T}(X_h, Y_h) = \pi(Y_h)X_h - \pi(X_h)Y_h$ .

This completes the proof of Theorem 2.8.  $\square$

In a basis  $\{e_i\}$ , (6) can be rewritten as

$$\bar{\Gamma}_{ij}^k = \{^k_{ij}\} + p_i \delta_j^k + q_j \delta_i^k = \{^k_{ij}\} + \varphi_i \delta_j^k + \varphi_j \delta_i^k + \rho_j \delta_i^k - \rho_i \delta_j^k,$$

where  $\rho_i = \pi_i/2, p_i = \varphi_i - \rho_i, q_i = \varphi_i + \rho_i$ . The Schouten curvature tensor, Ricci tensor and sub-Weyl projective curvature tensor are given, respectively, as

$$\begin{cases} \bar{R}_{ijk}^h = R_{ijk}^h + \beta_{ij} \delta_k^h + \alpha_{ik} \delta_j^h - \alpha_{jk} \delta_i^h - \delta_k^h \Omega_{ij}^s p_s - \Omega_{ij}^h p_k, \\ \bar{R}_{jk} = R_{jk} + \beta_{kj} - (\ell - 1) \alpha_{jk} - \Omega_{kj}^s p_s - \Omega_{\varepsilon j}^\varepsilon p_k, \\ W_{ijk}^h = R_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h R_{ik} - \delta_i^h R_{jk}), \end{cases} \tag{9}$$

where

$$\begin{cases} \beta_{ij} = (\nabla_i p)(e_j) - (\nabla_j p)(e_i) = \varphi_{ij} - \varphi_{ji} + \rho_{ji} - \rho_{ij}, \\ \alpha_{ij} = (\nabla_i q)(e_j) - q(e_i)q(e_j) = \varphi_{ij} + \rho_{ij} - \varphi_i \rho_j - \varphi_j \rho_i, \\ \varphi_{ij} = e_i(\varphi_j) - \Gamma_{ij}^\varepsilon \varphi_\varepsilon - \varphi_i \varphi_j = \nabla_i \varphi_j - \varphi_i \varphi_j, \\ \rho_{ij} = e_i(\rho_j) - \Gamma_{ij}^\varepsilon \rho_\varepsilon - \rho_i \rho_j = \nabla_i \rho_j - \rho_i \rho_j, \\ R_{jk} = R_{\varepsilon jk}^\varepsilon, \bar{R}_{jk} = \bar{R}_{\varepsilon jk}^\varepsilon. \end{cases} \tag{10}$$

**Theorem 2.9.** *The tensor  $S_{ijk}^h$  is an invariant under a SSNH-projective transformation, where*

$$\begin{aligned} S_{ijk}^h &= R_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h R_{ik} - \delta_i^h R_{jk}) \\ &\quad + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \{ \delta_j^h (R_{ik} - R_{ki}) - \delta_i^h (R_{jk} - R_{kj}) - (\ell - 1) \delta_k^h (R_{ij} - R_{ji}) \} \\ &\quad + \frac{1}{\ell^2 - \ell - 2} \{ \delta_j^h A_{ik} - \delta_i^h A_{jk} - (\ell - 1) \delta_k^h A_{ij} \}, \end{aligned} \tag{11}$$

and  $A_{ij} = R_{ijs}^s$ .

*Proof.* For simplicity, we choose  $\{e_i\}$  as a local frame field such that  $[e_i, e_j] \in VM$ , and hence we have  $\Omega_{ij}^h = 0$ . Then the Schouten curvature tensors and Ricci curvature tensors can be written simply as

$$\bar{R}_{ijk}^h = R_{ijk}^h + \beta_{ij} \delta_k^h + \alpha_{ik} \delta_j^h - \alpha_{jk} \delta_i^h \text{ and } \bar{R}_{jk} = R_{jk} + \beta_{kj} - (\ell - 1) \alpha_{jk},$$

Let  $k = h = \varepsilon$ , and denote by  $A_{ij} = R_{ije}^\varepsilon, \bar{A}_{ij} = \bar{R}_{ije}^\varepsilon$ , one obtains

$$\bar{A}_{ij} = A_{ij} + \ell \beta_{ij} + \alpha_{ij} - \alpha_{ji},$$

hence one arrives at

$$\begin{aligned} \beta_{jk} &= \frac{1}{\ell^2 - \ell - 2} [(\bar{R}_{jk} - \bar{R}_{kj}) - (R_{jk} - R_{kj}) + (\ell - 1)(\bar{A}_{jk} - A_{jk})], \\ \alpha_{jk} &= \frac{1}{\ell - 1} (R_{jk} - \bar{R}_{jk}) - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} [(\bar{R}_{jk} - \bar{R}_{kj}) - (R_{jk} - R_{kj})] \\ &\quad - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} (\bar{A}_{jk} - A_{jk}). \end{aligned}$$

moreover one has

$$\begin{aligned} \bar{R}_{ijk}^h &= R_{ijk}^h + \frac{\delta_k^h}{\ell^2 - \ell - 2} [(\bar{R}_{ij} - \bar{R}_{ji}) - (R_{ij} - R_{ji}) + (\ell - 1)(\bar{A}_{ij} - A_{ij})] \\ &+ \frac{\delta_j^h}{\ell - 1} (R_{ik} - \bar{R}_{ik}) - \frac{\delta_j^h}{(\ell - 1)(\ell^2 - \ell - 2)} [(\bar{R}_{ik} - \bar{R}_{ki}) - (R_{ik} - R_{ki})] \\ &- \frac{\delta_j^h}{(\ell - 1)(\ell^2 - \ell - 2)} (\bar{A}_{ik} - A_{ik}) - \frac{\delta_i^h}{\ell - 1} (R_{jk} - \bar{R}_{jk}) \\ &+ \frac{\delta_i^h}{(\ell - 1)(\ell^2 - \ell - 2)} [(\bar{R}_{jk} - \bar{R}_{kj}) - (R_{jk} - R_{kj})] \\ &+ \frac{\delta_i^h}{(\ell - 1)(\ell^2 - \ell - 2)} (\bar{A}_{jk} - A_{jk}). \end{aligned}$$

Rewriting the above equation by

$$\begin{aligned} &\bar{R}_{ijk}^h - \frac{\delta_k^h}{\ell^2 - \ell - 2} (\bar{R}_{ij} - \bar{R}_{ji}) + \frac{\ell - 1}{\ell^2 - \ell - 2} \delta_k^h \bar{A}_{ij} + \frac{1}{\ell - 1} \delta_j^h \bar{R}_{ik} \\ &+ \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_j^h (\bar{R}_{ik} - \bar{R}_{ki}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_j^h \bar{A}_{ik} \\ &- \frac{1}{\ell - 1} \delta_i^h \bar{R}_{jk} - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_i^h (\bar{R}_{jk} - \bar{R}_{kj}) - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_i^h \bar{A}_{jk} \\ = &R_{ijk}^h - \frac{\delta_k^h}{\ell^2 - \ell - 2} (R_{ij} - R_{ji}) + \frac{\ell - 1}{\ell^2 - \ell - 2} \delta_k^h A_{ij} + \frac{1}{\ell - 1} \delta_j^h R_{ik} \\ &+ \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_j^h (R_{ik} - R_{ki}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_j^h A_{ik} \\ &- \frac{1}{\ell - 1} \delta_i^h R_{jk} - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_i^h (R_{jk} - R_{kj}) - \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \delta_i^h A_{jk}, \end{aligned}$$

that is

$$\begin{aligned} &\bar{R}_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h \bar{R}_{ik} - \delta_i^h \bar{R}_{jk}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \{ \delta_j^h (\bar{R}_{ik} - \bar{R}_{ki}) - \delta_i^h (\bar{R}_{jk} - \bar{R}_{kj}) \\ &- (\ell - 1) \delta_k^h (\bar{R}_{ij} - \bar{R}_{ji}) \} + \frac{1}{\ell^2 - \ell - 2} \{ \delta_j^h \bar{A}_{ik} - \delta_i^h \bar{A}_{jk} - (\ell - 1) \delta_k^h \bar{A}_{ij} \} \\ = &R_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h R_{ik} - \delta_i^h R_{jk}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \{ \delta_j^h (R_{ik} - R_{ki}) - \delta_i^h (R_{jk} - R_{kj}) \\ &- (\ell - 1) \delta_k^h (R_{ij} - R_{ji}) \} + \frac{1}{\ell^2 - \ell - 2} \{ \delta_j^h A_{ik} - \delta_i^h A_{jk} - (\ell - 1) \delta_k^h A_{ij} \}. \end{aligned}$$

Denote by

$$\begin{aligned} \bar{S}_{ijk}^h &= \bar{R}_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h \bar{R}_{ik} - \delta_i^h \bar{R}_{jk}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \{ \delta_j^h (\bar{R}_{ik} - \bar{R}_{ki}) - \delta_i^h (\bar{R}_{jk} - \bar{R}_{kj}) \\ &- (\ell - 1) \delta_k^h (\bar{R}_{ij} - \bar{R}_{ji}) \} + \frac{1}{\ell^2 - \ell - 2} \{ \delta_j^h \bar{A}_{ik} - \delta_i^h \bar{A}_{jk} - (\ell - 1) \delta_k^h \bar{A}_{ij} \}, \end{aligned}$$

and

$$\begin{aligned} S_{ijk}^h &= R_{ijk}^h + \frac{1}{\ell - 1} (\delta_j^h R_{ik} - \delta_i^h R_{jk}) + \frac{1}{(\ell - 1)(\ell^2 - \ell - 2)} \{ \delta_j^h (R_{ik} - R_{ki}) - \delta_i^h (R_{jk} - R_{kj}) \\ &- (\ell - 1) \delta_k^h (R_{ij} - R_{ji}) \} + \frac{1}{\ell^2 - \ell - 2} \{ \delta_j^h A_{ik} - \delta_i^h A_{jk} - (\ell - 1) \delta_k^h A_{ij} \}. \end{aligned}$$

then one obtains  $S_{ijk}^h = \bar{S}_{ijk}^h$ . This ends the proof of Theorem 2.9.  $\square$

We now similarly define the sub-Weyl projective curvature tensor of the SSNH-projective connection by

$$\bar{W}_{ijk}^h = \bar{R}_{ijk}^h + \frac{1}{\ell - 1}(\delta_j^h \bar{R}_{ik} - \delta_i^h \bar{R}_{jk}), \tag{12}$$

then we have

$$\begin{aligned} W_{ijk}^h &= W_{ijk}^h + \delta_k^h(\beta_{ij} - \Omega_{ij}^s p_s) + \frac{1}{\ell - 1}(\delta_j^h \beta_{ik} - \delta_i^h \beta_{jk}) \\ &\quad - \frac{1}{\ell - 1}(\delta_j^h \Omega_{ki}^s - \delta_i^h \Omega_{kj}^s) p_s - [\Omega_{ij}^h - \frac{1}{\ell - 1}(\delta_j^h \Omega_{ie}^e - \delta_i^h \Omega_{je}^e)] p_k. \end{aligned}$$

We denote by

$$\begin{aligned} B_{ijk}^h &= \delta_k^h(\beta_{ij} - \Omega_{ij}^s p_s) + \frac{1}{\ell - 1}(\delta_j^h \beta_{ik} - \delta_i^h \beta_{jk}) \\ &\quad - \frac{1}{\ell - 1}(\delta_j^h \Omega_{ki}^s - \delta_i^h \Omega_{kj}^s) p_s - [\Omega_{ij}^h - \frac{1}{\ell - 1}(\delta_j^h \Omega_{ie}^e - \delta_i^h \Omega_{je}^e)] p_k, \end{aligned}$$

then it is obvious that  $\bar{W}_{ijk}^h = W_{ijk}^h + B_{ijk}^h$ .

**Definition 2.10.** If the 1-form  $p$  and  $q$  in (6) are horizontally closed, that is,

$$\begin{aligned} dp(X_h, Y_h) &= X_h(p(Y_h)) - Y_h(p(X_h)) - p([X_h, Y_h]_h) = 0, \\ dq(X_h, Y_h) &= X_h(q(Y_h)) - Y_h(q(X_h)) - q([X_h, Y_h]_h) = 0, \end{aligned}$$

then we call a SSNH-projective connection  $\tilde{\nabla}$  the special SSNH-projective connection.

**Theorem 2.11.** The sub-Weyl projective curvature tensor is an invariant under a special SSNH-projective transformation.

*Proof.* If  $\tilde{\nabla}$  is a special SSNH-projective connection, then the 1-form  $p$  and  $q$  in (9) are all horizontally closed. Therefore there holds

$$\begin{aligned} 0 &= dp(e_i, e_j) = e_i(p_j) - e_j(p_i) - p([e_i, e_j]_h) = \varphi_{ij} - \varphi_{ji} + \rho_{ij} - \rho_{ji}, \\ 0 &= dq(e_i, e_j) = e_i(q_j) - e_j(q_i) - q([e_i, e_j]_h) = \varphi_{ij} - \varphi_{ji} + \rho_{ji} - \rho_{ij}. \end{aligned}$$

By adding above two equations one gets  $\varphi_{ij} = \varphi_{ji}$ , and  $\rho_{ij} = \rho_{ji}$  by subtracting these equations. Then one obtains  $\beta_{ij} = 0$  and

$$\bar{R}_{ijk}^h = R_{ijk}^h + \alpha_{ik} \delta_j^h - \alpha_{jk} \delta_i^h,$$

Contracting by  $i$  and  $h$ , one gets

$$\bar{R}_{jk} = R_{jk} - (\ell - 1)\alpha_{jk},$$

Therefore, one obtains

$$\begin{aligned} \bar{W}_{ijk}^h &= \bar{R}_{ijk}^h + \frac{1}{\ell - 1}(\delta_j^h \bar{R}_{ik} - \delta_i^h \bar{R}_{jk}) \\ &= R_{ijk}^h + \alpha_{ik} \delta_j^h - \alpha_{jk} \delta_i^h + \frac{1}{\ell - 1} \delta_j^h (R_{ik} - (\ell - 1)\alpha_{ik}) - \frac{1}{\ell - 1} \delta_i^h (R_{jk} - (\ell - 1)\alpha_{jk}) \\ &= W_{ijk}^h. \end{aligned}$$

The proof is finished.  $\square$

**Remark 2.12.** It is obvious that a projectively flat sub-Riemannina manifold is transformed to a projectively flat sub-Riemannina manifold by a SSNH-projective transformation.

**Theorem 2.13.** A sub-Riemannian manifold  $(M, \Delta, g_\Delta)$  is projective flat if and only if the Schouten curvature tensor  $\tilde{R}$  of the special SSNH-projective connection  $\tilde{D}$  is vanished.

*Proof.* If  $\tilde{\nabla}$  is a special SSNH-projective connection and

$$\tilde{R}_{ijk}^h = R_{ijk}^h + \beta_{ij}\delta_k^h + \alpha_{ik}\delta_j^h - \alpha_{jk}\delta_i^h = 0, \tag{13}$$

then by contracting (13) with  $i, h$ , we have  $\tilde{R}_{jk} = R_{jk} + \beta_{kj} - (\ell - 1)\alpha_{jk} = 0$ . Since  $\tilde{\nabla}$  is special, then the 1-form  $p$  is horizontally closed. Hence we get  $\beta_{ij} = 0$ , and

$$R_{ijk}^h = \alpha_{jk}\delta_i^h - \alpha_{ik}\delta_j^h, R_{ik} = (\ell - 1)\alpha_{ik}, \tag{14}$$

By substituting (14) into the following equation

$$W_{ijk}^h = R_{ijk}^h + \frac{1}{\ell - 1}(\delta_j^h R_{ik} - \delta_i^h R_{jk}),$$

we obtain  $W_{ijk}^h = 0$ , that is,  $M$  is projectively flat.

Conversely, if  $M$  is projectively flat, then  $W_{ijk}^h = 0$ , and  $R_{ijk}^h = \frac{1}{\ell-1}(\delta_j^h R_{ik} - \delta_i^h R_{jk})$ , namely,  $R_{ijkh} = \frac{1}{\ell-1}(g_{ih}R_{jk} - g_{jh}R_{ik})$ . Since  $R_{ijhh} = 0$ , we get  $R_{ik} = \frac{R}{\ell}g_{ik}$ . If the 1-form  $p$  is horizontally closed, then the equation  $\tilde{R}_{ij} = R_{ij} + \beta_{ij} - (\ell - 1)\alpha_{ij} = 0$  is equivalent to

$$(\nabla_i q)(e_j) - q_i q_j = \frac{R}{\ell(\ell - 1)}g_{ij}, \tag{15}$$

where  $(\nabla_i q)(e_j) - q_i q_j = \alpha_{ij}$ .

Now taking a covariant derivative of Equation (15), we get

$$\begin{aligned} & (\nabla_i \nabla_j q)(e_k) + (\nabla_j q)(\nabla_i e_k) - (\nabla_i q)(e_j)q(e_k) - q(\nabla_i e_j)q(e_k) - q(e_j)(\nabla_i q)(e_k) - q(\nabla_i e_k)q(e_j) \\ &= \frac{K}{\ell(\ell - 1)}(g(\nabla_i e_j, e_k) + g(e_j, \nabla_i e_k)) \\ &= (\nabla_{\nabla_i e_j} q)(e_k) - q(\nabla_i e_j)q(e_k) + (\nabla_j q)(\nabla_i e_k) - q(\nabla_i e_k)q(e_j), \end{aligned}$$

where the last equality follows from Equation (15). Namely,

$$(\nabla_i \nabla_j q)(e_k) - (\nabla_i q)(e_j)q(e_k) - q(e_j)(\nabla_i q)(e_k) = (\nabla_{\nabla_i e_j} q)(e_k). \tag{16}$$

Since the horizontal 1-form  $p$  is closed, then by (15), (16) and  $W_{ijk}^h = 0$ , we obtain

$$(\nabla_i \nabla_j q - \nabla_j \nabla_i q - \nabla_{[e_i, e_j]_h} q)(e_k) = -R_{ijk}^h q_h, \tag{17}$$

therefore there exists a solution  $q$  to Equation (15), let

$$\tilde{\Gamma}_{ij}^k = \{^k_{ij}\} + p_i \delta_j^k + q_j \delta_i^k, \tag{18}$$

where  $p$  is a closed horizontal 1-form.

By Theorem 2.8, we know  $\tilde{\nabla}$  whose connection coefficients are defined by (18) is a SSNH-projective connection. On the other hand,  $\alpha_{ij}$  is proportional to  $g_{ij}$  by (15), so it is symmetric and  $dq(e_i, e_j) = \alpha_{ij} - \alpha_{ji} = 0$ , which implies that the 1-form  $q$  is horizontally closed.

This completes the proof of Theorem 2.13.  $\square$

### 3. Example

**Example 3.1.** (Almost contact metric sub-Riemannian manifold)

Let  $(M, \Delta, g_\Delta)$  be a  $(2n + 1)$ -dimensional sub-Riemannian manifold, an almost contact structure is denoted by  $(\varphi, \xi, \eta)$ , where  $\varphi$  is a horizontal  $(1, 1)$ -tensor field (i.e.  $\varphi(X_h) \in \Delta$ ),  $\xi$  is a vector field and  $\eta$  is a 1-form such that

$$\varphi^2 = -I + \eta \otimes \xi, \eta(\xi) = 1, g(\varphi X_h, \varphi Y_h) = g(X_h, Y_h) - \eta(X_h)\eta(Y_h).$$

then  $(M, \Delta, g, \varphi, \xi, \eta)$  is called an almost contact metric sub-Riemannian manifold. In virtue of this 1-form  $\eta$ , one defines a metric connection,

$$\tilde{\nabla}_{X_h} Y_h = \nabla_{X_h} Y_h + \eta(X_h)Y_h + \eta(Y_h)X_h, \tag{19}$$

in local coordinate, that is,

$$\tilde{\Gamma}_{ij}^k = \Gamma_{ij}^k + \eta_i \delta_j^k + \eta_j \delta_i^k, \tag{20}$$

where  $\nabla$  is the sub-Riemannian connection, then  $\tilde{\nabla}$  is actually a SSNH-projective connection.

In fact, if  $\gamma : x^a = x^a(t)$  is a SR-parallel curve with respect to sub-Riemannian connection, then it satisfies Equations (2), substituting (20) into the above Equations, one obtains,

$$\frac{d^2 x^k}{dt^2} + \tilde{\Gamma}_{ij}^k \frac{dx^i}{dt} \frac{dx^j}{dt} = -2\eta_i \frac{dx^i}{dt} \frac{dx^k}{dt},$$

Now we introduce a new parameter  $s$  by the equation

$$s = \int e^{\int -2\eta_i dx^i} dt,$$

and obtain the following relations by straight-forward calculation,

$$\begin{aligned} \frac{ds}{dt} &= e^{\int -2\eta_j dx^j}, \frac{d^2 s}{dt^2} = e^{\int -2\eta_j dx^j} (-2\eta_j \frac{dx^j}{dt}), \\ \frac{dx^i}{dt} &= e^{\int -2\eta_j dx^j} \frac{dx^i}{ds}, \frac{d^2 x^i}{dt^2} = e^{2 \int -2\eta_j dx^j} \left( \frac{d^2 x^i}{ds^2} - 2\eta_j \frac{dx^j}{ds} \frac{dx^i}{ds} \right). \end{aligned}$$

hence, we have

$$\frac{d^2 x^k}{ds^2} + \tilde{\Gamma}_{ij}^k \frac{dx^i}{ds} \frac{dx^j}{ds} = 0.$$

that is  $\gamma : x^a = x^a(t)$  is also a SR-parallel curve associated with the connection (19). On the other hand, one can prove the converse statement is also true by the same method. Therefore, the metric connection (19) is a SSNH-projective connection.

### References

[1] N. S. Agashe and M. R. Chafle, A semi-symmetric non-metric connection in a Riemannian manifold, *Indian J. Pure Appl. Math.*, **23** (1992), 399–409.  
 [2] B. Barua and A. K. Ray, Some properties of semisymmetric metric connection in a Riemannian manifold, *Indian J. Pure Appl. Math.*, **16** (1985), 736–740.  
 [3] O. Constantinescu and M. Crasmareanu, Sub-Weyl geometry and its linear connections, *Int. J. Geom. Methods Mod. Phys.*, **9** (2012).  
 [4] U. C. De and S. C. Biswas, On a type of semi-symmetric metric connection on a Riemannian manifold, *Publ. Inst. Math. (Beograd) (N. S.)*, **61** (75) (1997), 90–96.  
 [5] U. C. De and D. Kamilya, Hypersurfaces of a Riemannian manifold with semi-symmetric non-metric connection, *J. Indian Inst. Sci.*, **75** (1995), 707–710.



- [6] U. C. De and G. C. Ghosh, On conformal flat special quasi-Einstein manifolds, *Publ. Math. Debrecen*, **57** (3-4) (2000), 297–306.
- [7] U. C. De and J. Sengupta, On a type of semi-symmetric non-metric connection on a Riemannian manifold, *Bull. Cal. Math. Soc.*, **92** (2000), 375–384.
- [8] A. Friedmann and J. A. Schouten, Über die Geometrie der Halbsymmetrischen. Übertragung. *Math. Z.*, **21** (1924), 211–233.
- [9] F. Y. Fu, X. P. Yang, and P. B. Zhao, Geometrical and physical characteristics of some class of conformal mappings. *J. Geom. Phys.*, **62** (6) (2012), 1467–1479.
- [10] F. Y. Fu and P. B. Zhao, A property on geodesic mappings of pseudosymmetric Riemannian manifolds, *Bull. Malays. Math. Sci. Soc.*, **33** (2) (2010), 265–272.
- [11] I. Hinterleitner, Geodesic mappings on a compact Riemannian manifolds with conditions on sectional curvature, *Publ. Inst. Math. (Beograd) (N.S.)*, **94** (108) (2013), 125–130.
- [12] I. Hinterleitner and J. Mikeš, Geodesic mappings and differentiability of metrics affine and projective connections, *Filomat*, **29** (6) (2015), 1245–1249.
- [13] I. Hinterleitner and J. Mikeš, Geodesic mappings onto Weyl manifolds, *8th Int. Conf. Appl. Math. (Aplimat 2009)*, Bratislava; *8th Int. Conf. Prol. (2009)*, 423–430.
- [14] Y. L. Han, F. Y. Fu and P. B. Zhao, On semi-symmetric metric connection in sub-Riemannian manifold (to appear in Tamkang Journal of Mathematics)
- [15] T. Imai, Notes on semi-symmetric metric connections, *Tensor*, **24** (1972), 293–296.
- [16] J. Mikeš etc. al., Differential geometry of special mappings, *Palacky Univ. Olomouc*, Olomouc, 2015, 568pp.
- [17] J. Mikeš, Geodesic and holomorphically projective mappings of special Riemannian spaces, PhD. Thesis, Odessa University (1979).
- [18] J. Mikeš, Geodesic mappings on semisymmetric spaces, (Russian) *Izv. Vyssh. Uchebn. Zaved. Mat.*, 1994, no. 2, 37–43; translation in *Russian Math. (Iz. VUZ)*, **38** (2) (1994), 35–41.
- [19] D. K. Sen and J. R. Vastane, On Weyl and Lyra manifolds, *J. Math. Phys.*, **13** (1972), 990–993.
- [20] M. M. Tripathi and N. Nakkar, On a semi-symmetric non-metric connection in a Kenmotsu manifold, *Bull. Cal. Math. Soc.*, **16** (4) (2001), 323–330.
- [21] K. Yano, On semi-symmetric metric connection, *Rev. Roum. Math. Pureset Appl.*, **15** (1970), 1579–1586.
- [22] H. Yılmaz, F. Zengin and S. Uysal, On a Semi Symmetric Metric Connection with a Special Condition on a Riemannian Manifold, *Eur. J. Pure Appl. Math.*, **4** (2011), 152–161.
- [23] P. B. Zhao and L. Jiao, Conformal transformations on Carnot Carathéodory spaces, *Nihonkai Math. J.*, **17** (2) 2006, 167–185.
- [24] M. Zlatanović, I. Hinterleitner and M. Najdanović, On equitorsion concircular tensor of generalized Riemannian spaces, *Filomat*, **28** (3) (2014), 463–471.
- [25] M. Zlatanović, L. S. Velimirović and M. S. Stanković, Necessary and sufficient conditions for equitorsion geodesic mapping, *J. Math. Anal. Appl.*, **435** (1) (2016), 578–592.
- [26] M. Zlatanović, S. M. Minčić and L. S. Velimirović, On integrability conditions of derivation equations in a subspace of asymmetric affine connection space, *Filomat*, **29** (10) (2015), 2421–2427.
- [27] P. B. Zhao and H. Z. Song, An invariant of the projective semi-symmetric connection, *Chinese Quarterly J of Math.*, **16** (4) (2001), 49–54.