

# A combinatorial proof of Postnikov's identity and a generalized enumeration of labeled trees

Seunghyun Seo\*

Department of Mathematics  
Brandeis University, Waltham, MA 02454, USA  
shseo@brandeis.edu

Submitted: Sep 16, 2004; Accepted: Dec 16, 2004; Published: Jan 24, 2005  
Mathematics Subject Classifications: 05A15, 05C05, 05C30

## Abstract


In this paper, we give a simple combinatorial explanation of a formula of A. Postnikov relating bicolored rooted trees to bicolored binary trees. We also present generalized formulas for the number of labeled  $k$ -ary trees, rooted labeled trees, and labeled plane trees.

## 1 Introduction

In Stanley's 60th Birthday Conference, Postnikov [3, p. 21] showed the following identity:

$$(n+1)^{n-1} = \sum_{\mathfrak{b}} \frac{n!}{2^n} \prod_{v \in V(\mathfrak{b})} \left(1 + \frac{1}{h(v)}\right), \quad (1)$$

where the sum is over unlabeled binary trees  $\mathfrak{b}$  on  $n$  vertices and  $h(v)$  denotes the number of descendants of  $v$  (including  $v$ ). The figure below illustrates all five unlabeled binary trees on 3 vertices, with the value of  $h(v)$  assigned to each vertex  $v$ . In this case, identity (1) says that  $(3+1)^2 = 3 + 3 + 4 + 3 + 3$ .



---

\*Research supported by the Post-doctoral Fellowship Program of Korea Research Foundation (KRF).

Postnikov derived this identity from the study of a combinatorial interpretation for mixed Eulerian numbers, which are coefficients of certain reparametrized *volume polynomials* which introduced in [3]. For more information, see [2, 3].

In the same talk, he also asked for a combinatorial proof of identity (1). Multiplying both sides of (1) by  $2^n$  and expanding the product in the right-hand side yields

$$2^n (n + 1)^{n-1} = \sum_{\mathbf{b}} n! \sum_{\alpha \subseteq V(\mathbf{b})} \prod_{v \in \alpha} \frac{1}{h(v)}. \quad (2)$$

Let  $\text{LHS}_n$  (resp.  $\text{RHS}_n$ ) denote the left-hand (resp. right-hand) side of (2).

The aim of this paper is to find a combinatorial proof of (2). In Section 2 we construct the sets  $\mathcal{F}_n^{\text{bi}}$  of labeled bicolored forests on  $[n]$  and  $\mathcal{D}_n$  of certain labeled bicolored binary trees, where the cardinalities equal  $\text{LHS}_n$  and  $\text{RHS}_n$ , respectively. In Section 3 we give a bijection between  $\mathcal{F}_n^{\text{bi}}$  and  $\mathcal{D}_n$ , which completes the bijective proof of (2). Finally, in Section 4, we present generalized formulas for the number of labeled  $k$ -ary trees, rooted labeled trees, and labeled plane trees.

## 2 Combinatorial objects for $\text{LHS}_n$ and $\text{RHS}_n$

From now on, unless specified, we consider trees to be *labeled* and *rooted*.

A *tree* on  $[n] := \{1, 2, \dots, n\}$  is an acyclic connected graph on the vertex set  $[n]$  such that one vertex, called the *root*, is distinguished. We denote by  $\mathcal{T}_n$  the set of trees on  $[n]$  and by  $\mathcal{T}_{n,i}$  the set of trees on  $[n]$  where vertex  $i$  is the root. A *forest* is a graph such that every connected component is a tree. Let  $\mathcal{F}_n$  denote the set of forests on  $[n]$ . There is a canonical bijection  $\gamma : \mathcal{T}_{n+1,n+1} \rightarrow \mathcal{F}_n$  such that  $\gamma(T)$  is the forest obtained from  $T$  by removing the vertex  $n + 1$  and letting each neighbor of  $n + 1$  be a root. A graph is called *bicolored* if each vertex is colored with the color  $\mathbf{b}$  (black) or  $\mathbf{w}$  (white). We denote by  $\mathcal{F}_n^{\text{bi}}$  the set of bicolored forests on  $[n]$ . From Cayley's formula [1] and the bijection  $\gamma$ , we have

$$|\mathcal{F}_n| = |\mathcal{T}_{n+1,n+1}| = (n + 1)^{n-1} \quad \text{and} \quad |\mathcal{F}_n^{\text{bi}}| = 2^n \cdot (n + 1)^{n-1}. \quad (3)$$

Thus  $\text{LHS}_n$  can be interpreted as the cardinality of  $\mathcal{F}_n^{\text{bi}}$ .

Let  $F$  be a forest and let  $i$  and  $j$  be vertices of  $F$ . We say that  $j$  is a *descendant* of  $i$  if  $i$  is contained in the path from  $j$  to the root of the component containing  $j$ . In particular, if  $i$  and  $j$  are joined by an edge of  $F$ , then  $j$  is called a *child* of  $i$ . Note that  $i$  is also a descendant of  $i$  itself. Let  $S(F, i)$  be the induced subtree of  $F$  on descendants of  $i$ , rooted at  $i$ . We call this tree the descendant subtree of  $F$  rooted at  $i$ . A vertex  $i$  is called *proper* if  $i$  is the smallest vertex in  $S(F, i)$ ; otherwise  $i$  is called *improper*. Let  $\text{pv}(F)$  denote the number of proper vertices in  $F$ .

A *plane tree* or *ordered tree* is a tree such that the children of each vertex are linearly ordered. We denote by  $\mathcal{P}_n$  the set of plane trees on  $[n]$  and by  $\mathcal{P}_{n,i}$  the set of plane trees on  $[n]$  where vertex  $i$  is the root. Define a *plane forest* on  $[n]$  to be a finite ordered sequence of non-empty plane trees  $(P_1, \dots, P_m)$  such that  $[n]$  is the disjoint union of the

sets  $V(P_r)$ ,  $1 \leq r \leq m$ . We denote by  $\mathcal{PF}_n$  the set of plane forests on  $[n]$  and by  $\mathcal{PF}_n^{\text{bi}}$  the set of bicolored plane forests on  $[n]$ . There is also a canonical bijection  $\bar{\gamma} : \mathcal{P}_{n+1, n+1} \rightarrow \mathcal{PF}_n$  such that  $\bar{\gamma}(P) = (S(P, j_1), \dots, S(P, j_m))$  where each vertex  $j_r$  is the  $r$ th child of  $n+1$  in  $P$ . It is well-known that the number of unlabeled plane trees on  $n+1$  vertices is given by the  $n$ th Catalan number  $C_n = \frac{1}{n+1} \binom{2n}{n}$  (see [4, ex. 6.19]). Thus we have

$$|\mathcal{PF}_n| = |\mathcal{P}_{n+1, n+1}| = n! \cdot C_n = 2n(2n-1) \cdots (n+2). \quad (4)$$

A *binary tree* is a tree in which each vertex has at most two children and each child of a vertex is designated as its left or right child. We denote by  $\mathcal{B}_n$  the set of binary trees on  $[n]$  and by  $\mathcal{B}_n^{\text{bi}}$  the set of bicolored binary trees on  $[n]$ .

For  $k \geq 2$ , a *k-ary tree* is a tree where each vertex has at most  $k$  children and each child of a vertex is designated as its first, second,  $\dots$ , or  $k$ th child. We denote by  $\mathcal{A}_n^k$  the set of  $k$ -ary trees on  $[n]$ . Clearly, we have that  $\mathcal{A}_n^2 = \mathcal{B}_n$ . Since the number of unlabeled  $k$ -ary trees on  $n$  vertices is given by  $\frac{1}{(k-1)n+1} \binom{kn}{n}$  (see [4, p. 172]), the cardinality of  $\mathcal{A}_n^k$  is as follows:

$$|\mathcal{A}_n^k| = n! \cdot \frac{1}{(k-1)n+1} \binom{kn}{n} = kn(kn-1) \cdots (kn-n+2).$$

Now we introduce a combinatorial interpretation of the number  $\text{RHS}_n$ . Let  $\mathfrak{b}$  be an unlabeled binary tree on  $n$  vertices and  $\omega : V(\mathfrak{b}) \rightarrow [n]$  be a bijection. Then the pair  $(\mathfrak{b}, \omega)$  is identified with a (labeled) binary tree on  $[n]$ . Let  $\Pi(\mathfrak{b}, \omega)$  be the set of vertices  $v$  in  $\mathfrak{b}$  such that  $v$  has no descendant  $v'$  satisfying  $\omega(v) > \omega(v')$ , i.e.,  $\omega(v)$  is proper.

Let  $\mathcal{D}_n$  be the set of bicolored binary trees on  $[n]$  such that each proper vertex is colored with  $\mathbf{b}$  or  $\mathbf{w}$  and each improper vertex is colored with  $\mathbf{b}$ .

**Lemma 1.** *The cardinality of  $\mathcal{D}_n$  is equal to  $\text{RHS}_n$ .*

*Proof.* Let  $\mathcal{D}'_n$  be the set defined as follows:

$$\mathcal{D}'_n := \{ (\mathfrak{b}, \omega, \alpha) \mid (\mathfrak{b}, \omega) \in \mathcal{B}_n \text{ and } \alpha \subseteq \Pi(\mathfrak{b}, \omega) \}.$$

There is a canonical bijection from  $\mathcal{D}'_n$  to  $\mathcal{D}_n$  as follows: Given  $(\mathfrak{b}, \omega, \alpha) \in \mathcal{D}'_n$ , if a vertex  $v$  of  $\mathfrak{b}$  is contained in  $\alpha$  then color  $v$  with  $\mathbf{w}$ ; otherwise color  $v$  with  $\mathbf{b}$ . Thus it suffices to show that the cardinality of  $\mathcal{D}'_n$  equals  $\text{RHS}_n$ .

Given an unlabeled binary tree  $\mathfrak{b}$  and a subset  $\alpha$  of  $V(\mathfrak{b})$ , let  $l(\mathfrak{b}, \alpha)$  be the number of labelings  $\omega$  satisfying  $\alpha \subseteq \Pi(\mathfrak{b}, \omega)$ . Then for each  $v \in \alpha$ , the label  $\omega(v)$  of  $v$  should be the smallest among the labels of the descendants of  $v$ . If we pick a labeling  $\omega$  uniformly at random, the probability that  $\omega(v)$  is the smallest among the labels of the descendants of  $v$  is  $1/h(v)$ . So the number of possible labelings  $\omega$  is  $n! / \prod_{v \in \alpha} h(v)$ . Thus we have

$$\begin{aligned} |\mathcal{D}'_n| &= \sum_{\mathfrak{b}} \sum_{\alpha \subseteq V(\mathfrak{b})} l(\mathfrak{b}, \alpha) \\ &= \sum_{\mathfrak{b}} \sum_{\alpha \subseteq V(\mathfrak{b})} n! \prod_{v \in \alpha} \frac{1}{h(v)}, \end{aligned}$$

which coincides with  $\text{RHS}_n$ . □




Figure 1: The map  $\mathbf{f}$ . (Right improper vertices are in italics.)

### 3 A bijection

In this section we construct a bijection between  $\mathcal{F}_n^{\text{bi}}$  and  $\mathcal{D}_n$ , which gives a bijective proof of (2).

Given a vertex  $v$  of a bicolored binary tree  $B$ , let  $L(B, v)$  (resp.  $R(B, v)$ ) be the descendant subtree of  $B$ , which is rooted at the left (resp. right) child of  $v$ . Note that  $L(B, v)$  and  $R(B, v)$  may be empty, but  $L(B, v)$  or  $R(B, v)$  is nonempty when  $v$  is improper. For any kind of tree  $T$ , let  $m(T)$  be the smallest vertex in  $T$ . By convention, we put  $m(\emptyset) = \infty$ . For an improper vertex  $v$  of  $B$ , if  $m(L(B, v)) > m(R(B, v))$ , then we say that  $v$  is *right improper*; otherwise *left improper*.

For a vertex  $v$  of  $B$ , define the *flip* on  $v$ , which will be denoted by  $f_v$ , by swapping  $L(B, v)$  and  $R(B, v)$  and changing the color of  $v$ . Note that the flip satisfies  $f_v \circ f_v = \text{id}$  and  $f_v \circ f_w = f_w \circ f_v$ . For a bicolored binary tree  $D$  in  $\mathcal{D}_n$ , let  $\mathbf{f}$  be the map defined by

$$\mathbf{f}(D) := (f_{v_1} \circ \dots \circ f_{v_k})(D),$$

where  $\{v_1, \dots, v_k\}$  is the set of right improper vertices in  $D$ . (See Figure 1.)

Let  $\mathcal{E}_n$  be the set of bicolored binary trees  $E$  on  $[n]$  such that every improper vertex  $v$  is left improper, i.e.,  $m(L(E, v)) < m(R(E, v))$ .

**Lemma 2.** *The map  $\mathbf{f}$  is a bijection from  $\mathcal{D}_n$  to  $\mathcal{E}_n$ .*

*Proof.* For a bicolored binary tree  $E$  in  $\mathcal{E}_n$ , let  $\mathbf{f}'$  be the map defined by  $\mathbf{f}'(E) := (f_{u_1} \circ \dots \circ f_{u_j})(E)$ , where  $\{u_1, \dots, u_j\}$  is the set of white-colored improper vertices in  $E$ . Then the map  $\mathbf{f}'$  is the inverse of  $\mathbf{f}$ .  $\square$

Let  $\mathcal{G}_n$  (resp.  $\mathcal{Q}_n$ ) be the set of bicolored trees (resp. bicolored plane trees) on  $[n+1]$  such that  $n+1$  is the root colored with  $\mathbf{b}$ . Note that the map  $\gamma$  (resp.  $\bar{\gamma}$ ) given at the beginning of Section 2 can be regarded as a bijection  $\gamma : \mathcal{G}_n \rightarrow \mathcal{F}_n^{\text{bi}}$  (resp.  $\bar{\gamma} : \mathcal{Q}_n \rightarrow \mathcal{PF}_n^{\text{bi}}$ ). For a vertex  $v$  of  $Q \in \mathcal{Q}_n$ , let  $(w_1, \dots, w_r)$  be the children of  $v$ , in order. Then for each  $i = 1, \dots, r-1$ , we say that  $w_{i+1}$  is the *right sibling* of  $w_i$ . The set  $\mathcal{G}_n$  can be viewed as a subset of  $\mathcal{Q}_n$  satisfying the following condition: Suppose that  $v$  is the right sibling of  $u$  in  $Q \in \mathcal{Q}_n$ . Then  $m(S(Q, u)) < m(S(Q, v))$  holds.

Recall that  $\mathcal{B}_n^{\text{bi}}$  denotes the set of bicolored binary trees on  $[n]$ . Clearly we have  $\mathcal{E}_n \subseteq \mathcal{B}_n^{\text{bi}}$ . Let  $\Phi$  be a bijection from  $\mathcal{B}_n^{\text{bi}}$  to  $\mathcal{Q}_n$ , which maps  $B$  to  $Q$  as follows:




Figure 2: The bijection  $\Phi$ .

1. The root of  $B$  is the first child of  $n + 1$  in  $Q$ .
2.  $v$  is the first child of  $u$  in  $Q$  iff  $v$  is a left child of  $u$  in  $B$ .
3.  $v$  is the right sibling of  $u$  in  $Q$  iff  $v$  is a right child of  $u$  in  $B$ .
4. The color of  $v$  in  $Q$  is the same as the color of  $v$  in  $B$ .

Note that here  $\Phi$  is essentially an extension of a well-known bijection, which is described in [5, p. 60], from binary trees to plane trees.

**Lemma 3.** *The restriction  $\phi$  of  $\Phi$  to  $\mathcal{E}_n$  is a bijection from  $\mathcal{E}_n$  to  $\mathcal{G}_n$ .*

*Proof.* For any improper vertex  $v$  of  $E \in \mathcal{E}_n$ , we have  $m(L(E, v)) < m(R(E, v))$ . This guarantees that  $m(S(G, v)) < m(S(G, w))$  in  $G = \Phi(E)$ , where  $w$  (if it exists) is the right sibling of  $v$  in  $G$ . Thus  $\Phi(E) \in \mathcal{G}_n$ , i.e.,  $\Phi(\mathcal{E}_n) \subseteq \mathcal{G}_n$ . Similarly we can show that  $\Phi^{-1}(\mathcal{G}_n) \subseteq \mathcal{E}_n$ . So we have  $\Phi(\mathcal{E}_n) = \mathcal{G}_n$ , which implies  $\phi$  is bijective.  $\square$

From Lemma 3, we easily get that  $\gamma \circ \phi$  is a bijection from  $\mathcal{E}_n$  to  $\mathcal{F}_n^{\text{bi}}$ . Combining this result with Lemma 2 yields the following consequence.

**Theorem 4.** *The map  $\gamma \circ \phi \circ \mathbf{f}$  is a bijection from  $\mathcal{D}_n$  to  $\mathcal{F}_n^{\text{bi}}$ .*

Figure 3 shows how the bijection in Theorem 4 maps a bicolored binary tree  $D$  in  $\mathcal{D}_{11}$  to a bicolored forest  $F$  on [11]. From equation (3) the cardinality of  $\mathcal{F}_n^{\text{bi}}$  equals  $\text{LHS}_n$  and from Lemma 1 the cardinality of  $\mathcal{D}_n$  equals  $\text{RHS}_n$ . Thus Theorem 4 is a combinatorial explanation of identity (2).

## 4 Generalized formulas

Theorem 4 implies the set  $\mathcal{D}_n$  of binary trees on  $[n]$  such that each proper vertex is colored with the color  $\mathbf{b}$  or  $\mathbf{w}$  and each improper vertex is colored with the color  $\mathbf{b}$  has cardinality  $|\mathcal{D}_n| = 2^n (n + 1)^{n-1}$ . In this section we give a generalization of this result.




Figure 3: The bijection from  $\mathcal{D}_n$  to  $\mathcal{F}_n^{\text{bi}}$ .

For  $n \geq 1$ , let  $a_{n,m}$  denote the number of  $k$ -ary trees on  $[n]$  with  $m$  proper vertices. By convention, we put  $a_{0,m} = \delta_{0,m}$ . Let

$$a_n(t) = \sum_{m \geq 0} a_{n,m} t^m = \sum_{T \in \mathcal{A}_n^k} t^{\text{pv}(T)},$$

where  $\text{pv}(T)$  is the number of proper vertices of  $T$ . It is clear that for a positive integer  $t$  the number  $a_n(t)$  is the number of  $k$ -ary trees on  $[n]$  such that each proper vertex is colored with the color  $\bar{1}, \bar{2}, \dots$ , or  $\bar{t}$  and each improper vertex has one color  $\bar{1}$ . Let  $A(x)$  be denote the exponential generating function for  $a_n(t)$ , i.e.,

$$A(x) = \sum_{n \geq 0} a_n(t) \frac{x^n}{n!}.$$

**Lemma 5.** *The generating function  $A = A(x)$  satisfies the following differential equation:*

$$A' = kx A^{k-1} A' + t A^k, \tag{5}$$

where the prime denotes the derivative with respect to  $x$ .

*Proof.* Let  $T$  be an  $k$ -ary tree on  $[n] \cup \{0\}$ . Delete all edges going from the root  $r$  of  $T$ . Then  $T$  is decomposed into  $T' = (r; T_1, \dots, T_k)$  where each  $T_i$  is a  $k$ -ary tree and  $[n] \cup \{0\}$  is the disjoint union of  $V(T_1), \dots, V(T_k)$  and  $\{r\}$ . Consider two cases: (i) For some  $1 \leq i \leq k$ ,  $T_i$  has the vertex  $0$ ; (ii)  $r = 0$ . Then we have

$$\begin{aligned} a_{n+1}(t) &= \sum_{i=1}^k \sum_{n_1 + \dots + n_k = n-1} \binom{n}{1, n_1, \dots, n_k} a_{n_1}(t) \cdots a_{n_{i+1}}(t) \cdots a_{n_k}(t) \\ &\quad + t \sum_{n_1 + \dots + n_k = n} \binom{n}{n_1, \dots, n_k} a_{n_1}(t) \cdots a_{n_k}(t). \end{aligned}$$

Multiplying both sides by  $x^n/n!$  and summing over  $n$  yields (5). □

To compute  $a_n(t)$  from (5) we need the following theorem.

**Theorem 6.** Fix positive integers  $a$  and  $b$ . Let  $u = 1 + \sum_{n=1}^{\infty} u_n x^n/n!$  be a formal power series in  $x$  satisfying

$$u' = a x u^b u' + t u^{b+1}. \quad (6)$$

Then  $u_n$  is given by

$$u_n = t \prod_{i=1}^{n-1} ((bi + 1)t + a(n - i)), \quad n \geq 1.$$

*Proof.* Adding  $(bt - a) x u^b u'$  to both sides of (6) yields

$$(1 + (bt - a) x u^b) u' = t (b x u^{b-1} u' + u^b) u.$$

Since  $(1 + (bt - a) x u^b)' = (bt - a) (b x u^{b-1} u' + u^b)$ , we have

$$(bt - a) \log u = t \log(1 + (bt - a) x u^b).$$

Taking the exponential of both sides and the substitutions  $x = y^b$  and  $yu(y^b) = \hat{u}(y)$  yield

$$\hat{u}(y) = y (1 + (bt - a) \hat{u}(y)^b)^{t/(bt-a)}. \quad (7)$$

Applying the Lagrange Inversion Formula (see [4, p. 38]) to (7) yields that

$$\begin{aligned} [y^{bn+1}] \hat{u}(y) &= \frac{1}{bn + 1} [y^{bn}] (1 + (bt - a) y^b)^{\frac{t(bn+1)}{bt-a}} \\ &= \frac{1}{bn + 1} (bt - a)^n \binom{\frac{t(bn+1)}{bt-a}}{n} \\ &= \frac{t}{n!} \prod_{i=1}^{n-1} (t(bn + 1) - (bt - a)i). \end{aligned}$$

Since  $u_n = n! [y^{bn+1}] \hat{u}(y)$ , we obtain the desired result.  $\square$

Since (5) is a special case of (6) ( $a = k$ ,  $b = k - 1$ ), we can deduce a formula for  $a_n(t)$ .

**Corollary 7 ( $k$ -ary trees).** For  $n \geq 1$ ,  $a_n(t)$  is given by

$$a_n(t) = t \prod_{i=1}^{n-1} ((ki - i + 1)t + k(n - i)). \quad (8)$$

Clearly, substituting  $t = 1$  in (8) yields the number of  $k$ -ary trees on  $[n]$ , i.e.,

$$a_n(1) = kn(kn - 1) \cdots (kn - n + 2) = |\mathcal{A}_n^k|.$$

For some values of  $k$ , we can get interesting results. In particular when  $k = 2$  we have

$$a_n(t) = t \prod_{i=1}^{n-1} ((i+1)t + 2(n-i)) \xrightarrow{t=2} 2^n(n+1)^{n-1},$$

so this is a generalization of  $|\mathcal{D}_n| = 2^n(n+1)^{n-1}$ , i.e., identity (2).

In fact Theorem 6 has more applications. For  $n \geq 1$ , let  $f_{n,m}$  denote the number of forests on  $[n]$  with  $m$  proper vertices and let  $p_{n,m}$  denote the number of plane forests on  $[n]$  with  $m$  proper vertices. Let

$$f_n(t) = \sum_{m \geq 1} f_{n,m} t^m \quad \text{and} \quad p_n(t) = \sum_{m \geq 1} p_{n,m} t^m.$$

Let  $F(x)$  and  $P(x)$  be the exponential generating function for  $f_n(t)$  and  $p_n(t)$ , respectively, i.e.,

$$F(x) = 1 + \sum_{n \geq 1} f_n(t) \frac{x^n}{n!} \quad \text{and} \quad P(x) = 1 + \sum_{n \geq 1} p_n(t) \frac{x^n}{n!}.$$

Similarly to Lemma 5, we can get two differential equations:

$$F' = x F F' + t F^2, \tag{9}$$

$$P' = x P^2 P' + t P^3. \tag{10}$$

Since (9) and (10) are special cases of (6) ( $a = b = 1$  and  $a = 1, b = 2$ , respectively), we have the following results.

**Corollary 8.** *Suppose  $f_n(t)$  and  $p_n(t)$  are defined as above. Then we have*

1. For  $n \geq 1$ ,  $f_n(t)$  is given by

$$f_n(t) = t \prod_{i=1}^{n-1} ((i+1)t + (n-i)). \tag{11}$$

2. For  $n \geq 1$ ,  $p_n(t)$  is given by

$$p_n(t) = t \prod_{i=1}^{n-1} ((2i+1)t + (n-i)). \tag{12}$$

Note that (11) and (12) are generalizations of (3) and (4), respectively. Moreover, from these formulas, we can easily get

$$\begin{aligned} \sum_{T \in \mathcal{T}_{n+1}} t^{\text{pv}(T)} &= t \prod_{i=0}^{n-1} ((i+1)t + (n-i)), \\ \sum_{P \in \mathcal{P}_{n+1}} t^{\text{pv}(P)} &= t \prod_{i=0}^{n-1} ((2i+1)t + (n-i)), \end{aligned}$$

which are generalizations of  $|\mathcal{T}_{n+1}| = (n+1)^n$  and  $|\mathcal{P}_{n+1}| = (n+1)! C_n$ .

**Remark.** In spite of the simple expressions, we have not proved (8), (11) and (12) in a bijective way. Also a direct combinatorial proof of Theorem 6 would be desirable.



## Acknowledgment

The author thanks Ira Gessel for his helpful advice and encouragement. The author also thanks the referees for useful comments and suggestions.

## References

- [1] A. Cayley, A theorem on trees, *Quart. J. Math.* **23** (1889), 376–378.
- [2] J. Pitman and R. P. Stanley, A polytope related to empirical distributions, plane trees, parking functions, and the associahedron, *Discrete Comput. Geom.* **27** (2002), no. 4, 603–634.
- [3] A. Postnikov, Permutohedra, associahedra, and beyond, *Retrospective in Combinatorics: Honoring Richard Stanley's 60th Birthday*, Massachusetts Institute of Technology, Cambridge, Massachusetts, June 22–26, 2004, slide available at: <http://www-math.mit.edu/~apost/talks/perm-slides.pdf>
- [4] R. P. Stanley, *Enumerative Combinatorics*, Vol. 2, Cambridge University Press, Cambridge, 1999.
- [5] D. Stanton and D. White, *Constructive Combinatorics*, Springer-Verlag, New York, 1986.