

**A Common Challenge in Older Adults:  
Classification, Overlap, and Therapy of Depression and Dementia**

Thomas Leyhe<sup>a\*</sup>, Charles F. Reynolds III<sup>b</sup>, Tobias Melcher<sup>a</sup>, Christoph Linnemann<sup>a</sup>,  
Stefan Klöppel<sup>c</sup>, Kaj Blennow<sup>d</sup>, Henrik Zetterberg<sup>d,e</sup>,  
Bruno Dubois<sup>f</sup>, Simone Lista<sup>g,h</sup>, and Harald Hampel<sup>f,h</sup>

<sup>a</sup>Center of Old Age Psychiatry, Psychiatric University Hospital, Basel, Switzerland

<sup>b</sup>Western Psychiatric Institute and Clinic, Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

<sup>c</sup>Department of Psychiatry and Psychotherapy, Center for Geriatric Medicine and Gerontology, Department of Neurology, University Medical Center Freiburg, Germany

<sup>d</sup>Institute of Neuroscience and Physiology, Department of Psychiatry and Neurochemistry, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

<sup>e</sup>University College London Institute of Neurology, Queen Square, London, UK

<sup>f</sup>Sorbonne Universités, Université Pierre et Marie Curie, Paris 06, Institut de la Mémoire et de la Maladie d'Alzheimer (IM2A) & Institut du Cerveau et de la Moelle épinière (ICM), Département de Neurologie, Hôpital de la Pitié-Salpêtrière, Paris, France

<sup>g</sup>IHU-A-ICM – Paris Institute of Translational Neurosciences, Pitié-Salpêtrière University Hospital, Paris, France

<sup>h</sup>AXA Research Fund & UPMC Chair, Paris, France

\*Corresponding author

Prof. Dr. Thomas Leyhe

Center of Old Age Psychiatry

Psychiatric University Hospital

Wilhelm Klein-Strasse 27

4012 Basel, Switzerland

Tel.: +41(0)61 325 53 53

Fax: +41(0)61 325 55 85

E-mail: thomas.leyhe@upkbs.ch

**Abstract**

Late-life depression is frequently associated with cognitive impairment. Depressive symptoms are often associated with or even precede a dementia syndrome. Moreover, depressive disorders increase the risk of persistence for mild cognitive impairment and dementia. Here, we present both the current state of evidence and future perspectives regarding the integration and value of clinical assessments, neuropsychological, neurochemical, and neuroimaging biomarkers for the etiological classification of the dementia versus the depression syndrome and for the prognosis of depression relating to dementia risk. Finally, we summarize the existing evidence for both pharmacotherapy and psychotherapy of depression in demented patients. There is an urgent need for large-scale collaborative research to elucidate the role and interplay of clinical and biological features in elderly individuals with depressive disorders who are at elevated risk for developing dementia. To overcome barriers for successful drug development, we propose the introduction of the precision medicine paradigm to this research field.

**Keywords:** Depression, dementia, differential diagnosis, classification, prognosis, biomarkers, neuroimaging, pharmacotherapy, psychological intervention, precision medicine.

## 1. Introduction

Dementia and depression are the most common psychiatric syndromes in older age. Although early identification of underlying causes and subsequent treatment are essential, the accurate differential diagnosis and discrimination (classification) remain clinically extremely challenging [1]. Late-life depression is frequently associated with cognitive impairment. In turn, dementia has been related to an increased risk of depressive symptoms.

Dementia, a major psychopathological syndrome, is traditionally diagnosed according to very slowly evolving operationalized criteria manuals, including the International Statistical Classification of Diseases and Related Health Problems (10<sup>th</sup> revision) (ICD-10) [2] and the Diagnostic and Statistical Manual of Mental Disorders (5<sup>th</sup> Edition) (DSM-5), that have only been updated after decade long intervals [3]. Contrary to expectations, the recently published DSM-5 did not yet integrate any biological information (biological markers) into the diagnostic armamentarium. The conservative primarily symptom-based, descriptive approach has been maintained for the neurocognitive disorders categories as well [4]. However, advanced expert consensus criteria have attempted classification approaches grounded on clinical, biological, and etiological factors.

The most common underlying causes of dementia in the elderly include Alzheimer's disease (AD) [5], vascular dementia [6], mixed dementia, dementia with Lewy bodies [7] dementia in Parkinson's disease [8] and frontotemporal dementia [9]. Notably, depressive symptoms have been reported in 30-50% of patients with AD dementia and are especially common at the prodromal stage [10]. Overt major depression can be diagnosed in >10% of AD patients, mostly during the early to moderately impaired stage [11], and in up to 50% of patients with vascular dementia [12–14]. Moreover, approximately 50% of patients with dementia with Lewy bodies show depressive symptoms [15].

Late-life depression is also generally diagnosed according to the ICD-10 [2] or DSM-5 criteria [3]. In addition to standard clinical assessment, psychometric indexes, such as the Geriatric Depression Scale [16], are frequently administered in the elderly. Late-life depression is common in patients with chronic physical illnesses. Age- and disease-related changes, including arteriosclerosis, chronic inflammation, hormonal and immune modifications, may affect the integrity of frontostriatal circuits as well as the amygdala and

the hippocampus, ultimately increasing the vulnerability to depression [17]. In addition, age-related psychosocial stressors including poor socio-economic status, disability, and social isolation are significant risk factors for depression [18]. Vegetative symptoms and impairments of executive functions, attention, information processing, psychomotor speed, and working memory are common. In particular, subcortical vascular changes play a major role in the pathophysiology of late-life depression [1] leading to the conceptualization of vascular depression, as defined by the results of magnetic resonance imaging (MRI) [19–22]. The risk of suicide is approximately 2-fold higher in the elderly, especially in older males, compared with the general population [23]. Overall, late-life depression has distinctive features that allow its differentiation from depressive disorders occurring at a younger age [24].

## **2. Increased dementia risk in depression**

To date, most of the published studies have focused on late-life depression – i.e., depression in subjects aged 60 years and older – and the risk of dementia, as well as the link between depression and dementia; in contrast, relatively few studies have been conducted in patients with earlier-life depression, i.e., in subjects younger than 60 years. Since (I) depression onset shows a high degree of variability, (II) both young adulthood and middle age are characterized by a high incidence of depression, and (III) dementia is characterized by a long asymptomatic preclinical phase, the examination of earlier-life depression might represent an opportunity to examine whether depression is a risk factor for dementia many years prior to the advent of clinical signs. It should be acknowledged, in any case, that the association between late-life depression and dementia might allow for a more in-depth analysis of depression as part of the prodromal stage of dementia. Therefore, a careful analysis of both earlier-life and late-life depression is necessary to attain complementary evidence [25].

The risk of developing dementia later in life increases 2-fold in presence of a positive history of depression at younger age. In presence of recurrent depressive disorders, a monotonic rise in the risk of dementia can be observed with an estimated 14% increase with each episode [26]. Although the available findings remain partly inconsistent, it can be assumed that late-life depression leads to a substantially increased risk of dementia. In this setting, depression can be a risk factor, a prodrome, or a consequence of dementia [25]. A recent study suggested that chronic depression during life may be aetiologically associated with an increased risk for

developing dementia, particularly vascular dementia, whereas depression occurring for the first time in late life may reflect a prodromal stage of dementia, in particular AD [27].

Currently, various mechanisms have been proposed to explicate the association between depression and dementia. First, there is significant evidence indicating that vascular disease is the primary link between depression and dementia, which is substantiated by the “vascular depression hypothesis” [28,29]. This pathophysiological theory states that cerebrovascular disease is a risk factor, a trigger, or a perpetuating factor for depressive syndromes in the elderly [18,30]. In particular, vascular changes in the frontostriatal brain regions have been linked to both depressive symptoms and cognitive impairment [31-33].

In addition, increased cortisone levels, a biochemical alteration frequently observed in depressive disorders [34] can lead to worsening hippocampal atrophy associated with cognitive deficits [31,35]. Notably, atrophy of the hippocampus is a well-characterized brain alteration detected both in AD [36] and in patients with depression [37,38].

In the conceptual framework of protein misfolding disorders, the presence of accumulating brain amyloid beta ( $A\beta$ ) plaques represents a key pathological hallmark of AD. It is well-known that both  $A\beta$  peptides and hyperphosphorylated tau proteins accumulate significantly in AD brains, leading to the formation of neuritic plaques and neurofibrillary tangles, respectively [39,40]. Interestingly, evidence indicates that depression might lead to an increased disequilibrium in terms of  $A\beta$  production and/or clearance. This effect is mediated by the depression-related stress response and the resulting hypercortisolemia, as well as the direct impact on  $A\beta$  processing, probably due to alterations at the level of the serotonergic system [41–44]. Notably, depressed AD patients have a higher burden of  $A\beta$  plaques and neurofibrillary tangles in the hippocampus than AD patients without depression [45–48].

During the last decade, chronic inflammatory processes have been also implicated in both depression and dementia [49–51]. On the one hand, a subtle and chronic brain inflammatory state, inducing cellular dysbalance, the activation of microglia and reactive astrocytes, resulting into increased concentrations of brain cytokines detected in depression and dementia, may result in a reduced modulation of anti-inflammatory and immunosuppressive mechanisms, increased acute-phase and proinflammatory regionally spreading alterations in the central nervous system (CNS), and, ultimately, in a non-linear progressive fashion

inducing neural network dysbalance, decompensation and breakdown, cognitive deficits and subsequent dementia [52]. Moreover, proinflammatory cytokines overexpression are supposed to interfere with the serotonin metabolism, thereby decreasing both synaptic plasticity and hippocampal neurogenesis [42,49].

Another mechanism that may link depression with dementia is represented by decreased levels of circulating neurotrophic factors, mainly the brain-derived neurotrophic factor (BDNF). BDNF modulates neuronal structure and function and plays an important role in synapse development and plasticity [53]. Reduced plasma BDNF levels have been observed both in animal models of depression [54] as well as in patients with depression [55,56] and AD [57,58].

Recent data have revealed that depressed patients face an accelerated cellular aging. In particular, those with the most severe and chronic major depressive disorder displayed the shortest telomere length and participants with remitted major depressive disorders had shorter telomere length than controls [59].

A more comprehensive systems-based neurobiological approach – larger genetic and epigenetic studies, analyses of gene and biomarker expression pattern, as well as innovative multi-modal structural, functional and metabolic neuroimaging – is needed to shed more light on the pathophysiology of late-life depression. To this aim, much can be learned both conceptually and methodologically from recent discoveries in the field of AD [60-62].

### **3. Common occurrence of depression and mild cognitive impairment as a risk condition**

A recent large cohort study conducted in Northern Manhattan (New York, NY, USA), including more than 2,000 individuals aged 65 or older, has shown that late-life depression is associated with both an increased risk of prevalent mild cognitive impairment (MCI), an established risk factor for the progression and occurrence of dementia, as well as overt dementia itself. Depression was also associated with an increased risk of incident dementia but not incident MCI. Notably, individuals presenting a concomitant depressive disorder and MCI showed a significantly increased risk of developing dementia, in particular vascular dementia, compared to those with MCI without depression [63].

A concomitant diagnosis of MCI has been reported in 25% to 50% of patients with late-life depression [64–66], compared to a 3% to 6% prevalence of MCI in community-based samples [11,67]. In addition, cognitive impairments emerging during a depressive episode can persist even after the remission of depressive symptoms [64,68]. The extent of cognitive impairment identified in elderly depressed patients prior to treatment seems to predict cognitive outcome after therapy [65,69-71].

Another recent study has demonstrated that cognitive impairment in late life depression might be related to greater cerebrovascular disease along with abnormalities in the immune–inflammatory control, cell survival, intracellular signalling, protein and lipid homeostasis, and clotting processes. As a result, individuals with late life depression and cognitive impairment seem to be more susceptible to accelerated brain aging processes at the cellular and molecular levels [72].

#### **4. Classification – differential diagnosis – of dementia and depression**

The concomitant occurrence of a depressive disorder and cognitive impairment should always be carefully investigated from a diagnostic viewpoint. Olin and colleagues [73,74] have proposed criteria to discriminate major depression and depression in AD. Accordingly, depression due to AD can be diagnosed when all criteria of dementia of Alzheimer type are fulfilled and three (or more) typical depressive symptoms have been detected during the same two-week period and represent a perturbation from previous physiological activity. At least one of the symptoms consists of either depressed mood or decreased positive affect or pleasure. Symptoms that are clearly due to a medical condition other than AD, or are the direct result of non-mood-related dementia symptoms (e.g. loss of weight due to difficulties with food intake) should not be included. Clinical signs are often less severe and pervasive than in major depression. Age of onset, rate and course of cognitive change, subjective memory complaints and typical sleep-wake cycle disturbances can aid in differential diagnosis.

##### *4.1. The emerging role of specific memory testing for the differential diagnosis*

AD is currently designated as a clinical entity typically characterized by a progressive amnesic syndrome with appearance of other cognitive, behavioural and neuropsychiatric

changes [75]. The episodic memory disorder, in the typical form of AD, shows a specific pattern which is the expression of hippocampal dysfunction and can be identified by tests including word list learning. This amnesic syndrome of the hippocampal type [76] is defined by (I) low free recall as for any brain related memory disorder; and (II) a low total recall performance, despite retrieval facilitation with cueing, due to hippocampal damage which affects the ability to store new information. Therefore, information cannot be retrieved even after facilitation procedures. Such a pattern demonstrates excellent specificity for AD [77].

In case of a pure depressive disorder, there is no genuine storage deficit; rather, attention difficulties that impair encoding or retrieval strategies can be observed [78]. Therefore, the differential diagnosis between AD and a pure depressive disorder can be improved by using a test paradigm that provides encoding specificity (with semantic cues) [79] and a retrieval facilitation (with the same cues), such as the free and cued selective reminding test (FCSRT) [80]. Regarding the memory domain, an improvement with repeated exposure and a normal recall with both control of encoding and retrieval cues are typically found in major depression, while a flat learning curve despite repeated exposure, a rapid forgetting, the inefficacy of cueing for recall and intrusions are typical for AD.

#### *4.2. The role of biomarkers for the differential diagnosis*

Whilst there are no specifically established fluid biomarkers for depression, three relevant biomarkers have been detected in the cerebrospinal fluid (CSF) for the key neuropathological alterations in AD. Actually, owing to its contiguity to the brain parenchyma and the free exchange with the brain extracellular space, the biochemical composition of CSF is able to provide information on the brain chemistry. The “core”, feasible biomarkers are: (I) total tau (T-tau, a marker reflecting cortical axonal degeneration), (II) phospho-tau (P-tau, a marker reflecting tau phosphorylation and AD-type neurofibrillary tangle pathology), both tau-related markers are the best indicators for disease progression, and (III) the 42 amino acid long form of A $\beta$  (A $\beta$ <sub>1-42</sub>, a marker of senile plaque pathology) [81]. This biomarker “triad” can be used to examine if patients with dementia-like depressive symptoms have AD pathological changes. Depression *per se* does not result in an AD-like CSF biomarker pattern, *i.e.*, increased T-tau and P-tau concentrations, and reduced levels of A $\beta$ <sub>1-42</sub> [82] although marginally decreased CSF A $\beta$ <sub>1-42</sub> concentrations have been reported [83]. A positive AD biomarker pattern is around 90% specific for AD neuropathology but does not exclude co-



morbidity in AD and depression. As demonstrated by Ewers and colleagues [84] in a large-scale multi-centric study, CSF A $\beta_{1-42}$  best discriminates AD dementia from frontotemporal dementia, non-neurodegenerative neurological diseases and depression, but shows significant overlap with other non-AD forms of dementia, possibly reflecting the underlying mixed pathologies. The combination of the three gold-standard markers provides added diagnostic value; in particular, at a fixed sensitivity of 85%, the specificity was >85% [84].

Notably, several other molecular changes have been examined in CSF in relation to depression and dementia disorders; nevertheless, none of them has yet shown potential clinical utility for differential diagnosis and classification. However, depression is characterized by slightly increased CSF concentrations of several pro-inflammatory cytokines, which are most often disregarded in dementia-causing diseases [85]. Pro-inflammatory cytokines enhance the activity of the indoleamine 2,3-dioxygenase enzyme that is the first rate-limiting enzyme of the tryptophan degradation pathway, the kynurenine pathway. Increased tryptophan degradation may induce serotonin depletion and depression, which is reflected by low CSF concentrations of kynurenic acid [86]. This alteration is not found in AD or dementia with Lewy bodies [87], which suggests that CSF kynurenic acid may be a promising biomarker to explore further for differential diagnosis. It is important to note that altered immune function is by no means specific to depression; any inflammatory biomarker evaluated as a potential tool to differentiate depression from dementia disorders has to be carefully examined in relation to diabetes, obesity, and a range of other disorders.

At present, there are no established blood (plasma/serum)-based biomarkers for AD or other dementia-causing diseases. However, mounting evidence suggests that depression is associated with a pro-inflammatory cytokine response in serum. Studies have shown elevated serum concentrations of interleukins (ILs) such as IL-1 and IL-6, the tumour necrosis factors alpha (TNF- $\alpha$ ), the C-reactive protein (CRP), and the Monocyte Chemoattractant Protein-1 (MCP-1) in depressed patients with mixed results for IL-8 [88]. There is no study so far assessing the potential diagnostic and prognostic utility of this molecular pattern.

#### *4.3. The role of imaging for the differential diagnosis*

To some extent, CSF biomarkers listed in the previous section can also be quantified using positron emission tomography (PET) of the brain. The use of tracers for tau is still restricted

to research studies but amyloid PET is now an established diagnostic tool in many specialised memory clinics. Cross-sectional studies report increased amyloid depositions in cognitively normal depressed older individuals [89]. It seems most likely that these are individuals with very early stage AD which manifests with depression as first symptom. Longitudinal studies or studies focusing on individuals with early onset depression could provide further insights. A recent study did not find any significant difference in terms of brain  $\beta$ -amyloid deposition (and gray matter volume) between elderly patients with remitted major depression and persistent MCI compared with elderly patients with remitted major depression and normal cognitive function [72].

Although patients with major depression may also show an altered regional brain glucose metabolism as measured using [ $^{18}\text{F}$ ] Fluorodeoxyglucose – PET (FDG-PET) [90] this is far less consistent and pronounced and affects different regions than hypometabolism observed in overt clinical dementia or MCI.

Besides PET, quantifying hippocampus atrophy and brain white matter lesion load are relevant to distinguish late-life depression from mild dementia. Although both can be assessed using computed tomography (CT), MRI is substantially more sensitive. Recurrent depressive episodes lead to hippocampus atrophy while a high number of white matter lesions is a common risk factor for late-onset depression and dementia, particularly of vascular origin [28, 29].

Studies comparing the extent of hippocampus atrophy in late-onset depression and AD typically report substantially more pronounced atrophy in AD [91]. This result is not surprising given the extent of neurodegeneration observed in manifest dementia. A current large meta analyses indicates areas such as cingulate cortex and precuneus may serve best to separate AD from late-life depression [92]. The degree of hippocampal atrophy may be similar between subjects with MCI and those with a high number of depressive episodes; however, such studies are missing.

On the other hand, the first appearance of depression in late life is often associated with increased white matter hyperintensities and clinically with lack of initiative and cognitive slowing. Consequently, global brain atrophy and white matter hyperintensities contribute to a diagnostic separation between late-life depression and healthy aging [93]. However,

separating patients with late-onset depression from those in the initial stages of vascular dementia can be difficult, particularly since both can manifest in the same individual, with accelerated cognitive decline and poorer response to antidepressants.

## **5. Pharmacotherapy of depression in dementia**

The design and evaluation of pharmacological interventions of depression in dementia is challenging for several methodical reasons. The large range of published prevalence rates from under 5 % to nearly 50 % for major depression in AD [94,95] foreshadows the problems of the correct detection and diagnosis of depression in dementia. It is a well-known fact that cognitive decline and dementia can progressively limit language skills and self-awareness of depressive symptoms. The difficult differential diagnosis of apathy and depression contributes to the complexity.

Clinical trials assessing the effects of pharmacotherapy of depression in dementia rest upon rating scales. The Montgomery-Asberg Depression Rating Scale (MADRS) [96] is not validated for demented patients [97,98]. The Hamilton Depression Rating Scale (HAM-D) [99] is another widely used scale that has not been validated in patients with severe dementia [100]. However, the Cornell Scale for Depression in Dementia (CSDD) [101] and the Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change (ADCS-CGIC) [102] are validated in demented patients [102,103].

As reviewed recently by Leong [104] several randomized placebo-controlled interventions reported negative outcomes for antidepressant efficacy in dementia: with sertraline (95 mg daily) and mirtazapine (30 mg daily) for 13 weeks, assessed by CSDD in 326 patients [105]; sertraline (93 mg daily) for 24 weeks, assessed by CSDD and a modified ADCS-CGIC in 117 patients [106]; venlafaxine (75 mg daily) for six weeks, assessed by MADRS in 31 patients [107] fluoxetine (maximum 40 mg daily) for six weeks, assessed by HAM-D in 41 patients [108]; sertraline (maximum 100 mg daily) for eight weeks, assessed by CSDD in 31 patients [109], imipramine (83 mg daily) for eight weeks, assessed by HAM-D in 61 patients [110]. In contrast, positive outcomes were reported for the following randomized placebo controlled trials: with sertraline (95 mg daily) for twelve weeks, assessed by CSDD and HAM-D in 44 patients [111]; moclobemide (maximum 400 mg daily) for six weeks, assessed by HAM-D in 511 demented patients [112]; clomipramine (maximum 100 mg daily) for six weeks, assessed

by HAM-D in 21 patients [113]; and citalopram (maximum 30 mg daily) for four weeks in 98 patients [114].

In summary, well-controlled studies, systematic reviews and meta-analyses [104,105,115,116] have shown no reliable and convincing efficacy of antidepressants in patients with dementia and co-occurring depressive disorders. Even the addition of a cholinesterase inhibitor in depressed patients showed only a small effect on concurrent cognitive impairment and on the conversion rate to dementia syndrome, but with increased risk of recurrence of depression [117].

Given the fact that pharmacological interventions based on serotonergic and noradrenergic abnormalities have shown disappointing results, the discussion of novel strategies becomes pertinent. The role of glutamatergic signalling, especially the dysfunction of N-methyl-D-aspartate (NMDA) receptor complex signalling, might be a promising overlap in the pathology of AD and major depression in late life. NMDA receptor antagonists have been demonstrated to feature anti-dementia as well as antidepressant potential [for review see 118]. These joint NMDA receptor-regulated signalling pathways in depression and dementia might represent a point of action in - so far treatment-resistant - depression in dementia.

The sobering results of pharmacological trials also highlight the need to consider alternative non-pharmacological treatments; for instance, psychological and behavioural interventions.

## **6. Psychological intervention for depression in dementia**

Systematic reviews provide a strong evidence base for psychotherapy as a first-line treatment in older depressive patients, with effect sizes comparable to those of antidepressant pharmacotherapy [119–122]. Moreover, the level of efficacy of psychotherapy in geriatric depression appears to be similar to that in younger patients [e.g. 119,123–125]. Evidence-based recommendations have been made especially for the cognitive-behavioural therapy (CBT), interpersonal therapy (IPT), and problem solving therapy (PST), while the literature provides no profound evidence that psychodynamic therapies were less efficient compared to other types of psychotherapy in the treatment of geriatric depression [c.f. 121,126–128]. On the whole, psychotherapy in late-life could evidently use all approaches established in younger patients, which, however, always needs to be reviewed before application for

necessary modifications to accommodate age-related mental changes and relevant contextual characteristics in the life of elderly people [129].

A necessity for technical or procedural adaptations is particularly appropriate in older depressed patients with dementia, who are easily overstrained by standard psychotherapeutic procedures. The progressive course of disease inevitably requires permanent adaptation of therapeutic actions to accommodate the advancing loss of mental resources. Concrete modifications of psychotherapeutic procedures for elderly patients have been described in the literature [e.g. 130-132] and essentially comprise an increase of structure and redundancy, i.e. rehearsal of important issues or messages, combined with a slowing of the conversational flow (to accommodate slowed cognitive processing) and a flexible organization of session lengths, which together should preserve the patients' involvement and comprehension in the therapeutic process. Moreover, and partially evident from the former issues, psychotherapy in the elderly needs an especially active and supportive (i.e. not neutral) therapeutic stance combined with a limited changing intent, i.e. to adaptively shift the emphasis of therapeutic interventions from changing (cognition or behaviour) to insight and/or acceptance, at least in some domains [cf.133]. More specific modifications concern the therapeutic content and include a stronger interdisciplinary focus, i.e. an intensified collaboration between the psychotherapist on the one hand and physicians, social workers and ambulatory services on the other, as well as a strengthened therapeutic focus and goal-orientation such as an emphasized selection, prioritization, and pursuit of therapeutic goals [cf.132–134].

The necessity for major adaptations to therapeutic procedures contradicts, at least in part, the assumption that psychotherapy in late-life has similar outcomes, i.e. produces an equal amount of “positive change” as in younger patients. This claim arguably does not apply to the “older old” and dementia patients [cf.125,135]. Moreover, the adaptability of specific psychotherapies could be outpaced by a level of mental decline. Accordingly, different psychotherapies can be construed to have different indication areas depending on their individual adaptive flexibility. Concretely, modified psychodynamic therapy has been described to be restricted to early stages of dementia or to MCI, whereas modified behavioural therapy may be still practicable and effective even in severely demented patients [136,137].

Despite the evident limitations and barriers, there is a constantly growing evidence-base for psychotherapy in dementia patients, data which evidently provide a valuable alternative or complementary treatment option to the pharmacotherapy of neuropsychiatric and particularly depressive symptoms. More specifically, a series of studies show that the adapted application of standard psychotherapies like CBT, PST, and IPT (compared to usual community or residential care) has a moderate effect on the level of depression in dementia and, hence, the potential to significantly improve the patients' psychological well-being [138,139; for review, see 135,140].

Of note, these results compare quite favourably with the limited effects of psychopharmacological interventions on depression in dementia patients, who in addition are more susceptible to adverse side effects and interaction effects than younger patients [cf.141]. As a restriction, however, participants of such studies mostly have mild dementia so that it remains unclear as to whether similar positive effects of nonpharmacological interventions can be achieved in more severe forms, too. Problem Adaptation Therapy (PATH) [142] is a relative novel psychotherapeutic intervention specifically designed for the treatment of depression in dementia. This therapy, which can be applied from mild cognitive deficits to moderate dementia, specifically aims to strengthen the patient's emotional regulation and for this purpose integrates a problem solving approach with environmental adaptation, compensatory strategies, and caregiver participation. In a recent efficacy evaluation study [143] PATH has been compared to supportive therapy for cognitively impaired patients and, thereby, has been shown to be superior in the reduction of depressive symptoms. The authors conclude that PATH may provide a valuable relief to a large group of dementia patients.

Other psychological interventions which have been evaluated in dementia patients are not specifically or uniquely aimed at the reduction of depressive symptoms, but follow more general or nonspecific goals like e.g. to increase social interaction, to stimulate memory, and/or to stabilize the sense of identity. These therapies have been developed specifically for the treatment of dementia patients, so-called "dementia-specific therapies", and, therefore, can be generally applied without adaptation, even in advanced disease stages. Even though dementia-specific therapies are not specifically aimed at the reduction of depressive symptoms, they still may improve the patients' mood and reduce the emotional stress related to dementia. Reminiscence therapy and Validation therapy, for instance, are "emotion-oriented" psychological interventions, which can be applied to treatment for MCI and all

stages of dementia. Reminiscence therapy [144] consists of different techniques to stimulate memories of personal history and thereby to reactivate life experiences, particularly positive ones, and their related mood states. Techniques to make these memories more meaningful are e.g. asking deepening questions, which suggest the importance of the life event or the allocation of historical materials like vintage photographs, documents or auditory records. Validation therapy [e.g.145] was originated and further developed by gerontologist Naomi Feil. This therapy focuses on the comprehension of emotional messages, which are assumed to lie behind the partly confused speech and behaviour of the dementia patient. According to the therapy rationale, dementia patients actively retreat to an inner reality, which is based on feelings rather than intellect, because they cannot tolerate their present reality. Based on this assumption, validation therapists prioritize the emotional content over the person's orientation to the present, so to speak, and accordingly validate (i.e. mirror and confirm) the communicative efforts of the patients rather to correct them in their confused expressions. This procedure is expected to reduce negative affect and consequent behavioural disturbances.

Unfortunately, dementia-specific therapies still have a rather weak or inconclusive evidence base [e.g.146–148], while, however, a lack of evidence can and should not be interpreted as proof of inefficacy. Validation therapy and Reminiscence therapy, among other dementia-specific psychotherapies, are clinically relevant and an inherent part of the dementia care practice [e.g.146].

## **7. Conclusions and future directions**

Both depression and cognitive decline impose a significant burden on public health. It has been shown that depressive syndromes have distinct features in the elderly. Close monitoring of cognitive disorders in elderly people showing depressive symptoms is of paramount importance and the presence of dementia should be excluded through extensive neuropsychological investigations. At present, although core, feasible AD neuroimaging modalities and neurochemical CSF biomarkers show evolving evidence, they have not yet been generalized world-wide and implemented in guidelines for clinical practice and are not routinely utilized in patients with late-life depression for differential diagnosis/classification and progression/prognosis.

Whereas psychological interventions have shown promising beneficial results, the clinical utility of approved antidepressant pharmacotherapies in patients with dementia and depression remains questionable. Actually, the understanding of the biological mechanisms underlying both dementia and depressive disorders as well as their genetically-biologically determined endophenotypes deserve further scrutiny and may pave the way for the development of novel diagnostic strategies and primary therapeutic targets for both dementia and depression. In this scenario, future studies should investigate clinical and biological features that may predict the development of persistent cognitive impairment in both earlier-life and late-life depression.

Preventive strategies against cognitive decline should take into account the patients' affective vulnerability. Moreover, much research effort should be devoted to the understanding of the neurobiology of depression with concomitant cognitive decline, with special reference to the involvement of networks and neurotransmitter systems that may differ from those affected in depression without cognitive decline.

It appears that there is an urgent need for large-scale collaborative research to shed more light on the role of clinical, neuroimaging, neuropsychological, genetic, neurochemical, and environmental characteristics of patients with late-life depression. In addition, studies on their significance for the prediction of cognitive decline and dementia development need to be performed.

In general, depressive syndromes present a highly heterogeneous spectrum of clinical phenotypes and are characterized by a still largely unclarified, intricate genetics and biology reflecting a partly overlapping variety of underlying aetiologies. In addition, pathological findings in different brain areas show a high degree of interindividual variability. In order to take these factors into adequate account, further research is needed into the phenomenology and neurobiology of depression in different ages of life, in patients with neurodegenerative and vascular diseases as well as in those showing pathologically mixed forms. In order to achieve this goal, a large amount of demographic, psychopathological, genetic, biomarker, and imaging data should be gathered in a standardized fashion. All these variables need to be analysed both cross-sectionally and longitudinally. These results would be invaluable for refining research hypotheses and designing further studies aimed at developing novel interventional strategies in preclinical models and, subsequently, in clinical cohorts.



However, despite decade-long research in the fields of neurobiology and neuropsychology of depression and dementia translated clinical progress seems still very limited. A more progressive conceptual paradigm shift inspired by other more advanced fields in healthcare – such as oncology – should transfertilize research in neuroscience providing better solutions also for these complex neuropsychiatric disorders. As a consequence of the prevailing conceptual traditionalism and stagnation in Neuropsychiatry, both dementia and depression are still considered, since approximately a century, as clinically descriptive categorical entities and continue to be classified according to descriptive operationalized criteria catalogues. The assumption that a few drug classes will fit all genetic and biological heterogeneous clinically defined target populations (syndromes and “disease” entities) is treacherous and unlikely to lead to success.

The precision medicine paradigm [149,150], adopted from oncology (e.g. in patients with adenocarcinomas and lung cancer) may offer a way out. To this aim, we need to accept the notion that patients can indeed be stratified by complex genetic, epigenetic, and genomic patterns, by subsequent molecular and cellular pathways that can be tracked and that serve as targets for intervention. These biological conditions emerge (often) long before first clinical symptoms arise, which suggests the exploration of “silent” preclinical stages before symptoms and syndromes emerge as late stage signs of underlying disease. Biological markers reflecting specific biochemistry and mechanisms of action as well as pathophysiology need to systematically be discovered and validated. In this regard, the systems biology paradigm [61,151] - representing an integrated investigation of interacting biomolecules within cells and organisms and allowing comprehensive exploratory biomarker studies – offers the adequate toolset for supporting precision medicine.

Actually, the evolving hypothesis-free exploratory paradigm of systems biology, also referred to as integrative biology or network biology [152,153] is an integrative interdisciplinary strategy exploiting advances in multimodal high-throughput technological platforms enabling the investigation of networks of biological pathways where elevated amounts of structurally/functionally different molecules are simultaneously explored over time at a system level (i.e., at the level of cells, tissues, organs, apparatuses, or even whole organism). This approach requires the comprehensive enumeration and quantification of biological processes, followed by efficient data analysis and integration, to allow the generation of hypotheses that need to be validated at a system level [151].

Technologies utilized in systems biology are the high-throughput screening methods typical of the omic sciences, namely: genomics/epigenomics, transcriptomics, proteomics/peptidomics, and metabolomics/lipidomics. Omics-based data provide a comprehensive picture on complex systems at molecular level, which can be used to systematically elucidate molecular mechanisms of living organisms. Thus, omics sciences can inform a more definite prediction of the risk of developing the disease, its progression, the severity of symptoms, personalized to a specific individual [61,62]. This information is necessary to tailor precisely both prevention strategies and therapeutic approaches to that subject as well as to provide suitable decisions regarding lifestyle and preventative treatments.

In order to develop targeted therapeutic strategies in the field of depression, depression in dementia and dementia in depression, it is mandatory to integrate cutting-edge multi-modal biomarker technologies and transfertilization from more matured translational research fields, such as the previously mentioned area of oncology. These progresses will lead to a radical paradigm shift: from the traditional reductionistic “one-drug-fits-all” approach to the concept of precision medicine enabling the identification of patients who would likely benefit the most from a treatment and suffer the least side effects [154]. For instance, the genetic makeup most likely associated with the molecular mechanisms of action of a drug can be revealed, thus increasing the probability of a patient’s response to the therapy. This helps to identify better matches for existing and novel drugs, with the aim of ensuring that the “right drug” is delivered to the “right patient” at the “right time”, with the highest possible success at lowest risk [155].

A precision medicine and, consequently, more comprehensive systems-based neurobiological approach – requiring larger genetic/genomic (e.g. whole genome and exome sequencing) and epigenomic studies, analyses of gene and biomarker expression patterns, as well as with innovative multi-modal structural, functional and metabolic neuroimaging – is therefore needed to gain more insights into late-life depression. To this aim, much can be learned both conceptually and methodologically from recent discoveries in the field of AD [61,62].

In summary, the aim of precision medicine is specifically targeting the molecular and clinical heterogeneity by identifying an individual’s comprehensive and specific pattern of risk factors, by defining the precise underlying molecular pathophysiologic processes, and, lastly,

by aiming to administer a preventive or therapeutic intervention specifically “customized”, i.e. adapted, to the identified molecular pattern of risk and disease processes

Finally, in the context of the precision medicine paradigm, the putative application of a multimodal model that may integrate into the pathophysiology of depression both imaging and biomarker data, which are well established for dementia, can ultimately reduce the burden of late-life depression on healthcare systems. Similarly, the burden would be decreased on the patients and their families, as a result of most appropriate prevention and clinical management. The availability of risk information assessed and specified for individuals with depression would facilitate clinicians’ informed decision-making on cognitive outcomes, basically targeting a modifiable risk factor (i.e., depressed mood) for dementia.

### **Acknowledgements**

Harald Hampel is supported by the AXA Research Fund, the Fondation Université Pierre et Marie Curie and the “*Fondation pour la Recherche sur Alzheimer*”, Paris, France. The research leading to these results has received funding from the program “*Investissements d’avenir*” ANR-10-IAIHU-06 (*Agence Nationale de la Recherche-10-IA Agence Institut Hospitalo-Universitaire-6*).

Charles F. Reynolds III is supported from P30 MH90333 and from the UPMC Endowed Chair in Geriatric Psychiatry. The corresponding author confirms that he had full access to all the data in the study and had final responsibility for the decision to submit for publication.

### **References**

- [1] Steffens DC, Potter GG. Geriatric depression and cognitive impairment. *Psychol Med* 2008;38 (2):163–75.
- [2] World Health Organisation. International statistical classification of diseases and related health problems. 10<sup>th</sup> rev. Genf, 2007.
- [3] American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 5<sup>th</sup> edition. Washington, DC, 2013.

- [4] Möller HJ, Bandelow B, Bauer M, Hampel H, Herpertz SC, Soyka M, et al. DSM-5 reviewed from different angles: goal attainment, rationality, use of evidence, consequences—part 2: bipolar disorders, schizophrenia spectrum disorders, anxiety disorders, obsessive-compulsive disorders, trauma- and stressor-related disorders, personality disorders, substance-related and addictive disorders, neurocognitive disorders. *Eur Arch Psychiatry Clin Neurosci* 2015;265(2):87-106.
- [5] McKhann GM., Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7(3):263–69.
- [6] Román GC, Tatemichi TK, Erkinjuntti T, Cummings JL, Masdeu JC, Garcia JH, et al. Vascular dementia: diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology* 1993;43(2):250–60.
- [7] McKeith IG, Dickson DW, Lowe J, Emre M, O'Brien JT, Feldman H, et al. Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology* 2005;65:1863–872.
- [8] Goetz CG, Emre M, Dubois B. Parkinson's disease dementia: definitions, guidelines, and research perspectives in diagnosis. *Ann Neurol* 2008;64,Suppl 2:S81-92.
- [9] Neary D, Snowden JS, Gustafson L, Passant U, Stuss D, Black S, et al. Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 1998;51(6):1546-1554.
- [10] Zuberko GS, Zuberko WN, McPherson S, Spoor E, Marin DB, Farlow MR, et al. A collaborative study of the emergence and clinical features of the major depressive syndrome of Alzheimer's disease. *Am J Psychiat* 2003;160:857–66.
- [11] Lopez OL, Becker JT, Sweet RA, Klunk W, Kaufer DI, Saxton J, et al. Psychiatric symptoms vary with the severity of dementia in probable Alzheimer's disease. *J Neuropsych Clin Neurosci* 2003;15:346–53.

- [12] Park JH, Lee SB, Lee TJ, Lee DY, Jhoo JH, Youn JC, et al. Depression in vascular Dementia is quantitatively and qualitatively different from depression in Alzheimer's Disease. *Dement Geriatr Cogn Disord* 2007;23:67–73.
- [13] Ballard C, Bannister C, Solis M, Oyeboode F, Wilcock G. The prevalence, associations and symptoms of depression amongst dementia sufferers. *J Affect Disord* 1996;36:135–144.
- [14] Ballard C, Neill D, O'Brien J, McKeith IG, Ince P, Perry R. Anxiety, depression and psychosis in vascular dementia: prevalence and associations. *J Affect Disord* 2000;59:97–106.
- [15] Ballard C, Holmes C, McKeith I, Neill D, O'Brien J, Cairns N, et al. Psychiatric morbidity in dementia with Lewy bodies: a prospective clinical and neuropathological comparative study with Alzheimer's disease. *Am J Psychiat* 1999;156:1039–45.
- [16] Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res* 1982-1983;17(1):37-49.
- [17] Sexton CE, Mackay CE, Ebmeier KP. A systematic review and meta-analysis of magnetic resonance imaging studies in late-life depression. *Am J Geriatr Psychiatry*. 2013;21(2):184-95.
- [18] Alexopoulos GS: Depression in the elderly. *Lancet* 2005;365:1961–970.
- [19] Krishnan KR, Taylor WD, McQuoid DR, MacFall JR, Payne ME, Provenzale JM, Steffens DC. Clinical characteristics of magnetic resonance imaging-defined subcortical ischemic depression. *Biol Psych* 2004;55:390–7.
- [20] Krishnan KRR, Hays JC, Blazer DG. MRI-defined vascular depression. *Am J Psychiat* 1997;154:497–501.

- [21] Steffens DC, Krishnan KRR Structural neuroimaging and mood disorders: recent findings, implications for classification, and future directions. *Biol Psych* 1998;43:705–12.
- [22] Pimontel MA, Reinlieb ME, Johnert LC, Garcon E, Sneed JR, Roose SP. The external validity of MRI-defined vascular depression. *Int J Geriatr Psychiatry* 2013;28(11):1189-96.
- [23] Miniño AM, Arias E, Kochanek KD, Murphy SL, Smith BL. Deaths: final data for 2000. *Natl Vital Stat Rep* 2002;50:1–119.
- [24] Naismith SL1, Norrie LM, Mowszowski L, Hickie IB. The neurobiology of depression in later-life: clinical, neuropsychological, neuroimaging and pathophysiological features. *Prog Neurobiol* 2012;98(1):99-143.
- [25] Byers AL, Yaffe K. Depression and risk of developing dementia. *Nat Rev Neurol*. 2011;7(6):323–31.
- [26] Naismith SL1, Norrie LM, Mowszowski L, Hickie IB. Recurrent depressive symptoms and the incidence of dementia and mild cognitive impairment. *Neurology* 2010;75:27-34.
- [27] Barnes DE1, Yaffe K, Byers AL, McCormick M, Schaefer C, Whitmer RA. Midlife vs late-life depressive symptoms and risk of dementia: Differential effects for Alzheimer Disease and vascular dementia. *Arch Gen Psychiatry* 2012;69(5):493-98.
- [28] Alexopoulos GS, Meyers BS, Young RC, Campbell S, Silbersweig D, Charlson M. “Vascular depression” hypothesis. *Arch. Gen. Psychiatry* 1997;54:915-922.
- [29] Krishnan KR, Hays JC, George LK, Blazer DG. Six-month outcomes for MRI-related vascular depression. *Depress Anxiety*. 1998;8(4):142-6.
- [30] Alexopoulos GS. Vascular disease, depression, and dementia. *J Am Geriatr Soc* 2003;51:1178-80.

- [31] Butters MA, Young JB, Lopez O, Aizenstein HJ, Mulsant BH, Reynolds CF 3<sup>rd</sup>, et al. Pathways linking late-life depression to persistent cognitive impairment and dementia. *Dialogues Clin Neurosci* 2008;10:345–57.
- [32] Alexopoulos GS: The vascular depression hypothesis: 10 years later. *Biol Psychiatry* 2006;60:1304–05.
- [33] Sheline YI, Price JL, Vaishnavi SN, Mintun MA, Barch DM, Epstein AA, et al. Regional white matter hyperintensity burden in automated segmentation distinguishes late-life depressed subjects from comparison subjects matched for vascular risk factors. *Am J Psychiatry* 2008;165:524–32.
- [34] Wolkowitz OM1, Epel ES, Reus VI, Mellon SH. Depression gets old fast: do stress and depression accelerate cell aging? *Depress Anxiety* 2010;27:327–38.
- [35] Sierksma AS, van den Hove DL, Steinbusch HW, Prickaerts J. Major depression, cognitive dysfunction and Alzheimer's disease: is there a link? *Eur J Pharmacol* 2010;626:72–82.
- [36] van de Pol LA, Hensel A, Barkhof F, Gertz HJ, Scheltens P, van der Flier WM. Hippocampal atrophy in Alzheimer disease: age matters. *Neurology* 2006;66:236-38.
- [37] Videbech P, Ravnkilde B. Hippocampal volume and depression: a meta-analysis of MRI studies. *Am J Psychiatry* 2004;161:1957-66.
- [38] Colla M, Kronenberg G, Deuschle M, Meichel K, Hagen T, Bohrer M, Heuser I. Hippocampal volume reduction and HPA-system activity in major depression. *J Psychiatr Res* 2007;41:553-60.
- [39] Blennow K, de Leon MJ, Zetterberg H. Alzheimer's disease. *Lancet*. 2006;368(9533):387-403.

- [40] Blennow K, Hampel H, Weiner M, Zetterberg H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* 2010;6(3):131-44.
- [41] Caraci F1, Copani A, Nicoletti F, Drago F. Depression and Alzheimer's disease: neurobiological links and common pharmacological targets. *Eur J Pharmacol* 2010;626:64–71.
- [42] Cho S, Hu Y. Activation of 5-HT<sub>4</sub> receptors inhibits secretion of beta-amyloid peptides and increases neuronal survival. *Exp Neurol* 2007;203:274–78.
- [43] Lezoualc'h F. 5-HT<sub>4</sub> receptor and Alzheimer's disease: the amyloid connection. *Exp Neurol* 2007;205:325–29.
- [44] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002;297:353–56.
- [45] Leinonen V, Koivisto AM, Savolainen S, Rummukainen J, Tamminen JN, Tillgren T, et al. Amyloid and tau proteins in cortical brain biopsy and Alzheimer's disease. *Ann Neurol* 2010;68:446–53.
- [46] Maes M, Yirmiya R, Norberg J, Brene S, Hibbeln J, Perini G, et al. The inflammatory & neurodegenerative (I&ND) hypothesis of depression: leads for future research and new drug developments in depression. *Metab Brain Dis* 2009;24:27–53.
- [47] Rapp MA, Schnaider-Beerl M, Purohit DP, Perl DP, Haroutunian V, Sano M.. Increased neurofibrillary tangles in patients with Alzheimer disease with comorbid depression. *Am J Geriatr Psychiatry* 2008; 16: 168-74.
- [48] Rapp MA<sup>1</sup>, Schnaider-Beerl M, Grossman HT, et al. Increased hippocampal plaques and tangles in patients with Alzheimer disease with a lifetime history of major depression. *Arch. Gen. Psychiatry* 2006;63:161-67.
- [49] Leonard BE: Inflammation, depression and dementia: are they connected? *Neurochem Res* 2007;32:1749–56.



- [50] Rojo LE, Fernández JA, Maccioni AA, Jimenez JM, Maccioni RB. Neuroinflammation: implications for the pathogenesis and molecular diagnosis of Alzheimer's disease. *Arch Med Res* 2008;39:1–16.
- [51] Sorrells SF, Sapolsky RM.: An inflammatory review of glucocorticoid actions in the CNS. *Brain Behav Immun* 2007;21:259–72.
- [52] Yaffe K, Lindquist K, Penninx BW, Simonsick EM, Pahor M, Kritchevsky S, et al. Inflammatory markers and cognition in well-functioning African-American and white elders. *Neurology* 2003;61:76–80.
- [53] Fumagalli F, Molteni R, Calabrese F, Maj PF, Racagni G, Riva MA. Neurotrophic factors in neurodegenerative disorders: potential for therapy. *CNS Drugs* 2008;22:1005–19.
- [54] Krishnan V, Nestler EJ.: The molecular neurobiology of depression. *Nature* 2008;455:894–902.
- [55] Karege F1, Vaudan G, Schwald M, Perroud N, La Harpe R. Neurotrophin levels in post-mortem brains of suicide victims and the effects of antemortem diagnosis and psychotropic drugs. *Brain Res Mol Brain Res* 2005;136:29–37.
- [56] Angelucci F, Brenè S, Mathé AA. BDNF in schizophrenia, depression and corresponding animal models. *Mol Psychiatry* 2005;10:345–52.
- [57] Murer MG, Yan Q, Raisman-Vozari R. Brain-derived neurotrophic factor in the control human brain, and in Alzheimer's disease and Parkinson's disease. *Prog Neurobiol* 2001;63:71–124.
- [58] Cotman CW: The role of neurotrophins in brain aging: a perspective in honor of Regino Perez-Polo. *Neurochem Res* 2005;30:877–81.

- [59] Verhoeven JE, Révész D, Epel ES, Lin J, Wolkowitz OM, Penninx BWJH: Major depressive disorder and accelerated cellular aging: results from a large psychiatric cohort study. *Mol Psychiatry* 2014;19:895–901.
- [60] Hampel H, Frank R, Broich K, Teipel SJ, Katz RG, Hardy J, et al. Biomarkers for Alzheimer's disease: academic, industry and regulatory perspectives. *Nat Rev Drug Discov* 2010;9(7):560-574.
- [61] Hampel H, Lista S, Khachaturian ZS. Development of biomarkers to chart all Alzheimer's disease stages: the royal road to cutting the therapeutic Gordian Knot. *Alzheimers Dement* 2012;8(4):312-36.
- [62] Lista S, Khachaturian ZS, Rujescu D, Garaci F, Dubois B, Hampel H. Application of systems theory in longitudinal studies on the origin and progression of Alzheimer's Disease. *Methods Mol Biol* 2016;1303: 9-67.
- [63] Richard E, Reitz C, Honig LH, Schupf N, Tang MX, Manly JJ, et al. Late-life depression, mild cognitive impairment, and dementia. *JAMA Neurol* 2013;70(3):374-82.
- [64] Lopez OL, Jagust WJ, DeKosky ST, Becker JT, Fitzpatrick A, Dulberg C, et al. Prevalence and classification of mild cognitive impairment in the Cardiovascular Health Study Cognition Study: Part 1. *Arch Neurol* 2003;60:1385–89.
- [65] Adler G, Chwalek K, Jajcevic. A: Six-month course of mild cognitive impairment and affective symptoms in late-life depression. *Eur Psychiat* 2004;19:502–5.
- [66] Lee JS, Potter GG, Wagner HR, Welsh-Bohmer KA, Steffens DC. Persistent mild cognitive impairment in geriatric depression. *Int Psychoger* 2007;19:125–35.
- [67] Ganguli M, Dodge HH, Shen C, DeKosky ST. Mild cognitive impairment, amnesic type: an epidemiologic study. *Neurology* 2004;63:115–21.

- [68] Reischies FM, Neu P. Comorbidity of mild cognitive disorder and depression – a neuropsychological analysis. *Eur Arch Psy Clin N* 2000;50:186–93.
- [69] Butters MA, Becker JT, Nebes RD, Zmuda MD, Mulsant BH, Pollock BG, Reynolds CF 3rd. Changes in cognitive functioning following treatment of late-life depression. *Am J Psychiat* 2000;157: 949–54.
- [70] Nebes RD, Pollock BG, Houck PR, Butters MA, Mulsant BH, Zmuda MD, Reynolds CF 3rd. Persistence of cognitive impairment in geriatric patients following antidepressant treatment: a randomized, double-blind clinical trial with nortriptyline and paroxetine. *J Psychiat Res* 2003;37:99–108.
- [71] Murphy CF, Alexopoulos GS: Longitudinal association of initiation/perseveration and severity of geriatric depression. *Am J Geriat Psychiat* 2004;12:50–6.
- [72] Diniz BS, Sibille E, Ding Y, Tseng G, Tseng G, Aizenstein HJ, Lotrich F, et al. Plasma biosignature and brain pathology related to persistent cognitive impairment in late-life depression. *Mol Psychiatry* 2015;20:594–601.
- [73] Olin JT, Katz IR, Meyers BS, Schneider LS, Lebowitz BD. Provisional diagnostic criteria for depression of Alzheimer disease: rationale and background. *Am J Geriat Psychiat* 2002;10:129–41.
- [74] Olin JT, Schneider LS, Katz IR, Meyers BS, Alexopoulos GS, Breitner JC et al. Provisional diagnostic criteria for depression of Alzheimer disease. *Am J Geriat Psychiat* 2002;10:125–28.
- [75] Dubois B, Feldman HH, Jacova C, Dekosky ST, Barberger-Gateau P, Cummings J, et al.. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol* 2007;6(8):734-46.
- [76] Dubois B, Albert ML. Amnesic MCI or prodromal Alzheimer's disease? *Lancet Neurol* 2004;3:246–48.

- [77] Tounsi H, Deweer B, Ergis AM, Van der Linden M, Pillon B, Michon A, Dubois B. Sensitivity to semantic cuing: an index of episodic memory dysfunction in early Alzheimer disease. *Alzheimer Dis Assoc Disord* 1999;13:38–46.
- [78] Fossati P, Harvey PO, Le BG, Ergis AM, Jouvent R, Allilaire JF. Verbal memory performance of patients with a first depressive episode and patients with unipolar and bipolar recurrent depression. *J Psychiatr Res* 2004;38:137–44.
- [79] Grober E, Buschke H. Genuine memory deficit in dementia. *Dev Neuropsychol* 2006;3:13–36.
- [80] Buschke H, Sliwinski MJ, Kuslansky G, Lipton RB. Diagnosis of early dementia by the Double Memory Test: encoding specificity improves diagnostic sensitivity and specificity. *Neurology* 1997;48:989–97.
- [81] Rosén C, Hansson O, Blennow K, Zetterberg H. Fluid biomarkers in Alzheimer's disease - current concepts. *Mol Neurodegener* 2013;8:20. doi:10.1186/1750-1326-8-20.
- [82] Gudmundsson P, Skoog I, Waern M, Blennow K, Zetterberg H, Rosengren L, Gustafson D. Is there a CSF biomarker profile related to depression in elderly women? *Psychiatry Res* 2010;176(2-3):174-8.
- [83] Nascimento KK, Silva KP, Malloy-Diniz LF, Butters MA, Diniz BS. Plasma and cerebrospinal fluid amyloid- $\beta$  levels in late-life depression: A systematic review and meta-analysis. *J Psychiatr Res* 2015;69:35-41.
- [84] Ewers M, Mattsson N, Minthon L, Molinuevo JL, Antonell A, Popp J, et al. CSF biomarkers for the differential diagnosis of Alzheimer's disease: A large-scale international multicenter study. *Alzheimers Dement* 2015;11(11):1306-15.
- [85] Raedler TJ. Inflammatory mechanisms in major depressive disorder. *Curr Opin Psychiatry* 2011;24(6):19-25.

- [86] Bay-Richter C, Linderholm KR, Lim CK, Samuelsson M, Träskman-Bendz L, Guillemin GJ, et al. A role for inflammatory metabolites as modulators of the glutamate N-methyl-D-aspartate receptor in depression and suicidality. *Brain Behav Immun* 2015;43:110-7.
- [87] Wennström M, Nielsen HM, Orhan F, Londos E, Minthon L, Erhardt S. Kynurenic Acid levels in cerebrospinal fluid from patients with Alzheimer's disease or dementia with lewy bodies. *Int J Tryptophan Res* 2014;28(7):1-7.
- [88] Young JJ, Bruno D, Pomara N. A review of the relationship between proinflammatory cytokines and major depressive disorder. *J Affect Disord* 2014;169:15-20.
- [89] Harrington KD, Lim YY, Gould E, Maruff P. Amyloid-beta and depression in healthy older adults: a systematic review. *Aust N Z J Psychiatry* 2015;49(1):36-46.
- [90] Su L, Cai Y, Xu Y, Dutt A, Shi S, Bramon E. Cerebral metabolism in major depressive disorder: a voxel-based meta-analysis of positron emission tomography studies. *BMC Psychiatry* 2014;14:321. doi:10.1186/s12888-014-0321-9.
- [91] Schmaal L, Veltman DJ, van Erp TG, Sämann PG, Frodl T, Jahanshad N, et al. Subcortical brain alterations in major depressive disorder: findings from the ENIGMA Major Depressive Disorder working group. *Mol Psychiatry* 2016;21(6):724-5.
- [92] Boccia M, Acierno M, Piccardi L. Neuroanatomy of Alzheimer's Disease and late-life depression: A coordinate-based meta-analysis of MRI studies. *J Alzheimers Dis* 2015;46(4):963-70.
- [93] Shimoda K, Kimura M, Yokota M, Okubo Y. Comparison of regional gray matter volume abnormalities in Alzheimer's disease and late life depression with hippocampal atrophy using VSRAD analysis: a voxel-based morphometry study. *Psychiatry Res* 2015;232(1):71-5.

- [94] Weiner MF, Doody RS, Sairam R, Foster B, Liao TY. Prevalence and incidence of major depressive disorder in Alzheimer's disease: findings from two databases. *Dement Geriatr Cogn Disord* 2002;13:8-12.
- [95] Weiner MF, Edland SD, Luszczynska H. Prevalence and incidence of major depression in Alzheimer's disease. *Am J Psychiatry* 1994;151(7):1006-9.
- [96] Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry* 1979;134:382-89.
- [97] Conn D, Thorpe L. Assessment of behavioural and psychological symptoms associated with dementia. *Can J Neurol Sci* 2007;34 Suppl 1:S67-71.
- [98] Holroyd S, Clayton AH: Measuring Depression in the Elderly: Which Scale is Best? *Medscape General Medicine* 2000;2(4).
- [99] Hamilton M: A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56-62.
- [100] Lichtenberg PA, Marcopulos BA, Steiner DA, Tabscott JA. Comparison of the Hamilton Depression Rating Scale and the Geriatric Depression Scale: detection of depression in dementia patients. *Psychol Rep* 1992;70:515-21.
- [101] Alexopoulos GS, Abrams RC, Young RC, Shamoian CA. Cornell Scale for Depression in Dementia. *Biol Psychiatry* 1988;23:271-84.
- [102] Schneider LS, Olin JT, Doody RS, et al.: Validity and reliability of the Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997; 11 Suppl 2: S22-32.
- [103] Kørner A, Lauritzen L, Abelskov K, Clark CM, Morris JC, Reisberg B, et al. The Geriatric Depression Scale and the Cornell Scale for Depression in Dementia. A validity study. *Nord J Psychiatry* 2006;60:360-4.

- [104] Leong C. Antidepressants for depression in patients with dementia: a review of the literature. *Consult Pharm* 2014;29:254-63.
- [105] Banerjee S, Hellier J, Dewey M, Romeo R, Ballard C, Baldwin R, et al. Sertraline or mirtazapine for depression in dementia (HTA-SADD): a randomised, multicentre, double-blind, placebo-controlled trial. *Lancet*. 2011 30;378(9789):403-11.
- [106] Rosenberg PB, Drye LT, Martin BK, Frangakis C, Mintzer JE, Weintraub D, et al. Sertraline for the treatment of depression in Alzheimer's disease. *Am J Geriatr Psychiatry* 2010;18:136-45.
- [107] De Vasconcelos Cunha UG, Lopes Rocha F, Alves Valle E, de Souza Neto JJ, Mendes Brega R, et al.: A placebo-controlled double-blind randomized study of venlafaxine in the treatment of depression in dementia. *Dement Geriatr Cogn Disord* 2007;24:36-41.
- [108] Petracca GM, Chemerinski E, Starkstein SE. A double-blind placebo-controlled study of fluoxetine in depressed patients with Alzheimer's disease. *Int Psychogeriatr* 2001;13:233-40.
- [109] Magai C, Kennedy G, Cohen CI, Gomberg D. A controlled clinical trial of sertraline in the treatment of depression in nursing home patients with late-stage Alzheimer's disease. *Am J Geriatr Psych* 2000;8:66-74.
- [110] Reifler BV, Teri L, Raskind M, Veith R, Barnes R, White E, McLean P. Double-blind trial of imipramine in Alzheimer's disease patients with and without dementia. *Am J Psychiatry* 1989;146:45-9.
- [111] Lyketsos CG, DelCampo L, Steinberg M, Miles Q, Steele CD, Munro C, et al. Treating depression in Alzheimer disease: efficacy and safety of sertraline therapy, and the benefits of depression reduction: the DIADS. *Arch Gen Psychiatry* 2003;60:737-46.

- [112] Roth M, Mountjoy CQ, Amrein R. Moclobemide in elderly patients with cognitive decline and depression: an international double-blind placebo-controlled trial. *Br J Psychiatry* 1996;168:149-57.
- [113] Petracca G, Tesón A, Chemerinski E, Leiguarda R, Starkstein SE. A double-blind placebo-controlled study of clomipramine in depressed patients with Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1996;8:270-5.
- [114] Nyth AL, Gottfries CG: The clinical efficacy of citalopram in the measurement of emotional disturbances in dementia disorders. *Br. J. Psychiatry* 1990;157:894-901.
- [115] Nelson JC, Devanand DP: A systematic review and meta-analysis of placebo-controlled antidepressant studies in people with depression and dementia. *J Am Geriatr Soc* 2011;59:577-85.
- [116] Sepehry AA, Lee PE, Hsiung GY, Beattie BL, Jacova C. Effect of selective serotonin reuptake inhibitors in Alzheimer's disease with comorbid depression: a meta-analysis of depression and cognitive outcomes. *Drugs Aging* 2012;29:793-806.
- [117] Reynolds CF 3rd, Butters MA, Lopez O, Pollock BG, Dew MA, Mulsant BH, et al. Maintenance treatment of depression in old age: a randomized, double-blind, placebo-controlled evaluation of the efficacy and safety of donepezil combined with antidepressant pharmacotherapy. *Arch Gen Psychiatry* 2011;68:51-60.
- [118] Khundakar AA, Thomas AJ: Neuropathology of depression in Alzheimer's Disease: Current knowledge and the potential for new treatments. *J Alzheimers Dis* 2015;44:27-41.
- [119] Cuijpers P, Sijbrandij M, Sander L, Koole SL, Andersson G, Beekman AT, Reynolds CF. Adding psychotherapy to antidepressant medication in depression and anxiety disorders: a meta-analysis. *World Psychiatry* 2014;13(1):56-67.



- [120] Pincus M, Duberstein PR, Lyness JM. Treatments for later-life depressive conditions: a meta-analytic comparison of pharmacotherapy and psychotherapy. *Am J Psychiatry* 2006;163(9):1493-501.
- [121] Wilson KCM, Mottram PG, Vassilas CA. Psychotherapeutic treatments for older depressed people. *Cochrane Database Syst Rev* 2008;(1):CD004853. doi:10.1002/14651858.CD004853.pub2.
- [122] Peng XD, Huang CQ, Chen LJ, Lu ZC. Cognitive behavioural therapy and reminiscence techniques for the treatment of depression in the elderly: a systematic review. *J Int Med Res* 2009;37(4):975-82.
- [123] Robinson LA, Berman JS, Neimeyer RA. Psychotherapy for the treatment of depression: a comprehensive review of controlled outcome research. *Psychol Bull* 1990;108(1):30-49.
- [124] Cuijpers P, van Straten A, Andersson G, van Oppen P. Psychotherapy for depression in adults: a meta-analysis of comparative outcome studies. *J Consult Clin Psychol* 2008;76(6):909-22.
- [125] Cuijpers P, van Straten A, Smit F, Andersson G. Is psychotherapy for depression equally effective in younger and older adults? A meta-regression analysis. *Int Psychogeriatr* 2009;21(1):16-24.
- [126] Hollon SD, Jarrett RB, Nierenberg AA, Thase ME, Trivedi M, Rush AJ: Psychotherapy and medication in the treatment of adult and geriatric depression: which monotherapy or combined treatment? *J Clin Psychiatry* 2005;66(4):455-68.
- [127] Pincus M, Duberstein PR, Lyness JM: Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: a meta-analysis. *Aging Ment Health* 2007;11(6):645-57.

- [128] Kirkham JG, Choi N, Seitz DP. Meta-analysis of problem solving therapy for the treatment of major depressive disorder in older adults. *Int J Geriatr Psychiatry* 2016;31(5):526-35.
- [129] Laidlaw K: An empirical review of cognitive therapy for late life depression: does research evidence suggest adaptations are necessary for cognitive therapy with older adults? *Clin Psychol Psychother* 2001;8(1):1–14.
- [130] Qualls SH, Knight BG, eds. *Psychotherapy for depression in older adults* Hoboken, N.J., Wiley Series in Clinical Geropsychology. John Wiley & Sons, Inc., 2006.
- [131] Knight B., Poon CYM. Contextual adult lifespan theory for adapting psychotherapy with older adults. *J Ration Emot Cogn Behav Ther* 2008;26:232–49.
- [132] Laidlaw K, McAlpine S. Cognitive-behaviour therapy: How is it different with older people? *J Ration Emot Cogn Behav Ther* 2008;26(4):250-262.
- [133] Frank E, Frank N, Cornes C, Imber SD, Miller MD, Morris SM, Reynolds CF. Interpersonal psychotherapy in the treatment of late-life depression. In: Klerman GL, Weissman MM, eds. *New applications of interpersonal psychotherapy*. Washington, DC: American Psychiatric Press; 1993. 167–98.
- [134] Chand SP, Grossberg GT. How to adapt cognitive-behavioral therapy for older adults. *Curr Psychiatr* 2013;12(3):10-5.
- [135] Orgeta V, Qazi A, Spector AE, Orrell M. Psychological treatments for depression and anxiety in dementia and mild cognitive impairment. *Cochrane Database Syst Rev* 2014;22(1):CD009125. doi:10.1002/14651858.CD009125.pub2.
- [136] Wilkins VM et al.: Late-life depression with comorbid cognitive impairment and disability: nonpharmacological interventions. *Clin Interv Aging*. 2010;5:323–31.
- [137] Hirsch R.: Psychotherapy with people afflicted with dementia. *Psychotherapie* 2009; 14(2):317-31.

- [138] Burgener SC, Yang Y, Gilbert R, Marsh-Yant S. The effects of a multimodal intervention on outcomes of persons with early-stage dementia. *Am J Alzheimers Dis Other Demen* 2008;23(4):382-94.
- [139] Burns A, Guthrie E, Marino-Francis F, et al. Brief psychotherapy in Alzheimer's disease: randomised controlled trial. *Br J Psychiatry* 2005;187:143-7.
- [140] Apóstolo J, Queirós P, Rodrigues M, Castro I, Cardoso D. The effectiveness of nonpharmacological interventions in older adults with depressive disorders: a systematic review. *JBIC Database System Rev Implement Rep* 2015;13(6):220-78.
- [141] Bains J, Birks J, Denning T. Antidepressants for treating depression in dementia. *Cochrane Database Syst Rev* 2002;(4):CD003944.
- [142] Kiosses DN, Teri L, Velligan DI, Alexopoulos GS. A home-delivered intervention for depressed, cognitively impaired, disabled elders. *Int J Geriatr Psychiatry* 2011;26(3):256-62.
- [143] Kiosses DN, Ravdin LD, Gross JJ, Raue P, Kotbi N, Alexopoulos GS: Problemadaptation therapy for older adults with major depression and cognitive impairment: a randomized clinical trial. *JAMA Psychiatry*. 2015;72(1):22-30.
- [144] Webster J. *Critical Advances in Reminiscence Work: From Theory to Application*. New York, NY: Springer 2002.
- [145] Douglas S, James I, Ballard C. Non-pharmacological interventions in dementia. *Adv Psychiatr Treat* 2004;10:71–79.
- [146] Olazarán J, Reisberg B, Clare L. Nonpharmacological therapies in Alzheimer's disease: a systematic review of efficacy. *Dement Geriatr Cogn Disord*. 2010;30(2):161-78.

- [147] Livingston G: Systematic review of psychological approaches to the management of neuropsychiatric symptoms of dementia. *Am J Psychiatry* 2005;162(11):1996-2021.
- [148] Neal M, Barton W.P: Validation therapy for dementia. *Cochrane Database Syst Rev.* 2003;(3):CD001394.
- [149] Collins FS, Varmus H. A new initiative on precision medicine. *N Engl J Med* 2015;372:793–795.
- [150] Kelloff GJ, Sigman CC. Cancer biomarkers: selecting the right drug for the right patient. *Nat Rev Drug Discov* 2012;11(3):201-214.
- [151] Noorbakhsh, F., Overall, C.M., Power, C. Deciphering complex mechanisms in neurodegenerative diseases: the advent of systems biology. *Trends Neurosci* 2009;32:88–100.
- [152] Bensimon A, Heck AJ, Aebersold R. Mass spectrometry based proteomics and network biology. *Annu Rev Biochem* 2012;81:379–405.
- [153] Sabido E, Selevsek N, Aebersold R. Mass spectrometry based proteomics for systems biology. *Curr Opin Biotechnol* 2012;23:591–597.
- [154] Reitz C. Toward precision medicine in Alzheimer's disease. *Ann Transl Med* 2016;4(6):107
- [155] Sperling RA, Jack CR Jr, Aisen PS. Testing the right target and right drug at the right stage. *Sci Transl Med.* 2011;3(111):111cm33.