

A COMMON RECURSION FOR LAPLACIANS OF
MATROIDS AND SHIFTED SIMPLICIAL COMPLEXES

ART M. DUVAL

Received: June 10, 2004

Revised: March 31, 2005

Communicated by Günter Ziegler

ABSTRACT. A recursion due to Kook expresses the Laplacian eigenvalues of a matroid M in terms of the eigenvalues of its deletion $M - e$ and contraction M/e by a fixed element e , and an error term. We show that this error term is given simply by the Laplacian eigenvalues of the pair $(M - e, M/e)$. We further show that by suitably generalizing deletion and contraction to arbitrary simplicial complexes, the Laplacian eigenvalues of shifted simplicial complexes satisfy this exact same recursion. We show that the class of simplicial complexes satisfying this recursion is closed under a wide variety of natural operations, and that several specializations of this recursion reduce to basic recursions for natural invariants.

We also find a simple formula for the Laplacian eigenvalues of an arbitrary pair of shifted complexes in terms of a kind of generalized degree sequence.

2000 Mathematics Subject Classification: Primary 15A18; Secondary 05B35, 05E99

Keywords and Phrases: Laplacian, spectra, matroid complex, shifted simplicial complex, Tutte polynomial

1. INTRODUCTION

The independence complex of matroids and shifted simplicial complexes are two of only four types of simplicial complexes whose combinatorial Laplacians $L = \partial\partial^* + \partial^*\partial$ are known to have only integer eigenvalues (see Kook, Reiner, and Stanton [27], and [16], respectively). The other two types, which will not concern us further, are matching complexes of complete graphs [14] and chessboard complexes [21]. More information and background about the combinatorial Laplacian and its eigenvalues may be found in Section 2 and [16, 20, 27]. Our main result (Theorems 3.18 and 4.23) is another, more striking, similarity

between the Laplacian eigenvalues of matroids and shifted complexes: they satisfy the exact same recursion, which we call the *spectral recursion*, equation (2). This recursion is stated in terms of the *spectrum polynomial*, a natural generating function for Laplacian eigenvalues, defined in equation (1).

The Tutte polynomial T_M of a matroid M satisfies the recursion $T_M = T_{M-e} + T_{M/e}$, when e is neither a loop nor an isthmus, and where $M-e$ and M/e denote the deletion and contraction, respectively, of M with respect to ground element e . When Kook, Reiner, and Stanton proved that the Laplacian spectrum of a matroid is integral, they also speculated on the existence of a Tutte polynomial-like recursion for the spectrum polynomial of a matroid M , though possibly with a third “error” term, besides the deletion and contraction, on the right-hand side [27, Question 3]. Kook [26] found such a recursion, but the error term in his formulation is somewhat complicated to state, with two cases depending on whether or not the ground element e is a closed element in M . Subsequently, Kook and Reiner (private communication) asked if this error term might be just the spectrum polynomial of the matroid pair $(M - e, M/e)$.

One of our main results (Theorem 3.18) is that Kook and Reiner’s conjecture is true, that is, the spectrum polynomial of M can be expressed simply in terms of the spectrum polynomials of $M - e$, M/e , and $(M - e, M/e)$. This is the spectral recursion. We show, furthermore, by suitably generalizing the definitions of deletion and contraction from matroids to arbitrary simplicial complexes (Section 2), that shifted complexes also satisfy the spectral recursion (Theorem 4.23).

This raises the natural question: What is the largest class of simplicial complexes, necessarily a common generalization of matroids and shifted complexes, satisfying the spectral recursion? We will see that this class is closed under the operations of join, skeleta, Alexander dual, and disjoint union (Corollaries 4.5, 4.19, 6.8, and 6.11, respectively). We might hope that it is closed also under deletion and contraction, as matroids and shifted complexes each are. In the same vein, it may be worthwhile to restrict our attention to those complexes that are also Laplacian integral. Unfortunately, no hint to determining this common generalization is apparent in the proofs of either Laplacian integrality or the spectral recursion, which are each rather different for matroids and shifted complexes.

Jarrah and Laubenbacher [23] examined another property shared by matroids and shifted complexes. Klivans [24] has characterized simplicial complexes that are simultaneously shifted *and* the matroid complex of some matroid; this is, in some sense, the reverse of finding a natural common generalization of matroids and shifted complexes.

The common generalization includes neither of the other known types of Laplacian integral simplicial complexes. Direct computations show that the matching complex of the complete graph on 5 vertices and the 2×3 chessboard complex both fail to satisfy the spectral recursion with respect to any vertex. Also excluded is the 3-edge path (Example 2.5), which rules out as the common

generalization such otherwise likely candidates as vertex-decomposable [32][9, Section 11] or shellable complexes [8, 9].

A key piece of the proof that matroids satisfy the spectral recursion is a decomposition of the Laplacian of $(M - e, M/e)$ into a direct sum of Laplacians of M/C 's for all circuits C containing e (Lemma 3.3). We may combine this with the spectral recursion to express the spectrum polynomial of a matroid completely in terms of spectrum polynomials of smaller matroids (with no matroid pairs), which permits a truly recursive way of computing Laplacian eigenvalues for matroids (Remark 3.19).

Unfortunately, we are unable to state any formula for the Laplacian eigenvalues of an *arbitrary* matroid pair (*i.e.*, besides $(M - e, M/e)$). We are able, however, to use tools developed in the proof of the spectral recursion for shifted complexes to find a simple formula for the Laplacian eigenvalues of an arbitrary shifted simplicial pair (Theorem 5.7). This naturally generalizes a formula for a single shifted complex [16]; the graph case goes back to Merris [29]. Similarly, we generalize a related conjectured inequality on the Laplacian spectrum of an arbitrary simplicial complex [16] to an arbitrary simplicial pair (Conjecture 5.8); the graph case was conjectured by Grone and Merris [22]. Passing from graphs to simplicial complexes in [16] required generalizing the well-known notion of degree sequences for graphs. Now passing to simplicial *pairs*, we introduce a less than obvious, but perfectly natural, further generalization of degree sequence (Subsection 5.2).

The Tutte polynomial is arguably the most important invariant of matroid theory (see, *e.g.*, [12]). The spectrum polynomial shares several nice features with the Tutte polynomial, such as being well-behaved under join (Corollary 4.3), disjoint union (Lemma 6.9), and several dual operators (equations (29) and (32)). Furthermore, specializations obtained by plugging in particular values for one or the other of the variables of the spectrum polynomial reduce it to well-known invariants. Consequently (and now going beyond matroids and the Tutte polynomial), in each of these specializations, the spectral recursion holds for all simplicial complexes Δ (not just matroids and shifted complexes), because it reduces to a basic recursion expressing the relevant invariant for Δ in terms of that invariant for $\Delta - e$ and Δ/e (Theorem 2.4 and Corollary 4.8). In contrast to the Tutte polynomial recursion, the spectral recursion does not need to exclude loops and isthmuses as special cases. Indeed, the spectral recursion holds for all complexes (not just matroids and shifted complexes) when e is a loop (Proposition 2.3) or an isthmus (Proposition 2.2 and Theorem 2.4).

Section 2 contains more information about Laplacians and the spectral recursion, including some special cases. Sections 3 and 4 are devoted to the proofs that matroids and shifted complexes, respectively, satisfy the spectral recursion. The formula for eigenvalues of arbitrary shifted simplicial pairs is developed in Section 5. Finally, in Section 6, we show that disjoint union and several duality operators, including Alexander duality, all preserve the property of satisfying the spectral recursion.

2. LAPLACIANS OF SIMPLICIAL PAIRS

For further background on simplicial complexes, their boundary maps and homology groups, see, *e.g.*, [30, Chapter 1]. If Δ and Δ' are simplicial complexes on the same ground set of vertices, then we will say (Δ, Δ') is a *simplicial pair*, but we set $(\Delta, \Delta') = (\Gamma, \Gamma')$ when the set differences $\Delta \setminus \Delta'$ and $\Gamma \setminus \Gamma'$ are equal as subsets of the power set of the ground set of vertices (here $A \setminus B$ denotes the set difference $\{a \in A : a \notin B\}$ between sets A and B); more formally, then, a simplicial pair is an equivalence class on ordered pairs of simplicial complexes. In all cases, definitions applying to a simplicial pair (Δ, Δ') may be specialized to a single simplicial complex Δ , by letting $\Delta' = \emptyset$, the empty simplicial complex.

As usual, let $C_i = C_i(\Delta, \Delta'; \mathbb{R}) := C_i(\Delta; \mathbb{R})/C_i(\Delta'; \mathbb{R})$ denote the i -dimensional oriented \mathbb{R} -chains of (Δ, Δ') , *i.e.*, the formal \mathbb{R} -linear sums of oriented i -dimensional faces $[F]$ such that $F \in \Delta_i \setminus \Delta'_i$, where Δ_i denotes the set of i -dimensional faces of Δ . Let $\partial_{(\Delta, \Delta'); i} = \partial_i : C_i \rightarrow C_{i-1}$ denote the usual (signed) *boundary operator*. Via the natural bases $\Delta_i \setminus \Delta'_i$ and $\Delta_{i-1} \setminus \Delta'_{i-1}$ for $C_i(\Delta, \Delta'; \mathbb{R})$ and $C_{i-1}(\Delta, \Delta'; \mathbb{R})$, respectively, the boundary map ∂_i has an adjoint map $\partial_i^* : C_{i-1}(\Delta, \Delta'; \mathbb{R}) \rightarrow C_i(\Delta, \Delta'; \mathbb{R})$; *i.e.*, the matrices representing ∂ and ∂^* in the natural bases are transposes of one another.

DEFINITION. Let $L'_i = \partial_{i+1} \partial_{i+1}^*$ and $L''_i = \partial_i^* \partial_i$. Then the (*i -dimensional Laplacian*) of (Δ, Δ') is the map $L_i(\Delta, \Delta') : C_i(\Delta, \Delta'; \mathbb{R}) \rightarrow C_i(\Delta, \Delta'; \mathbb{R})$ defined by

$$L_i = L_i(\Delta, \Delta') := L'_i + L''_i = \partial_{i+1} \partial_{i+1}^* + \partial_i^* \partial_i.$$

For more information, see, *e.g.*, [16, 20, 27]. Laplacians of pairs of graphs were considered in [13]. Each of L'_i and L''_i is positive semidefinite, since each is the composition of a linear map and its adjoint. Therefore, their sum L_i is also positive semidefinite, and so has only non-negative real eigenvalues. (See also Proposition 4.6 and [20, Proposition 2.1].) These eigenvalues do not depend on the arbitrary ordering of the vertices of Δ , and are thus invariants of (Δ, Δ') ; see, *e.g.*, [16, Remark 3.2]. Define $\mathbf{s}_i(\Delta, \Delta')$ to be the multiset of eigenvalues of $L_i(\Delta, \Delta')$, and define $m_\lambda(L_i(\Delta, \Delta'))$ to be the multiplicity of λ in $\mathbf{s}_i(\Delta, \Delta')$. The single complex case ($\Delta' = \emptyset$) of the following proposition is the first result of combinatorial Hodge theory, which goes back to Eckmann [18].

PROPOSITION 2.1. *The multiplicity of 0 as an eigenvalue of the i -dimensional Laplacian L_i of (Δ, Δ') is the i th reduced Betti number of (Δ, Δ') , *i.e.*,*

$$m_0(L_i(\Delta, \Delta')) = \tilde{\beta}_i(\Delta, \Delta') = \dim_{\mathbb{R}} \tilde{H}_i(\Delta, \Delta'; \mathbb{R}).$$

Proof. A nice summary is given in the proof of [20, Proposition 2.1]. The usual setup is for just a single simplicial complex (*i.e.*, the special case $\Delta' = \emptyset$), but only depends on the C_i 's and ∂_i 's forming a chain complex ($\partial^2 = 0$), which still holds even when $\Delta' \neq \emptyset$. (*Cf.* Proposition 4.6.) \square

A natural generating function for the Laplacian eigenvalues of a simplicial pair (Δ, Δ') is

$$(1) \quad S_{(\Delta, \Delta')}(t, q) := \sum_{i \geq 0} t^i \sum_{\lambda \in \mathbf{s}_{i-1}(\Delta, \Delta')} q^\lambda = \sum_{i, \lambda} m_\lambda(L_{i-1}(\Delta, \Delta')) t^i q^\lambda.$$

We call $S_{(\Delta, \Delta')}$ the *spectrum polynomial* of (Δ, Δ') . Although $S_{(\Delta, \Delta')}$ is defined for any simplicial pair (Δ, Δ') , it is only truly a polynomial when the Laplacian eigenvalues are not only non-negative, but integral as well. This will be true for the cases we are concerned with, primarily matroids [27], shifted complexes [16], and shifted simplicial pairs (Theorem 5.7 and Remark 5.9). For the special case of a matroid, a “spectrum polynomial” *Spec* was defined, differently, in [27], but we will see later that the two definitions agree in this case up to simple changes in indexing (see Lemma 3.6 and [27, Corollary 18]). Letting $\lambda \in \mathbf{s}_{i-1}$ instead of $\lambda \in \mathbf{s}_i$ simplifies the statement of some later results, notably Corollary 4.3. Recall (e.g., [5, Section 7.3]) the *independence complex* $IN(M)$ of a matroid M on ground set E is the simplicial complex whose faces are the independent sets of M and whose vertex set is E . (For background about matroids, see, e.g., [31, 34, 35].) We will sometimes use M and $IN(M)$ interchangeably, so, for instance, $L_i(M) := L_i(IN(M)) = L_i(IN(M), \emptyset)$ and $S_M := S_{IN(M)} = S_{(IN(M), \emptyset)}$. Similarly, if N is another matroid on the same ground set such that $IN(N) \subseteq IN(M)$ (i.e., $N \leq M$ in the weak order on matroids), then $L_i(M, N) = L_i(IN(M), IN(N))$ and $S_{(M, N)} = S_{(IN(M), IN(N))}$. In this case, we say (M, N) is a *matroid pair*.

We now naturally generalize the notion of deletion and contraction for matroids (see e.g., [11]) to arbitrary simplicial complexes.

DEFINITION. Let Δ be a simplicial complex on vertex set V , and $e \in V$. Then the *deletion* of Δ with respect to e is the simplicial complex

$$\Delta - e = \{F \in \Delta : e \notin F\}$$

on vertex set $V - e$, and the *contraction* of Δ with respect to e is the simplicial complex

$$\Delta / e = \{F - e : F \in \Delta, e \in F\}$$

on vertex set $V - e$. Note that $\Delta / e = \text{lk}_\Delta e$, the usual simplicial complex link [30, Section 2]; we use the term “contraction” to highlight similarities to matroid theory.

It is easy to verify that $IN(M - e) = IN(M) - e$ as long as e is not an isthmus of M , and that $IN(M/e) = IN(M)/e$ as long as e is not a loop of M . There is thus no confusion in the notational shortcuts $S_{M-e} := S_{IN(M-e)} = S_{IN(M)-e}$ and $S_{M/e} := S_{IN(M/e)} = S_{IN(M)/e}$ as long as e is not an isthmus or a loop, respectively.

Since e is an isthmus of M precisely when e is a vertex of every facet of $IN(M)$, define e to be an *isthmus* of a simplicial complex Δ if e is a vertex of every facet of Δ (so Δ is a cone with apex e – see Subsection 4.1). Similarly, since e is a loop of M precisely when e is not a vertex of any face of $IN(M)$, define

e to be a *loop* of a simplicial complex Δ if e is in the vertex set of Δ , but in no face of Δ (even the singleton $\{e\}$ is not a face, contrary to usual simplicial complex conventions).

Our definitions mean that if e is an isthmus of simplicial complex Δ , then the deletion $\Delta - e$ equals Δ/e . (When e is an isthmus of a matroid M , the *matroid* deletion $M - e$ is left undefined in *e.g.*, Brylawski [11], though $M - e = M/e$ in Welsh [34, Section 4.2] and Oxley [31, Corollary 3.1.25].) If e is a loop of simplicial complex Δ , then the contraction Δ/e is \emptyset , the empty simplicial complex. (When e is a loop of a matroid M , the *matroid* contraction M/e equals $M - e$.)

DEFINITION. We will say that a simplicial complex Δ *satisfies the spectral recursion with respect to e* if e is a vertex of Δ and

$$(2) \quad S_{\Delta}(t, q) = qS_{\Delta-e}(t, q) + qtS_{\Delta/e}(t, q) + (1 - q)S_{(\Delta-e, \Delta/e)}(t, q).$$

We will say Δ *satisfies the spectral recursion* if Δ satisfies the spectral recursion with respect to every vertex in its vertex set. (Note that Proposition 2.3 below means we need not be too particular about the vertex set of Δ .)

Our main result is that Δ satisfies the spectral recursion when Δ is either the independence complex of a matroid (Theorem 3.18) or a shifted simplicial complex (Theorem 4.23), and e is any vertex of Δ . We illustrate now a few special cases of the spectral recursion, which are easy to verify, and some of which are used in later sections.

PROPOSITION 2.2. *The simplicial complex whose sole facet is a single vertex satisfies the spectral recursion.*

PROPOSITION 2.3. *If e is a loop of simplicial complex Δ , then Δ satisfies the spectral recursion with respect to e .*

Proposition 2.2 and Theorem 4.4 will show that, if e is an isthmus of Δ , then Δ satisfies the spectral recursion with respect to e .

THEOREM 2.4. *If Δ is any simplicial complex, and e is any vertex of Δ , then the spectral recursion holds when $q = 0$, $q = 1$, $t = 0$, or $t = -1$.*

Proof. Plugging $q = 0$ into S immediately yields $S_{(\Delta, \Delta')}(t, 0) = \sum_i t^i \tilde{\beta}_{i-1}(\Delta, \Delta')$, by Proposition 2.1. Proving the spectral recursion in this case then reduces to showing

$$(3) \quad \tilde{\beta}_{i-1}(\Delta) = \tilde{\beta}_{i-1}(\Delta - e, \Delta/e),$$

for all i . This, in turn, is a consequence of the basic topology facts $\tilde{\beta}_{i-1}(\Delta) = \tilde{\beta}_{i-1}(\Delta, \text{st}_{\Delta} e)$ and $(\Delta, \text{st}_{\Delta} e) = (\Delta - e, \Delta/e)$, where $\text{st}_{\Delta} e$ denotes the usual star of e in Δ , the simplicial complex whose facets are the facets of Δ containing e . Setting $q = 1$, we see $S_{(\Delta, \Delta')}(t, 1) = \sum_i (f_{i-1}(\Delta) - f_{i-1}(\Delta'))t^i$, where f_i is the number of i -dimensional faces of Δ , since there are as many eigenvalues of $L_{i-1}(\Delta, \Delta')$ as there are faces in $\Delta_{i-1} \setminus \Delta'_{i-1}$ (assuming $\Delta' \subseteq \Delta$). It is then

an easy exercise to verify that, when $q = 1$, the t^{i+1} coefficient of the spectral recursion reduces to the easy observation

$$(4) \quad f_i(\Delta) = f_i(\Delta - e) + f_{i-1}(\Delta/e).$$

If we set $t = 0$, it is easy to see that $S_\Delta(0, q) = q^{v(\Delta)}$, where $v(\Delta)$ denotes the number of non-loop vertices of Δ . The spectral recursion in this case reduces to the trivial observation that $v(\Delta) = 1 + v(\Delta - e)$ if e is not a loop, but $v(\Delta) = v(\Delta - e)$ if e is a loop.

We will also see in Corollary 4.8 that, when $t = -1$, the spectral recursion reduces to an easy identity about Euler characteristic. \square

In the special case where Δ is a near-cone (see Subsection 4.5) and e is its apex, it is not hard to verify that the $t^{\dim \Delta + 1}$ coefficient of the spectral recursion reduces to [16, Lemma 5.3].

The following complex is the simplest and smallest counterexample to both Laplacian integrality and the spectral recursion.

EXAMPLE 2.5. Let Δ be the 1-dimensional simplicial complex with vertices a, b, c, d and facets (maximal faces) $\{a, b\}, \{b, c\}$, and $\{c, d\}$. It is easy to check directly that $\Delta - e$, Δ/e , and $(\Delta - e, \Delta/e)$ are all Laplacian integral for any choice of e , while Δ is not integral. It then follows immediately that Δ does not satisfy the spectral recursion for any choice of e .

3. MATROIDS

In this section, we show that the independence complex of a matroid satisfies the spectral recursion, equation (2). The key step of the section is a simple trick in Subsection 3.1 to reduce the problem of computing $S_{(M-e, M/e)}$ to computing $S_{M/C}$ for all circuits C containing e . Subsection 3.2 shows how an algorithm due to Kook, Reiner, and Stanton [27] allows us to compute the spectrum polynomial of a matroid from its combinatorial information; we also compare what this algorithm computes for M , $M - e$, M/e , and M/C . The final steps of the calculation, which largely consist of translating to generating functions the results of the previous subsections, are in Subsection 3.3.

We first set our notation for matroids; for further background, and any terms not defined here, see [35]. Let $M = M(E)$ be a matroid on ground set E . We will let $\mathcal{B} = \mathcal{B}(M)$, $\mathcal{I} = \mathcal{I}(M)$, $\mathcal{C} = \mathcal{C}(M)$, and $\mathcal{F} = \mathcal{F}(M)$ denote the sets of bases, independent sets, circuits, and flats (closed sets) of M , respectively. If $A \subseteq E$, let $\text{rk}_M(A) = \text{rk}(A)$ denote the *rank* of A (with respect to M), and let $\bar{A} = \text{cl}_M(A)$ denote the *closure* of A (with respect to M). We will often write V for $M(V)$ in the special case when V is a flat of M . When $A \subseteq V$, the set $V - A$ may be considered to be the *matroid* V/A in matroid M/A , but considered to be the *matroid* $V - A$ in matroid $M - A$. We will also use the notions of internal and external activity as in, e.g., [5].

3.1. A PARTITION. If Δ is a simplicial complex and A is a set disjoint from the vertices of Δ , then let $A \circ \Delta$ denote

$$A \circ \Delta := \{A \dot{\cup} F : F \in \Delta\}.$$

It will soon be important to note that $A \circ \Delta$ is a simplicial pair; in fact $A \circ \Delta = (2^A * \Delta, (2^A \setminus \{A\}) * \Delta)$, where 2^A denotes the simplicial complex consisting of all subsets of A , and $*$ denotes the usual join, as defined in Section 4.

LEMMA 3.1. *If Δ is a simplicial complex and A a finite set disjoint from the vertices of Δ , then*

$$S_{A \circ \Delta}(t, q) = t^{|A|} S_{\Delta}(t, q).$$

Proof. Under the natural bijection between Δ and $A \circ \Delta$, given by $\phi : F \mapsto A \dot{\cup} F$, the boundary operators ∂_{Δ} and $\partial_{A \circ \Delta}$ are the same. That is, $\partial_{A \circ \Delta}[A \dot{\cup} F] = [A] \partial_{\Delta}[F]$, simply by numbering the vertices of $A \circ \Delta$ so that the elements of A all come last. Since the boundary operators are the same, so are the Laplacians, but the dimension shift in ϕ means $s_i(\Delta) = s_{i+|A|}(A \circ \Delta)$. The lemma now follows readily. \square

If I is independent in M and $p \in \bar{I} - I$, we will let $ci(p, I) = ci_M(p, I) = ci_{\bar{I}}(p, I)$ be the unique circuit of \bar{I} contained in $I \dot{\cup} p$. Dually, if $b \in I$, we will let $bo(b, I) = bo_M(b, I) = bo_{\bar{I}}(b, I)$ be the unique bond of \bar{I} contained in $(\bar{I} - I) \dot{\cup} b$. It is easy to see that if $p \notin \bar{I}$, then $I \in \mathcal{I}(M/p)$. Therefore we may safely refer to $ci_M(p, I)$ for any $I \in \mathcal{I}(M - p) - \mathcal{I}(M/p)$.

LEMMA 3.2. *If $I', I \in \mathcal{I}(M - e) - \mathcal{I}(M/e)$ and $I' \subseteq I$, then $ci_M(e, I') = ci_M(e, I)$.*

Proof. From $ci_M(e, I') \subseteq I' \dot{\cup} e \subseteq I \dot{\cup} e$ it follows that $ci_M(e, I')$ is a circuit in $I \dot{\cup} e$, and thus the unique circuit in $I \dot{\cup} e$, i.e., $ci_M(e, I)$. \square

The following lemma is the key step to proving that matroids satisfy the spectral recursion.

LEMMA 3.3. *Let $M(E)$ be a matroid, and $e \in E$. If e is not a loop, then*

$$L_i(M - e, M/e) = \bigoplus_{\substack{C \in \mathcal{C}(M) \\ e \in C}} L_i((C - e) \circ IN(M/C)).$$

Proof. For any $C \in \mathcal{C}(M)$ such that $e \in C$, let

$$M_C = \{I \in \mathcal{I}(M - e) - \mathcal{I}(M/e) : ci_M(e, I) = C\};$$

we will see shortly that this is a simplicial pair. By Lemma 3.2,

$$\partial_{(M - e, M/e)}[I] = \partial_C[I]$$

for any $I \in \mathcal{I}(M - e) - \mathcal{I}(M/e)$, where $C = ci_M(e, I)$. Thus removing M/e from $M - e$ partitions $L_i(M - e, M/e)$ into

$$L_i(M - e, M/e) = \bigoplus_{\substack{C \in \mathcal{C}(M) \\ e \in C}} L_i(M_C).$$

Furthermore, it is easy to see that

$$\begin{aligned} M_C &= \{I \in \mathcal{I}(M - e) : C - e \subseteq I\} = (C - e) \circ IN((M - e)/(C - e)) \\ &= (C - e) \circ IN(M/C). \end{aligned}$$

□

3.2. THE KOOK-REINER-STANTON ALGORITHM. The decomposition in Proposition 3.4 below was first discovered by Etienne and Las Vergnas [19, Theorem 5.1], but we will rely upon Algorithm 3.5, due to Kook, Reiner, and Stanton [27, proof of Theorem 1], for producing this decomposition.

PROPOSITION 3.4. *Given a base B of matroid M , there is a unique disjoint decomposition $B = B_1 \dot{\cup} B_2$ into two (necessarily) independent sets such that:*

- B_1 has internal activity 0; and
- B_2 has external activity 0, with respect to the matroid M/V , where $V = \overline{B_1}$.

ALGORITHM 3.5. This algorithm produces the decomposition guaranteed by the previous theorem. It takes the base B as input, and outputs the pair (B_1, B_2) .

Step 1: Set $B_1 = B, B_2 = \emptyset$.

Step 2: Let $V = \overline{B_1}$.

Step 3: Find an internally active element b for B_1 as a base of the flat V .

- If no such element b exists, then stop and output the pair (B_1, B_2) .
- If such a b exists, then set $B_1 := B_1 - b, B_2 := B_2 \dot{\cup} b$ (we call this step a *removal*), and return to Step 2.

NOTATION. If the decomposition of base B in matroid M produced by the above algorithm is $B = B_1 \dot{\cup} B_2$, then let $\pi(B) = \pi_M(B) = B_1$. If $I \in \mathcal{I}(M)$, then let $\overline{\pi}_M(I) = \text{cl}_V(\pi_V(I)) = \text{cl}_M(\pi_V(I))$, where $V = \text{cl}_M(I)$. If W is any closed set containing I (equivalently, containing $V = \text{cl}_M(I)$), then $\text{cl}_W(I) = \text{cl}_M(I) = V$, and so $\overline{\pi}_W(I) = \text{cl}_V(\pi_V(I)) = \overline{\pi}_M(I)$. In particular, $\overline{\pi}_V(I) = \overline{\pi}_M(I)$.

The following lemma, which is little more than a recasting of [27, Corollary 18] in language tailored to our purposes, reduces computations of the spectrum polynomial to computations of $\overline{\pi}$.

LEMMA 3.6. *For any matroid $M(E)$,*

$$S_M(t, q) = q^{|E|} \sum_{I \in \mathcal{I}(M)} t^{\text{rk}(\overline{I})} (q^{-1})^{|\overline{\pi}_M(I)|} = q^{|E|} \sum_{V \in \mathcal{F}(M)} t^{\text{rk}(V)} \sum_{I \in \mathcal{B}(V)} (q^{-1})^{|\overline{\pi}_V(I)|}.$$

Let $\tilde{\chi}(\Delta) := \sum (-1)^i f_i(\Delta)$ denote the (reduced) Euler characteristic of simplicial complex Δ ; we also use the shorthand $\tilde{\chi}(M) = \tilde{\chi}(IN(M))$. If $V \subseteq W$ are flats of matroid M , let $\mu(W, V) = \mu_M(W, V)$ denote the Möbius function of

the sublattice $[W, V]$ in the lattice of flats of M . The proof of [27, equation (2.2)] shows that

$$(5) \quad \sum_{B \in \overline{\mathcal{B}}(M)} x^{|\overline{\pi}_M(B)|} = \sum_{V \in \mathcal{F}(M)} |\tilde{\chi}(V)| |\mu(V, M)| x^{|V|}.$$

We use the same techniques to do something similar.

LEMMA 3.7. *For any matroid $M(E)$, and any $e \in E$,*

$$\sum_{\substack{B \in \overline{\mathcal{B}}(M) \\ e \in \overline{\pi}_M(B)}} x^{|\overline{\pi}_M(B)|} = \sum_{\substack{V \in \mathcal{F}(M) \\ e \in V}} |\tilde{\chi}(V)| |\mu(V, M)| x^{|V|}.$$

In particular, this sum is independent of the linear order on E .

Proof. By Algorithm 3.5 (see also its proof in [27]), there is a bijection between:

- the set \mathcal{V} of triples (V, B_1, B_2) where V is a flat of M , B_1 is a base of internal activity 0 for V (in particular, $V = \overline{B_1}$), and B_2 is a base of external activity 0 for M/V ; and
- the set \mathcal{B} of bases B of M .

Furthermore, $B = B_1 \dot{\cup} B_2$ and $\pi_M(B) = B_1$. Thus

$$\sum_{\substack{B \in \overline{\mathcal{B}}(M) \\ e \in \overline{\pi}_M(B)}} x^{|\overline{\pi}_M(B)|} = \sum_{\substack{(V, B_1, B_2) \in \mathcal{V} \\ e \in V}} x^{|V|}.$$

We must then determine how many triples (V, B_1, B_2) there are in \mathcal{V} for a fixed flat V . Mimicking an argument from the proof of [27, Theorem 1], we recall from [5, Theorem 7.8.4] that there are $|\tilde{\chi}(V)|$ bases of internal activity 0 for V , and from [5, Proposition 7.4.7] that there are $|\mu(V, M)|$ bases of external activity 0 for M/V . So for every V , there are $|\tilde{\chi}(V)|$ choices for B_1 , and, independently, $|\mu(V, M)|$ choices for B_2 . Thus,

$$\sum_{\substack{(V, B_1, B_2) \in \mathcal{V} \\ e \in V}} x^{|V|} = \sum_{\substack{V \in \mathcal{F}(M) \\ e \in V}} |\tilde{\chi}(V)| |\mu(V, M)| x^{|V|},$$

completing the proof. □

We now see how Algorithm 3.5 works on $M - e$ (Lemma 3.11) and M/e (Lemma 3.13), and on M/C when C is a circuit containing e (Lemma 3.15). We first need three technical lemmas whose easy proofs are omitted. We abuse set difference notation slightly to let $A \setminus x$ denote $\{a \in A : a \neq x\}$, when A is a set that may or may not contain element x .

LEMMA 3.8. *Let I be an independent set in matroid M , let e be last in the linear order, and assume that $e \notin I$ and that e is not an isthmus of M . Then b is internally active in I (with respect to M) iff b is internally active in I (with respect to $M - e$).*

LEMMA 3.9. *Let I be an independent set in matroid M , and let $e, b \in I$. Then b is internally active in I (with respect to M) iff b is internally active in $I - e$ (with respect to M/e).*

LEMMA 3.10. *Let I be an independent set in matroid M , and let i be an isthmus in \bar{I} . Then $b \neq i$ is internally active in I (with respect to M) iff b is internally active in $I - i$ (with respect to M).*

LEMMA 3.11. *Let B be a base of $M - e$, so B is also a base of M and $e \notin B$. Also assume e is last in the linear order. Then $\pi_{M-e}(B) = \pi_M(B)$.*

Proof. Use Algorithm 3.5 to compute $\pi_M(B)$. By Lemma 3.8, every step of the algorithm can be copied in $M - e$; that is, when element b is removed from B_1 in M , we can remove b from B_1 in $M - e$. And also by Lemma 3.8, when there are no more elements to remove from B_1 in M , then there are also no more elements to remove from B_1 in $M - e$. \square

COROLLARY 3.12. *Let I be an independent set of $M - e$, so I is also independent in M and $e \notin I$. Also assume e is last in the linear order. Then*

$$\bar{\pi}_{M-e}(I) = \bar{\pi}_M(I) \setminus e.$$

LEMMA 3.13. *Let B be a base of M such that $e \in B$, so $B - e$ is a base of M/e . Also assume e is last in the linear order. Then*

$$\pi_{M/e}(B - e) = \pi_M(B) \setminus e$$

Proof. Again use Algorithm 3.5 to compute $\pi_M(B)$, except do not remove e unless it is the only element that can be removed. As in Lemma 3.11, every step can be copied in M/e , this time by Lemma 3.9, as long as we are not removing e , and have not yet removed e . Also by Lemma 3.9, if we never remove e , then when there are no more elements to remove in M , there are no more elements to remove in M/e . Thus, if e is never removed (i.e., if $e \in \pi_M(B)$), then $\pi_{M/e}(B - e) = \pi_M(B) - e$.

If e is eventually removed in M , it must be when e is an isthmus, since e is ordered last (so it can be the minimal element of $\text{bo}(e, I)$ only if it is the only element - i.e., if it is an isthmus). Since we put off removing e until there were no other possible removals, Lemma 3.10 guarantees that there are no new removals possible after e is removed. Since the removals were identical in M and M/e until e was removed, $\pi_{M/e}(B - e) = \pi_M(B)$. \square

COROLLARY 3.14. *Let I be an independent set of M such that $e \in I$, so $I - e$ is independent in M/e . Also assume e is last in the linear order. Then*

$$\bar{\pi}_{M/e}(I - e) = \bar{\pi}_M(I) \setminus e.$$

Proof. Let $V = \text{cl}_M(I)$. Then $\text{cl}_{M/e}(I - e) = V - e$ as sets, so $\text{cl}_{M/e}(I - e) = V/e$ as matroids. Thus, by the definition of $\bar{\pi}$, we have

$$\bar{\pi}_{M/e}(I - e) = \text{cl}_{M/e}(\pi_{V/e}(I - e)).$$

If $e \in \pi_V(I)$, then simply $\text{cl}_{M/e}(\pi_{V/e}(I - e)) = \text{cl}_{M/e}(\pi_V(I) - e) = \text{cl}_M(\pi_V(I)) - e = \bar{\pi}_M(I) \setminus e$; the first equality is by Lemma 3.13, the second equality is a

routine exercise using $e \in \pi_V(I)$, and the last equality is from the definition of $\bar{\pi}$.

If $e \notin \pi_V(I)$, then the proof of Lemma 3.13 shows that e is an isthmus in $\text{cl}_V(\pi_V(I) \cup e)$. Then, since $\text{cl}(A \dot{\cup} i) = (\text{cl } A) \dot{\cup} i$ for any A and any isthmus $i \notin A$,

$$(6) \quad \text{cl}_M(\pi_V(I) \cup e) = \text{cl}_V(\pi_V(I) \cup e) = \text{cl}_V(\pi_V(I)) \cup e = \bar{\pi}_M(I) \cup e.$$

Now, also in this case,

$$\begin{aligned} \text{cl}_{M/e}(\pi_{V/e}(I - e)) &= \text{cl}_{M/e}(\pi_V(I)) = \text{cl}_M(\pi_V(I) \cup e) - e = (\bar{\pi}_M(I) \cup e) - e \\ &= \bar{\pi}_M(I) \setminus e; \end{aligned}$$

the first equality is by Lemma 3.13, the second equality is from the definition of $\text{cl}_{M/e}$, and the third equality is equation (6). \square

LEMMA 3.15. *Let B be a base of matroid $M(E)$, let e be first in the linear order on E , and assume that $e \notin B$ and e is not a loop. Let $C = \text{ci}(e, B)$, so $B - (C - e)$ is a base of M/C . Then*

$$\pi_{M/C}(B - (C - e)) = \pi_M(B) - (C - e).$$

Proof. It is an easy exercise to check that $\text{bo}_{M/C}(b, B - (C - e)) = \text{bo}_M(b, B)$ for any $b \in B - (C - e)$. It then follows that b is internally active in B (with respect to M) iff b is minimal in $\text{bo}_M(b, B) = \text{bo}_M(b, B - (C - e))$ iff b is internally active in $B - (C - e)$ (with respect to M/C).

Now, as in Lemmas 3.11 and 3.13, use Algorithm 3.5 to compute $\pi_{M/C}(B - (C - e))$. Once again, every step can be copied in M , computing $\pi_M(B)$. Furthermore, when there are no more elements in $B - (C - e)$ to remove in computing $\pi_{M/C}(B - (C - e))$, the only elements of B that could possibly be removed in computing $\pi_M(B)$ must be in $C - e$. We now show that any $c \in C - e$ is not internally active, and thus that the removals in M and M/C are identical, which will complete the proof.

It is easy to see that $C = \text{ci}_{B_1}^{-1}(e, B_1)$, where B_1 is what remains of B after performing all the removals in M corresponding to the removals in M/C . Thus $c \in C - e \subseteq \text{ci}_{B_1}^{-1}(e, B_1)$ implies, by *e.g.*, [5, Lemma 7.3.1], that $e \in \text{bo}_{B_1}(c, B_1)$. Since e is first in the linear order, c is, as desired, not internally active. \square

3.3. THE SPECTRAL RECURSION FOR MATROIDS. We now prove that matroids satisfy the spectral recursion (Theorem 3.18), by comparing $qtS_{M/e} + qS_{M-e} - S_M$ and $S_{(M-e, M/e)}$. In each case, we get two expressions, one in terms of $\tilde{\chi}$ and μ , the other in terms of $\bar{\pi}$. The expressions in terms of $\tilde{\chi}$ and μ lead to a quick proof, by reducing a key piece of the equation to the $q = 0$ case for a flat. The expressions in terms of $\bar{\pi}$ suggest a more bijective proof, which is not hard to prove either. Both proofs are given.

LEMMA 3.16. *If $M(E)$ is a matroid, and $e \in E$ is neither an isthmus nor a loop, then*

$$\begin{aligned} qS_{M-e}(t, q) + qtS_{M/e}(t, q) - S_M(t, q) &= (q-1)q^{|E|} \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{\substack{I \in \mathcal{B}(V) \\ e \in \bar{\pi}_V(I)}} (q^{-1})^{|\bar{\pi}_V(I)|} \\ &= (q-1) \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{\substack{W \in \mathcal{F}(V) \\ e \in W}} |\tilde{\chi}(W)| |\mu(W, V)| q^{|E|-|W|}. \end{aligned}$$

Proof. We compute each of S_{M-e} and $S_{M/e}$ using Lemma 3.6. First,

$$\begin{aligned} S_{M-e}(t, q) &= q^{|E-e|} \sum_{I \in \mathcal{I}(M-e)} t^{\text{rk}_{M-e}(\text{cl}_{M-e}(I))} (q^{-1})^{|\bar{\pi}_{M-e}(I)|} \\ (7) \quad &= q^{|E-e|} \sum_{\substack{I \in \mathcal{I}(M) \\ e \notin I}} t^{\text{rk}_M(\text{cl}_M(I))} (q^{-1})^{|\bar{\pi}_M(I) \setminus e|}, \end{aligned}$$

since: $I \in \mathcal{I}(M-e)$ iff $I \in \mathcal{I}(M)$ and $e \notin I$; $|\bar{\pi}_{M-e}(I)| = |\bar{\pi}_M(I) \setminus e|$, by Corollary 3.12; and $\text{rk}_{M-e}(\text{cl}_{M-e}(I)) = \text{rk}_M(\text{cl}_M(I) \setminus e)$ is an easy matroid exercise. Similarly,

$$\begin{aligned} S_{M/e}(t, q) &= q^{|E-e|} \sum_{I' \in \mathcal{I}(M/e)} t^{\text{rk}_{M/e}(\text{cl}_{M/e}(I'))} (q^{-1})^{|\bar{\pi}_{M/e}(I')|} \\ (8) \quad &= q^{|E-e|} \sum_{\substack{I \in \mathcal{I}(M) \\ e \in I}} t^{\text{rk}_M(\text{cl}_M(I)) - 1} (q^{-1})^{|\bar{\pi}_M(I) \setminus e|}, \end{aligned}$$

where $I = I' \dot{\cup} e$ for $I' \in \mathcal{I}(M/e)$, since: $|\bar{\pi}_{M/e}(I')| = |\bar{\pi}_{M/e}(I-e)| = |\bar{\pi}_M(I) \setminus e|$, by Corollary 3.14; and $\text{rk}_{M/e}(\text{cl}_{M/e}(I')) = \text{rk}_M(\text{cl}_M(I)) - 1$ is a routine exercise, using $e \in \text{cl}_M(I)$.

Combining equations (7) and (8), and then sorting independent sets by their closures, we get

$$\begin{aligned} qS_{M-e}(t, q) + qtS_{M/e}(t, q) &= q^{|E|} \sum_{I \in \mathcal{I}(M)} t^{\text{rk}_M(\text{cl}_M(I))} (q^{-1})^{|\bar{\pi}_M(I) \setminus e|} \\ (9) \quad &= q^{|E|} \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{I \in \mathcal{B}(V)} (q^{-1})^{|\bar{\pi}_V(I) \setminus e|}. \end{aligned}$$

Furthermore

$$\begin{aligned} \sum_{I \in \mathcal{B}(V)} (q^{-1})^{|\bar{\pi}_V(I) \setminus e|} &= \sum_{\substack{I \in \mathcal{B}(V) \\ e \notin \bar{\pi}_V(I)}} (q^{-1})^{|\bar{\pi}_V(I) \setminus e|} + \sum_{\substack{I \in \mathcal{B}(V) \\ e \in \bar{\pi}_V(I)}} (q^{-1})^{|\bar{\pi}_V(I) \setminus e|} \\ (10) \quad &= \sum_{I \in \mathcal{B}(V)} (q^{-1})^{|\bar{\pi}_V(I)|} + (q-1) \sum_{\substack{I \in \mathcal{B}(V) \\ e \in \bar{\pi}_V(I)}} (q^{-1})^{|\bar{\pi}_V(I)|}; \end{aligned}$$

plugging into equation (10) into equation (9) readily leads to the first equation of the lemma. The second equation then follows directly from Lemma 3.7. \square

LEMMA 3.17. *If $M(E)$ is a matroid, and $e \in E$ is neither an isthmus nor a loop, then*

$$\begin{aligned} S_{(M-e, M/e)}(t, q) &= q^{|E|} \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{\substack{C \in \mathcal{C}(V) \\ e \in C}} q^{-|C|} \sum_{I \in \mathcal{B}(V/C)} (q^{-1})^{|\bar{\pi}_{V/C}(I)|} \\ &= \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{\substack{W \in \mathcal{F}(V) \\ e \in W}} \sum_{\substack{C \in \mathcal{C}(W) \\ e \in W}} |\tilde{\chi}(W/C)| |\mu(W, V)| q^{|E|-|W|}. \end{aligned}$$

Proof. By Lemmas 3.6, 3.1, and 3.3,

$$\begin{aligned} S_{(M-e, M/e)}(t, q) &= \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} t^{\text{rk}_M(C)} S_{M/C}(t, q) \\ &= \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} t^{\text{rk}_M(C)} q^{|E-C|} \sum_{W \in \mathcal{F}(M/C)} t^{\text{rk}_{M/C}(W)} \sum_{I \in \mathcal{B}(W)} (q^{-1})^{|\bar{\pi}_W(I)|}. \end{aligned}$$

Now, the flats of M/C are $V - C$ as sets, and thus V/C as matroids, for all flats V of M containing C . Therefore,

$$\begin{aligned} S_{(M-e, M/e)}(t, q) &= q^{|E|} \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} t^{\text{rk}_M(C)} q^{-|C|} \sum_{\substack{V \in \mathcal{F}(M) \\ C \subseteq V}} t^{\text{rk}_M(V) - \text{rk}_M(C)} \sum_{I \in \mathcal{B}(V/C)} (q^{-1})^{|\bar{\pi}_{V/C}(I)|} \\ &= q^{|E|} \sum_{V \in \mathcal{F}(M)} t^{\text{rk}_M(V)} \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C \subseteq V}} q^{-|C|} \sum_{I \in \mathcal{B}(V/C)} (q^{-1})^{|\bar{\pi}_{V/C}(I)|}, \end{aligned}$$

which is the first equation of the lemma, once we note that $C \in \mathcal{C}(V)$ iff $C \in \mathcal{C}(M)$ and $C \subseteq V$.

The second equation of the lemma then follows from

$$\begin{aligned} &\sum_{\substack{C \in \mathcal{C}(V) \\ e \in C}} q^{-|C|} \sum_{I \in \mathcal{B}(V/C)} (q^{-1})^{|\bar{\pi}_{V/C}(I)|} \\ &= \sum_{\substack{C \in \mathcal{C}(V) \\ e \in C}} q^{-|C|} \sum_{W/C \in \mathcal{F}(V/C)} |\tilde{\chi}(W/C)| |\mu_{V/C}(W/C, V/C)| (q^{-1})^{|W/C|} \\ &= \sum_{\substack{C \in \mathcal{C}(V) \\ e \in C}} \sum_{\substack{W \in \mathcal{F}(V) \\ C \subseteq W}} |\tilde{\chi}(W/C)| |\mu(W, V)| (q^{-1})^{|W|} \\ &= \sum_{\substack{W \in \mathcal{F}(V) \\ e \in W}} \sum_{\substack{C \in \mathcal{C}(V) \\ e \in C \subseteq W}} |\tilde{\chi}(W/C)| |\mu(W, V)| (q^{-1})^{|W|}. \end{aligned}$$

The first equation above is from equation (5); we are also using the same characterization of flats of a contraction as in the previous paragraph. The second equation is since the interval $[W/C, V/C]$ in the lattice of flats of V/C is isomorphic to the interval $[W, V]$ in the lattice of flats of V , again by that same characterization of flats in a contraction. It only remains to again note that $C \in \mathcal{C}(W)$ iff $C \in \mathcal{C}(M)$ and $C \subseteq W$. \square

THEOREM 3.18. *If M is a matroid, then its independence complex $IN(M)$ satisfies the spectral recursion, equation (2).*

Proof. By Proposition 2.3, we may assume e is not a loop. By Lemma 2.2 and Theorem 4.4 below (which does not depend on anything in this section), we may assume e is not an isthmus. As discussed at the beginning of the subsection, there are now two ways to finish off the proof, one using the $q = 0$ case, the other using a bijection.

$q = 0$ *proof.* By Theorem 2.4, we know that the spectral recursion holds, for any matroid, with $q = 0$. By Lemmas 3.16 and 3.17, this means

$$(11) \quad |\tilde{\chi}(M)| = \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} |\tilde{\chi}(M/C)|$$

for any matroid M , since only terms with $W = E$ survive when $q = 0$. (Equation (11) is also, as noted by Kook [25], dual to Crapo’s complementation theorem (*e.g.*, [1, Theorem 4.33]) applied to the dual matroid of M .) Thus, simply by plugging in the flat W , as a matroid, for the matroid M in equation (11),

$$|\tilde{\chi}(W)| = \sum_{\substack{C \in \mathcal{C}(W) \\ e \in C}} |\tilde{\chi}(W/C)|$$

whenever W is a flat of M containing e . By Lemmas 3.16 and 3.17 again, we are done.

Bijjective proof. By Lemmas 3.16 and 3.17, it suffices to show

$$(12) \quad \sum_{\substack{I \in \mathcal{B}(M) \\ e \in \bar{\pi}_W(I)}} (q^{-1})^{|\bar{\pi}_M(I)|} = \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} \sum_{I \in \mathcal{B}(M/C)} (q^{-1})^{|\bar{\pi}_{M/C}(I)| + |C|}.$$

Further, Lemma 3.7 shows that the sum on the left-hand side of equation (12) is independent of the ordering of the ground set. Similarly, Lemma 3.17 itself shows the same thing for the sum on the right-hand side. So we now assume, for the remainder of this proof, that e is ordered first in the linear order on E . Equation (12) would follow naturally from a bijection

$$\phi: \{B \in \mathcal{B}(M) : e \in \bar{\pi}_M(B)\} \rightarrow \{(C, I) : C \in \mathcal{C}(M), I \in \mathcal{B}(M/C), e \in C\}$$

such that

$$(13) \quad \bar{\pi}_{M/C}(I) \dot{\cup} C = \bar{\pi}_M(B),$$

where $\phi(B) = (C, I)$. Such a bijection is given by, as we now show, $C = \text{ci}(e, B)$ and $I = B - (C - e)$ in one direction, and $B = I \dot{\cup} C - e$ in the other.

First note that, since e is ordered first, if $e \in B$ then e is internally active in B , and so $e \notin \pi_M(B)$. It is then easy to see in this case that $e \notin \bar{\pi}_M(B)$. We may therefore safely assume $e \notin B$, and so $C = \text{ci}(e, B)$ is well-defined. It then follows that ϕ is well-defined.

It is easy to see that ϕ is injective. Showing that ϕ is surjective reduces to verifying that $e \in \bar{\pi}_M(B)$ when $B = I \dot{\cup} C - e$; by Lemma 3.15, $C - e \subseteq \pi_M(B)$, so $e \in C = \overline{C - e} \subseteq \bar{\pi}_M(B)$.

Finally, to verify equation (13), by Lemma 3.15 and the definition of closure in a matroid contraction, $\bar{\pi}_{M/C}(B - (C - e)) = \text{cl}_M(\pi_M(B) \cup e) - C$. Also, $\text{cl}_M(\pi_M(B) \cup e) = \text{cl}_M(\pi_M(B)) = \bar{\pi}_M(B)$, since $e \in \bar{\pi}_M(B)$, which completes the proof of equation (13). \square

REMARK 3.19. The spectral recursion does not provide a truly recursive way to compute S_M , due to the presence of $S_{(M-e, M/e)}$, since the recursion only applies to a single matroid, and not a matroid pair like $(M - e, M/e)$. We can however, combine it with Lemmas 3.1 and 3.3 for a recursion that is truly recursive, albeit with more terms than the spectral recursion:

$$S_M(t, q) = qS_{M-e}(t, q) + qtS_{M/e}(t, q) + (1 - q) \sum_{\substack{C \in \mathcal{C}(M) \\ e \in C}} t^{\text{rk}_M(C)} S_{M/C}(t, q).$$

I am grateful to E. Babson for this observation.

4. SHIFTED COMPLEXES

We postpone until Subsection 4.5 the actual definition of shifted complexes, but we will see there that a shifted complex is a skeleton of a cone of a smaller shifted complex (Lemmas 4.21–4.22). To prove that shifted complexes satisfy the spectral recursion, equation (2), then, it suffices to show that taking skeleta and taking cones each preserve the property of satisfying the spectral recursion – which are interesting results in their own right.

We will prove in Subsection 4.1 that the property of satisfying the spectral recursion is preserved by taking joins (Corollary 4.5), and thus by taking cones (*cf.* Proposition 2.2). The key step is that a simple formula [16, Theorem 4.10] for the eigenvalues of the join generalizes straightforwardly from single simplicial complexes to simplicial pairs (Corollaries 4.2 and 4.3).

Proving that taking skeleta preserves the property of satisfying the spectral recursion is harder, and is the focus of Subsections 4.2–4.4. The key facts about Laplacians, established in Subsections 4.2 and 4.3, respectively, are that the non-zero eigenvalues come in pairs in consecutive dimensions (Lemma 4.7), and that taking $(d-1)$ -skeleta preserves non-zero eigenvalues of the finer Laplacians in dimension $d-1$ and below (Lemma 4.11).

The only eigenvalues in dimension $d-1$ and below that are changed by taking $(d-1)$ -skeleta, then, are some $(d-1)$ -dimensional eigenvalues that become 0 when their counterparts (in the sense of Lemma 4.7) in dimension d are removed. It is auspicious that these replaced $(d-1)$ -dimensional eigenvalues must line up properly in the spectral recursion (since their counterparts in

dimension d , the only non-zero eigenvalues in that dimension, do as well) and that the 0's that replace them also line up properly (since the spectral recursion is true with $q = 0$ for both the original complex and its skeleton, by Theorem 2.4). But it turns out that we are better off with f -vectors ($q = 1$, also a good case by Theorem 2.4) than with homology ($q = 0$), in part because the change in f -vectors resulting from taking skeleta is much easier to describe than the change in homology.

In Subsection 4.4, we will see that the difference between the spectrum polynomials of the skeleton and the original complex can be described largely in terms of the f -vector (Lemma 4.14), allowing us to describe the difference in the spectral recursion between the skeleton and the original complex in a particularly useful form (Lemma 4.15). From there, simple generating function manipulations lead to Theorem 4.18, which states that a d -dimensional simplicial complex satisfies the spectral recursion with respect to a vertex if and only if its $(d - 1)$ -skeleton and pure d -skeleton (the complex generated by its facets) do as well.

4.1. JOINS AND CONES. Define the *join* $(\Delta, \Delta') * (\Gamma, \Gamma')$ of two simplicial pairs on disjoint vertex sets to be

$$(\Delta, \Delta') * (\Gamma, \Gamma') := \{F \dot{\cup} G : F \in \Delta \setminus \Delta', G \in \Gamma \setminus \Gamma'\}$$

(here, $\dot{\cup}$ denotes disjoint union), which equals the simplicial pair

$$(14) \quad (\Delta * \Gamma, (\Delta' * \Gamma) \cup (\Delta * \Gamma')).$$

When $\Delta' = \Gamma' = \emptyset$, this reduces to the usual join $\Delta * \Gamma$. When, further, Δ is a single vertex, say v , the join is written as $v * \Gamma$, the *cone* over Γ with *apex* v . The proofs of the following two results on simplicial pairs are identical (modulo some indexing changes) to those of the analogous statements for single simplicial complexes [16, Section 4].

PROPOSITION 4.1. *For any two simplicial pairs (Δ, Δ') and (Γ, Γ') and every k , the map defined \mathbb{R} -linearly by $[F] \otimes [G] \mapsto [F \dot{\cup} G]$ identifies the vector spaces*

$$\bigoplus_{i+j=k} C_{i-1}((\Delta, \Delta'); \mathbb{R}) \otimes C_{j-1}((\Gamma, \Gamma'); \mathbb{R}) \cong C_{k-1}((\Delta, \Delta') * (\Gamma, \Gamma'); \mathbb{R})$$

and has the following property with respect to the Laplacians L of the appropriate dimensions in (Δ, Δ') , (Γ, Γ') , and $(\Delta, \Delta') * (\Gamma, \Gamma')$:

$$(15) \quad L((\Delta, \Delta') * (\Gamma, \Gamma')) = L(\Delta, \Delta') \otimes \text{id} + \text{id} \otimes L(\Gamma, \Gamma').$$

COROLLARY 4.2. *If (Δ, Δ') and (Γ, Γ') are two simplicial pairs, then*

$$\mathbf{s}_{k-1}((\Delta, \Delta') * (\Gamma, \Gamma')) = \bigcup_{\substack{i+j=k \\ \lambda \in \mathbf{s}_{i-1}(\Delta, \Delta'), \mu \in \mathbf{s}_{j-1}(\Gamma, \Gamma')}} \lambda + \mu.$$

It is then an easy exercise in generating functions to verify the following corollary.

COROLLARY 4.3. *If (Δ, Δ') and (Γ, Γ') are two simplicial pairs, then*

$$S_{(\Delta, \Delta')*(\Gamma, \Gamma')} = S_{(\Delta, \Delta')}S_{(\Gamma, \Gamma')}.$$

THEOREM 4.4. *If Δ satisfies the spectral recursion with respect to e , and Γ is any simplicial complex whose vertex set is disjoint from the vertex set of Δ , then the join $\Delta * \Gamma$ satisfies the spectral recursion with respect to e .*

Proof. By Corollary 4.3 twice, and our hypothesis,

$$\begin{aligned} S_{\Delta * \Gamma} &= S_{\Delta}S_{\Gamma} = (qS_{\Delta - e} + qtS_{\Delta/e} + (1 - q)S_{(\Delta - e, \Delta/e)})S_{\Gamma} \\ &= qS_{(\Delta - e)*\Gamma} + qtS_{(\Delta/e)*\Gamma} + (1 - q)S_{(\Delta - e, \Delta/e)*\Gamma}. \end{aligned}$$

This last expression is exactly what we need, since it is easy to verify that join commutes with deletion and contraction, *i.e.*, $(\Delta - e) * \Gamma = (\Delta * \Gamma) - e$ and $(\Delta/e) * \Gamma/e = (\Delta * \Gamma)/e$, and also since equation (14) with $\Delta' = \emptyset$ then yields

$$(\Delta - e, \Delta/e) * \Gamma = ((\Delta - e) * \Gamma, (\Delta/e) * \Gamma) = ((\Delta * \Gamma) - e, (\Delta * \Gamma)/e).$$

□

COROLLARY 4.5. *If Δ and Γ each satisfy the spectral recursion, then so does their join $\Delta * \Gamma$.*

4.2. FINER LAPLACIANS. Recall from Section 2 that $L'_i = L'_i(\Delta, \Delta') := \partial_{i+1}\partial_{i+1}^*$ and $L''_i = L''_i(\Delta, \Delta') := \partial_i^*\partial_i$, so that $L_i = L'_i + L''_i$. Define $\mathbf{s}'_i(\Delta, \Delta')$ and $\mathbf{s}''_i(\Delta, \Delta')$ to be the multiset of eigenvalues of $L'_i(\Delta, \Delta')$ and $L''_i(\Delta, \Delta')$, respectively, arranged in weakly decreasing order.

Following [16], let the equivalence relation $\lambda \overset{\circ}{=} \mu$ on multisets λ and μ denote that λ and μ agree in the multiplicities of all of their *non-zero parts*, *i.e.*, that they coincide except for possibly their number of zeroes. Also let $\lambda \cup \mu$ denote the $\overset{\circ}{=}$ -equivalence class whose non-zero parts are the multiset union of the non-zero parts of λ and μ .

PROPOSITION 4.6. *If (Δ, Δ') is a simplicial pair, then*

$$\mathbf{s}_i(\Delta, \Delta') \overset{\circ}{=} \mathbf{s}''_i(\Delta, \Delta') \cup \mathbf{s}''_{i+1}(\Delta, \Delta').$$

Proof. The proof is identical to the single simplicial complex ($\Delta' = \emptyset$) case in [16, Equation (3.6)], and depends only upon $\partial^2 = 0$ and routine eigenvalue calculations involving adjoints. □

If (Δ, Δ') is a simplicial pair, let

$$\begin{aligned} S''_{(\Delta, \Delta'), i}(q) &:= \sum_{\substack{\lambda \in \mathbf{s}''_i(\Delta, \Delta') \\ \lambda \neq 0}} q^\lambda, \quad \text{and} \\ S''_{(\Delta, \Delta')}(t, q) &:= \sum_i S''_{(\Delta, \Delta'), i-1}(q)t^i. \end{aligned}$$

Zero eigenvalues are omitted from these definitions of S'' in order to more naturally encode Proposition 4.6 into the language of generating functions, in Lemma 4.7, below. Also let

$$B_{(\Delta, \Delta')}(t) := \sum_i \tilde{\beta}_{i-1}(\Delta, \Delta') t^i = \sum_i m_0(L_{i-1}(\Delta, \Delta')) t^i = S_{(\Delta, \Delta')}(t, 0).$$

These three definitions of B are equivalent by Proposition 2.1.

From now on, when there is no confusion about the variables t and q , we will often omit them for clarity.

LEMMA 4.7. *If (Δ, Δ') is a simplicial pair, then*

$$S_{(\Delta, \Delta')} = (1 + t^{-1}) S''_{(\Delta, \Delta')} + B_{(\Delta, \Delta')}.$$

Proof. Combine Propositions 4.6 and 2.1. □

COROLLARY 4.8. *If Δ is any simplicial complex, and e is any vertex of Δ , then the spectral recursion holds when $t = -1$.*

Proof. By Lemma 4.7, for any simplicial pair (Δ, Δ') ,

$$S_{(\Delta, \Delta')}(-1, q) = B_{(\Delta, \Delta')}(-1, q) = \sum_i (-1)^i \tilde{\beta}_i(\Delta, \Delta') = \chi(\Delta, \Delta'),$$

where $\chi(\Delta, \Delta')$ denotes the Euler characteristic of the simplicial pair (Δ, Δ') (see e.g., [30]). The identity $\chi(\Delta, \Delta') = \chi(\Delta) - \chi(\Delta')$, which holds as long as $\Delta' \subseteq \Delta$, immediately reduces the $t = -1$ instance of the spectral recursion to $\chi(\Delta) = \chi(\Delta - e) - \chi(\Delta/e)$. This, in turn, follows from $\chi(\Delta) = \sum_i (-1)^i f_i(\Delta)$ and equation (4). □

If Δ is a simplicial complex, define

$$F_{\Delta}(t) := \sum_i f_{i-1}(\Delta) t^i.$$

If $\phi(q)$ is a function of q , define

$$D_q \phi := \phi(q) - \phi(1).$$

The point of D_q is that it helps us convert from B and homology (the effect on which of taking skeleta is hard to describe) to F and f -vectors (the effect on which of taking skeleta is easy to describe) in the following lemma.

LEMMA 4.9. *If $\Delta \subseteq \Delta'$ are simplicial complexes, then*

$$S_{(\Delta, \Delta')} = (1 + t^{-1}) D_q S''_{(\Delta, \Delta')} + F_{\Delta} - F_{\Delta'}.$$

Proof. By Lemma 4.7,

$$F_{\Delta}(t) - F_{\Delta'}(t) = S_{(\Delta, \Delta')}(t, 1) = (1 + t^{-1}) S''_{(\Delta, \Delta')}(t, 1) + B_{(\Delta, \Delta')}(t).$$

Thus

$$B_{(\Delta, \Delta')}(t) = -(1 + t^{-1}) S''_{(\Delta, \Delta')}(t, 1) + F_{\Delta}(t) - F_{\Delta'}(t),$$

which, when plugged back into Lemma 4.7, yields the desired result. □

4.3. SKELETA. Recall the s -skeleton of a simplicial complex Δ is

$$\Delta^{(s)} := \{F \in \Delta : \dim F \leq s\}.$$

Also recall that a simplicial complex is *pure* if all its facets have the same dimension. The *pure s -skeleton* of a simplicial complex Δ is

$$\Delta^{[s]} := \{F \in \Delta : F \subseteq G, G \in \Delta, \dim G = s\}.$$

In other words, $\Delta^{[s]}$ is the subcomplex of Δ consisting of the s -dimensional faces of Δ , and all their subfaces. (See [8, Definition 2.8].) The results of the following lemma are easy exercises.

LEMMA 4.10. *If Δ is a simplicial complex and e is a vertex of Δ , then*

- (1) $(\Delta - e)^{(s)} = \Delta^{(s)} - e$;
- (2) $(\Delta/e)^{(s-1)} = \Delta^{(s)}/e$;
- (3) $\Delta_s = (\Delta^{[s]})_s$;
- (4) $(\Delta - e)_s = (\Delta^{[s]} - e)_s$; and
- (5) $(\Delta/e)_{s-1} = (\Delta^{[s]}/e)_{s-1}$.

LEMMA 4.11. *If $\dim \Delta' \leq d - 1$, then*

$$\mathbf{s}''_{d-1}(\Delta, \Delta') \stackrel{\circ}{=} \mathbf{s}''_{d-1}(\Delta^{(d-1)}, \Delta'^{(d-2)}).$$

Proof. Since Δ and $\Delta^{(d-1)}$ agree in dimensions $d - 1$ and below,

$$\mathbf{s}''_{d-1}(\Delta, \Delta') = \mathbf{s}''_{d-1}(\Delta^{(d-1)}, \Delta').$$

Next, replacing Δ' by $\Delta'^{(d-2)}$ in $(\Delta^{(d-1)}, \Delta')$ has the effect of adding $(d - 1)$ -dimensional faces (in fact, all the $(d - 1)$ -dimensional faces of Δ') to the simplicial pair, all of whose boundary faces are still not present in the simplicial pair, since $\dim \Delta' \leq d - 1$. Thus

$$\partial_{(\Delta^{(d-1)}, \Delta'^{(d-2)}); d-1} = \partial_{(\Delta^{(d-1)}, \Delta'); d-1} \oplus 0$$

(equivalently, the matrices representing the two boundary operators differ only in some additional zero columns); *cf.* proof of Lemma 5.1. It is then easy to check that, since $L''_{d-1} = \partial_{d-1}^* \partial_{d-1}$,

$$L''_{d-1}(\Delta^{(d-1)}, \Delta'^{(d-2)}) = L''_{d-1}(\Delta^{(d-1)}, \Delta') \oplus 0,$$

and so

$$\mathbf{s}''_{d-1}(\Delta^{(d-1)}, \Delta'^{(d-2)}) \stackrel{\circ}{=} \mathbf{s}''_{d-1}(\Delta^{(d-1)}, \Delta').$$

□

COROLLARY 4.12. *If $\dim \Delta' \leq d - 1$, then*

$$S''_{(\Delta, \Delta'), d-1} = S''_{(\Delta^{(d-1)}, \Delta'^{(d-2)}), d-1}$$

COROLLARY 4.13. *If $\dim \Delta \leq d$ and $\dim \Delta' \leq d - 1$, then*

$$S''_{(\Delta^{(d-1)}, \Delta'^{(d-2)})} = S''_{(\Delta, \Delta')} - S''_{(\Delta, \Delta'), d} t^{d+1}.$$

Proof. Clearly, (Δ, Δ') and $(\Delta^{(d-1)}, \Delta'^{(d-2)})$ agree in dimensions $d - 2$ and below. Corollary 4.12 thus ensures $S''_{(\Delta^{(d-1)}, \Delta'^{(d-2)})} = \sum_{i \leq d} S''_{(\Delta, \Delta'), i-1} t^i$. Then simply note, since $\dim \Delta \leq d$, that $S''_{(\Delta, \Delta'), d} t^{d+1}$ is the only remaining term from $S''_{(\Delta, \Delta')}$ not found in $S''_{(\Delta^{(d-1)}, \Delta'^{(d-2)})}$. \square

4.4. THE SPECTRAL RECURSION AND SKELETA.

LEMMA 4.14. *If $\dim \Delta \leq d$, $\dim \Delta' \leq d - 1$, and $\Delta' \subseteq \Delta$, then*

$$\begin{aligned} & S_{(\Delta^{(d-1)}, \Delta'^{(d-2)})} \\ &= S_{(\Delta, \Delta')} - (f_d(\Delta) + f_{d-1}(\Delta))t^{d+1} - (D_q S''_{(\Delta, \Delta'), d} - f_{d-1}(\Delta'))(t^{d+1} + t^d). \end{aligned}$$

Proof. First use the definition of D_q and Corollary 4.13 to get

$$(16) \quad D_q S''_{(\Delta^{(d-1)}, \Delta'^{(d-2)})} = D_q S''_{(\Delta, \Delta')} - (D_q S''_{(\Delta, \Delta'), d})t^{d+1}.$$

Then apply Lemma 4.9 (twice) and equation (16) to compute

$$\begin{aligned} (17) \quad & S_{(\Delta^{(d-1)}, \Delta'^{(d-2)})} \\ &= S_{(\Delta, \Delta')} - (F_\Delta - F_{\Delta'}) - (t^d + t^{d+1})D_q S''_{(\Delta, \Delta'), d} \\ &\quad + (F_{\Delta^{(d-1)}} - F_{\Delta'^{(d-2)}}) \\ (18) \quad &= S_{(\Delta, \Delta')} - (f_d(\Delta)t^{d+1} - f_{d-1}(\Delta')t^d) - (t^d + t^{d+1})D_q S''_{(\Delta, \Delta'), d}. \end{aligned}$$

The lemma now follows by adding the quantity $(t^d + t^{d+1})f_{d-1}(\Delta')$ to the middle term of the right hand side of equation (18), while subtracting it from the last term. \square

If Δ is a simplicial complex and e is a vertex of Δ , let

$$\begin{aligned} \mathcal{S}_{\Delta, e} &:= S_\Delta - (qS_{\Delta-e} + qtS_{\Delta/e} + (1-q)S_{(\Delta-e, \Delta/e)}), \\ \mathcal{S}_{\Delta, e}^d &:= S''_{\Delta, d} - (qS''_{\Delta-e, d} + qS''_{\Delta/e, d-1} + (1-q)S''_{(\Delta-e, \Delta/e), d}), \text{ and} \\ \mathcal{D}_{\Delta, e}^d &:= D_q \mathcal{S}_{\Delta, e}^d + (1-q)f_{d-1}(\Delta/e). \end{aligned}$$

We have defined $\mathcal{S}_{\Delta, e}$ precisely so that Δ satisfies the spectral recursion with respect to e if and only if $\mathcal{S}_{\Delta, e} = 0$, and we have defined $\mathcal{S}_{\Delta, e}^d$ to be the d -dimensional finer Laplacian version of $\mathcal{S}_{\Delta, e}$. The significance of \mathcal{D} is made apparent by the next lemma, which is the last key step to proving Theorem 4.18.

LEMMA 4.15. *If $\dim \Delta \leq d$ and e is a vertex of Δ , then*

$$\mathcal{S}_{\Delta^{(d-1)}, e} = \mathcal{S}_{\Delta, e} - (t^d + t^{d+1})\mathcal{D}_{\Delta, e}^d.$$

Proof. Since $\dim \Delta \leq d$, then $\dim \Delta - e \leq d$ and $\dim \Delta/e \leq d - 1$. Therefore

$$\begin{aligned} & \mathcal{S}_{\Delta^{(d-1)},e} \\ &= \mathcal{S}_{\Delta^{(d-1)}} - q\mathcal{S}_{(\Delta-e)^{(d-1)}} - qt\mathcal{S}_{(\Delta/e)^{(d-2)}} - (1-q)\mathcal{S}_{((\Delta-e)^{(d-1)},(\Delta/e)^{(d-2)})} \\ &= \mathcal{S}_{\Delta} - q\mathcal{S}_{\Delta-e} - qt\mathcal{S}_{\Delta/e} - (1-q)\mathcal{S}_{(\Delta-e,\Delta/e)} \\ &\quad - f_d(\Delta)t^{d+1} + qf_d(\Delta-e)t^{d+1} + qtf_{d-1}(\Delta/e)t^d \\ &\quad + (1-q)(f_d(\Delta-e) + f_{d-1}(\Delta/e))t^{d+1} \\ &\quad - D_q\mathcal{S}''_{\Delta,d}(t^{d+1} + t^d) + qD_q\mathcal{S}''_{\Delta-e,d}(t^{d+1} + t^d) + qtD_q\mathcal{S}''_{\Delta/e,d-1}(t^d + t^{d-1}) \\ &\quad + (1-q)(D_q\mathcal{S}''_{(\Delta-e,\Delta/e),d} - f_{d-1}(\Delta/e))(t^{d+1} + t^d). \end{aligned}$$

The first equation above is by the definition of \mathcal{S} and Lemma 4.10. The second equation involves expanding each term of the left-hand side by Lemma 4.14, and then regrouping like terms. Now, the second line and third lines of this last expression add up to zero, by equation (4). The lemma then follows from the definitions of \mathcal{S} and \mathcal{S}^d . □

LEMMA 4.16. *If $\dim \Delta \leq d$ and e is a vertex of Δ , then $\mathcal{S}_{\Delta,e} = 0$ implies $\mathcal{D}^d_{\Delta,e} = 0$.*

Proof. It is easy to see that $\mathcal{S}_{\Delta^{(d-1)},e}$ has no power of t higher than $\dim \Delta^{(d-1)} + 1 = d$. But since $\mathcal{S}_{\Delta,e} = 0$, Lemma 4.15 implies that $0 = [t^{d+1}]\mathcal{S}_{\Delta^{(d-1)},e} = \mathcal{D}^d_{\Delta,e}$. Here, we are using the coefficient notation $[t^i](\sum_j a_j t^j) := a_i$. □

LEMMA 4.17. *If Δ is a simplicial complex and e is a vertex of Δ , then*

$$\mathcal{D}^d_{\Delta,e} = \mathcal{D}^d_{\Delta^{[d]},e}.$$

Proof. By expanding $\mathcal{D}^d_{\Delta,e}$ we need only show that we may replace Δ by $\Delta^{[d]}$ in each of $\mathcal{S}''_{\Delta,d}$, $\mathcal{S}''_{\Delta-e,d}$, $\mathcal{S}''_{\Delta/e,d-1}$, $\mathcal{S}''_{(\Delta-e,\Delta/e),d}$, and $f_{d-1}(\Delta/e)$. But this follows from Lemma 4.10 and the definition of \mathcal{S}'' . □

THEOREM 4.18. *If $\dim \Delta \leq d$, and e is a vertex of Δ , then Δ satisfies the spectral recursion with respect to e iff $\Delta^{(d-1)}$ and $\Delta^{[d]}$ do as well.*

Proof. First assume Δ satisfies the spectral recursion with respect to e . Then $0 = \mathcal{S}_{\Delta,e}$. By Lemma 4.16, then, $\mathcal{D}^d_{\Delta,e} = 0$. And then by Lemma 4.15, $\mathcal{S}_{\Delta^{(d-1)},e} = 0$. Furthermore, $\mathcal{D}^d_{\Delta^{[d]},e} = \mathcal{D}^d_{\Delta,e} = 0$, by Lemma 4.17.

Conversely, assume $\Delta^{(d-1)}$ and $\Delta^{[d]}$ satisfy the spectral recursion with respect to e . By Lemma 4.16, then $\mathcal{D}^d_{\Delta^{[d]},e} = 0$. And then by Lemmas 4.17 and 4.15,

$$\mathcal{S}_{\Delta,e} = \mathcal{S}_{\Delta^{(d-1)},e} + (t^d + t^{d+1})\mathcal{D}^d_{\Delta,e} = \mathcal{S}_{\Delta^{(d-1)},e} + (t^d + t^{d+1})\mathcal{D}^d_{\Delta^{[d]},e} = 0.$$

□

COROLLARY 4.19. *If $\dim \Delta \leq d$, then Δ satisfies the spectral recursion iff $\Delta^{(d-1)}$ and $\Delta^{[d]}$ do as well.*

4.5. **SHIFTED COMPLEXES.** Recall a k -set is a set with k elements, and a k -family over ground set E is a collection of k -subsets of E . For a k -set F , let $\text{bd } F$ denote the $(k-1)$ -family of all $(k-1)$ -subsets of F . For a k -family \mathcal{K} , its *unsigned boundary* $\text{bd } \mathcal{K}$ is the $(k-1)$ -family $\cup_{F \in \mathcal{K}} \text{bd } F$.

If $F = \{f_1 < \dots < f_k\}$ and $G = \{g_1 < \dots < g_k\}$ are k -subsets of integers, then $F \leq_P G$ under the *componentwise partial order* if $f_p \leq g_p$ for all p . A k -family \mathcal{K} is *shifted* if $F \leq_P G$ and $G \in \mathcal{K}$ together imply that $F \in \mathcal{K}$. A simplicial complex Δ is *shifted* if Δ_i is shifted for every i . The useful properties of shifted families in the following lemma are easy to verify.

LEMMA 4.20. *If \mathcal{K}_1 and \mathcal{K}_2 are shifted families, then so are $\text{bd } \mathcal{K}_1$ and $\mathcal{K}_1 \cap \mathcal{K}_2$.*

We say that Δ is a *near-cone with apex 1* if $\text{bd}(\Delta-1) \subseteq \Delta/1$, where bd denotes the usual unsigned boundary complex consisting of all faces that are not facets. Equivalently, Δ is a near-cone with apex 1 if $F - v \cup 1 \in \Delta$ whenever $F \in \Delta$, $1 \notin F$, and $v \in F$. (See, e.g., [7] for more on near-cones.) We omit the easy proofs of the following two lemmas.

LEMMA 4.21. *Let Δ be a simplicial complex on $[n]$. Then Δ is shifted if and only if Δ is a near-cone with apex 1, and both $\Delta-1, \Delta/1$ are shifted with respect to the ordered vertex set $[2, n]$.*

LEMMA 4.22. *If Δ is a pure d -dimensional near-cone with apex 1, then*

$$\Delta = (1 * (\Delta - 1))^{(d)}$$

THEOREM 4.23. *If Δ is a shifted simplicial complex, then Δ satisfies the spectral recursion, equation (2).*

Proof. The proof is by induction on the dimension and number of vertices of Δ . The base cases, when $\dim \Delta = 0$ or Δ has one vertex (a special case of $\dim \Delta = 0$, anyway) are easy to check.

Assume $\dim \Delta = d \geq 1$. By induction, $\Delta^{(d-1)}$ satisfies the spectral recursion. By Corollary 4.19, it remains to show that $\Delta^{[d]}$ satisfies the spectral recursion as well.

To this end, first note that Δ_d , the family of facets of $\Delta^{[d]}$, is shifted; then, by Lemma 4.20 and reverse induction on dimension, $\Delta^{[d]}$ is shifted. By definition, $\Delta^{[d]}$ is also pure, so Lemma 4.22 implies

$$\Delta^{[d]} = (1 * (\Delta^{[d]} - 1))^{(d)}.$$

Since $\Delta^{[d]}$ is shifted, $\Delta^{[d]} - 1$ is also shifted, with one less vertex, and so satisfies the spectral recursion, by induction. Thus $1 * (\Delta^{[d]} - 1)$ also satisfies the spectral recursion by Proposition 2.2 and Corollary 4.5. Then Corollary 4.19 guarantees that $\Delta^{[d]}$ satisfies the spectral recursion. \square

5. ARBITRARY SHIFTED SIMPLICIAL PAIRS

Merris [29] found a simple description of the Laplacian spectrum of a shifted graph (2-family), in terms of the degree sequence of the graph. This was generalized in [16] to shifted families, by suitably generalizing the notion of

degree sequence. In this section, we extend both the theorem, and the notion of degree sequence, to shifted family *pairs* (Theorem 5.7). As in [16], the technique is to find identical recursive formulas, similar to those in [16], for the Laplacian spectrum (Corollary 5.4) and the generalized degree sequence (Lemma 5.6), in Subsections 5.1 and 5.2, respectively. The two threads are tied together with the proof of Theorem 5.7 in Subsection 5.3. Along the way, we rely upon tools developed in Section 4.

Groene and Merris [22] conjectured that Merris' description of the spectrum of a shifted graph becomes a majorization inequality for an *arbitrary* graph. This was also generalized from graphs to families (though still not proved) in [16]. In Subsection 5.3, we also further extend this conjecture from families to family pairs (Conjecture 5.8).

5.1. LAPLACIANS. Recall the definition of family in Subsection 4.5. If (for some k), \mathcal{K} and \mathcal{K}' are a k -family and $(k-1)$ -family, respectively, on the same ground set of vertices, then we will say $(\mathcal{K}, \mathcal{K}')$ is a *family pair*, but we set $(\mathcal{K}, \mathcal{K}') = (\mathcal{K}, \mathcal{K}'')$ when $(\text{bd } \mathcal{K}) \cap \mathcal{K}' = (\text{bd } \mathcal{K}) \cap \mathcal{K}''$ (more formally, then, a family pair is an equivalence class on ordered pairs of families). We will say $(\mathcal{K}, \mathcal{K}')$ is a *shifted family pair* when \mathcal{K} is shifted and $(\mathcal{K}, \mathcal{K}') = (\mathcal{K}, \mathcal{K}'')$ for some \mathcal{K}'' that is shifted on the same ordered ground set as \mathcal{K} .

Let $C(\mathcal{K}; \mathbb{R})$ denote the *oriented chains* of k -family \mathcal{K} , *i.e.*, the formal \mathbb{R} -linear sums of oriented faces $[F]$ such that $F \in \mathcal{K}$. If $(\mathcal{K}, \mathcal{K}')$ is a family pair, then the *boundary operator* $\partial_{(\mathcal{K}, \mathcal{K}')}: C(\mathcal{K}; \mathbb{R}) \rightarrow C((\text{bd } \mathcal{K}) \setminus \mathcal{K}'; \mathbb{R})$ is defined as it is for simplicial complexes, except that the sum is now restricted to faces in $(\text{bd } \mathcal{K}) \setminus \mathcal{K}'$. Equivalently, $\partial_{(\mathcal{K}, \mathcal{K}')} = \partial_{(\Delta(\mathcal{K}), \Delta(\mathcal{K}'))}; k-1$, when \mathcal{K} is a k -family and \mathcal{K}' is a $(k-1)$ -family. As with simplicial complexes, the boundary operator has an adjoint $\partial_{(\mathcal{K}, \mathcal{K}')}^*$, so the matrices representing ∂ and ∂^* in the natural bases are transposes of one another.

DEFINITION. The *Laplacian* of $(\mathcal{K}, \mathcal{K}')$ is the map $L(\mathcal{K}, \mathcal{K}') : C(\mathcal{K}; \mathbb{R}) \rightarrow C(\mathcal{K}; \mathbb{R})$ defined by

$$L(\mathcal{K}, \mathcal{K}') := \partial_{(\mathcal{K}, \mathcal{K}')}^* \partial_{(\mathcal{K}, \mathcal{K}')}$$

It immediately follows that

$$(19) \quad L(\mathcal{K}, \mathcal{K}') = L''_{k-1}(\Delta(\mathcal{K}), \Delta(\mathcal{K}')),$$

where $\Delta(\mathcal{K})$ denote the pure $(k-1)$ -dimensional simplicial complex whose facets are the members of k -family \mathcal{K} .

It should be clear that $\partial_{(\mathcal{K}, \mathcal{K}')}$, and hence $L(\mathcal{K}, \mathcal{K}')$, is well-defined on family pairs; that is, $\partial_{(\mathcal{K}, \mathcal{K}')} = \partial_{(\mathcal{K}, \mathcal{K}'')}$ and $L(\mathcal{K}, \mathcal{K}') = L(\mathcal{K}, \mathcal{K}'')$, when $(\mathcal{K}, \mathcal{K}') = (\mathcal{K}, \mathcal{K}'')$. Of course, we may always specialize to a single family by letting $\mathcal{K}' = \emptyset$.

Recall that Δ_i denotes the $(i+1)$ -family of i -dimensional faces of simplicial complex Δ .

LEMMA 5.1. *If $\dim \Delta' \leq d-1$, then*

$$L''_d(\Delta, \Delta') = L(\Delta_d, \Delta'_{d-1}).$$

Proof. The boundary maps $\partial_{(\Delta, \Delta'); d}$ and $\partial_{(\Delta_d, \Delta'_{d-1})}$ used to define $L''_d(\Delta, \Delta')$ and $L(\Delta_d, \Delta'_{d-1})$, respectively, both act on $C_d(\Delta; \mathbb{R})$. By the definitions of L and L''_d , then, it will suffice to show that, for any $F \in \Delta_d$,

$$(20) \quad \partial_{(\Delta, \Delta'); d}[F] = \partial_{(\Delta_d, \Delta'_{d-1})}[F].$$

Now, the only difference between the left-hand and right-hand sides of this equation is that the left-hand side is a sum restricted to faces in the set difference $\Delta_{d-1} \setminus \Delta'_{d-1}$, and the right-hand side is a sum restricted to faces in $(\text{bd } \Delta_d) \setminus \Delta'_{d-1}$. Since Δ is a simplicial complex, $\text{bd } \Delta_d \subseteq \Delta_{d-1}$, so the only difference between the two sums is provided by faces in $\Delta_{d-1} \setminus (\text{bd } \Delta_d)$. But any such face will not be in $\text{bd } F$, the unsigned boundary of F , and thus not appear in the expression for the signed boundary map, anyway. (Equivalently, the matrices representing $\partial_{(\Delta, \Delta'); d}$ and $\partial_{(\Delta_d, \Delta'_{d-1})}$ differ only in extra 0 rows indexed by $(d - 1)$ -dimensional faces of Δ not contained in any d -dimensional face of Δ , and these extra 0 rows do not affect $L = \partial^* \partial$.) This establishes equation (20), and hence the lemma. (Cf. the proof of Lemma 4.11). \square

Lemma 5.1, Proposition 4.6, and equation (19) allow us to go back and forth between families and complexes.

LEMMA 5.2. *If (Γ, Γ') is a simplicial pair, then*

$$qtS_{(\Gamma, \Gamma')} = S''_{(1*\Gamma, 1*\Gamma')}.$$

Proof. We compute $S_{(1*\Gamma, 1*\Gamma')}$ in two different ways. Since $1*\Gamma$ and $1*\Gamma'$ are cones, $(1*\Gamma, 1*\Gamma')$ has trivial homology, so $B_{(1*\Gamma, 1*\Gamma')} = 0$. Thus, by Lemma 4.7,

$$S_{(1*\Gamma, 1*\Gamma')} = (1 + t^{-1})S''_{(1*\Gamma, 1*\Gamma')} + B_{(1*\Gamma, 1*\Gamma')} = t^{-1}(1 + t)S''_{(1*\Gamma, 1*\Gamma')}.$$

On the other hand, by Corollary 4.3,

$$S_{(1*\Gamma, 1*\Gamma')} = S_{1*(\Gamma, \Gamma')} = q(1 + t)S_{(\Gamma, \Gamma')}.$$

The lemma now follows immediately. \square

Define $\mathbf{s}(\mathcal{K}, \mathcal{K}')$ to be the multiset of eigenvalues of $L(\mathcal{K}, \mathcal{K}')$, arranged in weakly decreasing order. When $\mathbf{s}(\mathcal{K}, \mathcal{K}')$ consists of non-negative integers, it is a partition. We will use the notation of [28] for partitions, except that we will denote the *conjugate* or *transpose* of partition λ by λ^T . In particular, $1^m = (m)^T$ denotes the partition consisting of m 1's. Recall from Subsection 4.2 the definitions of $\overset{\circ}{=}$ and \cup for multisets, which apply equally well to partitions and weakly decreasing sequences.

Recall the definition of near-cone from subsection 4.5.

LEMMA 5.3. *If $\Delta' \subseteq \Delta$ are pure near-cones with apex 1, and $\dim \Delta = d$ and $\dim \Delta' = d - 1$, then, as partitions,*

$$\mathbf{s}''_d(\Delta, \Delta') \overset{\circ}{=} 1^{f_{d-1}(\Delta/1) - f_{d-1}(\Delta'^{-1})} + (\mathbf{s}''_d(\Delta - 1, \Delta' - 1) \cup \mathbf{s}''_{d-1}(\Delta/1, \Delta'/1)).$$

Proof. Recall the coefficient notation $[t^i](\sum_j a_j t^j) := a_i$. First note

$$(21) \quad [t^i]S''_{(\Gamma, \Gamma')} = S''_{(\Gamma, \Gamma'), i-1}$$

$$(22) \quad [t^i]B_{(\Gamma, \Gamma')} = \tilde{\beta}_{i-1}(\Gamma, \Gamma')$$

for any simplicial pair (Γ, Γ') and for any i . Then, by Lemmas 4.12, 4.22, 5.2, and equation (21),

$$S''_{(\Delta, \Delta'), d} = q[t^d]S_{(\Delta-1, \Delta'-1)},$$

so $s''_d(\Delta, \Delta')$ has just as many non-zero parts as there are terms in $q[t^d]S_{(\Delta-1, \Delta'-1)}$. Lemma 4.7 and equations (21) and (22) now imply

$$\begin{aligned} S''_{(\Delta, \Delta'), d} &= q[t^d]S_{(\Delta-1, \Delta'-1)} \\ &= q(S''_{(\Delta-1, \Delta'-1), d-1} + S''_{(\Delta-1, \Delta'-1), d} + \tilde{\beta}_{d-1}(\Delta-1, \Delta'-1)), \end{aligned}$$

so the non-zero parts of $s''_d(\Delta, \Delta')$ are given by adding 1 to every element of the multiset union of three partitions: $s''_{d-1}(\Delta-1, \Delta'-1)$; $s''_d(\Delta-1, \Delta'-1)$; and the partition consisting of $\tilde{\beta}_{d-1}(\Delta-1, \Delta'-1)$ zeros. This means

$$(23) \quad s''_d(\Delta, \Delta') \doteq 1^m + (s''_{d-1}(\Delta-1, \Delta'-1) \cup s''_d(\Delta-1, \Delta'-1)),$$

where m is the number of terms in $q[t^d]S_{(\Delta-1, \Delta'-1)}$, since we established above that $s''_d(\Delta, \Delta')$ has m non-zero parts. But $\Delta' \subseteq \Delta$ easily implies $\Delta'-1 \subseteq \Delta-1$, and so there are

$$m = f_{d-1}(\Delta-1) - f_{d-1}(\Delta'-1)$$

terms in $q[t^d]S_{(\Delta-1, \Delta'-1)}$.

It is easy to verify that, since Δ and Δ' are pure near-cones (of dimensions d and $d-1$, respectively) with apex 1,

$$(24) \quad (\Delta-1)^{(d-1)} = \Delta/1; \text{ and}$$

$$(25) \quad (\Delta'-1)^{(d-2)} = \Delta'/1.$$

From equation (24), we conclude

$$(26) \quad m = f_{d-1}(\Delta-1) - f_{d-1}(\Delta'-1) = f_{d-1}(\Delta/1) - f_{d-1}(\Delta'/1).$$

From equations (24) and (25), and Lemma 4.11, we conclude

$$(27) \quad s''_{d-1}(\Delta-1, \Delta'-1) \doteq s''_{d-1}(\Delta/1, \Delta'/1).$$

The lemma now follows from equations (23), (26), and (27). □

DEFINITION. Let \mathcal{K} be a k -family on ground set E , and $e \in E$. Then the *deletion* of \mathcal{K} with respect to e is the k -family

$$\mathcal{K} - e = \{F \in \mathcal{K} : e \notin F\}$$

on ground set $E - e$, and the *contraction* of \mathcal{K} with respect to e is the $(k-1)$ -family

$$\mathcal{K}/e = \{F - e : F \in \mathcal{K}, e \in F\}$$

on ground set $E - e$.

The following identities are immediate: $(\Delta(\mathcal{K}) - e)_{k-1} = \mathcal{K} - e$, $(\Delta(\mathcal{K})/e)_{k-2} = \mathcal{K}/e$, and $\Delta(\mathcal{K})_{k-1} = \mathcal{K}$.

Define a k -family to be a *near-cone with apex 1* when $\text{bd}(\mathcal{K} - 1) \subseteq \mathcal{K}/1$. It is an easy exercise to verify that \mathcal{K} is a near-cone iff $\Delta(\mathcal{K})$ is a near-cone. Also, as with simplicial complexes (Lemma 4.21), \mathcal{K} is shifted iff \mathcal{K} is a near-cone with apex 1 such that $\mathcal{K} - 1$ and $\mathcal{K}/1$ are shifted. The following corollary generalizes [16, Lemma 5.3].

COROLLARY 5.4. *If \mathcal{K} and \mathcal{K}' are near-cone families with apex 1 such that $\mathcal{K}' \subseteq \text{bd} \mathcal{K}$, then*

$$s(\mathcal{K}, \mathcal{K}') \doteq 1^{|\mathcal{K}/1| - |\mathcal{K}' - 1|} + (s(\mathcal{K} - 1, \mathcal{K}' - 1) \cup s(\mathcal{K}/1, \mathcal{K}'/1)).$$

Proof. Say \mathcal{K} is a k -family, so \mathcal{K}' is a $(k - 1)$ -family. Let $\Delta = \Delta(\mathcal{K})$ and $\Delta' = \Delta(\mathcal{K}')$. From $\mathcal{K}' \subseteq \text{bd} \mathcal{K}$, it follows that $\Delta' \subseteq \Delta$. Then, by Lemmas 5.1 and 5.3,

$$\begin{aligned} s(\mathcal{K}, \mathcal{K}') &= s(\Delta_{k-1}, \Delta'_{k-2}) = s''_{k-1}(\Delta, \Delta') \\ &\doteq 1^{f_{k-2}(\Delta/1) - f_{k-2}(\Delta' - 1)} + (s''_{k-1}(\Delta - 1, \Delta' - 1) \cup s''_{k-2}(\Delta/1, \Delta'/1)) \\ &= 1^{f_{k-2}(\Delta/1) - f_{k-2}(\Delta' - 1)} \\ &\quad + (s((\Delta - 1)_{k-1}, (\Delta' - 1)_{k-2}) \cup s((\Delta/1)_{k-2}, (\Delta'/1)_{k-3})) \\ &= 1^{|\mathcal{K}/1| - |\mathcal{K}' - 1|} + (s(\mathcal{K} - 1, \mathcal{K}' - 1) \cup s(\mathcal{K}/1, \mathcal{K}'/1)). \end{aligned}$$

□

5.2. DEGREE SEQUENCES.

NOTATION. We will write $F - \lambda$ to denote the set difference $F \setminus \{\lambda\}$, with the implicit assumption that $\lambda \in F$, just as writing $F \dot{\cup} \mu$ carries the implicit assumption that $\mu \notin F$. For instance, $\{F \in \mathcal{K} : F - \lambda \notin \mathcal{K}'\}$ in the following definition is shorthand for $\{F \in \mathcal{K} : \lambda \in F, F \setminus \{\lambda\} \notin \mathcal{K}'\}$.

DEFINITION. Let $(\mathcal{K}, \mathcal{K}')$ be a family pair on ground set E . Define the *degree* of λ in $(\mathcal{K}, \mathcal{K}')$ by

$$d_\lambda(\mathcal{K}, \mathcal{K}') := |\{F \in \mathcal{K} : F - \lambda \notin \mathcal{K}'\}|.$$

It is easy to see that d_λ is well-defined on family pairs; that is, $d_\lambda(\mathcal{K}, \mathcal{K}') = d_\lambda(\mathcal{K}, \mathcal{K}'')$ when $(\mathcal{K}, \mathcal{K}') = (\mathcal{K}, \mathcal{K}'')$. The *degree sequence* $\mathbf{d} = \mathbf{d}(\mathcal{K}, \mathcal{K}')$ is the partition whose parts are $\{d_\lambda : \lambda \in E\}$.

In other words, to find the degree sequence of $(\mathcal{K}, \mathcal{K}')$, label all the edges in the Hasse diagram of $\Delta(\mathcal{K})$ in the natural way, by the vertex being added; then d_λ counts the number of edges in the Hasse diagram labelled λ , and connecting a face in \mathcal{K} with a face in $(\text{bd} \mathcal{K}) \setminus \mathcal{K}'$. When $\mathcal{K}' = \emptyset$, then $\mathbf{d}(\mathcal{K}) = \mathbf{d}(\mathcal{K}, \emptyset)$ is the generalized degree sequence of family \mathcal{K} defined in [16, Section 2]. It is also easy to see that $d_\lambda(\mathcal{K}) = |\mathcal{K}/\lambda|$. When \mathcal{K} is the set of edges of a graph, then $\mathbf{d}(\mathcal{K})$ is the usual degree sequence of a graph.

LEMMA 5.5. *If $(\mathcal{K}, \mathcal{K}')$ is a shifted family pair on $[1, n]$ and $1 \leq \lambda < \mu \leq n$, then $d_\lambda(\mathcal{K}, \mathcal{K}') \geq d_\mu(\mathcal{K}, \mathcal{K}')$; i.e.*

$$\mathbf{d}(\mathcal{K}, \mathcal{K}') = (d_1(\mathcal{K}, \mathcal{K}'), d_2(\mathcal{K}, \mathcal{K}'), \dots, d_n(\mathcal{K}, \mathcal{K}')).$$

In other words, the ordering of the degrees of the degree sequence of a shifted family pair is given by the linear ordering of their vertices.

Proof. It will suffice to find an injection from $\{F \in \mathcal{K} : F - \mu \notin \mathcal{K}'\}$, a set whose cardinality equals $d_\mu(\mathcal{K}, \mathcal{K}')$, into $\{F \in \mathcal{K} : F - \lambda \notin \mathcal{K}'\}$, a set whose cardinality equals $d_\lambda(\mathcal{K}, \mathcal{K}')$. It is easy to verify, using that \mathcal{K} and \mathcal{K}' are shifted, that such an injection ϕ is given by

$$\phi(F) = \begin{cases} F & \text{if } \lambda \in F \\ F - \mu \dot{\cup} \lambda & \text{if } \lambda \notin F. \end{cases}$$

□

The following lemma generalizes [16, Lemma 5.2]

LEMMA 5.6. *If \mathcal{K} and \mathcal{K}' are shifted families on ground set $[1, n]$, and $\mathcal{K}' \subseteq \text{bd } \mathcal{K}$, then, as partitions,*

$$\mathbf{d}(\mathcal{K}, \mathcal{K}')^T = 1^{|\mathcal{K}/1| - |\mathcal{K}' - 1|} + (\mathbf{d}(\mathcal{K} - 1, \mathcal{K}' - 1)^T \cup \mathbf{d}(\mathcal{K}/1, \mathcal{K}'/1)^T).$$

Proof. By standard partition arguments, this reduces to showing

$$\mathbf{d}(\mathcal{K}, \mathcal{K}') = (|\mathcal{K}/1| - |\mathcal{K}' - 1|) \cup (\mathbf{d}(\mathcal{K} - 1, \mathcal{K}' - 1) + \mathbf{d}(\mathcal{K}/1, \mathcal{K}'/1)),$$

which is a direct consequence of the following two facts:

- $d_1(\mathcal{K}, \mathcal{K}') = |\mathcal{K}/1| - |\mathcal{K}' - 1|$; and
- if $\lambda > 1$, then $d_\lambda(\mathcal{K}, \mathcal{K}') = d_\lambda(\mathcal{K} - 1, \mathcal{K}' - 1) + d_\lambda(\mathcal{K}/1, \mathcal{K}'/1)$.

The indexing on the second fact is indeed what is necessary, thanks to Lemma 5.5, because $\mathcal{K} - 1$, $\mathcal{K}' - 1$, $\mathcal{K}/1$, and $\mathcal{K}'/1$ each have ground set $[2, n]$. Each fact is an easy exercise, the first of which depends upon \mathcal{K} being shifted. □

5.3. A RELATIVE GENERALIZED MERRIS THEOREM. Merris [29, Theorem 2] showed that when \mathcal{K} is the 2-family of edges of a shifted graph, then $\mathbf{s}(\mathcal{K}) \doteq \mathbf{d}(\mathcal{K})^T$. This was generalized in [16, Theorem 1.1] to allow \mathcal{K} to be any shifted family. The main result of this section, below, further generalizes this to shifted family *pairs*. The proof is similar to that of [16, Theorem 1.1].

THEOREM 5.7. *If $(\mathcal{K}, \mathcal{K}')$ is a shifted family pair, then*

$$\mathbf{s}(\mathcal{K}, \mathcal{K}') \doteq \mathbf{d}(\mathcal{K}, \mathcal{K}')^T$$

Proof. By Lemmas 4.20 and 4.21, $(\mathcal{K} - 1, \mathcal{K}' - 1) = (\mathcal{K} - 1, (\mathcal{K}' - 1) \cap \text{bd}(\mathcal{K} - 1))$ and $(\mathcal{K}/1, \mathcal{K}'/1) = (\mathcal{K}/1, (\mathcal{K}'/1) \cap \text{bd}(\mathcal{K}/1))$ are shifted family pairs. Then the result is immediate from Corollary 5.4, Lemmas 4.21 and 5.6, and induction on the number of vertices. □

Grone and Merris [22, Conjecture 2] conjectured that when \mathcal{K} is the 2-family of edges of an *arbitrary* graph, then the equality (modulo zeros) $\mathbf{s}(\mathcal{K}) \doteq \mathbf{d}(\mathcal{K})^T$ above becomes a majorization inequality $\mathbf{s}(\mathcal{K}) \preceq \mathbf{d}(\mathcal{K})^T$, *i.e.*, $\sum_{j=1}^k s_j \leq \sum_{j=1}^k d_j^T$ for all k , where $\mathbf{s}(\mathcal{K}) = (s_1, s_2, \dots)$ and $\mathbf{d}(\mathcal{K})^T = (d_1^T, d_2^T, \dots)$ are written as weakly decreasing sequences. This majorization inequality was also conjectured (but not proved) to hold when \mathcal{K} is any family, in [16, Conjecture 1.2]. Based on no more than a few examples, and that [16, Theorem 1] successfully extends to pairs in Theorem 5.7 above, we extend this conjecture to family pairs as well.

CONJECTURE 5.8. *If $(\mathcal{K}, \mathcal{K}')$ is a family pair, then*

$$\mathbf{s}(\mathcal{K}, \mathcal{K}') \preceq \mathbf{d}(\mathcal{K}, \mathcal{K}')^T.$$

Stephen [33, Theorem 4.3.1] has shown that if the Grone-Merris conjecture is true for all graphs, then Conjecture 5.9 holds for graph pairs (\mathcal{K} is a 2-family and \mathcal{K}' is a 1-family).

REMARK 5.9. Theorem 5.7 suffices to find the spectrum of a shifted simplicial pair (that is, a simplicial pair (Δ, Δ') , where Δ and Δ' are each shifted on the same ordered ground set), not just a shifted family pair. To see this, first note that by Proposition 4.6, finding $\mathbf{s}_i''(\Delta, \Delta')$ for all i determines the spectrum of the simplicial pair (Δ, Δ') . Since \mathbf{s}_i'' depends only on i - and $(i-1)$ -dimensional faces, $\mathbf{s}_i''(\Delta, \Delta') = \mathbf{s}_i''(\Delta^{(i)}, \Delta'^{(i)})$. Finally, then, $\mathbf{s}_i''(\Delta, \Delta') = \mathbf{s}_i''(\Delta^{(i)}, \Delta'^{(i)}) = \mathbf{s}_i''(\Delta^{(i)}, \Delta'^{(i-1)}) = \mathbf{s}(\Delta_i, \Delta'_{i-1})$, by Lemmas 4.11 and 5.1.

6. OPERATIONS THAT PRESERVE THE SPECTRAL RECURSION

In this section, we see how the spectral recursion, equation (2), and the spectrum polynomial behave with respect to some natural operators on simplicial complexes. Each operator has significance for, or motivation from, matroids and/or shifted complexes. Our main results are that the property of satisfying the spectral recursion is preserved by disjoint union (Corollary 6.11), Alexander duality (Corollary 6.8), and, with a slight modification allowing order filters as well as simplicial complexes, two other dual operators (Theorems 6.3 and 6.6).

6.1. DUALS. The Tutte polynomial for matroids (see, *e.g.*, [12]) whose recursion ($T_M = T_{M-e} + T_{M/e}$) inspired and resembles the spectral recursion, is well-behaved with respect to matroid duals ($T_M(x, y) = T_{M^*}(y, x)$), so it is natural to ask what duality does to the spectrum polynomial and the spectral recursion. There are three natural involutions on simplicial complexes that are each appropriate generalizations of matroid duality. How these involutions affect the Laplacians of *families* has already been considered in [16, Section 4]. Recall that an *order filter* Ψ with vertices V is a collection of subsets of V , closed under taking supersets; that is, $F \in \Psi$ and $F \subseteq G \subseteq V$ together imply $G \in \Psi$.

DEFINITION. Let Δ be a simplicial complex (respectively, order filter) with vertex set V . The *dual* of Δ is the order filter (respectively, simplicial complex)

$$\Delta^* = \{V - F : F \in \Delta\}.$$

The *complement* of Δ is the order filter (respectively, simplicial complex)

$$\Delta^c = \{F \subseteq V : F \notin \Delta\}.$$

The *Alexander dual* of Δ is the simplicial complex (respectively, order filter)

$$\Delta^\vee = \Delta^{*c} = \Delta^{c*}.$$

The Alexander dual has received attention lately in combinatorial topology (see, *e.g.*, [2, 6]) and in combinatorial commutative algebra (see, *e.g.*, [3, 4, 10, 17]).

It is easy to see that $\Delta^{**} = \Delta^{cc} = \Delta^{\vee\vee} = \Delta$ for every simplicial complex Δ , and similarly for order filters. If we define an order filter Ψ to be shifted when its every family Ψ_i of i -dimensional faces is shifted, then it is easy to see that duality and complementation preserve being shifted, though with the reverse vertex order. Consequently, Alexander duality preserves being shifted.

If Ψ and Ψ' are order filters on the same ground set of vertices, we define the *order filter pair* (Ψ, Ψ') to be the simplicial pair (Ψ'^c, Ψ^c) , as defined in Section 2. (This means that, more formally, an order filter pair is an equivalence class on ordered pairs of order filters.) Thus $(\Psi, \Psi') = (\Omega, \Omega')$ when the set differences $\Psi \setminus \Psi'$ and $\Omega \setminus \Omega'$ are equal as subsets of the power set of the ground set of vertices. As with simplicial complexes, results and definitions about order filter pairs (Ψ, Ψ') may be specialized to a single order filter, by letting $\Psi' = \emptyset$, the empty order filter.

The definitions of deletion and contraction extend naturally to order filters. The deletion and contraction $\Psi - e$ and Ψ/e of an order filter Ψ on vertex set V are still order filters, though on vertex set $V - e$. In contrast to simplicial complexes, Ψ/e is not necessarily a subset of Ψ (though $\Psi - e \subseteq \Psi$, still), and $\Psi - e \subseteq \Psi/e$ (whereas, for simplicial complexes, $\Delta/e \subseteq \Delta - e$).

We now borrow a trick from [27, Proposition 6] (see also [16, Proposition 4.2]) to investigate how the dual affects Laplacians and the spectral recursion. Let (Δ, Δ') be a simplicial pair with vertex set $[n]$; it is easy to specialize from pairs of duals to a single dual, since the dual of the empty simplicial complex is again empty, so $\Delta^* = (\Delta^*, \emptyset) = (\Delta^*, \emptyset^*)$. Define $\phi_i(\Delta, \Delta') : C_i(\Delta, \Delta'; \mathbb{R}) \rightarrow C_{n-i-2}(\Delta^*, \Delta'^*; \mathbb{R})$ to be the \mathbb{R} -linear isomorphism induced by

$$\phi_i(\Delta, \Delta') : [F] \mapsto \sigma(F)[\overline{F}],$$

where $\sigma(F) = (-1)^{\sum_{j \in F} j}$, and $\overline{F} = [n] - F$.

LEMMA 6.1. *Let (Δ, Δ') be a simplicial pair with vertex set $[n]$, and let $\phi_j = \phi_j(\Delta, \Delta')$ for any j . Then*

- (1) $\phi_{i+1}^{-1} \partial_{(\Delta^*, \Delta'^*); n-i-2} \phi_i = -\partial_{(\Delta, \Delta'); i+1}^*$, and
- (2) $\phi_i \partial_{(\Delta, \Delta'); i+1} \phi_{i+1}^{-1} = -\partial_{(\Delta^*, \Delta'^*); n-i-2}^*$.

Proof. These are each a routine check of signs. □

COROLLARY 6.2. *Let (Δ, Δ') be a simplicial pair with vertex set $[n]$, and let $\phi_j = \phi_j(\Delta, \Delta')$ for any j . Then*

$$L_i(\Delta, \Delta') = \phi^{-1}L_{n-i-2}(\Delta^*, \Delta'^*)\phi.$$

An immediate corollary is that, as first conjectured by V. Reiner (personal communication),

$$(28) \quad \mathbf{s}_i(\Delta, \Delta') = \mathbf{s}_{n-i-2}(\Delta^*, \Delta'^*),$$

which translates into generating functions as

$$(29) \quad S_{(\Delta^*, \Delta'^*)}(t, q) = t^n S_{(\Delta, \Delta')}(t^{-1}, q).$$

We might hope that, if simplicial complex Δ satisfies the spectral recursion with respect to a vertex e , then Δ^* would, too, but this is not quite true. Routine calculations using equation (29), and duality identities $(\Delta - e)^* = \Delta^*/e$ and $(\Delta/e)^* = \Delta^* - e$, show that

$$(30) \quad S_{\Delta^*}(t, q) = qtS_{\Delta^*/e}(t, q) + qS_{\Delta^*-e}(t, q) + (1 - q)tS_{(\Delta^*/e, \Delta^*-e)}(t, q).$$

We thus call

$$(31) \quad S_{\Psi} = qS_{\Psi-e}(t, q) + qtS_{\Psi/e}(t, q) + (1 - q)tS_{(\Psi/e, \Psi-e)}(t, q)$$

the spectral recursion for order filters. Theorem 6.6 below provides further evidence that this is the right formulation for order filters. A unified approach to the spectral recursions for simplicial complexes and order filters is to develop a spectral recursion for simplicial complex pairs (which includes simplicial complexes and order filters as special cases), which is explored in [15].

THEOREM 6.3. *If Δ is a simplicial complex and e is an element of its vertex set, then Δ satisfies the spectral recursion with respect to e iff Δ^* satisfies the spectral recursion for order filters, equation (31), with respect to e .*

Proof. The forward implication follows from equation (30) above. The proof of the reverse implication is similar. □

The following proposition is a restatement of [16, Corollary 4.7].

PROPOSITION 6.4. *Let Δ be a simplicial complex with vertex set $[n]$. If $\lambda \neq n$, then $m_{\lambda}(L_i(\Delta)) = m_{\lambda}(L_{n-i-3}(\Delta^{\vee}))$.*

The following corollary was first conjectured by V. Reiner (personal communication).

COROLLARY 6.5. *If Δ is a simplicial complex with vertex set $[n]$, then $\mathbf{s}_{i-1}(\Delta)$ and $\mathbf{s}_i(\Delta^c)$ agree, except for the multiplicity of n .*

Proof. By equation (28), $\mathbf{s}_i(\Delta^c) = \mathbf{s}_i(\Delta^{\vee*}) = \mathbf{s}_{n-i-2}(\Delta^{\vee})$, so, if $\lambda \neq n$, then

$$m_{\lambda}(L_i(\Delta^c)) = m_{\lambda}(L_{n-i-2}(\Delta^{\vee})) = m_{\lambda}(L_{n-3-(n-i-2)}(\Delta)) = m_{\lambda}(L_{i-1})$$

by Proposition 6.4. □

The preceding proof is not as simple as it seems. The proof of Proposition 6.4 in [16, Corollary 4.7] is somewhat involved, and gets to the Alexander dual via the complement. Especially in light of the simplicity of the statement of Corollary 6.5, we might hope it would have a more direct proof that does not call upon the Alexander dual.

Corollary 6.5 translates into generating functions as

$$(32) \quad S_{\Delta^c}(t, q) = tS_{\Delta}(t, q) + q^n A_{\Delta}(t),$$

which we may rewrite as

$$(33) \quad S_{\Delta}(t, q) = t^{-1}S_{\Delta^c}(t, q) - q^n t^{-1}A_{\Delta}(t),$$

where $A_{\Delta}(t)$ is a polynomial in t that depends on Δ .

THEOREM 6.6. *If e is a vertex of simplicial complex Δ , then Δ satisfies the spectral recursion with respect to e iff Δ^c satisfies the spectral recursion for order filters, equation (31), with respect to e .*

Proof. First assume Δ^c satisfies the spectral recursion for order filters with respect to e . Then, we may use equations (32) and (33), and the complement identities $(\Delta - e)^c = \Delta^c - e$ and $(\Delta/e)^c = \Delta^c/e$, to compute

$$(34) \quad S_{\Delta} = qS_{\Delta-e} + qtS_{\Delta/e} + (1-q)S_{(\Delta-e, \Delta/e)} + q^n t^{-1}(A_{\Delta-e} + A_{\Delta/e} - A_{\Delta}).$$

By Lemma 2.4, all simplicial complexes satisfy the spectral recursion when $q = 1$, so plugging $q = 1$ into the above equation yields

$$S_{\Delta}(1, t) = S_{\Delta}(1, t) + 1^n t^{-1}(A_{\Delta-e} + A_{\Delta/e} - A_{\Delta}).$$

Therefore $A_{\Delta-e} + A_{\Delta/e} - A_{\Delta} = 0$, which, when plugged back into equation (34), proves Δ satisfies the spectral recursion with respect to e .

The reverse implication is proved similarly. \square

Theorems 6.3 and 6.6 together imply the corresponding result for Alexander duality:

THEOREM 6.7. *If e is a vertex of simplicial complex Δ , then Δ satisfies the spectral recursion with respect to e iff Δ^{\vee} satisfies the spectral recursion with respect to e .*

COROLLARY 6.8. *If Δ is a simplicial complex, then Δ satisfies the spectral recursion iff Δ^{\vee} does as well.*

6.2. UNION. The Tutte polynomial is well-behaved with respect to matroid direct sum ($T_{M \oplus N} = T_M + T_N$), which corresponds to the union of simplicial complexes with disjoint vertex sets ($IN(M \oplus N) = IN(M) \cup IN(N)$). So it is natural to ask what disjoint union does to the spectrum polynomial and the spectral recursion.

LEMMA 6.9. *If Δ and Γ are two non-empty simplicial complexes with disjoint vertex sets, then*

$$S_{\Delta \cup \Gamma} = S_{\Delta} + S_{\Gamma} + (t^0 + t^1)(q^{n+m} - (q^n + q^m)) + t^1 q^0,$$

where Δ and Γ have $n = f_0(\Delta)$ and $m = f_0(\Gamma)$ non-loop vertices, respectively.

Proof. For $i > 1$, it is clear that $L_{i-1}(\Delta \cup \Gamma) = L_{i-1}(\Delta) \oplus L_{i-1}(\Gamma)$, since no $(i - 1)$ -dimensional face of Δ has any boundary in Γ , and vice versa. Thus

$$\mathbf{s}_{i-1}(\Delta \cup \Gamma) = \mathbf{s}_{i-1}(\Delta) \cup \mathbf{s}_{i-1}(\Gamma)$$

for $i > 1$.

The vertices of Δ and Γ are disjoint, but they *share* the empty face in their boundary. It is easy to see that $\mathbf{s}_0''(\Sigma) \doteq (f_0(\Sigma))$ for any simplicial complex Σ , so $\mathbf{s}_0''(\Delta \cup \Gamma) = (n + m)$, while $\mathbf{s}_0''(\Delta) \cup \mathbf{s}_0''(\Gamma) = (n, m)$. Also, since $\dim L_0(\Delta \cup \Gamma) = f_0(\Delta \cup \Gamma) = n + m = f_0(\Delta) + f_0(\Gamma) = \dim L_0(\Delta) + \dim L_0(\Gamma)$, then $\mathbf{s}_0(\Delta \cup \Gamma)$ and $\mathbf{s}_0(\Delta) \cup \mathbf{s}_0(\Gamma)$ have the same number of parts. By Proposition 4.6, it then follows that

$$\mathbf{s}_0(\Delta \cup \Gamma) \cup (n, m) = \mathbf{s}_0(\Delta) \cup \mathbf{s}_0(\Gamma) \cup (n + m, 0).$$

(In other words, to change $\mathbf{s}_0(\Delta) \cup \mathbf{s}_0(\Gamma)$ into $\mathbf{s}_0(\Delta \cup \Gamma)$, replace (n, m) in $\mathbf{s}_0(\Delta) \cup \mathbf{s}_0(\Gamma)$ by $(n + m, 0)$ in $\mathbf{s}_0(\Delta \cup \Gamma)$.) Similarly, since $\Delta \cup \Gamma$, Δ , and Γ each have exactly one empty face, $\mathbf{s}_{-1}(\Delta \cup \Gamma)$ has one element, and $\mathbf{s}_{-1}(\Delta) \cup \mathbf{s}_{-1}(\Gamma)$ has two elements, and so

$$\mathbf{s}_{-1}(\Delta \cup \Gamma) = (n + m),$$

while

$$\mathbf{s}_{-1}(\Delta) \cup \mathbf{s}_{-1}(\Gamma) = (n, m).$$

The lemma now follows immediately. □

We continue to assume Δ and Γ are non-empty simplicial complexes with disjoint vertex sets, and that Γ has m non-loop vertices. By arguments similar to those in the proof of Lemma 6.9,

$$S_{(\Gamma, \emptyset)} = S_\Gamma - (t^0 + t^1)q^m + t^1q^0,$$

and so

$$(35) \quad S_{(\Delta \cup \Gamma, \Delta')} = S_{(\Delta, \Delta')} + S_\Gamma - (t^0 + t^1)q^m + t^1q^0.$$

THEOREM 6.10. *If Δ satisfies the spectral recursion with respect to e , and Γ is any simplicial complex whose vertex set is disjoint from the vertex set of Δ , then $\Delta \cup \Gamma$ satisfies the spectral recursion with respect to e .*

Proof. If $\Gamma = \emptyset$, then the theorem is trivially true. Otherwise, it is a routine calculation with Lemma 6.9 and equation (35). □

COROLLARY 6.11. *If Δ and Γ each satisfy the spectral recursion, then so does their disjoint union $\Delta \cup \Gamma$.*

The following example shows that the *arbitrary* union of two simplicial complexes satisfying the spectral recursion does not itself necessarily satisfy the spectral recursion, even if both complexes are pure.

EXAMPLE 6.12. Let Δ be the pure 1-dimensional simplicial complex on vertex set $\{a, b, c, d, e\}$ with facets $\{ab, ac, ad, ae, bc, bd\}$. (We omit brackets and commas from each face for clarity.) Let Γ be the pure 1-dimensional simplicial complex on the same vertex set with facets $\{ab, ac, ad, ae, de\}$. Now, Δ is

shifted with vertices ordered $a < b < c < d < e$, and Γ is shifted with vertices ordered $a < d < e < b < c$, so each satisfies the spectral recursion.

On the other hand, we can easily show $\Delta \cup \Gamma$ does not satisfy the spectral recursion with respect to vertex d . First check directly that $\Delta \cup \Gamma$ is not Laplacian integral. (Note that $\Delta \cup \Gamma$ is the 1-dimensional skeleton of the cone over Example 2.5.) Next, since $(\Delta \cup \Gamma) - d$ and $(\Delta \cup \Gamma)/d$ are each isomorphic to shifted complexes (with different vertex orders), they are each Laplacian integral. It is also easy to directly verify that $((\Delta \cup \Gamma) - d, (\Delta \cup \Gamma)/d)$ is Laplacian integral as well. Thus, the right-hand side of the spectral recursion in this instance has all integer exponents, but the left-hand side does not.

7. ACKNOWLEDGEMENTS

I am grateful to Vic Reiner, Eric Babson, Woong Kook, and Tamon Stephen for many helpful comments and suggestions, to Vic Reiner also for introducing me to this problem, and to several referees for their insightful advice.

REFERENCES

- [1] M. Aigner, *Combinatorial theory*, Grundlehren der Mathematischen Wissenschaften, vol. 234, Springer-Verlag, Berlin-New York, 1979.
- [2] E. Babson, A. Björner, S. Linusson, J. Shareshian, and V. Welker, “Complexes of not i -connected graphs”, *Topology* 38 (1999), 271–299.
- [3] D. Bayer, “Monomial ideals and duality”, unpublished lecture notes, 1996, available at http://www.math.columbia.edu/~bayer/papers/Duality_B96.pdf.
- [4] D. Bayer, H. Charalambous, and S. Popescu, “Extremal Betti numbers and applications to monomial ideals”, *J. Alg.* 221 (1999), 497–512.
- [5] A. Björner, “The homology and shellability of matroids and geometric lattices”, in *Matroid applications*, pp. 226–283, Encyclopedia Math. Appl., vol. 40, Cambridge Univ. Press, Cambridge, 1992.
- [6] A. Björner, L. M. Butler, and A. O. Matveev “Note on a combinatorial application of Alexander duality”, *J. Combin. Theory Ser. A* 80 (1997), 163–165.
- [7] A. Björner and G. Kalai, “An extended Euler-Poincaré theorem”, *Acta Math.* 161 (1988), 279–303.
- [8] A. Björner and M. L. Wachs, “Shellable nonpure complexes and posets. I”, *Trans. Amer. Math. Soc.* 348 (1996), 1299–1327.
- [9] A. Björner and M. L. Wachs, “Shellable nonpure complexes and posets. II”, *Trans. Amer. Math. Soc.* 349 (1997), 3945–3975.
- [10] W. Bruns and J. Herzog, “Semigroup rings and simplicial complexes”, *J. Pure Appl. Algebra* 122 (1997), 185–208.
- [11] T. Brylawski, “Constructions”, in *Theory of matroids*, pp. 127–223, Encyclopedia Math. Appl., vol. 26, Cambridge Univ. Press, Cambridge, 1986.

- [12] T. Brylawski and J. Oxley, “The Tutte polynomial and its applications”, in *Matroid applications*, pp. 123–225, Encyclopedia Math. Appl., vol. 40, Cambridge Univ. Press, Cambridge, 1992.
- [13] F. Chung and R. B. Ellis, “A chip-firing game and Dirichlet eigenvalues”, *Discrete Math.* 257 (2002), 341–355.
- [14] X. Dong and M. L. Wachs, “Combinatorial Laplacian of the matching complex”, *Electron. J. Combin.* 9 (2002), #R17, 11 pp.
- [15] A. M. Duval, “A Relative Laplacian spectral recursion”, in preparation.
- [16] A. M. Duval and V. Reiner, “Shifted simplicial complexes are Laplacian integral”, *Trans. Amer. Math. Soc.* 354 (2002), 4313–4344.
- [17] J. A. Eagon and V. Reiner, “Resolutions of Stanley-Reisner rings and Alexander duality”, *J. Pure Appl. Algebra* 130 (1998), 265–275.
- [18] B. Eckmann, “Harmonische Funktionen und Randwertaufgaben in einem Komplex”, *Comment. Math. Helv.* 17 (1945), 240–255.
- [19] G. Etienne and M. Las Vergnas, “External and internal elements of a matroid basis”, *Discrete Math.* 179 (1998), 111–119.
- [20] J. Friedman, “Computing Betti numbers via combinatorial Laplacian”, in *Proceedings of the Twenty-eighth Annual ACM Symposium on the Theory of Computing*, (Philadelphia, 1996), pp. 386–391, ACM, New York, 1996.
- [21] J. Friedman and P. Hanlon, “On the Betti numbers of chessboard complexes”, *J. Alg. Comb.* 8 (1998), 193–203.
- [22] R. Grone and R. Merris, “The Laplacian spectrum of a graph II”, *SIAM J. Disc. Math.* 7 (1994), 221–229.
- [23] A. S. Jarrah and R. Laubenbacher, “Generic Cohen-Macaulay monomial ideals”, *Ann. Comb.* 8 (2004), 45–61.
- [24] C. J. Klivans, “Shifted matroid complexes”, preprint, 2003.
- [25] W. Kook, “A circuit formula for homology of matroids”, preprint, 2000.
- [26] W. Kook, “Recurrence relations for the spectrum polynomial of a matroid”, *Disc. Appl. Math.* 143 (2004), 312–317.
- [27] W. Kook, V. Reiner, and D. Stanton, “Combinatorial Laplacians of matroid complexes”, *J. Amer. Math. Soc.* 13 (2000), 129–148.
- [28] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford University Press, New York, 1979.
- [29] R. Merris, “Degree maximal graphs are Laplacian integral”, *Lin. Alg. Appl.* 199 (1994), 381–389.
- [30] J. R. Munkres, *Elements of algebraic topology*, Addison-Wesley, Menlo Park CA, 1984.
- [31] J. G. Oxley, *Matroid theory*, Oxford University Press, New York, 1992.

- [32] J. S. Provan and L. J. Billera, “Decompositions of simplicial complexes related to diameters of convex polyhedra”, *Math. Oper. Res.* 5 (1980), 576–594.
- [33] T. Stephen, “A Majorization bound for the eigenvalues of some graph Laplacians”, preprint, 2004; [arXiv:math.CO/0411153](https://arxiv.org/abs/math.CO/0411153).
- [34] D. J. A. Welsh, *Matroid theory*, Academic Press, New York, 1976.
- [35] N. White (ed.), *Theory of matroids*, Encyclopedia Math. Appl., vol. 26, Cambridge Univ. Press, Cambridge, 1986.

Art M. Duval
Department of Mathematical Sciences
University of Texas at El Paso
El Paso, TX 79968-0514
USA
artduval@math.utep.edu