

A COMMUTATIVE NOETHERIAN HOPF ALGEBRA OVER A FIELD IS FINITELY GENERATED

RICHARD K. MOLNAR

ABSTRACT. Let k be an arbitrary field and H a commutative Hopf algebra over k . We give a short proof of the fact that H is Noetherian if and only if H is finitely generated as a k -algebra.

In [4] M. Takeuchi has shown that a commutative (or cocommutative) Hopf algebra H is faithfully flat over any sub-Hopf algebra K . He then uses this result to give a simple proof of the following:

(INJ) If H is a commutative Hopf algebra over a field k then the correspondence

$$(\text{sub-Hopf algebras}) \rightarrow (\text{Hopf ideals})$$

$$K \mapsto K^+H$$

is injective.

As a corollary to (INJ) Takeuchi shows [4, 3.11] that any sub-Hopf algebra of a finitely generated, commutative Hopf algebra over a field is finitely generated. The purpose of this note is to show that a commutative Noetherian Hopf algebra over a field is finitely generated, and that this fact follows from (INJ) with essentially the same proof as Takeuchi's corollary.

For any Hopf algebra H over the field k , we let $H^+ = \ker(\epsilon)$ where ϵ is the counit of H , and we denote the antipode of H by S . A commutative Hopf algebra is said to be Noetherian (resp. finitely generated) if it is Noetherian (resp. finitely generated) as a k -algebra. For the basic facts about Hopf algebras see [1] or [3]. Clearly the desired result follows from the following

Proposition. *Let H be a commutative Hopf algebra over a field k . If H^+ is finitely generated as an ideal then H is finitely generated.*

Proof. Let $\{f_1, \dots, f_n\}$ be generators of the ideal H^+ . If C is the subcoalgebra of H generated by the f_i , then C is finite dimensional [1,

Received by the editors December 2, 1974 and, in revised form, April 1, 1975.
AMS (MOS) subject classifications (1970). Primary 16A24; Secondary 13E05.
Key words and phrases. Commutative Hopf algebra, Noetherian algebra.

1.4.3]. Since the antipode of H has order 2, it follows that $D = C + S(C)$ is a finite dimensional subcoalgebra of H which is stable under the antipode, and $\{f_1, \dots, f_n\} \subset D$. If we let $[D]$ be the subalgebra of H generated by D , then, in fact, $[D]$ is a sub-Hopf algebra of H which is finitely generated.

Now $\{f_1, \dots, f_n\} \subset [D] \cap H^+ = [D]^+$, and since the f_i are generators of the ideal H^+ , we must have $H^+ \subset [D]^+H$. On the other hand, $[D]^+H$ is clearly contained in H^+ . Thus $[D]^+H = H^+ = H^+H$, and so by (INJ) we must have $[D] = H$, which gives the desired conclusion.

One may easily generalize this result. For example, by using K. Newman's version of (INJ) for cocommutative Hopf algebras [2, 2.5]¹ along with the proof of the Proposition (which essentially depends only on (INJ) and the fact that the antipode has finite order), one can show that a cocommutative Noetherian Hopf algebra over a field is finitely generated.

REFERENCES

1. R. G. Heyneman and M. E. Sweedler, *Affine Hopf algebras*. I, *J. Algebra* 13 (1969), 192–241. MR 39 #6876.
2. K. Newman, *A correspondence between bi-ideals and sub-Hopf algebras in cocommutative Hopf algebras* (preprint).
3. M. E. Sweedler, *Hopf algebras*, Math. Lecture Note Series, Benjamin, New York, 1969. MR 40 #5705.
4. M. Takeuchi, *A correspondence between Hopf ideals and sub-Hopf algebras*, *Manuscripta Math.* 7 (1972), 251–270. MR 48 #328.

DEPARTMENT OF MATHEMATICS, OAKLAND UNIVERSITY, ROCHESTER, MICHIGAN 48063

¹ We thank the referee for pointing this out.