

HÜLYA ATIL¹ and ADEL SALAH KHATTAB²

A comparison of different methods of estimating sire transmitting ability of some milk traits in a herd of Holstein Friesian cattle

Summary

A total of 1931 normal first lactation records of Holstein Friesian cows kept at Dena Farm in Egypt during the period from 1987 to 1994 were used to estimate phenotypic and genetic parameters for 90 day milk yield (90 dMY), 305 day milk yield (305 dMY) and lactation period (LP). In addition, 76 bulls with at least ten daughters were used to compare three methods of sire transmitting ability.

A least squares analysis of variance show significant effect of month and year of calving and age at first calving for different traits studied, except the effect of age at first calving on LP.

Heritability estimates for 90 dMY, 305 dMY and LP were 0.39 ± 0.08 , 0.27 ± 0.07 and 0.14 ± 0.05 , respectively. Genetic and phenotypic correlations between different traits were positive and significant.

Sires with at least ten daughters were evaluated by best linear unbiased prediction (BLUP), least squares means (LSM) and regression of the future daughters mean on the present daughters mean. The product moment correlations between different traits studied were positive and high ($= 0.96$).

Key Words: Sire transmitting ability, milk yield, lactation period, heritability, genetic and phenotypic correlation

Zusammenfassung

Titel der Arbeit: Vergleich verschiedener Methoden der Schätzung von Bullenzuchtwerten anhand einiger Milchleistungsmerkmale in einer ägyptischen Holstein Friesian Herde

Anhand von 1931 Erstlaktationen bei Holstein Friesian Kühen eines Betriebes in den Jahren 1987 bis 1994 wurden phänotypische und genetische Parameter für folgende Merkmale geschätzt: 90-Tage-Milchleistung, 305-Tage-Milchleistung und Laktationsperiode. Für den Vergleich der drei Schätzmethoden BLUP, LSM und REG standen 76 Bullen mit mindestens 10 Töchtern zur Verfügung. Die Varianzanalyse zeigte bei den einzelnen Merkmalen signifikante Effekte bei Kalbmonat und -jahr sowie Erstkalbealter, mit Ausnahme des Einflusses vom Erstkalbealter auf die Laktationsperiode. Die Heritabilitätsschätzwerte betragen für die Merkmale: 90-Tage-Milchleistung $0,39 \pm 0,08$, 305-Tage-Milchleistung $0,27 \pm 0,07$, Laktationsperiode $0,14 \pm 0,05$. Die genetischen und phänotypischen Korrelationen zwischen den betrachteten Merkmalen waren signifikant positiv. Die Ergebnisse des Vergleiches der Schätzmethoden werden diskutiert.

Schlüsselwörter: Bullenzuchtwerte, Milchleistung, Laktationsperiode, Heritabilität, genetische und phänotypische Korrelationen

Resumen

Titulo del Artículo: Una comparación de métodos diferentes de estimación de transmisión del padre capacidad para leche rasgos en un rebaño de holstein frisón

Un total de 1931 primeras lactaciones registradas normales de Holstein frisonas vacas que criaron al Dena Granga en Egipto durante los períodos de 1987 a 1994 se usaron a estimar parámetros fenotipos y genéticos por 90 día leche rendimiento, 305 día leche rendimiento y períodos del lactacion. Además, 76 toros con por lo menos 10 hijas se usaron a comparar tres métodos de transmisión del padre capacidad.

Análisis de cuadrados mínimos de variación muestra significativa el efecto del mes y el año del parto y la edad al primero parto para rasgos diferentes estudió, excepto el efecto de edad al primero parto sobre lactacion períodos.

Heritabilidad estima para 90 día leche rendimiento, 305 día leche rendimiento y períodos del lactacion eran 0.39 ± 0.08 , 0.27 ± 0.07 y 0.14 ± 0.05 respectivamente.

Correlaciones genéticos y fenotipos entre rasgos diferentes eran positivas y significativas.

Padres con por lo menos 10 hijas fueron evaluadas por predicción mejor lineal imparcial, medias de los cuadrados mínimos y regresión de media de las hijas futuras en media de las hijas presentes. Las correlaciones del producto entre rasgos diferentes estudiado era positiva y alta ($=0.96$).

Palabras claves: Capacida de transmision del padre, periodos del lactacion, correlacion genéticos y fenotipos

Introduction

Milk production can not be measured on males, thus sire genetic evaluation must be based either on records of female ancestors or on records of the progeny (progeny test). Evaluation on the basis of progeny test results has much greater accuracy than pedigree evaluation (VAN VLECK, 1979).

There are many different methods of sire breeding value has been estimated by different workers in different countries. EL-CHAIFE (1981), ABUBAKAR et al. (1986), VIJ and TIWANA (1988), ABDEL GLIL (1991, 1996), REGE (1991), KHATTAB (1992), KHATTAB and MOURAD (1992), METRY et al. (1994) and TRIVENI et al. (1996) used the best linear unbiased prediction (BLUP). FAHMY (1972), VIJ and TIWANA (1988) and KHATTAB (1992) used average of daughters or least squares means for sire effect. KHATTAB et al. (1987), VIJ and TIWANA (1988) estimated sire breeding values by the regression of the future daughters mean on the present daughters mean.

The objective of this study were: (1) to estimate genetic and phenotypic parameters for first milk traits; for example 90 day milk yield (90 dMY), 305 day milk yield (305 dMY) and lactation period (LP) and (2) to compare three methods of estimating sire transmitting ability for Holstein Friesian cattle in Egypt.

Material and Methods

Data

A total of 1931 first lactation milk records of Holstein Friesian cows maintained at Dena Farm in Egypt during the period 1987 to 1994 were used. The number of sires and average of daughters per sire were 76 and 24.79, respectively. Sires with less than ten daughters were excluded. Artificial insemination using frozen semen was used. Traits studied were first 90 day milk yield, 305 day milk yield and lactation period. More information of the data, feeding system and management of that herd were described by ATIL and KHATTAB (1998).

Estimating of genetic parameters

Data were analysed using Mixed Model Least Squares and Maximum Likelihood Computer Program of HARVEY (1987). The mixed model used for the analysis included the fixed effects of year and month of calving as a fixed effects, age at first calving (AFC) as a regression and sire as a random effect on each of 90 dMY, 305 dMY and LP. Estimates of sire and residual components of variances and covariances

were computed according to II of HENDERSON (1953). Estimates of heritability (h^2), genetic correlations (r_g) with standard errors (SE) and phenotypic correlation (r_p) were estimated according to HARVEY (1987).

Estimating of sire transmitting ability (ETA's)

ABUBAKAR et al. (1986) suggested that ten daughters per sire are considered a minimum number for the evaluation of sires using BLUP procedures in Tropical areas. KHATTAB et al. (1987) using another set of Friesian cattle in Egypt, reported that the accuracy of sire evaluation increases as the number of daughters per sire increased. They also suggested that it is necessary to estimate the sire evaluation based on a large number of daughters per sire. Therefore, transmitting ability of sires at least ten daughters will be examined.

Three methods of sire transmitting abilities were followed:

1- Best Linear Unbiased Prediction (BLUP)

The above model can be written in matrix notation as:

$$Y = Xf + Zs + Wb + e$$

where Y was a vector of observations for each trait, X was a known fixed design matrix, f was an unknown vector of fixed effects representing the mean, year and month of calving, Z was a known design matrix, s was an unobservable vector of random sire effect, W was a vector of covariate variable (independent variable), b was a vector of partial regression of Y on W and e was unobservable random vector of errors with mean zero and variance - covariance matrix $I\sigma^2e$.

The mixed model equations (HENDERSON, 1953) are

$$\begin{bmatrix} X'X & X'Z & X'W \\ Z'X & Z'Z + G & Z'W \\ W'X & W'Z & W'W \end{bmatrix} * \begin{bmatrix} f \\ s \\ b \end{bmatrix} = \begin{bmatrix} X'Y \\ Z'Y \\ W'Y \end{bmatrix}$$

where $G = (4 - h^2) / h^2$ for each trait was added to the diagonal of sire effects in the matrix.

2- Least Squares Means (LSM):

Constants of each bull were obtained by using a model including sire, year and month of calving as a fixed effects and age at first calving as a regression.

3- Regression Of The Future Daughters Mean On The Present Daughters Mean (REG):

$$REG = b \left(D - \bar{X} \right)$$

where $b = n / (n+k)$, n is the number of half sib daughters, k is equal $((4 - h^2) / h^2)$, D is the traits adjusted to year and month of calving as a fixed and age at first calving as a regression and \bar{X} is the least squares mean of the herd.

Product moment correlations for the three methods of sire transmitting ability for each bull were computed.

Results and Discussion

Least squares means of 90 dMY, 305 dMY and LP are 1911 ± 63 kg., 5139 ± 164 kg. and 281 ± 7 d, respectively. The present means of 90 dMY and were higher than those reported by HUSSEIN (1996) using another set Holstein Friesian cattle in Egypt, being 1748 kg and 4938 kg, respectively. Also, the present mean of LP was lower than that reported by KHATTAB and SULTAN (1991) (341 d). The differences between the present estimates and those reported by other workers could be due to different climatic and management. Herds could be possibly the genetically different on be caused by different models used.

Least squares analysis of variances for 90 dMY, 305 dMY and LP are presented in Table 1. Significant ($P < 0.01$) effects of month and year of calving on each of the traits studied were found. These results are agree with the findings were also reported on Friesian raised in other countries by MILAGRES et al. (1989). REGE (1991), KAYA (1996), MAKUZA and McDANIEL (1996) and KELM et al. (1997). Differences in milk yield and lactation period attributed to month and year of calving were interpreted to be due to climatic, nutritional and managerial conditions which changed from one year or month of calving to another.

Table 1

F ratios for factors affecting 90 day milk yield (90 dMY), 305 day milk yield (305 dMY) and lactation period (LP) (Varianzanalyse, geschätzte Effekte und Produktionsmerkmale unterschiedlicher Kontrollabschnitte)

Source of Variation	d.f.	90 dMY F	305 dMY F	LP F
Sire	75	3.64**	2.82**	1.91**
Month of calving	11	4.61**	0.91	2.66**
Year of calving	7	37.03**	37.93**	15.22**
Regressions				
AFC, linear	1	11.26**	5.62**	1.46
AFC, quadratic	1	94.65**	27.72**	2.27
Remainder mean squares	1835	153919	1374676	4201
Number of records = 1931				

**:($P < 0.01$)

Estimates of partial linear and quadratic regression coefficients of 90 dMY and 305 dMY an age at first calving were significant ($P < 0.01$), being 4.03 ± 1.20 kg/mo and -0.34 ± 0.036 kg/mo², respectively for 90 dMY and 8.49 ± 3.58 kg/mo and 0.56 ± 0.11 kg/mo², respectively for 305 dMY. The curvilinear relationships of milk yield on age at first calving were reported by SALLAM et al. (1990), KHATTAB and SULTAN (1991), MANSOUR (1992) and KHATTAB et al.(1994). The present results suggested that reduction in age at first calving is desirable to prolong the length of herd life and to economic the cost of rearing the heifers.

The partial linear and quadratic regression coefficients of LP on age at first calving were not significant. SALLAM et al. (1990) and KHATTAB and SULTAN (1991) found similar results.

The present results indicate the importance of considering the effect of year and season of calving and age at first calving in the model used for describing productive traits for unbiased sire evaluation.

Results obtained in the present study show that the sire of the heifers had a highly significant effect on 90 dMY, 305 dMY and LP ($P < 0.01$, Table 1), accounting for 9.62, 6.85 and 3.54%, of the total variance for 90 dMY, 305 dMY and LP, respectively. This is similar to results of BASU et al. (1982), ABUBAKAR et al. (1986), MILAGNES et al. (1989), ABDEL GLIL (1991 and 1996) and KHATTAB and SULTAN (1991), which ranged from 1.6% and 10%.

Heritability estimates (h^2), genetic correlations (r_g) and phenotypic correlations (r_p) between different traits studied are presented in Table 2. Estimates of h^2 for 90 dMY and 305 dMY were 0.39 ± 0.08 and 0.27 ± 0.07 (Table 2). The present means and moderate and similar to those estimates reported RAGAB et al. (1973), BASU et al. (1982), KHATTAB and SULTAN (1991), ABDEL GLIL (1996) and KAYGISIZ and VANLI (1997) which ranged from 0.26 to 0.48. The present results concluded that the genetic improvement in milk production traits could be achieved through selective breeding values.

Heritability estimate for LP was 0.14 ± 0.05 (Table 2). This estimate is in agreement with those reported by RAGAB et al. (1973) (0.21) and KHATTAB and SULTAN (1991) (0.18).

Genetic and phenotypic correlations between 90 dMY and 305 dMY were 0.56 ± 0.12 and 0.51, respectively (Table 2). The present results indicate that 90 dMY can be a good indicator for production in 305 dMY. These results are in agreement with other estimates found by (RAGAB et al., 1973; KHATTAB and SULTAN, 1991; ABDEL GLIL, 1991 and 1996; and KAYA, 1996).

Table 2

Estimates of heritability (on diagonal), genetic (below diagonal) and phenotypic (above diagonal) correlations for productive traits (Heritabilitätsschätzwerte, genetische (oberhalb Diagonale) und phänotypische Korrelationen für die Produktionsmerkmale)

Traits correlated	90 dMY	305 dMY	LP
90 dMY	0.39 ± 0.08	0.51	0.11
305 dMY	0.56 ± 0.12	0.27 ± 0.07	0.52
LP	0.92 ± 0.13	0.54 ± 0.15	0.14 ± 0.05

Genetic and phenotypic correlation between LP and each of 90 dMY and 305 dMY were positive and significant ($P < 0.01$, Table 2). The present results indicate that there was a positive phenotypic association between milk and lactation period.

Sire values for 90 dMY, 305 dMY and LP were estimated by using the procedure of best linear unbiased prediction (BLUP), least squares means (LSM) and the regression of the future daughters mean on the present daughters mean (REG). Number of daughters per sire ranged from 10 to 77.

Estimates of sire transmitting abilities as deviations from the mean for 90 dMY ranged from -205 to 245 kg for BLUP values, from -374 to 543 for LSM values and from -227 to 280 kg for REG values. With the range being 450, 917 and 507 kg, respectively. Similarly, EL-CHAIFE (1981) working on Friesian X Native cows in Egypt, found that the range of breeding values of 17 sires for 100-day milk yield ranged from -56 to 88 kg. KHATTAB and MOURAD (1992) working on Egyptian buffaloes found that the estimation of sire transmitting ability for 90 dMY ranged from -24 to 37 kg.

Estimates of sire transmitting abilities for 305 dMY ranged from -506 to 675 kg for BLUP values, from -964 to 895 kg for LSM values and from -531 to 492 for REG values, with the range being 1181, 1859 and 1023 kg, respectively. Similarly, ABUBAKAR et al. (1986) with 15512 lactation records of daughters of 138 sires (each with ten or more daughters) found that predicted sire values (BLUP) in 305 dMY ranged from -400 to 400 kg. ABDEL GLIL (1991) with 1653 lactation records for daughters of 163 sires (each with 5 or more daughters) found that predicted sire values (BLUP) ranged from -466 to 681 kg. Also, ZARNECKI et al. (1990) found that the expected breeding values for 305 dMY ranged from -481 to 571 kg. In Egyptian buffaloes, KHATTAB and MOURAD (1992) found that the estimation of sire transmitting ability for 305 dMY ranged from 147 kg to 154 kg. Also, METRY et al. (1994) working on 316 buffaloes cows sired by 77 sires, found that expected breeding values for 305 dMY ranged from -424 to 247 kg. They suggested that potential genetic improvement of milk yield by mass selection was estimated at 2.2 % per year.

The present estimates showed large genetic differences between sires for 90 dMY and 305 dMY, which indicate the high potential for rapid genetic improvement in milk production of Friesian cattle raised in Egypt through selection.

Estimates of sire transmitting ability for LP ranged from -16 to 19 d for BLUP values, -59 to 51 d for LSM values and from -18 to 21 d for REG, with the range being 35, 110 and 39 d, respectively. EL-CHAIFE (1981) working Friesian X Native cows in Egypt, found that sire values for LP ranged from -36 to 46 d. Also, KHATTAB (1992) working on Friesian cattle in Egypt found that the sire values for LP ranged from -29 to 29 d.

The similarity of ranges of BLUP and REG values show the suitability of these two methods for estimating sire transmitting ability for the three traits studied.

The product moment correlations between estimates of sire transmitting abilities for different traits (Table 3) were positive and high (0.96).

KHATTAB (1992) arrived at the same conclusion by using another herd of Friesian cattle.

The product moment correlations between BLUP and LSM estimates were higher than the estimates of 0.71 and 0.88 reported by GLIL and PARMAR (1988) and VIJ and TIWANA (1988) working on Red Dane X Sahiwal and Murrah buffaloes, respectively.

Table 3

Product moment correlations between estimates of different methods of estimating sire transmitting abilities for productive traits (Beziehungen zwischen den unterschiedlichen Schätzmethoden bei den Produktionsmerkmalen)

Methods	90 dMY			305 dMY			LP		
	BLUP	LSM	REG	BLUP	LSM	REG	BLUP	LSM	REG
BLUP									
LSM	0.98			0.98			0.96		
REG	1.00	1.00		1.00	0.98		1.00	0.96	

The high product moment correlations of ETA's between 90 dMY and 305 dMY for the three methods indicated that sire evaluation could be possible using initial milk yield in order to decrease the time required for progeny test. This will reduce the cost of evaluation, and decrease the generation interval to increase the annual genetic gain.

Also, the high product moment correlations between different methods for different traits, indicated that the three methods were reported to succeed for estimating sire transmitting ability.

References

- ABUBAKAR, B.Y.; McDOWELL, R.E.; WELLINGTON, K.E.; VAN VLECK, L.D.:
Estimating genetic values for milk production in the tropics. *J. Dairy Sci.* 69 (1986), 1087 - 1092
- ABDEL GLIL, M.F.:
Sire differences for milk production traits in Friesian cattle. Ph.D. Thesis Faculty of Agric. Zarazig Univ. Egypt, 1991
- ABDEL GLIL, M.F.:
Estimation of genetic parameters and trend of some milk traits in a herd of Friesian cows in Egypt. *Egypt J. Anim. Prod.* (in Press), (1996).
- ATIL, H. ; KHATTAB, A.S.:
Seasonal age correction factors for 305 day milk yield in Holstein cattle. *Pakistan J. Biological Sci.* 2 (1999), 296 - 300
- BASU, S.B.; BAHTRAGAR, D.S.; TANEJA, A.:
Estimates of genetic parameters for lactation performance in Tharparker cattle. *Indian J. Anim. Sci.*, 45 (1982), 279 - 283
- EL-CHAIFE, O.M.:
Studies on cattle (cows and buffaloes) sire evaluation and genetic parameters of partial lactation, dry period and service period. M.Sc. Thesis Faculty of Agric. Alex. Univ., Egypt, 1981
- FAHMY, S.K.:
The reliability of using 5 daughters in estimating the breeding values of dairy sires. *Agric. Research Review.*, 50 (1972), 13 - 17
- GLIL, G.S.; PARMAR, O.S.:
Genetic merits of Red Dane bulls used in a crossbreeding project. *Indian J. Anim. Sci.*, 58 (1988), 811 - 813
- HARVEY, W.R.:
User's guide for LSMLMW. Mixed model least squares and maximum likelihood computer program. The Ohio State Univ., USA, (1987)
- HENDERSON, C.R.:
Estimation of variance and covariance components. *Biometrics* 9 (1953), 226 - 252
- HUSSEIN, A.M.:
Age correction factors for some productive traits in a herd of Holstein Friesian cattle. M.Sc. Thesis. Fac. of Agric. Tanta Univ., Egypt, (1996)
- KAYA, I.:
Parameter estimates for persistency of lactation and relationship of persistency with milk yield in Holstein cattle. Ph.D. Thesis, Ege Univ. Izmir, Turkey, 1996
- KAYGISIZ, A.; VANLI, Y.:
Genetic analysis of the Brown cattle herd of Van Agricultural Vocational High School. *Tarim Bilimleri Dergisi*, 2 (1997), 111 - 115
- KELM, S.C.; FREEMAN, A.E.; KELLEY, D.H.:
Realized versus expected gains in milk and fat production of Holstein cattle, considering effects of days open. *J. Dairy Sci.* 80 (1997), 1786 - 1794
- KHATTAB, A.S.; MABROUK, M.M.S.; EL-HAKIM, A.M.:
Lactation records adjusted for days open in sire evaluation. *J. Agric. Res., Tanta Univ.*, 13 (1987), 286 - 297
- KHATTAB, A.S.; SULTAN, Z.A.:
A comparison of different selection indices for genetic improvement of some dairy traits in Friesian cattle in Egypt. *J. Anim. Breed. Genet.* 108 (1991), 349 - 354
- KHATTAB, A.S.:
A comparison between different methods of estimating sire transmitting ability of some milk traits in Friesian cows in Egypt. *J. Agric. Res. Tanta Univ.* 18 (1992), 569 - 576

- KHATTAB, A.S.; MOURAD, K.A.:
Estimation of genetic parameters and genetic trends for some milk traits in a herd of Egyptian buffaloes. *Egypt. J. Anim. Prod.* **29** (1992), 33 - 44
- KHATTAB, A.S.; ABDEL GLIL, M.F.; SALEM, A.Y.:
Estimation of genetic parameters and sire values for lifetime production and longevity traits in Friesian cows in Egypt. *J. Agric. Sci. Mansoura Univ.* **2** (1994), 483 - 498
- MAKUZA, S.M.; McDANIEL, T.C.:
Effects of days dry, previous days open and current days open on milk yield of cows in Zimbabwe and North Carolina. *J. Dairy Sci.* **79** (1996), 702 - 709
- MANSOUR, H.:
Some reproductive performance parameters of Holstein cattle in The Kingdom Saudi Arabia. *Annals Agric. Sci.*, **37** (1992), 85 - 94
- METRY, G.H.; WILK, J.C.; McDOWELL, R.E.; EL-RIGALATY, H.A.:
Factors affecting the performance of Egyptian buffaloes. *Annals of Agric. Sci., Moshtohor, Egypt.* **32** (1994), 827 - 841
- MILAGRES, J.C.,A.; ALVES, A.J.R.; PEREIRA, J.C.; TEIXEIRA, N.M.:
Effect of genetic and environmental factors on milk yield of crossbreed Holstein, Brown Swiss, Jersey and Zebu cows. (*A.B.A.*, **57**:4786), (1989)
- RAGAB, M.T.; ABDEL AZIZ, A.S.; MOURAD, H.M.:
Phenotypic and genetic parameters of some productive traits in a herd of Friesian cattle at Tahreer province. *Egypt. J. Anim. Prod.* **1** (1973), 3 - 8
- REGE, J.E.O.:
Genetic analysis of reproductive and productive performance of Friesian cattle in Kenya. *J. Anim. Breed. Genet.* **108** (1991), 412 - 423
- SALLAM, M.T.; EL-FEEL, F.M.; KHATTAB, A.S.:
Effect of genetic and nongenetic factors on productive and reproductive traits in Friesian cows in Egypt. *El-Minia. J. Agric. Res.* **12** (1990), 521 - 539
- TRIVENI, D.; RAHAJE, K.L.; TNEJA, V.K.; DUTT, T.:
Estimations of sire breeding values and genetic and phenotypic trends for various economic traits in Murrah buffaloes. *Indian. J. of Anim. Sci.* **66** (1996), 1145 - 1148
- VAN VLECK, D.:
Notes on theory and application of selection principles for the genetic improvement of animals. Department of Animal Sci., Cornell Univ., Ithaca, New York 14853, (1979)
- VII, R.K.; TIWANA, M.S.:
A note on evaluation of buffaloes sires. *Indian J. Dairy Sci.* **41** (1988), 500 - 503
- ZARNECKI, A.; NORMAN, H.D.; JAMNOZIK, J.:
Lifetime performance of ten Friesian strains in Poland. 4th World Congress on Genetic Applied Livestock Production. Edinburgh, 23-27 July, (1990)

Received: 1998-10-23

Accepted: 2000-02-23

Authors' addresses

Dr. HÜLYA ATIL
Ege University, Faculty of Agriculture, Animal Sciences Dept.,
Bornova, Izmir
Turkey

Prof. Dr. ADEL SALAH KHATTAB
Tanta University, Faculty of Agriculture, Animal Production Dept.,
Tanta
Egypt