

Dianne L. Watson · Euan S. Harvey  
Marti J. Anderson · Gary A. Kendrick


## A comparison of temperate reef fish assemblages recorded by three underwater stereo-video techniques

Published online: 9 February 2006  
© Springer-Verlag 2006

### Marine Biology (2005) 148:415–425

Figure 2a, b was incorrect. Please find the correct figure below.

**Fig. 2** The power of stereo-video transects, baited and unbaited remote stereo-video techniques to detect a 25% change in the mean total number of individuals ( $\alpha=0.05$ ) at **a** high-relief reef and **b** low-relief reef sites



The online version of the original article can be found at <http://dx.doi.org/10.1007/s00227-005-0090-6>

D. L. Watson (✉) · E. S. Harvey · G. A. Kendrick  
CRC for Coastal Zone, Estuary and Waterway Management,  
School of Plant Biology, The University of Western Australia,  
35 Stirling Hwy, Crawley, WA 6009, Australia  
E-mail: [dwatson@cyllene.uwa.edu.au](mailto:dwatson@cyllene.uwa.edu.au)  
Tel.: +61-8-64882246  
Fax: +61-8-64881001

M. J. Anderson  
Department of Statistics, University of Auckland,  
Private Bag 92019, Auckland, New Zealand