

A complete enzymatic capacity for long-chain polyunsaturated fatty acid biosynthesis is present in the Amazonian teleost tambaqui, *Colossoma macropomum*

Renato B. Ferraz^{a,b}, Naoki Kabeya^c, Mónica Lopes-Marques^a, André M. Machado^a,
Ricardo A. Ribeiro^d, Ana L. Salaro^e, Rodrigo Ozório^a, L. Filipe C. Castro^{a,ff} and Óscar
Monroig^{g,h†}

^a CIIMAR – Interdisciplinary Centre of Marine and Environmental Research, U. Porto – University of Porto, Portugal.

^b ICBAS - Institute of Biomedical Sciences Abel Salazar, University of Porto, Portugal

^c Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Japan

^d Federal University of Acre, Brazil

^e Department of Biology, Federal University of Viçosa, Brazil

^f Department of Biology, Faculty of Sciences, University of Porto, Portugal

^g Instituto de Acuicultura Torre de la Sal (IATS-CSIC), Ribera de Cabanes, Castellon, Spain

^h Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Scotland, UK

[†] joint last and corresponding authors: filipe.castro@ciimar.up.pt; oscar.monroig@csic.es

Postal address:

^a CIIMAR - Terminal De Cruzeiros De Leixões. Av. General Norton De Matos S/N4450-208 Matosinhos, Portugal.

^b ICBAS - Rua de Jorge Viterbo Ferreira, 228, 4050-313, Porto, Portugal.

^c Department of Aquatic Bioscience - The University of Tokyo 1-1-1, Yayoi, Bunkyo-ku, Tokyo, 113-865

^d Federal University of Acre - Rodovia BR 364, Km 04, s/n - Distrito Industrial, Rio Branco - AC, 69920-900, Brazil

^e Federal University of Viçosa - Avenida Peter Henry Rolfs, s/n - Campus Universitário, Viçosa - MG, 36570-900, Brazil

^f Department of Biology, Faculty of Sciences, University of Porto - Rua Campo Alegre s/n, 4169-007, Portugal

^g Instituto de Acuicultura Torre de la Sal (IATS-CSIC), Ribera de Cabanes 12595, Castellon, Spain

^h Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK94LA, Scotland, UK

Abstract

In vertebrates, the essential fatty acids (FA) that satisfy the dietary requirements for a given species depend upon its desaturation and elongation capabilities to convert the C₁₈ polyunsaturated fatty acids (PUFA), namely linoleic acid (LA, 18:2n-6) and α -linolenic acid (ALA, 18:3n-3), into the biologically active long-chain (C₂₀₋₂₄) polyunsaturated fatty acids (LC-PUFA), including arachidonic acid (ARA, 20:4n-6), eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3). Recent studies have established that tambaqui (*Colossoma macropomum*), an important aquaculture-produced species in Brazil, is a herbivorous fish that can fulfil its essential FA requirements with dietary provision C₁₈ PUFA LA and ALA, although the molecular mechanisms underpinning such ability remained unclear. The present study aimed at cloning and functionally characterizing genes encoding key desaturase and elongase enzymes, namely *fads2*, *elovl5* and *elovl2*, involved in the LC-PUFA biosynthetic pathways in tambaqui. First, a *fads2*-like desaturase was isolated from tambaqui. When expressed in yeast, the tambaqui Fads2 showed Δ 6, Δ 5 and Δ 8 desaturase capacities within the same enzyme, enabling all desaturation reactions required for ARA, EPA and DHA biosynthesis. Moreover, tambaqui possesses two elongases that are *bona fide* orthologs of *elovl5* and *elovl2*. Their functional characterization confirmed that they can operate towards a variety of PUFA substrates with chain lengths ranging from 18 to 22 carbons. Overall our results provide compelling evidence that demonstrates that all the desaturase and elongase activities required to convert LA and ALA into ARA, EPA and DHA are present in tambaqui within the three genes studied herein, i.e. *fads2*, *elovl5* and *elovl2*.

Keywords

Biosynthesis; *Colossoma macropomum*; Desaturase; Elongase; Essential fatty acids.

1. Introduction

Long-chain (C₂₀₋₂₄) polyunsaturated fatty acids (LC-PUFA) such as arachidonic acid (ARA, 20:4n-6), eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3), play many essential roles in growth, development and reproduction of vertebrates (Cook, 1996; Jaya-Ram et al., 2011; Innis, 2008; Vagner and Santigosa, 2011; Calder, 2014). In vertebrates, LC-PUFA can be obtained through the diet or endogenously synthesized from C₁₈ polyunsaturated fatty acid (PUFA) precursors, namely linoleic acid (LA, 18:2n-6) and α -linolenic acid (ALA, 18:3n-3), themselves being dietary essential compounds since they cannot be biosynthesized *de novo* by vertebrates (Castro et al., 2016). Biosynthesis of the physiologically important LC-PUFA from the dietary essential C₁₈ PUFA precursors requires the coordinated action of both fatty acid desaturases (Fads) and elongation of very long chain fatty acids (Elovl) proteins, membrane-bound enzymes localized in the endoplasmic reticulum (Guillou et al., 2010). The characterization of Fads and Elovl involved in LC-PUFA biosynthesis in fish has been extensively investigated (Castro et al., 2016; Kabeya et al., 2018; Monroig et al., 2018). This research has been mainly driven by the need to understand the endogenous ability that farmed species have to utilize commonly used ingredients such as vegetable oils (VO), devoid of LC-PUFA but often rich in their biosynthetic precursors LA and ALA (Turchini et al., 2009; Tocher, 2010). These studies revealed that the LC-PUFA biosynthetic capability varies remarkably among teleosts, depending upon the species-specific *fads* and *elovl* gene complement and function (Castro et al., 2016; Monroig et al., 2016, 2018).

Fads enzymes introduce a double bond (unsaturation) at a specific position of the fatty acid (FA) chain and are termed as “ Δ x desaturase”, with “x” denoting the position of the carbon from the carboxylic group at which the new double bond is introduced (Leonard et al., 2004). Vertebrate Fads are characterized by a cytochrome b5-like domain

containing a heme-binding motif (HPGG), three histidine boxes consisting HXXXH, HXXHH and QXXHH, and several membrane-spanning domains (Sperling et al., 2003). Unlike most vertebrates that have Fads1 and Fads2 with $\Delta 5$ and $\Delta 6$ desaturase activities, respectively (Guillou et al., 2010), teleost fish appear to have lost the *fads1* gene, presenting exclusively *fads2* genes (Castro et al., 2012, 2016), with the exception of the Japanese eel *Anguilla japonica*, a basal teleostei possessing a Fads1 with $\Delta 5$ activity (Lopes-Marques et al., 2018). Interestingly, teleostei Fads2 are functionally more diverse than their non-teleost counterparts (Fonseca-Madrugal et al., 2014; Castro et al., 2016). While the majority of teleost Fads2 enzymes functionally characterized to date present $\Delta 6$ desaturase activity (Zheng et al., 2004, 2009; González-Rovira et al., 2009; Mohd-Yusof et al., 2010; Monroig et al., 2013, 2010a; Kabeya et al., 2018), an increasingly high number of teleost Fads2 have alternative desaturase substrate specificities including both $\Delta 6$ and $\Delta 5$ activities within the same enzyme (Hastings et al., 2001; Tanomman et al., 2013; Kuah et al., 2016; Oboh et al., 2016), $\Delta 5$ (Abdul Hamid et al., 2016) and $\Delta 4$, the latter often exhibiting some residual activity as $\Delta 5$ desaturase (Li et al., 2010; Morais et al., 2012; Fonseca-Madrugal et al., 2014; Oboh et al., 2017). Consistent with the mammalian FADS2 (Park et al., 2009), fish Fads2 typically possess $\Delta 8$ desaturation capacity (Monroig et al., 2011a), enabling the “ $\Delta 8$ pathway”, an alternative route to the most prominent “ $\Delta 6$ pathway” leading to the biosynthesis of $20:3n-6$ and $20:4n-3$, immediate precursors of ARA and EPA, respectively (Castro et al., 2016).

Regarding Elovl, these enzymes catalyze the condensation of malonyl-CoA into an activated FA, a rate-limiting reaction in the elongation pathway resulting in the extension of the pre-existing FA in 2 carbons (Jakobsson et al., 2006). Several distinct Elovl (termed Elovl 1-7) have been described in vertebrates (Jakobsson et al., 2006; Guillou et al., 2010), each one having different substrate preferences. Interestingly, Elovl2, Elovl4 and

Elovl5 can elongate PUFA substrates and thus play major roles in LC-PUFA biosynthesis (Jakobsson et al., 2006; Guillou et al., 2010). In fish, the majority of Elovl5 orthologs investigated to date showed ability to efficiently elongate C₁₈ and C₂₀ PUFA substrates, with remarkably lower efficiency generally exhibited towards C₂₂ PUFA (Castro et al., 2016). Vertebrate Elovl2 can elongate C₂₀ and C₂₂ PUFA, but it has little ability to elongate C₁₈ PUFA. Loss of Elovl2 in recently emerged teleost lineages has been postulated to be partly compensated by Elovl4, an enzyme that, along its role in the biosynthesis of very long-chain (>C₂₄) PUFA (Monroig et al., 2010b), can operate towards C₂₂ PUFA like Elovl2 thus playing a role in LC-PUFA biosynthesis (Monroig et al., 2011b; Kabeya et al., 2015).

Overall, it has become clear that the endogenous capacity for LC-PUFA production in fish species cannot be anticipated from their genetic repertoire of *fads* and *elovl*. Rather, functional analyses of the gene products (i.e. enzymes) is required to fully understand the physiological capacities that farmed fish species have to biosynthesize ARA, EPA and DHA from C₁₈ precursors contained in VO. The *Colossoma macropomum*, popularly known as “tambaqui”, is a freshwater fish and is currently the main native species farmed in Brazil, representing 28% of the total of the national production. Importantly, tambaqui displays high growth rates, adaptation to intensive culture systems and good fillet quality (Guimarães and Martins, 2015). The adult fish are predominantly herbivorous, feeding abundantly on fruits and seeds during flooding episodes (Almeida et al., 2008). Such natural adaptation to low LC-PUFA diets suggests that tambaqui possess an active LC-PUFA biosynthetic capacity that enable them to thrive with little input of these essential nutrients. In order to test this hypothesis, the present study aimed at cloning and functionally characterize the key desaturase and elongase genes, namely *fads2*, *elovl5* and *elovl2*, involved in the LC-PUFA biosynthetic pathways in this species.

2. Materials and Methods

2.1. RNA sampling and cDNA synthesis

Liver samples from an adult tambaqui were preserved in an RNA stabilization buffer (3.6 M ammonium sulphate, 18 mM sodium citrate, 15 mM EDTA, pH 5.2) and stored at -80°C prior to RNA extraction. RNA extraction was performed using the Illustra RNA spin kit (GE Healthcare, Little Chalfont, UK) according to the manufacturer's guidelines including an on-column DNase treatment. Final RNA samples were run on a 1 % agarose TAE gel stained with GelRed™ nucleic acid stain (Biotium, Hayward, CA, USA) to evaluate integrity, while RNA quantification was performed using BiotTek® microplate reader to determine sample absorbance. Subsequently, cDNA synthesis was performed from 1 μg of total RNA and using iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA) according to manufacturer's recommendations. Additionally, 5' and 3' Rapid Amplification of cDNA Ends (RACE) cDNA was prepared using SMARTer® RACE 5'/3' Kit (Clontech, Mountain View, CA, USA).

2.2. Isolation of ORF *fads* and *elovl* genes from tambaqui

For amplification of the first fragments of the target genes *fads2*, *elovl5* and *elovl2*, polymerase chain reactions (PCR) were conducted using degenerated primers designed on conserved regions of teleosts orthologs of *fads2*, *elovl5* and *elovl2* using CODEHOP (<http://blocks.fhcrc.org/codehop.html>) (Rose et al., 2003). Amplifications of partial fragments of the target genes were carried out using a 1 μl of liver cDNA, 500 nM of sense and antisense primers, and Flash High-Fidelity PCR Master Mix (Thermo Fisher Scientific, Waltham, MA, USA), set for a final volume of 20 μl (see Supplementary Table 1 for primers, PCR conditions). Resulting PCR products were analyzed on a 1 % agarose gel, and fragments with the expected size were purified with NYZGelpure (NZYtech,

Lisbon, Portugal) and confirmed by DNA sequencing (GATC Biotech, Constance, Germany).

To obtain full-length cDNA sequences, RACE PCR were performed. Gene specific primers for RACE PCR were designed using the previously isolated first fragments, with adapter specific primers being provided with the kit (SMARTer® RACE 5'/3' Kit, Clontech, Mountain View, CA, USA). The RACE PCR were performed with Flash High-Fidelity PCR Master Mix (Thermo Fisher Scientific, Waltham, MA, USA) using as template 5' and 3' RACE cDNA prepared from liver RNA. Each RACE PCR contained 1 µl of gene specific primer combined with 2 µl Universal primer mix (Clontech, Mountain View, CA, USA) and corresponding RACE cDNA template (see Supplementary Table 1 for primers and PCR conditions). Resulting 5' and 3' RACE PCR products were confirmed by sequencing (GATC Biotech, Constance, Germany) and assembled with first fragments to obtain open reading frame (ORF) sequences for each target gene (Geneious V7.1.9).

2.3. Sequence collection and phylogenetic analysis

Blastp searches in the public database NCBI were performed using human Fads1, Fads2, Elovl2, Elovl5 and Elovl4, as queries. Fads and Elovl and amino acid sequences were collected from the major vertebrate lineages, namely Sarcopterygii (Mammalia, Reptilia, Aves, Amphibia, and Coelacanthiformes), Chondrichthyes (Elasmobranchii and Chimaera) and Actinopterygii (Elopomorpha, Osteoglossomorpha, Otomorpha, and Euteleostomorpha), as well as from invertebrate groups including Hemichordata (*Saccoglossus kowalevskii*), Cephalochordata (*Branchiostoma lanceolatum*), Urochordata (*Ciona inestinalis*) and Cephalopod (*Octopus vulgaris*) (see Supplementary Table 2 for accession numbers). Next, two phylogenetic trees were constructed containing Fads and Elovl sequences, with alignments for both Fads and Elovl performed

independently in MAFFT (Katoh and Toh, 2008) with the L-INS-i method. The resulting sequence alignments were inspected and stripped of 90% columns containing gaps leaving a total of 59 sequences with 467 positions for phylogenetic analysis for Fads, and 53 sequences and 318 positions for Elovl. Final sequence alignments were submitted for phylogenetic analysis to PhyML V3.0 (Guindon et al., 2010). For each analysis evolutionary model was determined using the smart model selection (SMS) option resulting in a JTT+G+I for Fads and JTT+G+I for Elovl and branch support in both runs was calculated using aBayes. The resulting trees were visualized and analyzed in Fig. Tree V1.3.1 available at <http://tree.bio.ed.ac.uk/software/figtree/>.

2.4. Functional characterization of *tambaqui fads2*, *elovl5* and *elovl2*

Functions of the *tambaqui* Fads2, Elovl5 and Elovl2 enzymes were characterized individually by heterologous expression in yeast, *Saccharomyces cerevisiae*. First, the ORF sequences of the *tambaqui fads2*, *elovl5* and *elovl2* were isolated and cloned into the yeast expression vector pYES2 (Thermo Fisher Scientific, Waltham, MA, USA) (Lopes-Marques et al., 2017). The ORF of the target genes were isolated by PCR (Flash High-Fidelity PCR Master Mix, Thermo Fisher Scientific, Waltham, MA, USA) using *C. macropomum* liver cDNA as template and primers containing appropriate restriction sites for further cloning into pYES2 (see Supplementary Table 1 for primers and PCR conditions). PCR products and pYES2 were subsequently digested with appropriate restriction enzymes (Promega, Madison, WI, USA) and ligated to produce the plasmid constructs pYES2-*fads2*, pYES2-*elovl5* and pYES2-*elovl2*. Sequences of inserts in each plasmid construct was confirmed (GATC Biotech, Constance, Germany) before being used to transform yeast competent cells.

One colony of transgenic yeast carrying either pYES2-*fads2*, pYES2-*elovl5* or pYES2-*elovl2* was grown in *S. cerevisiae* minimal (SCMM-uracil) medium lacking uracil

to produce a bulk culture and subsequently diluted to OD₆₀₀ of 0.4 in all the Erlenmeyer flasks used to test each potential FA substrate assayed. For *Fads2*, transgenic yeast were grown in the presence of $\Delta 6$ (18:2_{n-6} and 18:3_{n-3}), $\Delta 8$ (20:2_{n-6} and 20:3_{n-3}), $\Delta 5$ (20:3_{n-6} and 20:4_{n-3}) and $\Delta 4$ (22:4_{n-6} and 22:5_{n-3}) desaturase substrates. In addition, to determine the desaturase activity towards 24:5_{n-3}, the transgenic yeast co-expressing zebrafish *elovl2* and the tambaqui *fads2* were grown in the presence of 22:5_{n-3} as described previously by Oboh (Oboh et al., 2016). Here the conversion towards 18:3_{n-3} in the co-expression yeast system was also determined as positive control. With regards to tambaqui elongases, both *Elovl5* and *Elovl2* were functionally characterized by growing transgenic yeast in medium supplemented with one of the following PUFA substrates: 18:2_{n-6}, 18:3_{n-3}, 18:3_{n-6}, 18:4_{n-3}, 20:4_{n-6}, 20:5_{n-3}, 22:4_{n-6} and 22:5_{n-3}. Given that PUFA uptake by yeast decreases with increasing chain length, PUFA substrates were added to the yeast cultures from both desaturase and elongase assays at final concentrations of 0.5 mM (C₁₈), 0.75 mM (C₂₀) and 1.0 mM (C₂₂) (Lopes-Marques et al., 2017). Additionally, yeast transformed with empty pYES2 were also grown in presence of PUFA substrates as control treatments. After 2 days of culture at 30 °C, yeast were harvested, washed, and homogenized in chloroform/methanol (2:1, v/v) containing 0.01% (w/v) butylated hydroxytoluene (BHT) and kept -20 °C until further analysis. All PUFA substrates, except stearidonic acid (18:4_{n-3}), were from Nu-Chek Prep, Inc. (Elysian, MN, USA). Stearidonic acid and chemicals used to prepare SCMM-uracil were from Sigma-Aldrich (Darmstadt, Germany), except for the bacteriological agar obtained from Oxoid Ltd. (Hants, UK).

2.5. Fatty acid analysis of yeast

Total lipid extracted from yeast (Folch et al., 1957) was used to prepare fatty acyl methyl esters (FAME) that were further analyzed by gas chromatography as previously

described (Hastings et al., 2001; Li et al., 2010). FAME were identified based on retention times using an Fisons GC-8160 (Thermo Fisher Scientific, Waltham, MA, USA) gas chromatograph equipped with a 60 m x 0.32 mm i.d. x 0.25 μ m ZB-wax column (Phenomenex, Macclesfield, UK) and flame ionization detector. The desaturation or elongation conversion efficiencies from exogenously added PUFA substrates were calculated by the proportion of substrate FA converted to desaturated or elongated products as $[\text{all product areas}/(\text{all products areas} + \text{substrate area})] \times 100$. For the tambaqui elongases, “all product areas” include those of the initial elongation products, as well as those from stepwise elongations occurring subsequently (Monroig et al., 2012). Similarly, the tambaqui Fads2 exhibited multifunctional abilities, and thus the conversions on $\Delta 8$ substrates (20:2n-6 and 20:3n-3) include stepwise $\Delta 5$ desaturase reactions (Fonseca-Madrighal et al., 2014).

3. Results

3.1. Phylogenetic analysis of the tambaqui *fads* and *elovl* sequences

To determine orthology of the isolated sequences from tambaqui, two independent phylogenetic trees including *Fads* (Fig. 1) and *Elovl* (Fig. 2) were constructed. For *Fads*, the resulting phylogenetic tree showed two well supported vertebrate clades that contained either *Fads1* or *Fads2* sequences, and both being out-grouped by invertebrate *Fads* (Fig. 1). We found that the isolated *Fads* from tambaqui was placed within the *Fads2* clade together with orthologs from other teleost species such as *Danio rerio* and *Ictalurus punctatus*. This positioning indicates that newly cloned tambaqui *fads* is a *bona fide* *Fads2* desaturase, orthologous to previously characterized *Fads2* from other vertebrate species (Fig. 1).

Regarding the *Elovl* phylogenetic analysis, the phylogenetic tree revealed that vertebrate *Elovl4* sequences including the invertebrate elongase from *C. intestinalis* formed an independent monophyletic group positioned. The remaining vertebrate *Elovl2* and *Elovl5* sequences constituted two well supported independent clades with the invertebrate *B. lanceolatum* *Elovl2/5* sequence as an outgroup (Monroig et al., 2016). Regarding the herein isolated elongases from tambaqui, one clustered within the *Elovl2* clade and the other within the *Elovl5* clade, in both cases including characterized *Elovl* sequences from other teleosts and vertebrate species. These results indicate that the isolated tambaqui elongases are orthologs of *elovl2* and *elovl5* (Fig. 2).

3.2. Functional characterization of *C. macropomum fads2*, *elovl5* and *elovl2* in *S. cerevisiae*

Control yeast transformed with the empty vector pYES2 did not show activity towards any of the substrates tested (data not shown). Transgenic yeast expressing the tambaqui *fads2* exhibited $\Delta 6$ desaturase activity since $18:2n-6$ and $18:3n-3$ were converted to $18:3n-6$ (28.3 % conversion) and $18:4n-3$ (63.5 % conversion), respectively (Table 1).

The tambaqui Fads2 also showed $\Delta 5$ desaturase activity since transgenic yeast converted 20:3n-6 and 20:4n-3 were converted to 20:4n-6 (14.8 % conversion) and 20:5n-3 (17.8 % conversion), respectively (Table 1). These results indicate that the tambaqui Fads2 is a dual $\Delta 6\Delta 5$ desaturase. Moreover, this enzyme exhibited $\Delta 8$ desaturase activity, with exogenously added 20:2n-6 and 20:3n-3 being desaturated to 20:3n-6 and 20:4n-3, respectively (Table 1). In addition, the tambaqui Fads2 showed $\Delta 6$ desaturation capacity towards 24:5n-3, which was desaturated to 24:6n-3.

Functional assays of tambaqui elongases showed that yeast expressing Elov15 were able to elongate all C₁₈ (18:2n-6, 18:3n-3, 18:3n-6 and 18:4n-3) and C₂₀ (20:4n-6 and 20:5n-3) PUFA substrates, yet no elongation was detected for C₂₂ (22:4n-6 and 22:5n-6) substrates (Table 2). Moreover, yeast expressing the tambaqui Elov12 exhibited elongation ability for all PUFA including C₁₈, C₂₀ and C₂₂, with C₂₀ substrates appearing as preferred substrates for elongation (Table 2).

4. Discussion

The essential FA for a given species depends upon its desaturation and elongation capability to endogenously convert dietary C₁₈ PUFA LA and ALA into the biologically active LC-PUFA, namely ARA, EPA and DHA (Tocher, 2015). Recent investigations have shown that tambaqui juveniles exhibited no difference in growth performance when fed diets containing mixtures of VO (corn and linseed oils) in comparison to fish fed a fish oil (control) diet (Pereira et al., 2017; Paulino et al., 2018). Consequently it was established that tambaqui can fulfil its essential FA requirements with dietary provision C₁₈ PUFA (Paulino et al., 2018). The present study provides compelling molecular evidence that support previous findings and unequivocally demonstrates that all the desaturase and elongase activities required to convert C₁₈ PUFA into ARA, EPA and DHA exist in tambaqui within the three genes studied herein, i.e. *fads2*, *elovl5* and *elovl2* (Fig. 3).

The phylogenetic analysis confirmed that the *fads*-like desaturase isolated from tambaqui is an ortholog of *fads2*. Additionally, sequence alignment of the tambaqui desaturase with Fads2 from other teleost species showed a high degree of conservation of the classic desaturase motifs, namely histidine boxes and heme binding region (Supplementary Fig. 1). Recently Lopes-Marques et al. (2018) demonstrated that, along the formerly characterized Fads2 (Wang et al., 2014), *Anguilla japonica* possess a Fads1 with $\Delta 5$ activity similarly to chondrichthyes and mammals (Marquardt et al., 2000; Castro et al., 2012), being the first report of a non-*fads2* desaturase described in teleosts. Nevertheless, the presence of *fads1* among teleosts appears to be anecdotic and restricted to certain post-3R lineages like Elopomorpha (Lopes-Marques *et al.*, 2018) since the vast majority of teleosts possess only *fads2* as the sole *fads* desaturase found in their genomes (Castro et al., 2016). Indeed, loss of *fads1* has been often postulated as a major evolutionary driver partly explaining the unique functional plasticity of teleosts Fads2

(Fonseca-Madrugal et al., 2014; Castro et al., 2016). In agreement, the functional characterization of the tambaqui Fads2 demonstrated this is a functionalized desaturase that, in addition to the expected $\Delta 6$ activity, it also exhibited $\Delta 5$ desaturase capacity. Therefore, the tambaqui Fads2 can be categorized as a bifunctional or dual $\Delta 6\Delta 5$ desaturase, an enzyme type previously described in *D. rerio* (Hastings et al., 2001), *Siganus canaliculatus* (Li et al., 2010), *Oreochromis niloticus* (Tanomman et al., 2013), *Channa striata* (Kuah et al., 2016) and *Clarias gariepinus* (Obloh et al., 2016). Moreover, the tambaqui Fads2 also showed $\Delta 8$ desaturase activity as described in a wide range of teleosts (Monroig et al., 2011a; Obloh et al., 2016; Lopes-Marques et al., 2017; Kabeya et al., 2018). With the exception of the marine herbivore *S. canaliculatus*, all teleost species reported to have dual $\Delta 6\Delta 5$ desaturase inhabit freshwater ecosystems with limited abundance of LC-PUFA compared to marine environments (Colombo et al., 2016). Indeed, such dietary restriction of LC-PUFA has been hypothesized to account for the enhanced LC-PUFA biosynthesizing capacity of freshwater teleosts in comparison to their marine counterparts (Tocher et al., 2003; Leaver et al., 2008). Interestingly, the varied range of desaturase activities contained within the tambaqui Fads2 implies that this enzyme enables all the desaturation reactions required to biosynthesize ARA and EPA from the C₁₈ PUFA precursors LA and ALA, respectively (Fig. 3). Also, the analysis of critical residues involved in determining desaturation position preference in mammalian FADS desaturases (residues marked with “*” in Supplementary Fig. 1) (Watanabe et al., 2016) revealed that tambaqui Fads2 has a similar profile to that found in the bifunctional *D. rerio* Fads2 (red boxes Supplementary Fig. 1), presenting no additional distinctive residue replacements. Similarly, the elongation capacities involved in the biosynthesis of ARA and EPA were also demonstrated within the two newly cloned *elovl* from tambaqui.

Phylogenetic analysis of the two tambaqui elongases showed that these were orthologs of Elovl2 and Elovl5 confirming the presence of both types of elongases as anticipated by the phylogenetic position of tambaqui (Characiformes) (Ravi and Venkatesh, 2018). Certainly, whereas Elovl5 is present virtually in all teleosts (Castro et al., 2016), distribution of Elovl2 is more restricted, with absence in recently emerged teleosts (Morais et al., 2009; Monroig et al., 2016) and presence reported only in relatively ancient lineages such as Cypriniformes (Monroig et al., 2009), Siluriformes (Oboh et al., 2016) and Salmoniformes (Morais et al., 2009; Gregory and James, 2014). Importantly, functional characterization of the tambaqui elongases further demonstrated their role in LC-PUFA biosynthesis, with both enabling the elongation capacity required to produce ARA and EPA from their C₁₈ precursors (Fig. 3). Thus, the tambaqui Elovl5 showed elongation capacity towards C₁₈ and C₂₀ PUFA, while no activity detected towards C₂₂ substrates. Such elongation capacities exhibited by the tambaqui Elovl5 are generally consistent with those of other teleost species (Castro et al., 2016), with some interesting particularities. Thus, the tambaqui Elovl5 showed no activity towards the C₂₂ PUFA substrates 22:4n-6 and 22:5n-3, in contrast to Elovl5 from other species such as *S. canaliculatus* and *Thunnus thynnus*, with remarkably high C₂₂ elongation capacity when expressed in yeast (Morais et al., 2011; Monroig et al., 2012). On the other hand, the tambaqui Elovl2 showed activity towards all C₁₈, C₂₀ and C₂₂ PUFA substrates tested. Yet, certain C₁₈ substrates, namely LA (18:2n-6) and ALA (18:3n-3), were elongated at a comparatively lesser extent, suggesting that Elovl2 has a minor role in the Δ 6 desaturase – elongase - Δ 5 desaturase pathway leading to ARA and EPA biosynthesis from LA and ALA, respectively (Fig. 3). Unlike Elovl5 though, the tambaqui Elovl2 had efficiency to operate towards C₂₂ substrates. For instance, 22:5n-3 was efficiently elongated to 24:5n-3, a key intermediate component of the so-called “Sprecher pathway”, a metabolic route

that accounts for DHA biosynthesis in vertebrates (Buzzi et al., 1996, 1997; Sprecher, 2000) and recently demonstrated to be widespread among teleosts (Oboh et al., 2017). Certainly, our results show that tambaqui can also biosynthesize DHA through the Sprecher pathway since, in addition to the capacity of Elovl2 to produce 24:5n-3 mentioned above, its Fads2 has the ability to desaturate 24:5n-3 to 24:6n-3, a key desaturation reaction preceding a final β -oxidation step required to produce DHA. Similarly, to the enzymatic reactions required for ARA and EPA, our results confirm that both the elongation and desaturase capacities involved to biosynthesize DHA from EPA (elongase – elongase - Δ 6 desaturase) are present in tambaqui.

In conclusion, the combination of the enzymatic capacities demonstrated by the tambaqui Fads2, Elovl2 and Elovl5 allows this species to convert the dietary essential C₁₈ PUFA LA (18:2n-6) and ALA (18:3n-3) into the biologically active ARA, EPA, and DHA. These results confirm that tambaqui can fulfil its essential FA requirements with an adequate dietary provision of C₁₈ PUFA. Therefore, tambaqui emerges as a valuable species for the sustainable development of Brazilian aquaculture, given its ability to efficiently utilize alternative sources of essential FA such as VO as confirmed at molecular level by the enzymatic capabilities shown in this study.

Acknowledgements

This study was supported by CNPq, Conselho Nacional de Desenvolvimento Científico e Tecnológico – Brasil and by INNOVMAR - Innovation and Sustainability in the Management and Exploitation of Marine Resources (reference NORTE-01-0145-FEDER-000035, within Research Line INSEAFOD), supported by North Portugal Regional Operational Programme (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement, through the European Regional Development Fund (ERDF)”.

References

- Abdul Hamid, N.K., Carmona-Antonanzas, G., Monroig, O., Tocher, D.R., Turchini, G.M., Donald, J.A., 2016. Isolation and functional characterisation of a *fads2* in rainbow trout (*Oncorhynchus mykiss*) with $\Delta 5$ desaturase activity. PLoS One 11, e0150770.
- Almeida, N.M., Visentainer, J.V., Franco, M.R.B., 2008. Composition of total, neutral and phospholipids in wild and farmed tambaqui (*Colossoma macropomum*) in the Brazilian Amazon area. J. Sci. Food Agric. 88, 1739-1747.
- Buzzi, M., Henderson, R., Sargent, J., 1996. The desaturation and elongation of linolenic acid and eicosapentaenoic acid by hepatocytes and liver microsomes from rainbow trout (*Oncorhynchus mykiss*) fed diets containing fish oil or olive oil. Biochim. Biophys. Acta 1299, 235-244.
- Buzzi, M., Henderson, R., Sargent, J., 1997. Biosynthesis of docosahexaenoic acid in trout hepatocytes proceeds via 24-carbon intermediates. Comp. Biochem. Physiol. B 116, 263-267.
- Calder, P.C., 2014. Very long chain omega-3 (n-3) fatty acids and human health. Eur. J. Lipid Sci. Technol. 116, 1280-1300.
- Castro, L.F., Monroig, O., Leaver, M.J., Wilson, J., Cunha, I., Tocher, D.R., 2012. Functional desaturase Fads1 ($\Delta 5$) and Fads2 ($\Delta 6$) orthologues evolved before the origin of jawed vertebrates. PLoS One 7, e31950.
- Castro, L.F., Tocher, D.R., Monroig, O., 2016. Long-chain polyunsaturated fatty acid biosynthesis in chordates: Insights into the evolution of Fads and Elovl gene repertoire. Prog. Lipid Res. 62, 25-40.
- Colombo, S.M., Wacker, A., Parrish, C.C., Kainz, M.J., Arts, M.T., 2016. A fundamental dichotomy in long-chain polyunsaturated fatty acid abundance between and within marine and terrestrial ecosystems. Environ. Rev. 25, 163-174.
- Cook, H.W., 1996. Fatty acid desaturation and chain elongation in eukaryotes. In: Vance, D.E., Vance J.E. (Ed.), Biochemistry of Lipids, Lipoproteins and Membranes. Elsevier Science, Alberta Canada, pp. 129- 152.
- Folch, J., Lees, M., Sloane Stanley, G., 1957. A simple method for the isolation and purification of total lipids from animal tissues. J. Biol. Chem. 226, 497-509.
- Fonseca-Madrugal, J., Navarro, J.C., Hontoria, F., Tocher, D.R., Martinez-Palacios, C.A., Monroig, O., 2014. Diversification of substrate specificities in teleostei Fads2: characterization of $\Delta 4$ and $\Delta 6\Delta 5$ desaturases of *Chirostoma estor*. J. Lipid Res. 55, 1408-1419.
- González-Rovira, A., Mourente, G., Zheng, X., Tocher, D.R., Pendón, C., 2009. Molecular and functional characterization and expression analysis of a $\Delta 6$ fatty acyl desaturase cDNA of European sea bass (*Dicentrarchus labrax* L.). Aquaculture 298, 90-100.

- Gregory, M.K., James, M.J., 2014. Rainbow trout (*Oncorhynchus mykiss*) Elovl5 and Elovl2 differ in selectivity for elongation of omega-3 docosapentaenoic acid. *Biochim. Biophys. Acta* 1841, 1656-1660.
- Guillou, H., Zadavec, D., Martin, P.G., Jacobsson, A., 2010. The key roles of elongases and desaturases in mammalian fatty acid metabolism: Insights from transgenic mice. *Prog. Lipid Res.* 49, 186-199.
- Guimarães, I.G., Martins, G.P., 2015. Nutritional requirement of two Amazonian aquacultured fish species, *Colossoma macropomum* (Cuvier, 1816) and *Piaractus brachypomus* (Cuvier, 1818): a mini review. *J. Appl. Ichthyol.* 31, 57-66.
- Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307-321.
- Hastings, N., Agaba, M., Tocher, D.R., Leaver, M.J., Dick, J.R., Sargent, J.R., Teale, A.J., 2001. A vertebrate fatty acid desaturase with Delta5 and Delta6 activities. *Proc. Natl. Acad. Sci. U.S.A.* 98, 14304-14309.
- Innis, S.M., 2008. Dietary omega 3 fatty acids and the developing brain. *Brain Res.* 1237, 35-43.
- Jakobsson, A., Westerberg, R., Jacobsson, A., 2006. Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog. Lipid Res.* 45, 237-249.
- Jaya-Ram, A., Ishak, S.D., Enyu, Y.L., Kuah, M.K., Wong, K.L., Shu-Chien, A.C., 2011. Molecular cloning and ontogenic mRNA expression of fatty acid desaturase in the carnivorous striped snakehead fish (*Channa striata*). *Comp. Biochem. Physiol. A* 158, 415-422.
- Kabeya, N., Yamamoto, Y., Cummins, S.F., Elizur, A., Yazawa, R., Takeuchi, Y., Haga, Y., Satoh, S., Yoshizaki, G., 2015. Polyunsaturated fatty acid metabolism in a marine teleost, Nibe croaker *Nibea mitsukurii*: Functional characterization of Fads2 desaturase and Elovl5 and Elovl4 elongases. *Comp. Biochem. Physiol. B* 188, 37-45.
- Kabeya, N., Yevzelman, S., Oboh, A., Tocher, D.R., Monroig, O., 2018. Essential fatty acid metabolism and requirements of the cleaner fish, ballan wrasse *Labrus bergylta*: Defining pathways of long-chain polyunsaturated fatty acid biosynthesis. *Aquaculture* 488, 199-206.
- Katoh, K., Toh, H., 2008. Recent developments in the MAFFT multiple sequence alignment program. *Brief Bioinform.* 9, 286-298.
- Kuah, M.K., Jaya-Ram, A., Shu-Chien, A.C., 2016. A fatty acyl desaturase (*fads2*) with dual $\Delta 6$ and $\Delta 5$ activities from the freshwater carnivorous striped snakehead *Channa striata*. *Comp. Biochem. Physiol. A* 201, 146-155.
- Leaver, M.J., Bautista, J.M., Björnsson, B.T., Jönsson, E., Krey, G., Tocher, D.R., Torstensen, B.E., 2008. Towards fish lipid nutrigenomics: current state and prospects for fin-fish aquaculture. *Rev. Fish. Sci.* 16, 73-94.

- Leonard, A.E., Pereira, S.L., Sprecher, H., Huang, Y.-S., 2004. Elongation of long-chain fatty acids. *Prog. Lipid Res.* 43, 36-54.
- Li, Y., Monroig, O., Zhang, L., Wang, S., Zheng, X., Dick, J.R., You, C., Tocher, D.R., 2010. Vertebrate fatty acyl desaturase with $\Delta 4$ activity. *Proc. Natl. Acad. Sci. U.S.A.* 107, 16840-16845.
- Lopes-Marques, M., Ozorio, R., Amaral, R., Tocher, D.R., Monroig, O., Castro, L.F., 2017. Molecular and functional characterization of a *fads2* orthologue in the Amazonian teleost, *Arapaima gigas*. *Comp. Biochem. Physiol. B* 203, 84-91.
- Lopes-Marques, M., Kabeya, N., Qian, Y., Ruivo, R., Santos, M., Venkatesh, B., Tocher, D.R., Castro, L.F., Monroig, O., 2018. Retention of fatty acyl desaturase 1 (*fads1*) in Elopomorpha and Cyclostomata provides novel insights into the evolution of long-chain polyunsaturated fatty acid biosynthesis in vertebrates. *BMC Evolutionary Biology* (in press)
- Marquardt, A., Stöhr, H., White, K., Weber, B.H., 2000. cDNA cloning, genomic structure, and chromosomal localization of three members of the human fatty acid desaturase family. *Genomics* 66, 175-183.
- Mohd-Yusof, N.Y., Monroig, O., Mohd-Adnan, A., Wan, K.L., Tocher, D.R., 2010. Investigation of highly unsaturated fatty acid metabolism in the Asian sea bass, *Lates calcarifer*. *Fish Physiol. Biochem.* 36, 827-843.
- Monroig, O., Rotllant, J., Sanchez, E., Cerda-Reverter, J.M., Tocher, D.R., 2009. Expression of long-chain polyunsaturated fatty acid (LC-PUFA) biosynthesis genes during zebrafish *Danio rerio* early embryogenesis. *Biochim. Biophys. Acta* 1791, 1093-1101.
- Monroig, O., Zheng, X., Morais, S., Leaver, M.J., Taggart, J.B., Tocher, D.R., 2010a. Multiple genes for functional $\Delta 6$ fatty acyl desaturases (Fad) in Atlantic salmon (*Salmo salar* L.): gene and cDNA characterization, functional expression, tissue distribution and nutritional regulation. *Biochim. Biophys. Acta* 1801, 1072-1081.
- Monroig, O., Rotllant, J., Cerda-Reverter, J.M., Dick, J.R., Figueras, A., Tocher, D.R., 2010b. Expression and role of Elovl4 elongases in biosynthesis of very long-chain fatty acids during zebrafish *Danio rerio* early embryonic development. *Biochim. Biophys. Acta* 1801, 1145-1154.
- Monroig, O., Li, Y., Tocher, D.R., 2011a. Delta-8 desaturation activity varies among fatty acyl desaturases of teleost fish: high activity in delta-6 desaturases of marine species. *Comp. Biochem. Physiol. B* 159, 206-213.
- Monroig, Ó., Webb, K., Ibarra-Castro, L., Holt, G.J., Tocher, D.R., 2011b. Biosynthesis of long-chain polyunsaturated fatty acids in marine fish: Characterization of an Elovl4-like elongase from cobia *Rachycentron canadum* and activation of the pathway during early life stages. *Aquaculture* 312, 145-153.
- Monroig, Ó., Wang, S., Zhang, L., You, C., Tocher, D.R., Li, Y., 2012. Elongation of long-chain fatty acids in rabbitfish *Siganus canaliculatus*: Cloning, functional

- characterisation and tissue distribution of Elovl5- and Elovl4-like elongases. *Aquaculture* 350-353, 63-70.
- Monroig, Ó., Tocher, D.R., Hontoria, F., Navarro, J.C., 2013. Functional characterisation of a *Fads2* fatty acyl desaturase with $\Delta 6/\Delta 8$ activity and an Elovl5 with C₁₆, C₁₈ and C₂₀ elongase activity in the anadromous teleost meagre (*Argyrosomus regius*). *Aquaculture* 412-413, 14-22.
- Monroig, O., Lopes-Marques, M., Navarro, J.C., Hontoria, F., Ruivo, R., Santos, M.M., Venkatesh, B., Tocher, D.R., Castro, L.F., 2016. Evolutionary functional elaboration of the Elovl2/5 gene family in chordates. *Sci. Rep.* 6, 20510.
- Monroig, O., Tocher, D.R., Castro, L.F.C., 2018. Polyunsaturated fatty acid biosynthesis and metabolism in fish, Polyunsaturated Fatty Acid Metabolism. In: Burdge, G.C. (Ed.), Polyunsaturated Fatty Acid Metabolism. Academic Press, London, pp. 31-60.
- Morais, S., Monroig, O., Zheng, X., Leaver, M.J., Tocher, D.R., 2009. Highly unsaturated fatty acid synthesis in Atlantic salmon: characterization of Elovl5- and Elovl2-like elongases. *Mar. Biotechnol.* 11, 627-639.
- Morais, S., Mourente, G., Ortega, A., Tocher, J.A., Tocher, D.R., 2011. Expression of fatty acyl desaturase and elongase genes, and evolution of DHA:EPA ratio during development of unfed larvae of Atlantic bluefin tuna (*Thunnus thynnus* L.). *Aquaculture* 313, 129-139.
- Morais, S., Castanheira, F., Martinez-Rubio, L., Conceicao, L.E., Tocher, D.R., 2012. Long chain polyunsaturated fatty acid synthesis in a marine vertebrate: ontogenetic and nutritional regulation of a fatty acyl desaturase with $\Delta 4$ activity. *Biochim. Biophys. Acta* 1821, 660-671.
- Oboh, A., Betancor, M.B., Tocher, D.R., Monroig, O., 2016. Biosynthesis of long-chain polyunsaturated fatty acids in the African catfish *Clarias gariepinus*: Molecular cloning and functional characterisation of fatty acyl desaturase (*fads2*) and elongase (*elovl2*) cDNAs. *Aquaculture* 462, 70-79.
- Oboh, A., Kabeya, N., Carmona-Antoñanzas, G., Castro, L.F.C., Dick, J.R., Tocher, D.R., Monroig, O., 2017. Two alternative pathways for docosahexaenoic acid (DHA, 22: 6n-3) biosynthesis are widespread among teleost fish. *Sci. Rep.* 7, 3889.
- Park, W.J., Kothapalli, K.S., Lawrence, P., Tyburczy, C., Brenna, J.T., 2009. An alternate pathway to long-chain polyunsaturates: the *FADS2* gene product $\Delta 8$ -desaturates 20: 2n-6 and 20: 3n-3. *J. Lipid Res.* 50, 1195-1202.
- Paulino, R.R., Pereira, R.T., Fontes, T.V., Oliva-Teles, A., Peres, H., Carneiro, D.J., Rosa, P.V., 2018. Optimal dietary linoleic acid to linolenic acid ratio improved fatty acid profile of the juvenile tambaqui (*Colossoma macropomum*). *Aquaculture* 488, 9-16.
- Pereira, R.T., Paulino, R.R., de Almeida, C.A.L., Rosa, P.V., Orlando, T.M., Fortes-Silva, R., 2017. Oil sources administered to tambaqui (*Colossoma macropomum*): growth, body composition and effect of masking organoleptic properties and fasting on diet preference. *Appl. Anim. Behav. Sci.* 199, 103-110.

- Ravi, V., Venkatesh, B., 2018. The divergent genomes of teleosts. *Annu. Rev. Anim. Biosci.* 6, 47-68.
- Rose, T.M., Henikoff, J.G., Henikoff, S., 2003. CODEHOP (CONsensus-DEgenerate hybrid oligonucleotide primer) PCR primer design. *Nucleic Acids Res.* 31, 3763-3766.
- Sperling, P., Ternes, P., Zank, T.K., Heinz, E., 2003. The evolution of desaturases. *Prostaglandins Leukot. Essent. Fat. Acids* 68, 73–95.
- Sprecher, H., 2000. Metabolism of highly unsaturated n-3 and n-6 fatty acids. *Biochim. Biophys. Acta* 1486, 219-231.
- Tanomman, S., Ketudat-Cairns, M., Jangprai, A., Boonanuntanasarn, S., 2013. Characterization of fatty acid delta-6 desaturase gene in Nile tilapia and heterogenous expression in *Saccharomyces cerevisiae*. *Comp. Biochem. Physiol. B* 166, 148-156.
- Tocher, D.R., Agaba, M., Hastings, N., Teale, A.J., 2003. Metabolism and functions of lipids and fatty acids in teleost fish *Rev. Fish. Sci.*, 107-184.
- Turchini, G.M., Torstensen, B.E., Ng, W.K., 2009. Fish oil replacement in finfish nutrition. *Rev. Aquacult.* 1, 10-57.
- Tocher, D.R., 2010. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquac. Res.* 41, 717-732.
- Tocher, D.R., 2015. Omega-3 long-chain polyunsaturated fatty acids and aquaculture in perspective. *Aquaculture* 449, 94-107.
- Vagner, M., Santigosa, E., 2011. Characterization and modulation of gene expression and enzymatic activity of delta-6 desaturase in teleosts: A review. *Aquaculture* 315, 131-143.
- Wang, S., Monroig, Ó., Tang, G., Zhang, L., You, C., Tocher, D.R., Li, Y., 2014. Investigating long-chain polyunsaturated fatty acid biosynthesis in teleost fish: Functional characterization of fatty acyl desaturase (Fads2) and Elovl5 elongase in the catadromous species, Japanese eel *Anguilla japonica*. *Aquaculture* 434, 57-65.
- Watanabe, K., Ohno, M., Taguchi, M., Kawamoto, S., Ono, K., Aki, T., 2016. Identification of amino acid residues that determine the substrate specificity of mammalian membrane-bound front-end fatty acid desaturases. *J. Lipid Res.* 57, 89-99.
- Zheng, X., Seiliez, I., Hastings, N., Tocher, D.R., Panserat, S., Dickson, C.A., Bergot, P., Teale, A.J., 2004. Characterization and comparison of fatty acyl Δ 6 desaturase cDNAs from freshwater and marine teleost fish species. *Comp. Biochem. Physiol. B* 139, 269-279.
- Zheng, X., Ding, Z., Xu, Y., Monroig, O., Morais, S., Tocher, D.R., 2009. Physiological roles of fatty acyl desaturases and elongases in marine fish: Characterisation of cDNAs of fatty acyl Δ 6 desaturase and *elovl5* elongase of cobia (*Rachycentron canadum*). *Aquaculture* 290, 122-131.

Figure Captions

Figure 1. Maximum likelihood phylogenetic analysis of Fads amino acid sequences rooted with the invertebrate clade. Numbers at nodes indicate branch support in posterior probabilities calculated using aBayes. *C. macropomum* Fads2 studied herein is highlighted. Accession numbers for all Fads sequences are available in Supplementary Table 2.

Figure 2. Maximum likelihood phylogenetic analysis of Elovl amino acid sequences rooted with the Elovl4 sequences. Numbers at nodes indicate branch support in posterior probabilities calculated using aBayes. *C. macropomum* sequences (Elovl2 and Elovl5) studied herein are highlighted. Accession numbers for all Elovl sequences are available in Supplementary Table 3.

Figure 3. The pathways of biosynthesis long-chain (C₂₀₋₂₄) polyunsaturated fatty acids in *C. macropomum* predicted from activity of Fads2, Elovl2 and Elovl5 measured in yeast. Desaturation reactions are indicated as “Δx”, whereas elongation reactions are indicated as “Elo”. LA, linoleic acid (18:2n-6); ALA, α-linolenic acid (18:3n-3); ARA, arachidonic acid; EPA, eicosapentaenoic acid; DHA, docosahexaenoic acid. β-ox, beta-oxidation.

Figures

Figure 1




Figure 2





Figure 3



Tables

Table 1. Functional characterization of the tambaqui (*C. macropomum*) Fads2 in yeast *S. cerevisiae*. Yeast were transformed with the *C. macropomum fads2* ORF and grown in the presence of $\Delta 6$ (18:2n-6 and 18:3n-3), $\Delta 8$ (20:2n-6 and 20:3n-3), $\Delta 5$ (20:3n-6 and 20:4n-3), and $\Delta 4$ (22:4n-6 and 22:5n-3) fatty acid (FA) substrates. Desaturation of 24:5n-3 was tested by co-expressing both *D. rerio elovl2* and *C. macropomum fads2* in *S. cerevisiae* (Oboh et al., 2017). Conversions in both expression systems were calculated according to the formula [all product areas/(all product areas + substrate area)] \times 100.

FA substrate	FA product	Conversion (%)	Activity
18:2n-6	18:3n-6	28.3	$\Delta 6$
18:3n-3	18:4n-3	63.5	$\Delta 6$
20:2n-6	20:3n-6	2.8 ^a	$\Delta 8$
20:3n-3	20:4n-3	4.5 ^a	$\Delta 8$
20:3n-6	20:4n-6	14.8	$\Delta 5$
20:4n-3	20:5n-3	17.8	$\Delta 5$
22:4n-6	22:5n-6	Nd	$\Delta 4$
22:5n-3	22:6n-3	Nd	$\Delta 4$
Expressed with <i>D. rerio elovl2</i>			
18:3n-3	18:4n-3	20.5	$\Delta 6$
24:5n-3	24:6n-3	21.9	$\Delta 6$

^a Conversions of $\Delta 8$ substrates (20:2n-6 and 20:3n-3) include stepwise reactions due to multifunctional desaturation abilities. Thus, the conversions of the *C. macropomum* Fads2 on 20:2n-6 and 20:3n-3 include the $\Delta 8$ desaturation toward 20:3n-6 and 20:4n-3, respectively, and their subsequent $\Delta 5$ desaturations to 20:4n-6 and 20:5n-3, respectively.

Nd, not detected.

Table 2. Functional characterization of the tambaqui (*C. macropomum*) Elov15 and Elov12 elongases in yeast *S. cerevisiae*. Yeast transformed with the *C. macropomum* *elov15* and *elov12* ORF sequences and grown in the presence of exogenously added fatty acid (FA) substrates. Conversions by Elov15 and Elov12 were calculated according to the formula [all product areas/(all product areas + substrate area)] × 100.

Conversion (%)				
FA substrate	FA product	Elov15	Elov12	Activity
18:2n-6	20:2n-6	29.8	1.5	C ₁₈ →C ₂₀
18:3n-3	20:3n-3	47.7	7.6	C ₁₈ →C ₂₀
18:3n-6	20:3n-6	79.8	13.5	C ₁₈ →C ₂₀
18:4n-3	20:4n-3	79.0	26.3	C ₁₈ →C ₂₀
20:4n-6	22:4n-6	33.4	30.9	C ₂₀ →C ₂₂
20:5n-3	22:5n-3	53.1	66.5	C ₂₀ →C ₂₂
22:4n-6	24:4n-6	Nd	16.4	C ₂₂ →C ₂₄
22:5n-3	24:5n-3	Nd	21.8	C ₂₂ →C ₂₄

Nd, not detected.

Supplementary Material

Supplementary Table 1. Primer sets and corresponding PCR conditions used in the cloning of the *C. macropomum fads2*, *elovl5* and *elovl2* ORF sequences.

Supplementary Table 2. Accession numbers of all sequences used in phylogenetic analysis of Fads amino acid sequences.

Supplementary Table 3. Accession numbers of all sequences used in phylogenetic analysis of Elovl amino acid sequences.

Supplementary Figure 1. Sequence alignment and conservation analysis of desaturase sequence motifs of *C. macropomum* Fads2. *- indicates residues involved in the determination of the desaturation activity in mammalian desaturases (Watanabe et al., 2016). Red boxes highlight conserved residue replacements with *Danio rerio* $\Delta 5\Delta 6$ Fads2.

Supplementary Table 1. Primer sets and corresponding PCR conditions used in the cloning of the *C. macropomum fads2*, *elovl5* and *elovl2* ORF sequences.

	Primer set function	Primer name	Primer sequence	Cycles	TM	Extension (size bp)
FADS2	Degenerate primers	FADS2_degen_F	GCGCCTCCGCCAAygtggaayc	40	54 °C	72 °C/10 s
		FADS2_degen_R	TGGCCGGAGAACcartcrttraa			
	Gene specific race	Cma_FADS2_3RACE_F	GGAGTCTTCGGATCATTTGCGCTTC	45	65 °C	72 °C/15 s
		Cma_FADS2_5RACE_R	GCAGAGGAGGACCAATCAGGAAGAA			
	Full ORF	Cma_FADS2_ORF_F	TCATCAGAGAGAGCAGCGAG	35	59 °C	72 °C/24 s
		Cma_FADS2_ORF_R	CCAGCATAGATGGCAGAGGA			
Restriction site	Cma_FADS2_Pyes_KPNI_F	CCC <u>GGTACC</u> ATAATGGGTGGGGGCACTCAT	35	68 °C	72 °C/21 s	
	Cma_FADS2_Pyes_XBAL_R	CCCTCTAGATTATTTGTGGAGGTATGCGTCC				
ELOVL5	Degenerate primers	ELOVEL5_degen_F	TGAACGTCCTGTgtggtaytayt	45	52 °C	72 °C/5 s
		ELOVEL5_degen_R	GCTGCACCTGGGTGATGTACykyttccacca			
	Gene specific race	Cma_ELOVEL5_3RACE_F	TGGACACGTTCTTCTTCATCCTGCG	20	65 °C	72 °C/8 s
			nested Cma_ELOVEL5_3RACE_F2	GTCCTGTGCTGTAGTCTGGCCCTGC	25	65 °C
		Cma_ELOVEL5_5RC_R	CGCAGGATGAAGAAGAACGTGTCCA	45	65 °C	72 °C/5 s
	Restriction site	Cma_ELOVEL5_Pyes_KPNI_F	CCC <u>GGTACCA</u> AAGATGGAGGCCCTTTAATCACA	35	58 °C	72 °C/15 s
Cma_ELOVEL5_Pyes_XBAL_R		CCCTCTAGATCAATCTGCCCGCGGCTT				
ELOVL2	Degenerate primers	ELOVL2_degen_F	TACTTGGGACCAAAGTACATGA	45	60 °C	72 °C/7 s
		ELOVL2_degen_R	AGATAGCGTTTCCACCACAG			
	Gene specific race	Cma_ELOVL2_3RACE_F	ACCACGCCTCCATGTTCAATATCTGGTG	45	72 °C	72 °C/7 s
		Cma_ELOVL2_5RACE_R	CCCGCTGACCAGGTGTGTAATAA	45	69 °C	73 °C/7 s
	Full ORF	Cma_ELOVL2_ORF_F	GAGAGGCGCGGCGAGGAAACAAC	35	68 °C	73 °C/19s
		Cma_ELOVL2_ORF_R	GGCTGTGGTTGTGCATATGTCTCAG			
Restriction site	Cma_ELOVL2_Pyes_KPNI_F	CCC <u>GGTACC</u> ACCATGGAGCTCTTTAGCATGAA	35	62 °C	73 °C/14 s	
	Cma_ELOVL2_Pyes_XBAL_R	CCCTCTAGATTACTGTAGCTTATGTTTGGCTCC				

Supplementary Table 2. Accession numbers of all sequences used in phylogenetic analysis of Fads amino acid sequences.

Accession numbers	
Fads 1	
<i>Homo sapiens</i>	NP_037534.3
<i>Sus scrofa</i>	NP_001106512.1
<i>Mus musculus</i>	AAH26848.1
<i>Rattus norvegicus</i>	NP_445897.2
<i>Monodelphis domestica</i>	H9H609
<i>Ornithorhynchus anatinus</i>	XP_016084018.1
<i>Latimeria chalumnae</i>	XP_005988035.1
<i>Scyliorhinus canicula</i>	AEY94454.1
<i>Callorhynchus milii</i>	XP_007885635.1
<i>Xenopus tropicalis</i>	XP_002943012.2
<i>Xenopus laevis</i>	XP_018112981.1
<i>Xenopus laevis</i>	XP_018115603.1
<i>Anolis carolinensis a</i>	XP_003224167.1
<i>Anolis carolinensis c</i>	XP_003224188.1
<i>Anolis carolinensis d</i>	XP_003224187.1
<i>Gallus gallus a</i>	XP_421052.4
<i>Gallus gallus b</i>	XP_426408.2
<i>Gallus gallus c</i>	XP_421051.3
<i>Alligator mississippiensis</i>	XP_006274989.1
Fads 2	
<i>Homo sapiens</i>	AAG23121.1
<i>Mus musculus</i>	NP_062673.1
<i>Rattus norvegicus</i>	NP_112634.1
<i>Sus scrofa</i>	NP_001165221.1
<i>Anolis carolinensis</i>	XP_003224168.1
<i>Gallus gallus</i>	NP_001153900.1
<i>Alligator mississippiensis</i>	XP_006274951.1
<i>Xenopus tropicalis</i>	NP_001120262.1
<i>Salmo salar</i> (Fads2Δ6 _c)	NP_001165752.1
<i>Salmo salar</i> (Fads2Δ6 _b)	NP_001165251.1
<i>Salmo salar</i> (Fads2Δ6 _a)	NP_001117047.1
<i>Salmo salar</i> (Fads2Δ5)	NP_001117014.1
<i>Oncorhynchus masou</i>	ABU87822.1
<i>Oncorhynchus mykiss</i> (Fads2Δ6)	NP_001117759.1
<i>Oncorhynchus mykiss</i> (Fads2Δ5)	AFM77867.1
<i>Gadus morhua</i> (Fads2Δ6)	AAV46796.1
<i>Scatophagus argus</i> (Fads2Δ6)	AHA62794.1
<i>Oreochromis niloticus</i> (Fads2Δ6)	AGV52807.1
<i>Dicentrarchus labrax</i> (Fads2Δ6)	ACD10793.1

<i>Argyrosomus regius</i> (Fads2Δ6/ Δ8)	AGG69480.1
<i>Sparus aurata</i> (Fads2Δ6)	AAL17639.1
<i>Thunnus maccoyii</i> (Fads2Δ6)	ADG62353.1
<i>Scophthalmus maximus</i> (Fads2Δ6)	AAS49163.1
<i>Solea senegalensis</i> (Fads2Δ6)	AEQ92868.1
<i>Siganus canaliculatus</i> (Fads2Δ4)	ADJ29913.1
<i>Lates calcarifer</i> (Fads2Δ6)	ACS91458.1
<i>Rachycentron canadum</i> (Fads2Δ6)	ACJ65149.1
<i>Anguilla japonica</i> (Fads2Δ6/ Δ8)	AHY22375.1
<i>Arapaima gigas</i> (Fads2Δ6/ Δ8)	AOO19789.1
<i>Scleropages formosus</i>	XP_018585703.1
<i>Danio rerio</i> (Fads2Δ5/ Δ6)	NP_571720.2
<i>Colossoma macropomum</i>	MH734335
<i>Astyanax mexicanus</i>	XP_007235183.1
<i>Ictalurus punctatus</i>	XP_017341187.1
<i>Clarias gariepinus</i> (Fads2Δ5/ Δ6)	AMR43366.1
<i>Latimeria chalumnae</i>	XP_005988034.2
<i>Scyliorhinus canicula</i>	AEY94455.1
<i>Callorhynchus milii</i>	XP_007885636.1
invertebrates	Fads
<i>Saccoglossus kowalevskii</i>	XP_006822674.1
<i>Octopus vulgaris</i>	AEK20864.1

Supplementary Table 3. Accession numbers of all sequences used in phylogenetic analysis of Elovl amino acid sequences.

Accession numbers Elovl	
	Elovl2
<i>Homo sapiens</i>	NP_060240.3
<i>Bos taurus</i>	NP_001076986.1
<i>Mus musculus</i>	NP_062296.1
<i>Rattus norvegicus</i>	NP_001102588.1
<i>Gallus gallus</i>	NP_001184237.1
<i>Xenopus tropicalis</i>	NP_001016159.1
<i>Xenopus laevis</i>	NP_001087564.1
<i>Callorhinchus milii</i>	XP_007900820.1
<i>Latimeria chalumnae</i>	XP_006006450.1
<i>Esox lucius</i>	XP_010884057.1
<i>Salmo salar</i>	NP_001130025.1
<i>Oncorhynchus mykiss</i>	AIT56593.1
<i>Clupea harengus</i>	XP_012671565.1
<i>Danio rerio</i>	NP_001035452.1
<i>Colossoma macropomum</i>	MH734337
<i>Astyanax mexicanus</i>	XP_007260136.1
<i>Clarias gariepinus</i>	AOY10780.1
	Elovl5
<i>Homo sapiens</i>	NP_068586.1
<i>Mus musculus</i>	NP_599016.2
<i>Bos taurus</i>	NP_001040062.1
<i>Gallus gallus</i>	NP_001186126.1
<i>Xenopus laevis</i>	NP_001089883.1
<i>Sparus aurata</i>	AAT81404.1
<i>Dicentrarchus labrax</i>	CBX53576.1
<i>Rachycentron canadum</i>	ACJ65150.1
<i>Salmo salar</i> (Elovl5b)	NP_001130024.1
<i>Salmo salar</i> (Elovl5a)	NP_001117039.1
<i>Larimichthys crocea</i>	AFB81415.1
<i>Nibeia mitsukurii</i>	ACR47973.1
<i>Argyrosomus regius</i>	AGG69479.1
<i>Siganus canaliculatus</i>	ADE34561.1
<i>Clupea harengus</i>	XP_012695835.1
<i>Clarias gariepinus</i>	AAT81405.1
<i>Ictalurus punctatus</i>	NP_001188041.1
<i>Pangasius larnaudii</i>	AGR45586.1
<i>Danio rerio</i>	NP_956747.1
<i>Colossoma macropomum</i>	MH734336
<i>Branchiostoma lanceolatum</i>	ALZ50284.1

Elov14

<i>Homo sapiens</i>	NP_073563.1
<i>Mus musculus</i>	NP_683743.2
<i>Bos taurus</i>	NP_001092520.1
<i>Gallus gallus</i>	NP_001184238.1
<i>Danio rerio a</i>	NP_957090.1
<i>Danio rerio b</i>	NP_956266.1
<i>Takifugu rubripes a</i>	XP_003966009.1
<i>Takifugu rubripes b</i>	XP_003971605.1
<i>Salmo salar</i>	NP_001182481.1
<i>Rachycentron canadum</i>	ADG59898.1
<i>Ciona intestinalis</i>	NP_0010290

Supplementary Figure 1. Sequence alignment and conservation analysis of destaturase sequence motifs of *C. macropomum* Fads2. *- indicates residues involved in the determination of the destaturation activity in mammalian desaturases (Watanabe et al., 2016). Red boxes highlight conserved residue replacements with *Danio rerio* Δ5Δ6 Fads2.

