

A Complete Formulation of Baum-Connes' Conjecture for the Action of Discrete Quantum Groups

D. Goswami

Indian Statistical Institute

Kolkata, India

We formulate a version of Baum-Connes' conjecture for a discrete quantum group, building on our earlier work ([9]). Given such a quantum group \mathcal{A} , we construct a directed family $\{\mathcal{E}_F\}$ of C^* -algebras (F varying over some suitable index set), borrowing the ideas of [5], such that there is a natural action of \mathcal{A} on each \mathcal{E}_F satisfying the assumptions of [9], which makes it possible to define the "analytical assembly map", say $\mu_i^{r,F}$, $i = 0, 1$, as in [9], from the \mathcal{A} -equivariant K -homology groups of \mathcal{E}_F to the K -theory groups of the "reduced" dual $\hat{\mathcal{A}}_r$. As a result, we can define the Baum-Connes' maps $\mu_i^r : \varinjlim KK_i^{\mathcal{A}}(\mathcal{E}_F, \mathbb{C}) \rightarrow K_i(\hat{\mathcal{A}}_r)$, and in the classical case, i.e. when \mathcal{A} is $C_0(G)$ for a discrete group, the isomorphism of the above maps for $i = 0, 1$ is equivalent to the Baum-Connes' conjecture. Examples are discussed.

References

- [1] **P. Baum and A. Connes**, *Geometric K-theory for Lie groups and foliations*, Enseign. Math. **46** (2000), 3–42.
- [2] **P. Baum, A. Connes, and N. Higson**, *Classifying space for proper actions and K-theory of group C*-algebras*, in: *C*-algebras*, Contemp. Math., 167, Amer. Math. Soc., Providence, RI, 1994, 240–291
- [3] **S. Baaj and G. Skandalis**, *C*-algebres de Hopf et thorie de Kasparov èquivariante*, *K-Theory* **2** (1989), 686–721.
- [4] **A. Connes**, *Noncommutative Geometry*, Academic Press, San Diego, CA ,1994.
- [5] **J. Cuntz**, *Noncommutative Simplicial Complexes and the Baum-Connes Conjecture*, preprint, math.OA 0103219 (2001).
- [6] **V. G. Drinfeld**, *Quantum Groups*, Proc. I.C. M. (1,2), Berkeley, California, 1986, pp. 798-820.
- [7] **D. A. Ellwood**, *A New Characterisation of Principal Actions*, *J. Funct. Anal.* **173** (2000), 49-60.
- [8] **E. G. Effros and Z-J. Ruan**, *Discrete quantum groups. I. The Haar measure*, *Internat. J. Math.* **5**(1994), 681-723.
- [9] **D. Goswami and A. O. Kuku**, *Towards the Baum-Connes' Analytical Assembly Map for the Actions of Discrete Quantum Groups*, ICTP preprint, also available at math.KT 0207303.
- [10] **K. K. Jensen and K. Thomsen**, *Elements of KK-Theory*, in: *Mathematics: Theory and Applications*, Birkhäuser Boston, Inc., Boston, MA, 1991, pp. viii-202.
- [11] **M. Jimbo**, *A q-analogue of $U(\mathfrak{gl}(N + 1))$, Hecke algebra and Yang Baxter equation*, *Lett. Math. Phys.* **11** (1986), 247-252.

- [12] **J. Kustermans**, *The analytic structure of an algebraic quantum group*, preprint, available at funct-an/9707010.
- [13] **R. Meyer**, *Generalized fixed point algebras and square-integrable groups actions*, *J. Funct. Anal.* **186** (2001), 167–195.
- [14] **P. Podleś**, *Symmetries of quantum spaces. Subgroups and quotient spaces of quantum $SU(2)$ and $SO(3)$ groups*, *Comm. Math. Phys.* **170** (1995), 1–20.
- [15] **A. Van Daele**, *Discrete quantum groups*, *J. Algebra* **180** (1996), 431–444.
- [16] **A. Van Daele**, *Multiplier Hopf *-algebras with positive integrals: A laboratory for locally compact quantum groups*, preprint, available at math.OA/0205285.
- [17] **A. Van Daele**, *Multiplier Hopf algebras*, *Trans. Amer. Math. Soc.* **342** (1994), 917–932.
- [18] **A. Valette**, *Introduction to the Baum-Connes Conjecture*, ETH (Zurich) Lecture Notes, Birkhäuser, to appear (also available at <http://www.math.ethz.ch/indra/BC.dvi>.)
- [19] **A. Valette**, *On the Baum-Connes assembly map for discrete groups*, available at <http://www.unine.ch/math/preprints/preprints.html>.
- [20] **S. L. Woronowicz**, *Twisted $SU(2)$ group: an example of a noncommutative differential calculus*, *Publ. R.I.M.S.* **23**(1987), 117–181.