


Published in final edited form as:

J Org Chem. 2005 August 19; 70(17): 6907–6912. doi:10.1021/jo051002a.

A Convenient Preparation of Xanthene Dyes

Youjun Yang, Jorge O. Escobedo, Alexander Wong, Corin M. Schowalter, Michael C. Touchy, Lijuan Jiao, William E. Crowe, Frank R. Fronczek, and Robert M. Strongin*
 Department of Chemistry, Louisiana State University, Baton Rouge, Louisiana 70803

Abstract



A facile synthetic route utilizing readily available reagents affords a series of regioisomerically pure xanthene dye derivatives. Advantages include relatively mild conditions and good to excellent yields. Nonpolar, highly crystalline intermediates are isolable by standard chromatographic techniques. The intermediates are in the requisite xanthene oxidation state, thus avoiding the need for relatively inefficient oxidation chemistry and/or harsh conditions. During the course of this work, a new boron-mediated 1,2-aryl migration reaction was discovered.

Introduction

Fluorescein and fluorone are structurally related xanthene dyes (Figure 1). Fluorone is formally decarboxylated fluorescein. Fluorone derivatives have found numerous applications. They have been used in the detection of a variety of metal ions,¹ sugars,² phosphorylated molecules,³ HIV-1 nucleocapsid protein,⁴ reactive oxygen species,⁵ in screening assays for mitochondrial permeability,⁶ acetylcholinesterase inhibition,⁷ and telomerase inhibition.⁸ We have reported the use of fluorones as sialic acid⁹ and homocysteine probes.¹⁰

The initial fluorone synthesis was reported by Mohlau and Koch.¹¹ Typical syntheses include the following sequence: (i) formation of the leuco base via the condensation of resorcinol with aldehydes under thermal and acid-catalyzed conditions and (ii) the formation of dye via the oxidation of the leuco base. It has been reported that the oxidation step is often low-yielding. Additionally, there can be purification problems due to the formation of byproducts and the relatively polar nature of fluorone derivatives.¹²

Recently, a novel and innovative microwave protocol resulted in improved yields in related rosamine derivative syntheses.^{12d} The microwave-assisted condensation and oxidation gave relatively higher yields (typically ranging from 27 to 73%) than traditional thermal conditions (typical ranges from 8 to 35%).

Fluorone dye preparations have also been reported earlier by Neckers et al. as part of his extensive and pioneering work on xanthenes, involving heating intermediates in a sealed

tube at 200 °C.¹³ Benzophenones have also been used as condensation partners in high-temperature, acid-catalyzed syntheses of xanthenes.¹⁴ A novel oxidation reaction of substituted triphenylmethanes has been reported to occur via Br₂/CCl₄.¹⁵

Herein, we report a simple new protocol for attaining a series of fluorone derivatives. It is based on forming tertiary carbinol leuco bases via Grignard reactions. Treatment of carbinol leuco bases with BBr₃ affords the desired dyes. The carbinol intermediate is much less polar than the dye products. It can thus be purified via standard column chromatography. These intermediates are often highly crystalline (vide infra). The carbinol carbon is already in the correct oxidation state of the desired dyes. Therefore, potentially troublesome oxidations are avoided. The dye products can be purified by simple filtration methods without chromatography. The use of low-temperature, basic conditions may serve as an attractive alternative to more common, relatively harsh acid-catalyzed and oxidative methods. In addition, during the course of this work we discovered a novel 1,2-aryl migration mediated by boron that furnishes potentially useful dendrimer cores as well as dye architectures.

Results and Discussion

In our initial investigations, we use free radical bromination of appropriate triaryls as a means to obtain dye precursors with the requisite oxidation state. Tetramethyl ether **2** is obtained in nearly quantitative yield by stirring **1**¹⁶ via known procedures.¹⁷ Compound **2** (500 mg in 20 mL of CCl₄) is treated with NBS (1.05 equiv) in the presence of a catalytic amount of (PhCO₂)₂ in CCl₄ (Scheme 1).

Upon washing the crude brominated product with a saturated NH₄Cl solution, carbinol **4** is obtained in a 13% yield. Reverse condensation product **5** and brominated congener **6** are also obtained in 6% and 15% yields, respectively. The mechanism of resorcinarene reverse condensation has previously been studied in great detail.^{12c,18} Compounds **5** and **6** were synthesized independently.¹⁹ The structures of **1**, **2**, **4**, **5**, and **6** are verified by single-crystal X-ray structural analysis (Supporting Information).

Since solvolysis of bromide **3** occurs readily during workup,¹⁵ we decided to synthesize the carbinol precursor directly. It is conveniently accessed by the reaction of readily available Grignard reagents and esters. Compound **9** is obtained in one step in a yield of 96% by reacting 2,4-dimethoxybenzenemagnesium bromide **7** and methyl benzoate **8** (Scheme 2). The structure of **9** is confirmed by X-ray crystallography (Supporting Information). When **9** is treated with BBr₃ (6 equiv), monomethyl ether **10** is obtained in 64% yield. Using a greater excess of BBr₃ (16 equiv) furnishes fully deprotected **11** in a 65% yield. Compound **11** is conveniently isolated by filtration (Scheme 3).

It is straightforward to apply the above protocol to the synthesis of a series of regioisomerically pure fluorone dyes. When various methyl benzoates **12a–e** are used, the corresponding carbinols **13a–e** are obtained in excellent yields (>90%). The reaction of the carbinols **13a–d** with 16 equiv of BBr₃ furnishes the corresponding fluorone dyes with yields ranging from 70 to 88% while deprotection of **13e** affords methyl ether **15** (Table 1). Compound **15** is completely deprotected using 20 equiv of BBr₃ to furnish **14e** in a 76% yield (Scheme 4). In each case, the fluorone products are obtained without preparative chromatography. The structures of **13b** and **13e** are confirmed by single-crystal X-ray structure analysis (Supporting Information).

To prepare a potential sialic acid sensor **16**,⁹ carbinol **17** was targeted (Scheme 5).

The reaction of **7** (2.4 equiv) and **18** (1 equiv), however, affords compound **19** in 13% yield. When 3.6 equiv of **7** is used, the yield of **19** improves to 32%. In each run, 2,4-dimethoxyphenylboronic acid **20** is isolated in trace amounts (Scheme 6). Structures of **18**, **19** (Figure 2), and **20** were confirmed by single-crystal X-ray structure analysis (Supporting Information).

A proposed mechanism for the formation of compound **19** is shown in Scheme 7. The first equivalent of **7** reacts with boron to form the tetrahedral anionic boronate **21**. This is supported by the fact that **18** is quantitatively recovered when 1 equiv of **7** is added to a solution of **18** and stirred overnight before quenching with water. The second and third equivalents of **7** react with the ester to afford the tertiary oxide **22**. Upon workup, a 1,2-aryl shift mediated by boron affords **19**.

1,2-Alkyl shifts have been known for decades.²⁰ In 1963, Matteson described the prototypical 1,2-alkyl shift involving α -haloalkyl borates.²¹ Negishi and co-workers established the analogous 1,2-rearrangement of 1-chloroalkyl complexes of Al, Mg, Zn, Cd, Ti, Zr, Hf, V, Cr, Mn, Fe, Co, Ni, and Cu.²² It has also been found that organoboronate complexes bearing α,β -unsaturation and leaving groups at the α - and γ -carbons can undergo a 1,2-alkyl group migration to afford homologated organoboron compounds.²³ A recent article described a transformation whereby an organozincate bearing a leaving group at the remote benzylic position would rearrange with concomitant carbon-carbon bond formation to furnish the benzylzincate^{22j,24} (Scheme 8). In the current case (Scheme 7), the boronic acid is not found in the final product. This is the first observation of a 1,2-aryl group migration mediated by an arylboronate complex involving a leaving group at a remote benzylic position. Compound **19** and congeners may serve as functional dye and/or dendrimer substrates.

Conclusion

Carbinol leuco bases are easily prepared and purified precursors for fluorone dyes. A novel reaction that involves a boron-mediated 1,2-aryl shift reaction with a leaving group at a remote benzylic position is described. The methods described herein are now being utilized in our lab toward the synthesis of new naphthofluorescein dye architectures currently unattainable via other methodologies.

Experimental Section

5,5'-Dichloro-2,2',4,4'-tetramethoxytrityl Alcohol (**4**)

In a 50-mL round-bottom flask, compound **2** (0.500 g, 1.18 mmol) and NBS (0.315 g, 1.77 mmol) are dissolved in 20 mL of CCl₄. A catalytic amount of benzoyl peroxide (~10 mg) is added. The reaction mixture is heated at reflux for 1 h with vigorous stirring. It is cooled to room temperature and quenched with 50 mL of saturated NH₄Cl (aq). The mixture is extracted with CH₂Cl₂ (3 × 15 mL). The combined extracts are dried over MgSO₄ and filtered, and the filtrate is evaporated to dryness. The solid residue is purified by flash column chromatography (silica gel; CH₂Cl₂) to give 72 mg (14%) of **4** along with 18 mg (6%) of **5** and 43 mg (15%) of **6**. The structures of **5** and **6** were verified via independent syntheses (vide infra). Data for **4**: ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.23–7.16 (m, 5H), 7.05 (s, 2H), 6.76 (s, 2H), 5.52 (s, 1H), 3.88 (s, 6H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 156.6, 154.4, 146.0, 129.0, 127.3, 127.3, 127.0, 126.4, 111.3, 98.8, 78.0, 56.2, 55.9. MALDI-TOF *m/z* 431.517 [M – OH]⁺, 447.544 [M]⁺.

5-Chloro-2,4-dimethoxybenzophenone (5)

A 20-mL volume of CH₃CN solution containing 2,4-dimethoxybenzophenone (0.500 g, 2.07 mmol) and NCS (0.200 g, 2.17 mmol) is mixed with 100 mg of 60 Å silica gel. The reaction mixture is heated at reflux for 5 h with vigorous stirring and cooled to room temperature. Silica gel is removed via suction filtration, and CH₃CN is removed under reduced pressure. The residue is purified by flash column chromatography (silica gel; CH₂-Cl₂) to afford 0.470 g (82%) of **5**. ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.68–7.60 (m, 3H), 7.52–7.70 (m, 2H), 7.39 (s, 1H), 6.89 (s, 1H), 3.97 (s, 3H), 3.69 (s, 3H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 193.6, 157.7, 157.6, 137.6, 133.1, 130.1, 129.1, 128.5, 121.0, 112.5, 98.1, 56.6, 56.2. MALDI-TOF *m/z* 276.1 [M]⁺, 277.4 [M + H]⁺, 299.0 [M + Na]⁺, 314.9 [M + K]⁺.

1-Bromo-5-chloro-2,4-dimethoxy-benzene (6)

This compound was prepared following the procedures above for compound **5** except that 2,4-dimethoxybromobenzene (1.507 g, 6.9 mmol) is used instead of 2,4-dimethoxybenzophenone. The yield of **6** (1.22 g) is 70%. ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.55 (d, *J* = 1.2 Hz, 1H), 6.83 (d, *J* = 1.2 Hz, 1H), 3.89 (s, 3H), 3.88 (s, 3H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 155.4, 155.0, 132.1, 112.7, 100.5, 98.6, 56.6, 56.5. MALDI-TOF *m/z* 247.890 [M]⁺.

2,2',4,4'-Tetramethoxytrityl Alcohol (9)

Magnesium turnings (0.543 g, 22.3 mmol) and a few crystals of I₂ are placed in a 250-mL three-neck round-bottom flask fitted with a dropping funnel and a condenser. A solution of 2,4-dimethoxybromobenzene (5.0 g, 23.0 mmol) in 20 mL of anhydrous THF is added dropwise to the magnesium. The mixture is stirred for 20–30 min. The resulting Grignard reagent (2,4-dimethoxyphenyl-magnesium bromide) is cooled in a dry ice/acetone bath before a solution of methyl benzoate (1.25 g, 9.38 mmol) in 40 mL of dry THF is added dropwise. The mixture is stirred overnight and then quenched with 100 mL of distilled water and neutralized with 2 N HCl. The unreacted 2,4-dimethoxybromobenzene as well as THF are removed by steam distillation. The resulting mixture is extracted with CH₂Cl₂ (3 × 30 mL). The combined extracts are dried over MgSO₄ and filtered, and the filtrate is evaporated to dryness. The residue is purified by flash chromatography (silica gel; EtOAc–hexane, 20:80) to afford 3.49 g (96%) of **9**. Compound **9** was prepared previously via other methodology.²⁵ NMR data is in agreement with the assigned structure: ¹H NMR (DMSO-*d*₆, 300 MHz) δ 7.21–7.10 (m, 5H), 6.86 (d, *J* = 8.6 Hz, 2H), 6.52 (d, *J* = 2.2 Hz, 2H), 6.42 (dd, *J* = 8.6, 2.2 Hz, 2H), 5.17 (s, 1H), 3.73 (s, 6H), 3.40 (s, 6H); ¹³C NMR (DMSO-*d*₆, 75.5 MHz) δ 159.7, 157.9, 147.4, 129.2, 127.4, 126.7, 127.9, 104.1, 99.5, 79.1, 55.4, 55.1.

9-Phenyl-6-methoxy-3-fluorone (10)

A solution of **8** (0.178 g, 0.468 mmol) in 10 mL of dry CH₂Cl₂ is cooled to –78 °C using a dry ice/acetone bath before BBr₃ (0.935 g, 3.74 mmol) is added dropwise. The mixture is allowed to warm to room temperature gradually before being quenched with 20 mL of distilled H₂O. The mixture is extracted with CH₂Cl₂ (3 × 10 mL). The combined extracts are dried over MgSO₄ and filtered, and the filtrate is concentrated under reduced pressure. The residue is purified by flash chromatography (silica gel, EtOAc), affording 90 mg (64%) of **10**. ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.63–7.61 (m, 3H), 7.47–7.43 (m, 2H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.08 (d, *J* = 9.0 Hz, 1H), 7.01 (d, *J* = 9.7 Hz, 1H), 6.94 (dd, *J* = 9.0, 2.4 Hz, 1H), 6.42 (dd, *J* = 9.7, 1.8 Hz, 1H), 6.21 (d, *J* = 1.8 Hz, 1H), 3.90 (s, 3H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 183.8, 164.1, 158.4, 154.0, 149.2, 132.4, 130.8, 129.7, 129.6, 129.5, 129.4, 128.8, 117.0, 113.9, 113.6, 104.7, 100.6, 56.3. MALDI-TOF *m/z* 303.154 [M + H]⁺.

9-Phenyl-6-hydroxy-3-fluorone (**11**)

To a stirred solution of **8** (0.400 g, 1.05 mmol) in 10 mL of dry CH₂Cl₂ at -78 °C, BBr₃ (4.20 g, 16.8 mmol) is added dropwise. The mixture is warmed to room temperature gradually before being quenched with 20 mL of distilled H₂O. After being stirred for 20 min, filtration leads to a collection of a red precipitate (**11**). A sample for analytical purposes is obtained by flash chromatography (silica gel, EtOAc–MeOH 9:1). A quantity of 197 mg (65%) of **11** is collected. Compound **11** is known; however, no characterization data was previously reported via the older synthetic methods.²⁶ ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.63–7.60 (m, 3H), 7.46–7.43 (m, 2H), 7.01 (d, *J* = 9.2 Hz, 2H), 6.60 (dd, *J* = 9.2, 2.1 Hz, 2H) 6.60 (d, *J* = 2.1 Hz, 2H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 156.3, 149.8, 132.5, 130.5, 129.5, 129.3, 128.8, 114.6, 103.4.

2,2',4,4'-Tetramethoxy-4''-bromotrityl Alcohol (**13a**)

This compound was prepared following the protocol described above for compound **9** except that 4-bromo methyl benzoate **12a** (2.01 g, 9.38 mmol) was used as the ester. The yield of **13a** (3.97 g) is 92%. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 7.25 (d, *J* = 8.5 Hz, 2H), 6.94 (d, *J* = 8.5 Hz, 2H), 6.83 (d, *J* = 8.6 Hz, 2H), 6.42 (d, *J* = 2.0 Hz, 2H), 6.34 (dd, *J* = 8.6, 2.0 Hz, 2H), 5.17 (s, 1H), 3.63 (s, 6H), 3.32 (s, 6H); ¹³C NMR (DMSO-*d*₆, 75.5 MHz) δ 159.9, 157.8, 147.2, 129.7, 129.4, 129.1, 125.9, 118.9, 104.2, 99.4, 78.5, 55.3, 55.1. MALDI-TOF *m/z* 457.880 [M]⁺, 441.521 [M – OH]⁺, 481.532 [M + K]⁺.

2,2',4,4'-Tetramethoxy-4''-phenyltrityl Alcohol (**13b**)

This compound is prepared following the procedure described above for compound **9** except that 4-phenyl methyl benzoate **12b** (1.13 g, 5.33 mmol) is used. The yield of **13b** (2.43 g) is 99%. ¹H NMR (DMSO-*d*₆, 250 MHz) δ 7.64 (d, *J* = 7.6 Hz, 2H), 7.47 (d, *J* = 8.4 Hz, 2H), 7.43 (t, *J* = 7.8 Hz, 2H), 7.32 (t, *J* = 7.1 Hz, 1H), 7.19 (d, *J* = 8.4 Hz, 2H), 6.93 (d, *J* = 8.6 Hz, 2H), 6.55 (d, *J* = 2.3 Hz, 2H), 6.45 (dd, *J* = 8.6, 2.3 Hz, 2H); ¹³C NMR (DMSO-*d*₆, 62.9 MHz) δ 159.7, 157.9, 146.7, 140.0, 137.6, 129.2, 128.9, 128.1, 127.1, 126.6, 126.5, 125.0, 104.1, 99.5, 78.9, 55.4, 55.1. MALDI-TOF *m/z* 439.654 [M – OH]⁺, 479.599 [M + Na]⁺.

2,2',4,4'-Tetramethoxy-4''-methoxytrityl Alcohol (**13c**)

This compound is prepared following the procedures described above for compound **9** except that 4-methoxy methyl benzoate **12b** (1.55 g, 9.38 mmol) is used. The yield of **13c** (3.2 g) is 83%. ¹H NMR (CDCl₃, 300 MHz) δ 7.16 (d, *J* = 6.9 Hz, 2H), 6.83 (d, *J* = 8.6 Hz, 2H), 6.79 (d, *J* = 6.9 Hz, 2H), 6.49 (d, *J* = 2.4 Hz, 2H), 6.37 (dd, *J* = 8.6, 2.4 Hz, 2H), 3.79 (s, 9H), 3.54 (s, 6H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 160.3, 158.7, 158.3, 139.7, 130.4, 129.1, 127.6, 112.7, 103.8, 100.2, 80.6, 55.9, 55.5, 55.4. MALDI-TOF *m/z* 409.490 [M]⁺.

2,2',4,4'-Tetramethoxy-3''-nitrotrityl Alcohol (**13d**)

This compound is prepared following the procedure described above for compound **9** except that 3-nitro methyl benzoate **12d** (1.70 g, 9.38 mmol) is used. The yield of **13d** (3.86 g) is 95%. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 8.00–7.98 (m, 2H), 7.55–7.42 (m, 2H), 7.08 (d, *J* = 8.5 Hz, 2H), 6.54 (d, *J* = 1.9 Hz, 2H), 6.50 (dd, *J* = 8.5, 1.9 Hz, 2H), 5.59 (s, 1H), 3.75 (s, 6H), 3.39 (s, 6H); ¹³C NMR (DMSO-*d*₆, 75.5 MHz) δ 160.2, 157.6, 150.3, 146.7, 134.2, 129.1, 127.8, 125.0, 122.0, 120.7, 104.4, 99.4, 78.1, 55.1. MALDI-TOF *m/z* 424.481 [M]⁺, 448.565 [M + Na]⁺, 464.538 [M + K]⁺.

2,2',4,4'-Tetramethoxy-4''-nitrotrityl Alcohol (**13e**)

This compound is prepared following the procedure described above for compound **9** except that 4-nitro methyl benzoate **12e** (1.70 g, 9.38 mmol) is used. The yield of **13e** (3.51 g) is

91%. ^1H NMR (DMSO- d_6 , 300 MHz) δ 8.04 (d, J = 8.9 Hz, 2H), 7.35 (d, J = 8.9 Hz, 2H), 7.06 (d, J = 8.4 Hz, 2H), 6.54 (d, J = 2.4 Hz, 2H), 6.49 (dd, J = 8.4, 2.4 Hz, 2H), 5.53 (s, 1H), 3.75 (s, 6H), 3.40 (s, 6H); ^{13}C NMR (DMSO- d_6 , 75.5 MHz) δ 160.2, 157.6, 155.9, 145.4, 129.0, 128.6, 124.9, 121.7, 104.4, 99.4, 78.2, 55.1. MALDI-TOF m/z 424.701 [M] $^+$, 408.624 [M - OH] $^+$.

9-(4-Bromophenyl)-6-hydroxy-3-fluorone (14a)

This compound was prepared following the procedure above for compound **11** except that compound **13a** (0.400 g, 0.871 mmol) is used. The yield of **14a** (223 mg) is 70%. ^1H NMR (DMSO- d_6 , 250 MHz) δ 7.88 (d, J = 7.5 Hz, 2H), 7.47 (d, J = 7.5 Hz, 2H), 7.29 (d, J = 9.4 Hz, 2H), 6.90 (d, J = 2.1 Hz, 2H), 6.88 (dd J = 9.4, 2.1 Hz, 2H); ^{13}C NMR (DMSO- d_6 , 62.9 MHz) δ 172.1, 158.0, 132.5, 131.9, 131.7, 124.2, 120.9, 115.6, 102.8. MALDI-TOF m/z 367.228 [M + H] $^+$.

9-(4-Biphenyl)-6-hydroxy-3-fluorone (14b)

This compound is prepared following the procedure above for compound **11** except that compound **22** (0.500 g, 1.09 mmol) is used. The yield of **14b** (330 mg) is 87%. ^1H NMR (DMSO- d_6 , 250 MHz) δ 7.91 (d, J = 7.8 Hz, 2H), 7.79 (d, J = 7.9 Hz, 2H), 7.55–7.49 (m, 5H), 7.09 (d, J = 9.0 Hz, 2H), 6.59 (dd, J = 9.0, 2.2 Hz, 2H), 6.54 (d, J = 2.2 Hz, 2H); ^{13}C NMR (DMSO- d_6 , 62.9 MHz) δ 157.2, 150.4, 141.9, 140.1, 132.6, 131.4, 131.0, 130.0, 128.9, 127.8, 127.7, 122.6, 115.0, 104.3. MALDI-TOF m/z 365.312 [M + H] $^+$, 387.322 [M + Na] $^+$.

9-(4-Hydroxyphenyl)-6-hydroxy-3-fluorone (14c)

This compound is prepared following the procedure above for compound **11** except that compound **13c** (1.26 g, 3.07 mmol) is used. The yield of **14c** (900 mg) is 96%. ^1H NMR (CDCl $_3$, 250 MHz) δ 10.07 (s, 1H), 7.25 (d, J = 8.2 Hz, 2H), 7.14 (d, J = 9.2 Hz, 2H), 6.98 (d, J = 8.2 Hz, 2H), 6.59 (d, J = 9.4 Hz, 2H), 6.52 (s, 2H); ^{13}C NMR (CDCl $_3$, 62.9 MHz) δ 158.6, 156.4, 150.4, 131.1, 130.8, 122.6, 115.5, 114.6, 105.5, 103.3. MALDI-TOF m/z 305.127 [M + H] $^+$, 327.100 [M + Na] $^+$, 341.221 [M + K] $^+$.

9-(3-Nitro-phenyl)-6-hydroxy-3-fluorone (14d)

This compound is prepared following the procedure above for compound **11** except that compound **14d** (0.480 g, 1.13 mmol) is used. The yield of **14d** (274 mg) is 73%. ^1H NMR (DMSO- d_6 , 300 MHz) δ 8.42 (d, J = 7.0 Hz, 1H), 8.26 (s, 1H), 7.89 (m, 2H), 6.96 (d, J = 9.1 Hz, 2H), 6.55 (dd, J = 9.1, 2.9 Hz, 2H), 6.53 (d, J = 2.9 Hz, 2H); ^{13}C NMR (DMSO- d_6 , 62.9 MHz) δ 156.3, 138.0, 147.0, 136.0, 134.3, 130.5, 130.3, 124.3, 114.3, 103.5. MALDI-TOF m/z 334.186 [M + H] $^+$, 356.167 [M + Na] $^+$.

9-(4-Nitrophenyl)-6-hydroxy-3-fluorone (14e)

This compound was prepared following the procedure above for compound **11** except that compound **15** (30 mg) is used. The eluent for flash chromatography is EtOAc–MeOH 8:2. The yield of **14e** (21 mg) is 76%. ^1H NMR (DMSO- d_6 , 250 MHz) δ 8.44 (d, J = 8.6 Hz, 2H), 7.46 (d, J = 8.6 Hz, 2H), 6.90 (dd, J = 8.9, 1.1 Hz, 2H), 6.54 (d, J = 8.9 Hz, 2H), 6.52 (d, J = 1.1 Hz, 2H); ^{13}C NMR (DMSO- d_6 , 62.9 MHz) δ 174.8, 156.3, 148.0, 147.3, 139.7, 131.0, 130.1, 123.4, 122.1, 113.5, 103.6. MALDI-TOF m/z 334.273 [M + H] $^+$, 356.301 [M + Na] $^+$.

9-(4-Nitrophenyl)-6-methoxy-3-fluorone (15)

This compound is prepared following the procedure above for compound **11** except that compound **13e** (0.500 g, 1.18 mmol) is used. The yield of **15** (346 mg) is 85%. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 8.46(d, *J* = 8.6 Hz, 2H), 7.25 (d, *J* = 2.3 Hz, 1H), 7.07 (d, *J* = 9.4 Hz, 1H), 6.98(d, *J* = 9.8 Hz, 1H), 6.93 (dd, *J* = 9.4, 2.3 Hz, 1H), 6.44 (dd, *J* = 9.8, 1.8 Hz, 1H), 6.26 (d, *J* = 1.8 Hz, 1H); ¹³C NMR (DMSO-*d*₆, 75.5 MHz) δ 183.8, 164.3, 158.2, 154.0, 148.2, 139.3, 131.1, 130.5, 129.8, 129.5, 127.1, 124.0, 123.2, 117.2, 113.68, 113.4, 105.0, 100.8, 56.4. MALDI-TOF *m/z* 348.419 [M + H]⁺.

Bis-(2,4-Dimethoxyphenyl)-4-(2, 4-dimethoxyphenyl)-phenyl Methane (19)

This compound is prepared following the procedure described above for compound **9** except that **18** (1.64 g, 6.26 mmol) is used. The yield of **19** (1.00 g) is 32%. ¹H NMR (CDCl₃, 300 MHz) δ 7.37 (d, *J* = 8.1 Hz, 2H), 7.24 (d, *J* = 4.1 Hz, 1H), 7.05 (d, *J* = 8.1 Hz, 2H), 6.77 (d, *J* = 8.3 Hz, 2H), 6.55 (d, *J* = 4.1 Hz, 1H), 6.53 (s, 1H), 6.46 (d, *J* = 2.3 Hz, 2H), 6.38 (dd, *J* = 8.3, 2.3 Hz, 2H), 6.05 (s, 1H), 3.83 (s, 3H), 3.78 (s, 9H), 3.71 (s, 6H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 160.2, 159.3, 158.3, 157.6, 142.7, 135.6, 131.4, 130.7, 129.0, 129.0, 127.8, 104.7, 103.7, 99.1, 98.9, 55.9, 55.7, 55.6, 55.4, 41.8. MALDI-TOF *m/z*, 500.935 [M]⁺, 523.990 [M + Na]⁺.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

We thank the National Institutes of Health for funding this research via Grant R01 EB002044. The purchase of the Nonius Kappa CCD diffractometer was made possible by Grant No. LEQSF-(1999–2000)-ESH-TR-13, administered by the Louisiana Board of Regents.


References

- (1). (a) Hirano T, Kikichi K, Urano Y, Higuchi T, Nagono T. *Angew. Chem., Int. Ed.* 2000; 39:1052. (b) Liu S, Xie Y, Yong G, Dai Y. *J. Agric. Food. Chem.* 2000; 48:5860. [PubMed: 11141259] (c) Li Z, Tang J, Pan J. *Analyst.* 2001; 126:1154. [PubMed: 11478652] (d) Fujita Y, Mori I, Yamaguchi T, Hoshino M, Shigemura Y, Yasuyuki, Shimano M. *Anal. Sci.* 2001; 17:853. [PubMed: 11708118] (e) Gee KR, Zhou Z-L, Qian W-J, Kennedy R. *J. Am. Chem. Soc.* 2002; 124:776. [PubMed: 11817952] (f) Li Z, You F, Liu Z, Jian T. *Talanta.* 2004; 63:647. [PubMed: 18969482]
- (2). Turchini J. *Acta Histochem.* 1957; 4:15. [PubMed: 13457799]
- (3). U.S. Patent Appl. Publ. CODEN: USXXCO US 2004038306 A1 20040226 2004. p. 83
- (4). Stephen AG, Worthy KM, Towler E, Mikovits JA, Sei S, Roberts P, Yang Q, Akee RK, Klausmeyer P, McCloud TG, Henderson L, Rein A, Covell DG, Currens M, Shoemaker RH, Fisher RJ. *Biochem. Biophys. Res. Commun.* 2002; 296:1228. [PubMed: 12207905]
- (5). Nagano T, Urano Y. *Jpn. Kokai Tokyo Koho CODEN: JKXXAF JP 2000321262 A2 20001124.* 2000:14.
- (6). Blattner JR, He L, Lemasters JJ. *Anal. Biochem.* 2001; 295:220. [PubMed: 11488625]
- (7). Mizutani MY, Itai A. *J. Med. Chem.* 2004; 47:4818. [PubMed: 15369385]
- (8). Tolman RL, Gamsey S, Mehta S, Pongracz K. *Pct Int. Appl. CODEN: PIXXD2 WO 2002076397 A2 20021003.* 2002:41.
- (9). Yang Y, Lewis PT, Escobedo JO, St. Luce NN, Trealeaven WD, Cooks RL, Strongin RM. *Collect. Czech. Chem. Commun.* 2004; 69:1282.
- (10). Wang W, Escobedo JO, Lawrence CM, Strongin RM. *J. Am. Chem. Soc.* 2004; 126:3400. [PubMed: 15025448]


- (11). Mohlau R, Koch P. Ber. 1894; 27:2887.
- (12). (a) Swain CJ, Hedberg K. J. Am. Chem. Soc. 1950; 72:3373.(b) Tilak BD. Ind. Chim. Belge. 1967; 32:50.(c) He M, Johnson RJ, Escobedo JO, Beck PA, Kim KK, St. Luce NN, Davis CJ, Lewis PT, Fronczek FR, Melancon BJ, Mrse AA, Treleaven WD, Strongin RM. J. Am. Chem. Soc. 2002; 124:5000. [PubMed: 11982364] (d) Jiao G-S, Castro JC, Thoresen LH, Burgess K. Org. Lett. 2003; 5:3675. [PubMed: 14507202]
- (13). Shi J, Zhang X, Neckers DC. J. Org. Chem. 1992; 57:4418.
- (14). Chang CJ, Jaworski J, Nolan EM, Sheng M, Lippard SJ. Proc. Natl. Acad. Sci. U.S.A. 2004; 101:1129. [PubMed: 14734801]
- (15). Dinger MB, Scott MJ. J. Chem. Soc., Perkin Trans. 1. 2000; 11:1741.
- (16). Jpn. Kokai Tokkyo Koho CODEN: JKXXAF JP 07138200 A2 19950530 Heisei. 1995. p. 5
- (17). Heaney H, Hollinshead JH, Ley SV. Tetrahedron. 1995; 51:7741.
- (18). (a) Weinelt F, Schneider H-J. J. Org. Chem. 1991; 56:5527.(b) Fronczek FR, St. Luce NN, Strongin RM. Acta Crystallogr. 2001; C57:1423.
- (19). Detailed synthetic route is described in the Experimental Section.
- (20). Selected references:Brown HC. Organic Synthesis via Boranes. :1975.Wiley-InterscienceNew York. Negishi E. Wilkinson G, Stone FGA, Abel EW. Comprehensive Organometallic Chemistry. 1982; Vol. 5:255.Pergamon PressOxford. Pelter A, Smith K, Brown HS. Borane Reagents. 1988Academic PressLondon
- (21). Matteson DS, Mah RWH. J. Am. Chem. Soc. 1963; 85:2599.
- (22). (a) Kitatani K, Hiyama T, Nozaki H. Bull. Chem. Soc. Jpn. 1977; 50(b) Negishi E, Akiyoshi K. J. Am. Chem. Soc. 1988; 110:646.(c) Negishi E, Akiyoshi K, O'Connor B, Takagi K, Wu G. J. Am. Chem. Soc. 1989; 111:3089.(d) Kocienski P, Wadman S, Cooper K. J. Am. Chem. Soc. 1989; 111:2363.(e) Miller JA. J. Org. Chem. 1989; 54:998.(f) Knochel P, Jeong N, Rozema MJ, Yeh MCP. J. Am. Chem. Soc. 1989; 111:6474.(g) Stocks M, Kocienski P, Donald KK. Tetrahedron Lett. 1990; 31:1637.(h) AchyuthaRao S, Rosema MJ, Knochel P. J. Org. Chem. 1993; 58:2694.(i) Kakiya H, Inoue R, Shinokubo H, Oshima K. Tetrahedron Lett. 1997; 38:3275.(j) Harada T, Kaneko T, Fujiwara T, Oku A. J. Org. Chem. 1997; 62:8966.(k) Fillion E, Carson RJ, Trepanier VE, Goll JM, Remorova AA. J. Am. Chem. Soc. 2004; 126:15354. [PubMed: 15563142]
- (23). (a) Harada T, Hara D, Hattori K, Oku A. Tetrahedron Lett. 1988; 29:3821.(b) Harada T, Katsuhira T, Osada A, Iwazaki K, Maejima R, Oku A. J. Am. Chem. Soc. 1996; 118:11377.
- (24). (a) Harada T, Kaneko T, Fujiwara T, Oku A. Tetrahedron. 1998; 54:9317.(b) Harada T, Chiba M, Oku A. J. Org. Chem. 1999; 64:8210. [PubMed: 11674738]
- (25). Yonezawa N, Hino T, Matsuda K, Matsuki T, Narushima O, Kobayashi M, Ikeda T. J. Org. Chem. 2000; 65:941. [PubMed: 10814036]
- (26). Ramart-Lucas P. Bull. Soc. Chim. 1945; 12:477.




Scheme 1.




Scheme 2.




Scheme 3.




Scheme 4.




Scheme 5.



Scheme 6.



Scheme 7.



Scheme 8.
22j




Figure 1.
Structures of some common xanthene dyes.





Figure 2.
X-ray structure for **19**.

TABLE 1



entry	substrate	R ₁	R ₂	carbinol	carbinol yield (%)	fluorone	fluorone yield (%)
1	12a	Br	H	13a	92	14a	70
2	12b	Ph	H	13b	99	14b	87
3	12c	OMe	H	13c	83	14c	96
4	12d	H	NO ₂	13d	91	14d	73
5	12e	NO ₂	H	13e	95		