

A convergence result for the Emery topology and a variant of the proof of the Fundamental Theorem of Asset Pricing

Christa Cuchiero
(based on joint work with Josef Teichmann)

Technical University Vienna

Workshop on Mathematical Finance:
Arbitrage and Portfolio Optimization,
May 16, 2014

Problem formulation and motivation

- The question we deal with is to find conditions under which a sequence of semimartingales, which converges pathwise uniformly in probability (“up-convergence”), converges in the Emery topology.

Problem formulation and motivation

- The question we deal with is to find conditions under which a sequence of semimartingales, which converges pathwise uniformly in probability (“up-convergence”), converges in the Emery topology.
- The motivation for this question stems from the proof of the Fundamental Theorem of Asset Pricing (FTAP) in continuous time where one crucial part consists in proving exactly this statement under (NFLVR) and a certain maximality assumption.

Problem formulation and motivation

- The question we deal with is to find conditions under which a sequence of semimartingales, which converges pathwise uniformly in probability (“up-convergence”), converges in the Emery topology.
- The motivation for this question stems from the proof of the Fundamental Theorem of Asset Pricing (FTAP) in continuous time where one crucial part consists in proving exactly this statement under (NFLVR) and a certain maximality assumption.
- Is it possible to transform this to a criterion/theorem interesting in its own right?

Goal of this talk

- Present a general principle for sequences of semimartingales, the so-called **P-UT property**, which allows together with a **maximality assumption** to conclude the desired convergence in the Emery topology for up-convergent sequences of semimartingales.

Goal of this talk

- Present a general principle for sequences of semimartingales, the so-called **P-UT property**, which allows together with a **maximality assumption** to conclude the desired convergence in the Emery topology for up-convergent sequences of semimartingales.
- Application to the setting of FTAP, which allows to obtain a variant of one part of the proof by showing that the **P-UT property** is implied by (NFLVR).

Setting and notation

- We consider a finite time horizon $T = 1$ and a fixed filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0,1]}, P)$. For a process Z we write $|Z|_1^* = \sup_{t \in [0,1]} |Z_t|$.

Setting and notation

- We consider a finite time horizon $T = 1$ and a fixed filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0,1]}, P)$. For a process Z we write $|Z|_1^* = \sup_{t \in [0,1]} |Z_t|$.
- The set of semimartingales starting at 0 on the time interval $[0, 1]$ is denoted by \mathbb{S} .

Setting and notation

- We consider a finite time horizon $T = 1$ and a fixed filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0,1]}, P)$. For a process Z we write $|Z|_1^* = \sup_{t \in [0,1]} |Z_t|$.
- The set of semimartingales starting at 0 on the time interval $[0, 1]$ is denoted by \mathbb{S} .
- We equip \mathbb{S} with the Emery metric

$$\sup_{H \in b\mathcal{E}, \|H\|_\infty \leq 1} E[|(H \bullet (X - Y))|_1^* \wedge 1] = d_E(X, Y),$$

making it a complete metric space. Here, $b\mathcal{E}$ denotes the set of bounded simple predictable strategies.

Setting and notation

- We consider a finite time horizon $T = 1$ and a fixed filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0,1]}, P)$. For a process Z we write $|Z|_1^* = \sup_{t \in [0,1]} |Z_t|$.
- The set of semimartingales starting at 0 on the time interval $[0, 1]$ is denoted by \mathbb{S} .
- We equip \mathbb{S} with the Emery metric

$$\sup_{H \in b\mathcal{E}, \|H\|_\infty \leq 1} E[|(H \bullet (X - Y))|_1^* \wedge 1] = d_E(X, Y),$$

making it a complete metric space. Here, $b\mathcal{E}$ denotes the set of bounded simple predictable strategies.

- Pathwise uniform convergence in probability is metrized by

$$E[|X - Y|_1^* \wedge 1] = d(X, Y),$$

which makes the set of càdlàg processes a complete metric space.

Definition of the (P-UT) property

Let us introduce the (P-UT) property (cf. Jacod and Shiryaev (2003)), originally introduced by Stricker (1985) under Condition (*) and Jakubowski, Mémin and Pagès (1989) under the name (UT) (for “uniforme tension”).

Definition of the (P-UT) property

Let us introduce the (P-UT) property (cf. Jacod and Shiryaev (2003)), originally introduced by Stricker (1985) under Condition (*) and Jakubowski, Mémin and Pagès (1989) under the name (UT) (for “uniforme tension”).

Definition

We say that a sequence $(X^n)_{n \geq 0}$ of semimartingales satisfies the (P-UT) property (predictably uniformly tight) if for every $t > 0$ the family of random variables $\{(H \bullet X^n)_t : H \in b\mathcal{E}, \|H\|_\infty \leq 1, n \in \mathbb{N}\}$ is bounded in L^0 , that is,

$$\lim_{c \uparrow \infty} \sup_{H, n \in \mathbb{N}} P[|(H \bullet X^n)|_t > c] = 0.$$

Definition of the (P-UT) property

Let us introduce the (P-UT) property (cf. Jacod and Shiryaev (2003)), originally introduced by Stricker (1985) under Condition (*) and Jakubowski, Mémin and Pagès (1989) under the name (UT) (for “uniforme tension”).

Definition

We say that a sequence $(X^n)_{n \geq 0}$ of semimartingales satisfies the (P-UT) property (predictably uniformly tight) if for every $t > 0$ the family of random variables $\{(H \bullet X^n)_t : H \in b\mathcal{E}, \|H\|_\infty \leq 1, n \in \mathbb{N}\}$ is bounded in L^0 , that is,

$$\lim_{c \uparrow \infty} \sup_{H, n \in \mathbb{N}} P[|(H \bullet X^n)|_t > c] = 0.$$

- The (P-UT) property can be seen as boundedness in the Emery topology.
- Interpretation from mathematical finance: No unbounded profit or loss with simple, predictable, bounded by 1, long or short positions

Convergence of stochastic integrals under the (P-UT) property

“...the following theorem is in fact the very reason for introducing the (P-UT) property” (Jacod, Shirayev (2003))

Theorem (Jakubowski, Mémin and Pagès (1989))

Let $(H^n)_{n \geq 0}$ be a sequence of adapted càdlàg processes and $(X^n)_{n \geq 0}$ a sequence of semimartingales satisfying the (P-UT) property. If (H^n, X^n) converges pathwise uniformly in probability to (H, X) , then $(H_-^n \bullet X^n)$ converges to $(H_- \bullet X)$ pathwise uniformly in probability as well.

Convergence of stochastic integrals and the quadratic variation under the (P-UT) property

Corollary (Jakubowski, Mémin and Pagès (1989))

Let $(X^n)_{n \geq 0}$ be a sequence of semimartingales satisfying the (P-UT) property. If (X^n) converges pathwise uniformly in probability to X , then $[X^n, X^n]$ converges to $[X, X]$ pathwise uniformly in probability as well.

Sketch of the proof

This follows from the above theorem by Itô's integration by parts formula.

Emery convergence for the local martingale and the big jump part under (P-UT) and up-convergence

For a sequence of semimartingales (X^n) with $X_0^n = 0$ and some $C > 0$ we consider the following decomposition

$$X^n = B^{n,C} + M^{n,C} + \check{X}^{n,C},$$

where $\check{X}_t^{n,C} = \sum_{s \leq t} \Delta X_s^n \mathbf{1}_{\{|\Delta X_s| > C\}}$, $B^{n,C}$ is the predictable, finite variation part and $M^{n,C}$ the local martingale part of the canonical decomposition of the special semimartingale $X^n - \check{X}^{n,C}$.

Emery convergence for the local martingale and the big jump part under (P-UT) and up-convergence

For a sequence of semimartingales (X^n) with $X_0^n = 0$ and some $C > 0$ we consider the following decomposition

$$X^n = B^{n,C} + M^{n,C} + \check{X}^{n,C},$$

where $\check{X}_t^{n,C} = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s^n| > C\}}$, $B^{n,C}$ is the predictable, finite variation part and $M^{n,C}$ the local martingale part of the canonical decomposition of the special semimartingale $X^n - \check{X}^{n,C}$.

Theorem (Mémin and Slominski (1991))

Let (X^n) be a sequence of semimartingales with $X_0^n = 0$, which converges pathwise uniformly in probability to X and satisfies the (P-UT) property. Then there exists some $C > 0$ such that $M^{n,C} \rightarrow M^C$ and $\check{X}^{n,C} \rightarrow \check{X}^C$ in the Emery topology and $B^{n,C} \rightarrow B^C$ pathwise uniformly in probability.

Remarks on the proof

Preliminaries on Emery convergence

- A sequence of adapted finite variation processes (X^n) converges in the Emery topology to 0 if $\lim_{n \rightarrow \infty} TV(X^n)_1 \rightarrow 0$ in probability.
- A sequence of local martingale (X^n) with $|\Delta X^n|_1^* \leq C$ (uniformly in n) converges in the Emery topology to 0 if $\lim_{n \rightarrow \infty} [X^n, X^n]_1 \rightarrow 0$ in probability.

Remarks on the proof

Preliminaries on Emery convergence

- A sequence of adapted finite variation processes (X^n) converges in the Emery topology to 0 if $\lim_{n \rightarrow \infty} TV(X^n)_1 \rightarrow 0$ in probability.
- A sequence of local martingale (X^n) with $|\Delta X^n|_1^* \leq C$ (uniformly in n) converges in the Emery topology to 0 if $\lim_{n \rightarrow \infty} [X^n, X^n]_1 \rightarrow 0$ in probability.

Sketch of the proof (of the Mémin and Slominski Theorem)

The process $\check{X}^{n,C} - \check{X}^C$ converges in variation and thus in the Emery topology to 0. Moreover, (P-UT) and up convergence imply

$$[(X^n - \check{X}^{n,C}) - (X - \check{X}^C), (X^n - \check{X}^{n,C}) - (X - \check{X}^C)]_1 \rightarrow 0$$

in probability. From this $[M^{C,n} - M^C, M^{C,n} - M^C]_1 \rightarrow 0$ in probability and thus $M^{n,C} - M^C \rightarrow 0$ in the Emery topology can be deduced. Pathwise uniform convergence in probability of $B^{n,C} \rightarrow B^C$ is obtained by difference.

How to deal with the finite variation part (without big jumps)

- For a sequence of semimartingales (X^n) , consider for every $r \in \mathbb{N}$ the following sets

$$\mathcal{Y}^r = \{H \bullet X^n + G \bullet X^m \mid n, m \geq r, H, G \text{ bounded predictable} \\ \|H\|_\infty \leq 1, \|G\|_\infty \leq 1, HG = 0\},$$

and denote by $\mathcal{Z}^r = \text{conv}(\mathcal{Y}^r)$ its convex hull. Moreover, we denote $\mathcal{V}_1^r = \{Z_1 \mid Z \in \mathcal{Z}^r\}$ and by $\widehat{\mathcal{V}}_1^r$ its closure in L^0 . Define $\widehat{\mathcal{V}}_1 = \bigcap_{r \geq 0} \widehat{\mathcal{V}}_1^r$.

How to deal with the finite variation part (without big jumps)

- For a sequence of semimartingales (X^n) , consider for every $r \in \mathbb{N}$ the following sets

$$\mathcal{Y}^r = \{H \bullet X^n + G \bullet X^m \mid n, m \geq r, H, G \text{ bounded predictable} \\ \|H\|_\infty \leq 1, \|G\|_\infty \leq 1, HG = 0\},$$

and denote by $\mathcal{Z}^r = \text{conv}(\mathcal{Y}^r)$ its convex hull. Moreover, we denote $\mathcal{V}_1^r = \{Z_1 \mid Z \in \mathcal{Z}^r\}$ and by $\widehat{\mathcal{V}}_1^r$ its closure in L^0 . Define $\widehat{\mathcal{V}}_1 = \bigcap_{r \geq 0} \widehat{\mathcal{V}}_1^r$.

Lemma

Let (X^n) be a sequence of semimartingales satisfying the (P-UT) property. Then the set $\widehat{\mathcal{V}}_1$ is bounded in L^0 .

How to deal with the finite variation part (without big jumps)

- For a sequence of semimartingales (X^n) , consider for every $r \in \mathbb{N}$ the following sets

$$\mathcal{Y}^r = \{H \bullet X^n + G \bullet X^m \mid n, m \geq r, H, G \text{ bounded predictable} \\ \|H\|_\infty \leq 1, \|G\|_\infty \leq 1, HG = 0\},$$

and denote by $\mathcal{Z}^r = \text{conv}(\mathcal{Y}^r)$ its convex hull. Moreover, we denote $\mathcal{V}_1^r = \{Z_1 \mid Z \in \mathcal{Z}^r\}$ and by $\widehat{\mathcal{V}}_1^r$ its closure in L^0 . Define $\widehat{\mathcal{V}}_1 = \bigcap_{r \geq 0} \widehat{\mathcal{V}}_1^r$.

Lemma

Let (X^n) be a sequence of semimartingales satisfying the (P-UT) property. Then the set $\widehat{\mathcal{V}}_1$ is bounded in L^0 .

- From this it follows that $\widehat{\mathcal{V}}_1$ contains a maximal element.

A convergence result in the Emery topology

Theorem

Let (X^n) be a sequence of semimartingales satisfying the (P-UT) property. Take a sequence $(Z^n) \in \mathcal{Z}^1$ such that (Z_1^n) converges a.s. to a maximal element Z_1 in $\widehat{\mathcal{V}}_1$. Then (Z^n) converges in the Emery topology.

A convergence result in the Emery topology

Theorem

Let (X^n) be a sequence of semimartingales satisfying the (P-UT) property. Take a sequence $(Z^n) \in \mathcal{Z}^1$ such that (Z_1^n) converges a.s. to a maximal element Z_1 in $\widehat{\mathcal{V}}_1$. Then (Z^n) converges in the Emery topology.

Proof.

- Due to the P-UT property of (X^n) , \mathcal{Z}^1 is bounded in L^0 and the sequence (Z^n) satisfies the (P-UT) property.
- Moreover, as in the proof of FTAP by Delbaen and Schachermayer the maximality of the end value allows to conclude up-convergence.
- An application of Mémin and Slominski's theorem yields Emery convergence of the local martingale and the big jump part of Z^n in the Emery topology.
- Using this together with the maximality of Z_1 , an adaptation of the last part in the proof of FTAP yields convergence of the finite variation part in the Emery topology.

Towards the setting of FTAP

- The conclusion of the above result holds for sequences of portfolio wealth processes bounded from below by -1 under the (NUPBR) (No unbounded profit with bounded risk) condition, which means L^0 -boundedness of portfolio end values.

Towards the setting of FTAP

- The conclusion of the above result holds for sequences of portfolio wealth processes bounded from below by -1 under the (NUPBR) (No unbounded profit with bounded risk) condition, which means L^0 -boundedness of portfolio end values.
- The crucial issue consists in proving that (NUPBR) implies (P-UT).

The Fundamental Theorem of Asset pricing

- FTAP lays the foundations of mathematical finance and can therefore be considered as the main result in mathematical finance.

The Fundamental Theorem of Asset pricing

- FTAP lays the foundations of mathematical finance and can therefore be considered as the main result in mathematical finance.
- In continuous time it states the equivalence of an “absence of arbitrage” property (NFLVR) with the existence of an equivalent σ -martingale measure.

The Fundamental Theorem of Asset pricing

- FTAP lays the foundations of mathematical finance and can therefore be considered as the main result in mathematical finance.
- In continuous time it states the equivalence of an “absence of arbitrage” property (NFLVR) with the existence of an equivalent σ -martingale measure.
- The first complete proof has been presented by F. Delbaen and W. Schachermayer (1994, 1998).

The Fundamental Theorem of Asset pricing

- FTAP lays the foundations of mathematical finance and can therefore be considered as the main result in mathematical finance.
- In continuous time it states the equivalence of an “absence of arbitrage” property (NFLVR) with the existence of an equivalent σ -martingale measure.
- The first complete proof has been presented by F. Delbaen and W. Schachermayer (1994, 1998).
- Their beautiful and impressive proof builds on deep insights and is in some parts quite tricky. No essential simplification has been obtained since then.

The Fundamental Theorem of Asset pricing

- FTAP lays the foundations of mathematical finance and can therefore be considered as the main result in mathematical finance.
- In continuous time it states the equivalence of an “absence of arbitrage” property (NFLVR) with the existence of an equivalent σ -martingale measure.
- The first complete proof has been presented by F. Delbaen and W. Schachermayer (1994, 1998).
- Their beautiful and impressive proof builds on deep insights and is in some parts quite tricky. No essential simplification has been obtained since then.
- It was however realized soon that the presented proof can be transformed to a more abstract setting of portfolio wealth processes (Y. Kabanov (1996)).

Definition of portfolio wealth processes

Definition

- We consider a convex set $\mathcal{X}_1 \subset \mathbb{S}$ of semimartingales
 - ▶ starting at 0,
 - ▶ bounded from below by -1 ,
 - ▶ being closed in the Emery topology, and
 - ▶ satisfying the following concatenation condition: for all $X, Y \in \mathcal{X}_1$ and all bounded predictable strategies $H, G \geq 0$, with $HG = 0$ and $Z = (H \bullet X) + (G \bullet Y) \geq -1$, it holds that $Z \in \mathcal{X}_1$.

Definition of portfolio wealth processes

Definition

- We consider a convex set $\mathcal{X}_1 \subset \mathbb{S}$ of semimartingales
 - ▶ starting at 0,
 - ▶ bounded from below by -1 ,
 - ▶ being closed in the Emery topology, and
 - ▶ satisfying the following concatenation condition: for all $X, Y \in \mathcal{X}_1$ and all bounded predictable strategies $H, G \geq 0$, with $HG = 0$ and $Z = (H \bullet X) + (G \bullet Y) \geq -1$, it holds that $Z \in \mathcal{X}_1$.
- We denote by \mathcal{X} the set $\mathcal{X} = \cup_{\lambda > 0} \lambda \mathcal{X}_1$ and call its elements admissible portfolio wealth processes.

Definition of portfolio wealth processes

Definition

- We consider a convex set $\mathcal{X}_1 \subset \mathbb{S}$ of semimartingales
 - ▶ starting at 0,
 - ▶ bounded from below by -1 ,
 - ▶ being closed in the Emery topology, and
 - ▶ satisfying the following concatenation condition: for all $X, Y \in \mathcal{X}_1$ and all bounded predictable strategies $H, G \geq 0$, with $HG = 0$ and $Z = (H \bullet X) + (G \bullet Y) \geq -1$, it holds that $Z \in \mathcal{X}_1$.
- We denote by \mathcal{X} the set $\mathcal{X} = \cup_{\lambda > 0} \lambda \mathcal{X}_1$ and call its elements admissible portfolio wealth processes.
- We denote by K_0 , respectively K_0^1 the evaluations of elements of \mathcal{X} , respectively \mathcal{X}_1 , at final time $T = 1$.

Remarks

- The setting of portfolio wealth processes is an **abstract formulation of the classical setting of Delbaen and Schachermayer**, where trading (in an admissible way) with respect to a d -dimensional semimartingale S is considered.
- In particular, the above requirements (admissibility, concatenation, closedness in the Emery topology) are those which are satisfied in the case of $\mathcal{X}_1 = \{H \bullet S \mid H \text{ 1-admissible}\}$. Closedness in the Emery topology of this set was proved by Mémin (1980).

Notions of No Arbitrage

(NA) The set \mathcal{X} is said to satisfy **No Arbitrage** if $K_0 \cap L_{\geq 0}^0 = \{0\}$ which can be shown to be equivalent to $C \cap L_{\geq 0}^\infty = \{0\}$, with $C = (K_0 - L_{\geq 0}^0) \cap L^\infty$.

(NFLVR) The set \mathcal{X} is said to satisfy **No free lunch with vanishing risk** if

$$\bar{C} \cap L_{\geq 0}^\infty = \{0\},$$

where \bar{C} denotes the norm closure in L^∞ .

(NFL) The set \mathcal{X} is said to satisfy **No free lunch** if

$$\bar{C}^* \cap L_{\geq 0}^\infty = \{0\},$$

where \bar{C}^* denotes the weak- $*$ -closure in L^∞ .

Notions of No Arbitrage

(NUPBR) The set \mathcal{X}_1 is said to satisfy No unbounded profit with bounded risk if K_0^1 is bounded in L^0 .

Notions of No Arbitrage

(NUPBR) The set \mathcal{X}_1 is said to satisfy **No unbounded profit with bounded risk** if K_0^1 is bounded in L^0 .

- (NFLVR) \Leftrightarrow (NA) + (NUPBR) (D & S, Proposition 9.3.1, Corollary 9.3.9)

Notions of No Arbitrage

(NUPBR) The set \mathcal{X}_1 is said to satisfy **No unbounded profit with bounded risk** if K_0^1 is bounded in L^0 .

- (NFLVR) \Leftrightarrow (NA) + (NUPBR) (D & S, Proposition 9.3.1, Corollary 9.3.9)
- Both (NFLVR) and (NUPBR) are **economically convincing minimal requirement for models**, but only (NFL) allows to conclude relatively directly the existence of an equivalent separating measure, defined via

Notions of No Arbitrage

(NUPBR) The set \mathcal{X}_1 is said to satisfy **No unbounded profit with bounded risk** if K_0^1 is bounded in L^0 .

- (NFLVR) \Leftrightarrow (NA) + (NUPBR) (D & S, Proposition 9.3.1, Corollary 9.3.9)
- Both (NFLVR) and (NUPBR) are **economically convincing minimal requirement for models**, but only (NFL) allows to conclude relatively directly the existence of an equivalent separating measure, defined via

Definition

The set \mathcal{X} satisfies the (ESM) (**equivalent separating measure**) property if there exists an equivalent measure $Q \sim P$ such that $\mathbb{E}_Q[X_1] \leq 0$ for all $X \in \mathcal{X}$.

(NFLVR) \Leftrightarrow (ESM)

- It is a consequence of **Hahn-Banach's Theorem (the Kreps-Yan Theorem)** that (NFL) implies the existence of an equivalent measure $Q \sim P$ such that $E_Q[f] \leq 0$ for all $f \in C$ and hence for all $f \in K_0$.

$(NFLVR) \Leftrightarrow (ESM)$

- It is a consequence of **Hahn-Banach's Theorem (the Kreps-Yan Theorem)** that (NFL) implies the existence of an equivalent measure $Q \sim P$ such that $E_Q[f] \leq 0$ for all $f \in C$ and hence for all $f \in K_0$.
- We obviously have $(NFL) \Rightarrow (NFLVR) \Rightarrow (NA)$, but it is a deep insight that **under (NFLVR) it holds that $C = \overline{C}^*$** , i.e. the cone C is already weak-*closed and (NFL) holds.

(NFLVR) \Leftrightarrow (ESM)

- It is a consequence of **Hahn-Banach's Theorem (the Kreps-Yan Theorem)** that (NFL) implies the existence of an equivalent measure $Q \sim P$ such that $E_Q[f] \leq 0$ for all $f \in C$ and hence for all $f \in K_0$.
- We obviously have $(NFL) \Rightarrow (NFLVR) \Rightarrow (NA)$, but it is a deep insight that **under (NFLVR) it holds that $C = \overline{C}^*$** , i.e. the cone C is already weak- $*$ -closed and (NFL) holds.
- The goal is to show $(NFLVR) \Rightarrow C = \overline{C}^*$, which proves the Fundamental Theorem of Asset Pricing:

Theorem (Delbaen and Schachermayer (1994))

Under (NFLVR) the cone C is weak $$ -closed, hence (NFL) holds, which is equivalent to (ESM). In other words: (NFLVR) \Leftrightarrow (ESM).*

First part of the original proof by Delbaen and Schachermayer

- The convex cone $C = (K_0 - L_{\geq 0}^0) \cap L^\infty$ is closed with respect to the weak-* topology if and only if $C_0 = K_0 - L_{\geq 0}^0$ is Fatou-closed, i.e. for any sequence (f_n) in C_0 bounded from below and converging almost surely to f it holds that $f \in C_0$.

First part of the original proof by Delbaen and Schachermayer

- The convex cone $C = (K_0 - L_{\geq 0}^0) \cap L^\infty$ is closed with respect to the weak-* topology if and only if $C_0 = K_0 - L_{\geq 0}^0$ is Fatou-closed, i.e. for any sequence (f_n) in C_0 bounded from below and converging almost surely to f it holds that $f \in C_0$.
- Take now $-1 \leq f_n \in C_0$ converging almost surely to f . Then we can find $g_n \in K_0$, i.e. $g_n = Y_1^n$ with $Y^n \in \mathcal{X}$ such that $f_n \leq g_n = Y_1^n$.

First part of the original proof by Delbaen and Schachermayer

- The convex cone $C = (K_0 - L_{\geq 0}^0) \cap L^\infty$ is closed with respect to the weak-* topology if and only if $C_0 = K_0 - L_{\geq 0}^0$ is Fatou-closed, i.e. for any sequence (f_n) in C_0 bounded from below and converging almost surely to f it holds that $f \in C_0$.
- Take now $-1 \leq f_n \in C_0$ converging almost surely to f . Then we can find $g_n \in K_0$, i.e. $g_n = Y_1^n$ with $Y^n \in \mathcal{X}$ such that $f_n \leq g_n = Y_1^n$.
- By (NA) it follows that $Y^n \in \mathcal{X}_1$.

First part of the original proof by Delbaen and Schachermayer

- The convex cone $C = (K_0 - L_{\geq 0}^0) \cap L^\infty$ is closed with respect to the weak-* topology if and only if $C_0 = K_0 - L_{\geq 0}^0$ is **Fatou-closed**, i.e. for any sequence (f_n) in C_0 bounded from below and converging almost surely to f it holds that $f \in C_0$.
- Take now $-1 \leq f_n \in C_0$ converging almost surely to f . Then we can find $g_n \in K_0$, i.e. $g_n = Y_1^n$ with $Y^n \in \mathcal{X}$ such that $f_n \leq g_n = Y_1^n$.
- By **(NA)** it follows that $Y^n \in \mathcal{X}_1$.
- By **(NUPBR)** it follows that there are forward-convex combinations $\widetilde{Y}^n \in \text{conv}(Y^n, Y^{n+1}, \dots)$ such that $\widetilde{Y}_1^n \rightarrow \widetilde{h}_0 \geq f$ almost surely for some finite \widetilde{h}_0 . This implies that the set $\widehat{K}_0^1 \cap \{g \in L^0 \mid g \geq f\}$, where \widehat{K}_0^1 denotes the closure of K_0^1 in L^0 is **non-empty**. Since it is also **bounded and closed**, a maximal element h_0 exists.

First part of the original proof by Delbaen and Schachermayer

- Since $h_0 \in \widehat{K}_0^1$, we can find a sequence of semimartingales $X^n \in \mathcal{X}_1$ such that $X_1^n \rightarrow h_0$ almost surely.

First part of the original proof by Delbaen and Schachermayer

- Since $h_0 \in \widehat{K}_0^1$, we can find a sequence of semimartingales $X^n \in \mathcal{X}_1$ such that $X_1^n \rightarrow h_0$ almost surely.
- This “maximal” sequence of semimartingales $X^n \in \mathcal{X}_1$ converges pathwise uniformly in probability, i.e. $|X^n - X|_1^* \rightarrow 0$ in probability for some càdlàg process X .

First part of the original proof by Delbaen and Schachermayer

- Since $h_0 \in \widehat{K}_0^1$, we can find a sequence of semimartingales $X^n \in \mathcal{X}_1$ such that $X_1^n \rightarrow h_0$ almost surely.
- This “maximal” sequence of semimartingales $X^n \in \mathcal{X}_1$ converges pathwise uniformly in probability, i.e. $|X^n - X|_1^* \rightarrow 0$ in probability for some càdlàg process X .
- It is now the goal to show that indeed $X^n \rightarrow X$ in the Emery topology, an apparently much stronger statement. If we can show this, it follows that $h_0 = \lim X_1^n = X_1 \in K_0^1$ since $X \in \mathcal{X}_1$ by the closedness of \mathcal{X}_1 in the Emery topology. This then implies that $f \in C_0$ and we are done.

Comments on the original proof

- By performing an equivalent measure change to deal with special L^2 -semimartingales, the original proof proceeds by showing Emery convergence of the local martingale part and the finite variation part of forward convex combinations of (X^n) . This is achieved by a series of quite tricky lemmas.

Comments on the original proof

- By performing an equivalent measure change to deal with special L^2 -semimartingales, the original proof proceeds by showing Emery convergence of the local martingale part and the finite variation part of forward convex combinations of (X^n) . This is achieved by a series of quite tricky lemmas.
- Two questions come up, namely can the result be achieved without the change of measure and is it possible to do without the passage to forward convex combinations?

Comments on the original proof

- By performing an equivalent measure change to deal with special L^2 -semimartingales, the original proof proceeds by showing Emery convergence of the local martingale part and the finite variation part of forward convex combinations of (X^n) . This is achieved by a series of quite tricky lemmas.
- Two questions come up, namely can the result be achieved without the change of measure and is it possible to do without the passage to forward convex combinations?
- Is it possible to replace the series of lemmas by one theorem?

Comments on the original proof

- By performing an equivalent measure change to deal with special L^2 -semimartingales, the original proof proceeds by showing Emery convergence of the local martingale part and the finite variation part of forward convex combinations of (X^n) . This is achieved by a series of quite tricky lemmas.
- Two questions come up, namely can the result be achieved without the change of measure and is it possible to do without the passage to forward convex combinations?
- Is it possible to replace the series of lemmas by one theorem?

Goal: Show the Emery convergence of the “maximal” up-convergent sequence of semimartingales (X^n) directly via the (P-UT) property

Convergence in the Emery topology

The series of lemmas can be summarized by the following theorem.

Theorem

Let \mathcal{X}_1 satisfy (NUPBR) and let $(X^n) \in \mathcal{X}_1$ be a sequence of semimartingales, whose end values (X_1^n) converges almost surely to a maximal element in \widehat{K}_0^1 . Then $X^n \rightarrow X$ in the Emery topology.

Convergence in the Emery topology

The series of lemmas can be summarized by the following theorem.

Theorem

Let \mathcal{X}_1 satisfy (NUPBR) and let $(X^n) \in \mathcal{X}_1$ be a sequence of semimartingales, whose end values (X_1^n) converges almost surely to a maximal element in \widehat{K}_0^1 . Then $X^n \rightarrow X$ in the Emery topology.

Proof.

The maximality of the end value allows to conclude up-convergence. The crucial part then consists in proving ((NUPBR) \Rightarrow (P-UT)). The remaining part follows as before from Memin and Slominski's theorem and a modification of the last part of the proof of FTAP, namely Emery convergence of the finite variation part. □

Two proofs of the (P-UT) property under (NUPBR)

Two proofs of the (P-UT) property under (NUPBR)

- ① Approach from mathematical finance: Use the existence of a supermartingale deflator and the fact that sequences of supermartingales satisfy the (P-UT) property, which then easily translates to the original sequence of semimartingales.

Two proofs of the (P-UT) property under (NUPBR)

- ① **Approach from mathematical finance:** Use the existence of a supermartingale deflator and the fact that sequences of supermartingales satisfy the (P-UT) property, which then easily translates to the original sequence of semimartingales.
- ② **Direct Approach:** Mimic parts of the proof of one key lemma of the original proof by Delbaen and Schachermayer.

Two proofs of the (P-UT) property under (NUPBR)

- ① **Approach from mathematical finance:** Use the existence of a supermartingale deflator and the fact that sequences of supermartingales satisfy the (P-UT) property, which then easily translates to the original sequence of semimartingales.
- ② **Direct Approach:** Mimic parts of the proof of one key lemma of the original proof by Delbaen and Schachermayer.

Remark

Due to the concatenation property (NUPBR) allows to deduce directly that $\{(H \bullet X)_1 \mid X \in \mathcal{X}_1, H \geq 0 \text{ bounded predictable s.t. } H \bullet X \geq -1\}$ is bounded in L^0 , but in order to get it for all $H \in b\mathcal{E}$ with $\|H\|_\infty \leq 1$ more work has to be done.

Supermartingale deflator

Definition

A positive càdlàg adapted process D is called **supermartingale deflator** for $1 + \mathcal{X}_1$ if D is strictly positive, $D_0 \leq 1$ and $D(1 + X)$ is a supermartingale for all $X \in \mathcal{X}_1$.

Supermartingale deflator

Definition

A positive càdlàg adapted process D is called **supermartingale deflator** for $1 + \mathcal{X}_1$ if D is strictly positive, $D_0 \leq 1$ and $D(1 + X)$ is a supermartingale for all $X \in \mathcal{X}_1$.

Theorem (Karatzas and Kardaras (2007)/ Kardaras (2013))

Assume (NUPBR) for \mathcal{X}_1 , then there exists a supermartingale deflator D .

(P-UT) property for supermartingales

Lemma

Let (Z^n) be a sequence of *non-negative supermartingales* such that $Z_0^n \leq K$ for all $n \in \mathbb{N}$ and some $K > 0$. Then (Z^n) satisfies the *P-UT property*.

(P-UT) property for supermartingales

Lemma

Let (Z^n) be a sequence of *non-negative supermartingales* such that $Z_0^n \leq K$ for all $n \in \mathbb{N}$ and some $K > 0$. Then (Z^n) satisfies the *P-UT property*.

Proof.

By an inequality of Burkholder for non-negative supermartingales S and processes $H \in b\mathcal{E}$ with $\|H\| \leq 1$ it holds that

$$cP[|(H \bullet S)|_1^* \geq c] \leq 9\mathbb{E}[|S_0|]$$

for all $c \geq 0$. Applying this inequality to Z^n and letting $c \rightarrow \infty$ yields the P-UT property. □

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Let \mathcal{X}_1 satisfy (NUPBR) and let (X^n) be any sequence of semimartingales in \mathcal{X}_1 . Then (X^n) satisfies the P-UT property.

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Let \mathcal{X}_1 satisfy (NUPBR) and let (X^n) be any sequence of semimartingales in \mathcal{X}_1 . Then (X^n) satisfies the P-UT property.

Proof.

The (P-UT) property of the supermartingales $(Z^n) := (D(1 + X^n))$ can be easily transferred to the sequence (X^n) . It relies on Itô's integration by parts formula and the fact that $(H_-^n \bullet S^n)$ satisfies (P-UT), if (S^n) is a sequence of semimartingales satisfying (P-UT) and (H^n) a sequence of adapted càdlàg processes such that $(|H_{1-}^n|_1^*)_n$ is bounded in L^0 . \square

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $|X^n - X|_1^ \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .*

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $|X^n - X|_1^* \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .

- Then for every $C > 0$ there exists a decomposition $X^n = M^n + B^n + \check{X}^n$ into a local martingale M^n , a predictable, finite variation process B^n and the finite variation process of big jumps $\check{X}_t^n = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s| > C\}}$, for $n \geq 0$, such that jumps of M^n and B^n are bounded by $2C$ uniformly in n .

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $|X^n - X|_1^* \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .

- Then for every $C > 0$ there exists a decomposition $X^n = M^n + B^n + \check{X}^n$ into a local martingale M^n , a predictable, finite variation process B^n and the finite variation process of big jumps $\check{X}_t^n = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s| > C\}}$, for $n \geq 0$, such that jumps of M^n and B^n are bounded by $2C$ uniformly in n .
- The sequence $(|M^n|_1^*)_{n \geq 0}$ is bounded in L^0 and $(M^n)_{n \geq 0}$ satisfies P-UT.

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $|X^n - X|_1^* \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .

- Then for every $C > 0$ there exists a decomposition $X^n = M^n + B^n + \check{X}^n$ into a local martingale M^n , a predictable, finite variation process B^n and the finite variation process of big jumps $\check{X}_t^n = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s^n| > C\}}$, for $n \geq 0$, such that jumps of M^n and B^n are bounded by $2C$ uniformly in n .
- The sequence $(|M^n|_1^*)_{n \geq 0}$ is bounded in L^0 and $(M^n)_{n \geq 0}$ satisfies P-UT.
- The sequence $(\text{TV}(B^n)_1)_{n \geq 0}$ of total variations of B^n is bounded in L^0 and $(B^n)_{n \geq 0}$ satisfies P-UT.

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $\|X^n - X\|_1^* \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .

- Then for every $C > 0$ there exists a decomposition $X^n = M^n + B^n + \check{X}^n$ into a local martingale M^n , a predictable, finite variation process B^n and the finite variation process of big jumps $\check{X}_t^n = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s| > C\}}$, for $n \geq 0$, such that jumps of M^n and B^n are bounded by $2C$ uniformly in n .
- The sequence $(\|M^n\|_1^*)_{n \geq 0}$ is bounded in L^0 and $(M^n)_{n \geq 0}$ satisfies P-UT.
- The sequence $(\text{TV}(B^n)_1)_{n \geq 0}$ of total variations of B^n is bounded in L^0 and $(B^n)_{n \geq 0}$ satisfies P-UT.
- Then the sequence $(\text{TV}(\check{X}_1^n))_{n \geq 0}$ of total variations of \check{X}^n is bounded in L^0 and $(\check{X}^n)_{n \geq 0}$ satisfies the P-UT property.

(P-UT) property for sequences in \mathcal{X}_1

Proposition

Assume (NUPBR). Let $(X^n)_{n \geq 0}$ together with an adapted, càdlàg process X such that $\|X^n - X\|_1^* \rightarrow 0$ in probability as $n \rightarrow \infty$ be a sequence in \mathcal{X}_1 .

- Then for every $C > 0$ there exists a decomposition $X^n = M^n + B^n + \check{X}^n$ into a local martingale M^n , a predictable, finite variation process B^n and the finite variation process of big jumps $\check{X}_t^n = \sum_{s \leq t} \Delta X_s^n 1_{\{|\Delta X_s| > C\}}$, for $n \geq 0$, such that jumps of M^n and B^n are bounded by $2C$ uniformly in n .
- The sequence $(\|M^n\|_1^*)_{n \geq 0}$ is bounded in L^0 and $(M^n)_{n \geq 0}$ satisfies P-UT.
- The sequence $(\text{TV}(B^n)_1)_{n \geq 0}$ of total variations of B^n is bounded in L^0 and $(B^n)_{n \geq 0}$ satisfies P-UT.
- Then the sequence $(\text{TV}(\check{X}_1^n))_{n \geq 0}$ of total variations of \check{X}^n is bounded in L^0 and $(\check{X}^n)_{n \geq 0}$ satisfies the P-UT property.
- The sequence $(X^n)_{n \geq 0}$ satisfies P-UT.

Remarks

Remarks

- The (P-UT) property can be easily deduced from the existence of supermartingale deflators or...

Remarks

- The (P-UT) property can be easily deduced from the existence of supermartingale deflators or...
- ...from one key lemma proved by Delbaen and Schachermayer with an additional analysis of the finite variation part.

Remarks

- The (P-UT) property can be easily deduced from the existence of supermartingale deflators or...
- ...from one key lemma proved by Delbaen and Schachermayer with an additional analysis of the finite variation part.
- The P-UT property corresponds to boundedness in the Emery topology with an interpretation from mathematical finance. It is therefore natural to investigate this property.

Conclusions

- The (P-UT) property allows to conclude Emery convergence of sequences of semimartingales whose end values satisfy a certain maximality assumption.
- This allows to obtain a variant of the the second part of the proof of FTAP without performing a measure change and passing to forward convex combinations.

Conclusions

- The (P-UT) property allows to conclude Emery convergence of sequences of semimartingales whose end values satisfy a certain maximality assumption.
- This allows to obtain a variant of the the second part of the proof of FTAP without performing a measure change and passing to forward convex combinations.
- Given a supermartingale deflator, which is implied by (NUPBR), the P-UT property is an easy consequence of a Burkholder's inequality for supermartingales.

Conclusions

- The (P-UT) property allows to conclude Emery convergence of sequences of semimartingales whose end values satisfy a certain maximality assumption.
- This allows to obtain a variant of the the second part of the proof of FTAP without performing a measure change and passing to forward convex combinations.
- Given a supermartingale deflator, which is implied by (NUPBR), the P-UT property is an easy consequence of a Burkholder's inequality for supermartingales.
- The Emery convergence result for sequences in \mathcal{X}_1 can be summarized by

(P-UT) + maximal end value \Rightarrow Convergence in Emery

- Thank you for your attention!

