

A CONVERSE THEOREM ON h -STABILITY VIA IMPULSIVE VARIATIONAL SYSTEMS

SUNG KYU CHOI AND NAMJIP KOO

ABSTRACT. In this paper we develop useful relations which estimate the difference between the solutions of nonlinear impulsive differential systems with different initial values. Then we obtain the converse h -stability theorem of Massera's type for the nonlinear impulsive systems by employing the t_∞ -similarity of the associated impulsive variational systems and relations.

1. Introduction

The qualitative properties of solutions of differential equations with impulse effects have been developed by a large number of mathematicians due to the wide application of these systems to the control theory, biology, electronics, etc (see e.g. [2, 3, 13]).

Simeonov and Bainov [20] studied the exponential stability of the solutions for impulsive differential equations by using the comparison method and piecewise continuous auxiliary functions which are analogues to Lyapunov's functions. Also, Kulev and Bainov [12] introduced the notions of various types of uniform Lipschitz stability for impulsive differential systems and obtained sufficient conditions for these notions and their relations.

Pinto [17] introduced the notion of h -stability for differential systems without impulse effect with the intention of obtaining results about stability for weakly stable differential systems under some perturbations. The various notions of h -stability given in [17, 18] include several types of known stability properties as uniform stability, exponential asymptotic stability [15] and uniform Lipschitz stability [9].

Choi et al. [5, 6] studied h -stability for the linear impulsive differential equations by means of the notions of similarity, t_∞ -similarity, and impulsive integral

Received July 16, 2015; Revised October 29, 2015.

2010 *Mathematics Subject Classification.* 34A37, 34D20.

Key words and phrases. impulsive variational systems, h -stability, impulsive differential systems, t_∞ -similarity, converse stability theorem.

This work was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(NRF-2015R1D1A1A01060103).

inequality. In [4], we proved that two concepts of h -stability and h -stability in variation for nonlinear impulsive differential systems are equivalent via t_∞ -similarity of the associated variational impulsive systems and impulsive integral inequalities. Also, we characterized h -stability for nonlinear impulsive differential systems by using the notions of piecewise continuous auxiliary functions and impulsive differential inequalities. Many authors [2, 3, 5, 12, 13, 16, 20] have studied the various types of stability of solutions for impulsive differential equations. However, to the best of our knowledge, there are no papers published on the converse h -stability theorem for nonlinear impulsive differential systems.

Motivated by the above discussion, we develop useful relations which estimate the difference between the solutions of nonlinear impulsive differential systems with different initial values. Then we obtain the converse h -stability theorem of Massera's type for the nonlinear impulsive systems by employing the t_∞ -similarity of the associated impulsive variational systems and relations.

2. Preliminary notes and definitions

Let $\mathbb{R}_+ = [0, \infty)$ and \mathbb{R}^n be the n -dimensional Euclidean space with a convenient vector norm $|\cdot|$. We consider the impulsive differential system with impulses at fixed times

$$(2.1) \quad \begin{cases} x' &= f(t, x), \quad t \neq \tau_k, \\ \Delta x &= I_k(x), \quad t = \tau_k, \quad k = 1, 2, \dots, \end{cases}$$

where $\Delta x(t) = x(t+0) - x(t-0)$. Assume that the following basic conditions hold:

- (A1) $\{\tau_k\}$ is an unbounded increasing sequence satisfying $0 \leq t_0 = \tau_0 < \tau_1 < \tau_2 < \dots$ and $\lim_{k \rightarrow \infty} \tau_k = \infty$.
- (A2) The function $f : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and has a continuous partial derivative $f_x = \frac{\partial f}{\partial x}$ in $(\tau_{k-1}, \tau_k] \times \mathbb{R}^n$, $k = 1, 2, \dots$, and $f(t, 0) = 0$ for each $t \in \mathbb{R}_+$.
- (A3) For any $x \in \mathbb{R}^n$ and any $k = 1, 2, \dots$, the functions f and f_x have finite limits as $(t, y) \rightarrow (\tau_k, x)$, $t > \tau_k$.
- (A4) Each function $I_k : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuously differentiable in \mathbb{R}^n and there exist nonnegative constants l_k such that

$$|I_k(x) - I_k(y)| \leq l_k |x - y|, \quad k \in \mathbb{N}, x, y \in \mathbb{R}^n,$$

and $I_k(0) = 0$, $k = 1, 2, \dots$

- (A5) The solution $x(t, t_0, x_0)$ of system (2.1) which satisfies the initial condition $x(t_0+0, t_0, x_0) = x_0$ is defined in the interval (t_0, ∞) , and is left continuous. And at the moments τ_k the following relations hold

$$x(\tau_k-0) = x(\tau_k), \quad \Delta x(\tau_k) = x(\tau_k+0) - x(\tau_k-0).$$

Then it follows from the condition (A5) that the solution $x(t, t_0, x_0)$ of system (2.1) through the initial value (t_0, x_0) is described as follows:

$$x(t, t_0, x_0) = \begin{cases} x(t, t_0, x_0), & t_0 \leq t \leq \tau_1, \\ x(t, \tau_1, x_1^+), & \tau_1 < t \leq \tau_2, \\ \dots \\ x(t, \tau_k, x_k^+), & \tau_k < t \leq \tau_{k+1}, \\ \dots, \end{cases}$$

where $x_k^+ = x_k + I_k(x_k)$ and $x_k = x(\tau_k)$. For global existence of solutions for system (2.1), see [21]. The basic notions and important theory for system (2.1) were described in detail in [2, 3, 13].

Together with system (2.1), we consider the associated impulsive variational systems

$$(2.2) \quad \begin{cases} v' & = f_x(t, 0)v, \quad t \neq \tau_k, \\ \Delta v & = \frac{\partial I_k(0)}{\partial x} v, \quad t = \tau_k, \quad k \in \mathbb{N}, \\ v(t_0 + 0) & = v_0 \end{cases}$$

and

$$(2.3) \quad \begin{cases} z' & = f_x(t, x(t, t_0, x_0))z, \quad t \neq \tau_k, \\ \Delta z & = \frac{\partial I_k(x(\tau_k))}{\partial x} z, \quad t = \tau_k, \quad k \in \mathbb{N}, \\ z(t_0 + 0) & = z_0. \end{cases}$$

It follows from [13, Theorem 2.4.1] that the fundamental matrix solution $\Phi(t, t_0, 0)$ of system (2.2) is given by

$$\Phi(t, t_0, 0) = \frac{\partial x(t, t_0, 0)}{\partial x_0}, \quad t \neq \tau_k$$

and the fundamental matrix solution $\Phi(t, t_0, x_0)$ of system (2.3) is given by

$$(2.4) \quad \Phi(t, t_0, x_0) = \frac{\partial x(t, t_0, x_0)}{\partial x_0}, \quad t \neq \tau_k.$$

Let $M_n(\mathbb{R})$ be the set of all $n \times n$ matrices over \mathbb{R} and let $PC(\mathbb{R}_+, M_n(\mathbb{R}))$ denote the class of piecewise continuous functions from \mathbb{R}_+ to $M_n(\mathbb{R})$ with discontinuities of the first kind only at $t = \tau_k, k \in \mathbb{N}$.

We consider two linear homogeneous impulsive systems

$$(2.5) \quad \begin{cases} x' = A(t)x, & t \neq \tau_k, \\ \Delta x = A_k x, & t = \tau_k, \quad k \in \mathbb{N}, \end{cases}$$

and

$$(2.6) \quad \begin{cases} y' = B(t)y, & t \neq \tau_k, \\ \Delta y = B_k y, & t = \tau_k, \quad k \in \mathbb{N}, \end{cases}$$

under the assumption that the following conditions hold:

(B1) $A, B \in PC(\mathbb{R}_+, M_n(\mathbb{R}))$.

(B2) $A_k, B_k \in M_n(\mathbb{R}), \det(E + A_k) \neq 0$, and $\det(E + B_k) \neq 0, k \in \mathbb{N}$.

Then it follows from [2, p. 7] that the solutions of system (2.5) can be written in the form

$$(2.7) \quad x(t, t_0, x_0) = X(t, t_0)x_0,$$

where $X(t)$ is a fundamental matrix of system (2.5) and the Cauchy matrix $X(t, t_0)$ [2, 13] is given by

$$X(t, t_0) = \begin{cases} U_k(t, t_0), & 0 \leq t_0 \leq t \leq \tau_k, \\ U_{k+1}(t, \tau_k^+)(E + A_k)U_k(\tau_k, t_0), & t_0 \leq \tau_k < t \leq \tau_{k+1}, \\ U_{k+1}(t, \tau_k^+) \prod_{j=1}^k (E + A_j)U_j(\tau_j, \tau_{j-1}^+), & t_0 = \tau_0^+ \leq \tau_1 < \tau_k < t \leq \tau_{k+1}. \end{cases}$$

Here $U_k(t, s) (k \in \mathbb{N}, t, s \in (\tau_{k-1}, \tau_k])$ is the Cauchy matrix for the linear differential system

$$(2.8) \quad x' = A(t)x, \quad t \in (\tau_{k-1}, \tau_k], k = 1, 2, \dots$$

Denote by \mathfrak{S} the set of all matrix functions $S : \mathbb{R}_+ \rightarrow M_n(\mathbb{R})$ which belong to $PC(\mathbb{R}_+, M_n(\mathbb{R}))$ and are bounded in \mathbb{R}_+ together with their inverse $S^{-1}(t)$. Let \mathcal{M} be the set of the linear homogeneous impulsive systems as follows

$$\mathcal{M} := \{(A, A_k) \mid A \in PC(\mathbb{R}_+, M_n(\mathbb{R})), A_k \in M_n(\mathbb{R}), \det(E + A_k) \neq 0, k \in \mathbb{N}\}.$$

We recall the notion of t_∞ -similarity with impulse effect in \mathcal{M} which is analogue to the concept of t_∞ -similarity introduced by Conti [8].

Definition 2.1 ([3]). We say that $(A, A_k) \in \mathcal{M}$ is t_∞ -similar to $(B, B_k) \in \mathcal{M}$ if there exists a matrix-valued function $S \in \mathfrak{S}$ such that

$$(2.9) \quad S'(t) - A(t)S(t) + S(t)B(t) \equiv F_0 \in L_1, \quad t \neq \tau_k,$$

$$(2.10) \quad \Delta S(\tau_k) - A_k S(\tau_k) + S(\tau_k^+) B_k \equiv F_k \in l_1, \quad t = \tau_k, k \in \mathbb{N},$$

where $\Delta S(\tau_k) = S(\tau_k^+) - S(\tau_k)$. We say that system (2.5) is t_∞ -similar to system (2.6) if $(A, A_k) \in \mathcal{M}$ is t_∞ -similar to $(B, B_k) \in \mathcal{M}$ for each $k \in \mathbb{N}$. Note that the relation $(A, A_k) \sim (B, B_k) : S$ is an equivalence relation.

Remark 2.2. The notion of t_∞ -similarity preserves various concepts of stability: stability, uniform stability, uniform asymptotic stability and strict stability (see [3, Theorem 10.3]). In the case when $F_0 = 0$ in (2.9) and $F_k = 0$ in (2.10) for each $k \in \mathbb{N}$, then t_∞ -similarity reduces to kinematical similarity for impulsive linear equations [1].

Definition 2.3 ([20]). We say that the function $V : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ belongs to the class v_0 if

- (i) V is continuous in $G_k = \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^n : \tau_{k-1} < t < \tau_k\}$ for each $k \in \mathbb{N}$ and $V(t, 0) = 0$ for each $t \in \mathbb{R}_+$;

(ii) for any $k \in \mathbb{N}$ and $x \in \mathbb{R}^n$ there exist the finite limits

$$V(\tau_k - 0, x) = \lim_{\substack{(t,y) \rightarrow (\tau_k, x) \\ t < \tau_k}} V(t, y), \quad V(\tau_k + 0, x) = \lim_{\substack{(t,y) \rightarrow (\tau_k, x) \\ t > \tau_k}} V(t, y)$$

and the equality $V(\tau_k - 0, x) = V(\tau_k, x)$ holds.

We note that if $t \neq \tau_k$, then $V(t + 0, x)$ denotes $V(t, x)$.

Definition 2.4 ([20]). The function $V \in v_0$ belongs to the class v_1 if it is continuously differentiable in $G = \cup_{k=1}^\infty G_k$.

Remark 2.5 ([22]). For $(t, x) \in G$ we define the derivative of the function $V \in v_1$ with respect to system (2.1) by

$$\dot{V}_{(2.1)}(t, x) = \frac{\partial V(t, x)}{\partial t} + \frac{\partial V(t, x)}{\partial x} f(t, x)$$

and the upper right derivative of the function $V \in v_0$ with respect to the solutions of system (2.1) by

$$D_{(2.1)}^+ V(t, x) = \limsup_{s \rightarrow 0^+} \frac{1}{s} [V(t + s, x(t + s, t, x)) - V(t, x)].$$

We note that if the function $V \in v_0$ is locally Lipschitz continuous with respect to $x \in \mathbb{R}^n$, then for $(t, x) \in G_k$ we have

$$D_{(2.1)}^+ V(t, x) = \limsup_{s \rightarrow 0^+} \frac{1}{s} [V(t + s, x + sf(t, x)) - V(t, x)].$$

We recall some notions of various types of h -stability for the zero solution of system (2.1). For nonlinear differential systems without impulse effect, this notion was introduced by Pinto [17, 18].

Definition 2.6. The zero solution of system (2.1) is said to be

(hS) h -stable if there exist a positive bounded left-continuous function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$, $\delta > 0$, and a constant $c \geq 1$ such that

$$|x(t, t_0, x_0)| \leq c|x_0|h(t)h(t_0)^{-1}, \quad t \geq t_0$$

for $|x_0| \leq \delta$ (here $h(t)^{-1} = \frac{1}{h(t)}$);

(GhS) globally h -stable if system (2.1) is h -stable for every $x_0 \in D$, where $D \subset \mathbb{R}^n$ is a region which includes the origin;

(hSV) h -stable in variation if system (2.3) is h -stable;

(GhSV) globally h -stable in variation if system (2.3) is globally h -stable.

The various notions of h -stability given by Definition 2.6 include several notions of well-known stability such as uniform stability, uniform asymptotic stability, uniform Lipschitz stability, exponential stability, and non-uniform stability. For linear impulsive differential systems, we note that the notions of various types of the above h -stability are equivalent.

3. Main results

In this section we develop a useful inequality which estimates the difference between the solutions of nonlinear impulsive systems with different initial values by using fundamental solutions of the associated impulsive variational systems. Then we obtain the converse h -stability theorem of Massera's type for the nonlinear impulsive systems by employing the t_∞ -similarity of the associated impulsive variational systems and an inequality.

To do this, we need the following lemma for differential systems.

Lemma 3.1 ([14, Theorem 1.2.9]). *Let $g \in C(\mathbb{R}_+ \times \mathbb{R}^n, \mathbb{R}^n)$, and g_y exist and be continuous on $\mathbb{R}_+ \times \mathbb{R}^n$. Assume that $y(t, t_0, x_0)$ and $\hat{y}(t, t_0, y_0)$ are the solutions of differential system of the following form*

$$(3.1) \quad y' = g(t, y)$$

through (t_0, x_0) and (t_0, y_0) , respectively, existing for $t \geq t_0$, such that x_0 and y_0 belong to a convex subset of \mathbb{R}^n . Then

$$y(t, t_0, x_0) - \hat{y}(t, t_0, y_0) = \int_0^1 \Psi(t, t_0, y_0 + s(x_0 - y_0)) ds \cdot (x_0 - y_0)$$

holds for $t \geq t_0$, where $\Psi(t, t_0, x_0) = \frac{\partial y(t, t_0, x_0)}{\partial x_0}$.

We need some lemmas to obtain the converse h -stability theorem for system (2.1). We obtain the following result which estimates the difference between the solutions of nonlinear impulsive system (2.1) with different initial values in terms of the fundamental solutions of the associated impulsive variational systems.

Lemma 3.2. *Suppose that $D \subset \mathbb{R}^n$ is a convex subset and each map I_k in condition (A4) is Lipschitzian in x with nonnegative constants l_k , i.e.,*

$$(3.2) \quad |I_k(x) - I_k(y)| \leq l_k |x - y|, \quad k \in \mathbb{N}.$$

Then, for $t \geq t_0$,

$$\begin{aligned} & |x(t, t_0, x_0) - \hat{x}(t, t_0, y_0)| \\ & \leq \sup_{\eta \in D} |\Phi(t, \tau_k^+, \eta)| \prod_{i=1}^k (1 + l_i) |\Phi(\tau_i, \tau_{i-1}^+, \eta)| |x_0 - y_0|, \quad \tau_k < t \leq \tau_{k+1}, \quad k \in \mathbb{N}, \end{aligned}$$

where $x(t, t_0, x_0)$ and $\hat{x}(t, t_0, y_0)$ are the solutions of system (2.1) through (t_0, x_0) and (t_0, y_0) , respectively, existing for $t \geq t_0$, such that x_0 and y_0 belong to a convex subset of \mathbb{R}^n and $\tau_0^+ = t_0$. Moreover, $\Phi(t, t_0, x_0)$ is the fundamental matrix solution of system (2.3) given by (2.4).

Proof. If $t_0 \leq t \leq \tau_1$, then it follows from Lemma 3.1 that

$$(3.3) \quad x(t, t_0, x_0) - \hat{x}(t, t_0, y_0) = \int_0^1 \Phi(t, t_0, y_0 + s(x_0 - y_0)) ds \cdot (x_0 - y_0).$$

Thus, we have

$$(3.4) \quad |x(t, t_0, x_0) - \hat{x}(t, t_0, y_0)| \leq \int_0^1 |\Phi(t, t_0, y_0 + s(x_0 - y_0))| ds |x_0 - y_0| \\ \leq \sup_{\eta \in D} |\Phi(t, t_0, \eta)| |x_0 - y_0|, \quad t_0 \leq t \leq \tau_1,$$

where D is a convex subset of \mathbb{R}^n containing x_0 and y_0 .

If $\tau_1 < t \leq \tau_2$, then it follows from Lemma 3.1 and (3.3) that

$$x(t, \tau_1, x(\tau_1^+)) - \hat{x}(t, \tau_1, \hat{x}(\tau_1^+)) \\ = \int_0^1 \Phi(t, \tau_1^+, p_1(s)) ds [x(\tau_1^+) - \hat{x}(\tau_1^+)] \\ = \int_0^1 \Phi(t, \tau_1^+, p_1(s)) ds \left[\int_0^1 \Phi(\tau_1, t_0, p_0(s)) ds (x_0 - y_0) + I_1(x(\tau_1)) - I_1(\hat{x}(\tau_1)) \right],$$

where $p_0(s) = y_0 + s(x_0 - y_0)$ and $p_1(s) = \hat{x}(\tau_1^+) + s(x(\tau_1^+) - \hat{x}(\tau_1^+))$ and $0 \leq s \leq 1$. In view of (3.2) and (3.4), we obtain

$$(3.5) \quad |x(t, \tau_1, x(\tau_1^+)) - \hat{x}(t, \tau_1, \hat{x}(\tau_1^+))| \\ \leq \int_0^1 |\Phi(t, \tau_1^+, p_1(s))| ds \left[\int_0^1 |\Phi(\tau_1, t_0, p_0(s))| ds |x_0 - y_0| + |I_1(x(\tau_1)) - I_1(\hat{x}(\tau_1))| \right] \\ \leq \int_0^1 |\Phi(t, \tau_1^+, p_1(s))| ds \left[\int_0^1 |\Phi(\tau_1, t_0, p_0(s))| ds |x_0 - y_0| + l_1 |x(\tau_1) - \hat{x}(\tau_1)| \right] \\ \leq \int_0^1 |\Phi(t, \tau_1^+, p_1(s))| ds \int_0^1 |\Phi(\tau_1, t_0, p_0(s))| ds |x_0 - y_0| (1 + l_1) \\ \leq \sup_{\eta \in D} |\Phi(t, \tau_1^+, \eta)| (1 + l_1) |\Phi(\tau_1, t_0, \eta)| |x_0 - y_0|, \quad \tau_1 < t \leq \tau_2,$$

where $p_0(s) = y_0 + s(x_0 - y_0)$ and $p_1(s) = \hat{x}(\tau_1^+) + s(x(\tau_1^+) - \hat{x}(\tau_1^+))$ and $0 \leq s \leq 1$.

If $\tau_2 < t \leq \tau_3$, then it follows from Lemma 3.1 that

$$x(t, \tau_2, x(\tau_2^+)) - \hat{x}(t, \tau_2, \hat{x}(\tau_2^+)) = \int_0^1 \Phi(t, \tau_2^+, p_2(s)) ds [x(\tau_2^+) - \hat{x}(\tau_2^+)] \\ = \int_0^1 \Phi(t, \tau_2^+, p_2(s)) ds [x(\tau_2) - \hat{x}(\tau_2) + I_2(x(\tau_2)) - I_2(\hat{x}(\tau_2))],$$

where $p_2(s) = \hat{x}(\tau_2^+) + s(x(\tau_2^+) - \hat{x}(\tau_2^+))$ and $0 \leq s \leq 1$. In view of (3.2) and (3.5), we have

$$|x(t, \tau_2, x(\tau_2^+)) - \hat{x}(t, \tau_2, \hat{x}(\tau_2^+))| \\ \leq \int_0^1 |\Phi(t, \tau_2^+, p_2(s))| ds [|x(\tau_2) - \hat{x}(\tau_2)| + |I_2(x(\tau_2)) - I_2(\hat{x}(\tau_2))|]$$

$$\begin{aligned}
 &\leq \int_0^1 |\Phi(t, \tau_2^+, p_2(s))| ds [|x(\tau_2) - \hat{x}(\tau_2)| + l_2|x(\tau_2) - \hat{x}(\tau_2)|] \\
 &\leq \int_0^1 |\Phi(t, \tau_2^+, p_2(s))| ds (1 + l_2)|x(\tau_2) - \hat{x}(\tau_2)| \\
 &\leq \int_0^1 |\Phi(t, \tau_2^+, p_2(s))| ds (1 + l_1)(1 + l_2) \int_0^1 |\Phi(\tau_2, \tau_1^+, p_1(s))| ds \\
 &\quad \int_0^1 |\Phi(\tau_1, t_0, p_0(s))| ds |x_0 - y_0| \\
 &\leq \sup_{\eta \in D} |\Phi(t, \tau_2^+, \eta)| ds (1 + l_1)(1 + l_2) |\Phi(\tau_2, \tau_1^+, \eta)| |\Phi(\tau_1, t_0, \eta)| |x_0 - y_0|,
 \end{aligned}$$

where $\tau_2 < t \leq \tau_3$, $p_0(s) = y_0 + s(x_0 - y_0)$ and $p_i(s) = \hat{x}(\tau_i^+) + s(x(\tau_i^+) - \hat{x}(\tau_i^+))$, $i = 1, 2$ and $0 \leq s \leq 1$.

It follows from mathematical induction that

$$\begin{aligned}
 &x(t, \tau_k, x(\tau_k^+)) - \hat{x}(t, \tau_k, \hat{x}(\tau_k^+)) = \int_0^1 \Phi(t, \tau_k^+, p_k(s)) ds [x(\tau_k^+) - \hat{x}(\tau_k^+)] \\
 &= \int_0^1 \Phi(t, \tau_k^+, p_k(s)) ds [x(\tau_k) - \hat{x}(\tau_k) + I_k(x(\tau_k)) - I_k(\hat{x}(\tau_k))],
 \end{aligned}$$

where $\tau_k < t \leq \tau_{k+1}$, $k \in \mathbb{N}$, $p_k(s) = \hat{x}(\tau_k^+) + s(x(\tau_k^+) - \hat{x}(\tau_k^+))$, $0 \leq s \leq 1$. Thus it follows from (3.2) that

$$\begin{aligned}
 &|x(t, \tau_k, x(\tau_k^+)) - \hat{x}(t, \tau_k, \hat{x}(\tau_k^+))| \\
 &\leq \int_0^1 |\Phi(t, \tau_k^+, p_k(s))| ds [|x(\tau_k) - \hat{x}(\tau_k)| + |I_k(x(\tau_k)) - I_k(\hat{x}(\tau_k))|] \\
 &\leq \int_0^1 |\Phi(t, \tau_k^+, p_k(s))| ds (1 + l_k) |x(\tau_k) - \hat{x}(\tau_k)| \\
 &\leq \sup_{\eta \in D} |\Phi(t, \tau_k^+, \eta)| \prod_{i=1}^k (1 + l_i) |\Phi(\tau_i, \tau_{i-1}^+, \eta)| |x_0 - y_0|, \quad \tau_k < t \leq \tau_{k+1}, k \in \mathbb{N},
 \end{aligned}$$

where $p_k(s) = \hat{x}(\tau_k^+) + s(x(\tau_k^+) - \hat{x}(\tau_k^+))$, $0 \leq s \leq 1$, and $\tau_0^+ = t_0$. This completes the proof. \square

As a consequence of Lemma 3.2, we easily obtain the following result.

Corollary 3.3 ([4]). *Let $I_k(x) = d_k x$ in condition (A4), and let d_k be constants. Assume that $x(t, t_0, x_0)$ and $\hat{x}(t, t_0, y_0)$ are the solutions of system (2.1) through (t_0, x_0) and (t_0, y_0) , respectively, existing for $t \geq t_0$, such that x_0 and y_0 belong to a convex subset of \mathbb{R}^n . Then,*

$$\begin{aligned}
 &x(t, t_0, x_0) - \hat{x}(t, t_0, y_0) \\
 &= \int_0^1 \Phi(t, \tau_k^+, p_k(s)) ds \prod_{i=1}^k (1 + d_i) \int_0^1 \Phi(\tau_i, \tau_{i-1}^+, p_{i-1}(s)) ds \cdot (x_0 - y_0),
 \end{aligned}$$

holds for $\tau_k < t \leq \tau_{k+1}$, $t \geq t_0$, where $p_0(s) = y_0 + s(x_0 - y_0)$, $p_i(s) = \hat{x}(\tau_i^+) + s(x(\tau_i^+) - \hat{x}(\tau_i^+))$, $i = 1, \dots, k$, $0 \leq s \leq 1$, and $\tau_0^+ = t_0$.

We can obtain the following relation between the solution $x(t, t_0, x_0)$ of system (2.1) and $\Phi(t, t_0, x_0)$ given by (2.4).

Corollary 3.4. *Assume that $\hat{x}(t, t_0, y_0) = 0$ and $y_0 = 0$ in Corollary 3.3. Then,*

$$x(t, t_0, x_0) = \int_0^1 \Phi(t, \tau_k^+, sx(\tau_k^+))ds \prod_{i=1}^k (1 + d_i) \int_0^1 \Phi(\tau_i, \tau_{i-1}^+, sx(\tau_{i-1}^+))ds \cdot x_0,$$

holds for $\tau_k < t \leq \tau_{k+1}$, $k \in \mathbb{N}$, $t \geq t_0$, where $\tau_0 = t_0$.

Remark 3.5. Assume that $I_k(x) = 0$, $k \in \mathbb{N}$, in condition (A4). From Corollary 3.3 and the uniqueness of solutions, it follows that

$$x(t, t_0, x_0) = \int_0^1 \Phi(t, t_0, sx_0)ds \cdot x_0, \quad t \geq t_0.$$

From Theorems 3.9 and 3.11 in [4], we can show that two notions of hS and hSV for system (2.1) are equivalent.

Lemma 3.6 ([4, Theorem 3.11]). *Assume that impulsive variational system (2.2) is t_∞ -similar to impulsive variational system (2.3) and $\sup_{k \in \mathbb{N}} \frac{h(\tau_k)}{h(\tau_k^+)}$ is bounded. Then, the zero solution of system (2.3) is h -stable provided the zero solution of system (2.1) is h -stable.*

To prove our main result, we need the following stability criterion for linear impulsive systems which is adapted from Lemma 1 in [18]. We state the result of [4, Lemma 3.7] without its proof.

Lemma 3.7. *Suppose that the zero solution of system (2.3) is h -stable. Then, there exist a constant $c \geq 1$ and a positive bounded left-continuous function h defined on \mathbb{R}_+ such that for every $x_0 \in \mathbb{R}^n$,*

$$(3.6) \quad |\Phi(t, t_0, x_0)| \leq ch(t)h(t_0)^{-1}, \quad t \geq t_0 \geq 0,$$

where $\Phi(t, t_0, x_0)$ is the fundamental matrix solution of system (2.3).

Choi and Koo [4] provided some sufficient conditions for h -stability of solutions of system (2.1) by using the comparison method and piecewise continuous auxiliary functions which are analogues to Lyapunov functions. In order to obtain the main result (Theorem 3.8), Lemma 3.2 and the notion of t_∞ -similarity of the associated impulsive variational systems are used. Now, we can obtain the converse h -stability theorem of Massera's type for nonlinear impulsive differential systems which guarantee the existence of piecewise continuous Lyapunov's functions with certain properties. It is adapted from Lemma 3 in [20] and Theorem 15.3 in [2].

Theorem 3.8. Assume that $(f_x(t, 0), \frac{\partial I_k(0)}{\partial x}) \in \mathcal{M}$ is t_∞ -similar to

$$(f_x(t, x(t, t_0, x_0)), \frac{\partial I_k(x(\tau_k))}{\partial x}) \in \mathcal{M}$$

and $\sum_{k=1}^\infty l_k < \infty$ for nonnegative constants l_k of each map I_k in the condition (A4). Suppose further that the zero solution of system (2.1) is globally h -stable such that $h'(t)$ exists and is continuous on \mathbb{R}_+ . Then, there exist a positive constant c and a function $V : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

- (i) $V \in v_0$ and $V(t, x)$ is Lipschitzian in x with Lipschitz constant L for each $t \in \mathbb{R}_+$;
- (ii) $|x| \leq V(t, x) \leq c|x|$, $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^n$;
- (iii) $D_{(2.1)}^+ V(t, x) \leq \frac{h'(t)}{h(t)} V(t, x)$, $t \neq \tau_k$;
- (iv) $V(\tau_k^+, x + I_k(x)) \leq V(\tau_k, x)$, $x \in \mathbb{R}^n$, $k \in \mathbb{N}$,

where $L = c \exp(\sum_{i=1}^\infty l_i)$.

Proof. Since the zero solution of system (2.1) is globally h -stable, there exist a constant $c \geq 1$ and a positive bounded continuous function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that

$$|x(t, t_0, x_0)| \leq c|x_0|h(t)h(t_0)^{-1}, \quad t \geq t_0, \quad x_0 \in \mathbb{R}^n.$$

Define the function $V : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}_+$ by

$$V(t, x) = \sup_{\tau \geq 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t), \quad t \neq \tau_k,$$

$$V(\tau_k, x) = V(\tau_k^-, x), \quad k = 1, 2, \dots,$$

where $x(t + \tau, t, x)$ is a solution of system (2.1) through $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^n$. Then, we obtain

$$|x| = |x(t, t, x)| \leq \sup_{\tau \geq 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t)$$

and

$$V(t, x) \leq c|x|h(t + \tau)h(t)^{-1}h(t + \tau)^{-1}h(t) = c|x|.$$

Thus the property (ii) is proved for $t \neq \tau_k$ for each $k \in \mathbb{N}$.

From the definition of hS and uniqueness of solutions of system (2.1), it follows that $V(t, x)$ is defined on $\mathbb{R}_+ \times \mathbb{R}^n$.

We show that $V(t, x)$ is Lipschitzian in x for each $t \in \mathbb{R}_+$. Let $(t, x), (t, y) \in \mathbb{R}_+ \times \mathbb{R}^n$. From Lemmas 3.2, 3.6 and 3.7, we have

$$(3.7) \quad |V(t, x) - V(t, y)|$$

$$\leq \sup_{\tau \geq 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t) - \sup_{\tau \geq 0} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t)|$$

$$\leq \sup_{\tau \geq 0} |x(t + \tau, t, x) - x(t + \tau, t, y)|h(t + \tau)^{-1}h(t)$$

$$\leq |x - y| \sup_{\eta \in D} |\Phi(t + \tau, \tau_k^+, \eta)| \prod_{i=1}^k (1 + l_i) |\Phi(\tau_i, \tau_{i-1}^+, \eta)|h(t + \tau)^{-1}h(t)$$

$$\begin{aligned} &\leq |x - y|c \prod_{i=1}^{\infty} (1 + l_i)h(t + \tau)h(t)^{-1}h(t + \tau)^{-1}h(t) \\ &\leq |x - y|c \exp\left(\sum_{i=1}^{\infty} l_i\right) \\ &\leq L|x - y|, \tau_k < t + \tau \leq \tau_{k+1}, \end{aligned}$$

where D is a convex subset of \mathbb{R}^n containing x and y , and $L = c \exp(\sum_{i=1}^{\infty} l_i)$. This implies that $V(t, x)$ is Lipschitzian in x for each $t \in \mathbb{R}_+$. Let $x, y \in \mathbb{R}^n$, $\tau_{k-1} < t < \tau_k$ and $\delta > 0$ be such that $t + \delta < \tau_k$. Then we have

$$(3.8) \quad \begin{aligned} |V(t + \delta, x) - V(t, y)| &\leq |V(t + \delta, x) - V(t + \delta, y)| \\ &\quad + |V(t + \delta, y) - V(t + \delta, x(t + \delta, t, y))| \\ &\quad + |V(t + \delta, x(t + \delta, t, y)) - V(t, y)|. \end{aligned}$$

In view of (3.7) and $\lim_{\delta \rightarrow 0} |y - x(t + \delta, t, y)| = 0$ for $t \neq \tau_k$, then the first two terms in the right-hand side of estimate (3.8) are small if $|x - y|$ and δ are small.

Denote $a(\delta) = \sup_{\tau > \delta} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t)$. The function $a(\delta)$ is non-increasing for $\delta \geq 0$ and $\lim_{\delta \rightarrow 0+} a(\delta) = a(0)$ since $|x(t + \tau, t, x)|h(t + \tau)h(t)^{-1}$ is a bounded and piecewise continuous function for $\tau \geq 0$ and is continuous in some neighborhood of $\tau = 0$.

Then, for the third term in (3.8), we obtain

$$(3.9) \quad \begin{aligned} &|V(t + \delta, x(t + \delta, t, y)) - V(t, y)| \\ &= \left| \sup_{s > 0} |x(t + \delta + s, t + \delta, x(t + \delta, t, y))|h(t + \delta + s)^{-1}h(t + \delta) \right. \\ &\quad \left. - \sup_{\tau > 0} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t) \right| \\ &= \left| \sup_{\tau > \delta} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t + \delta) - \sup_{\tau > 0} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t) \right| \\ &= |a(\delta)h(t)^{-1}h(t + \delta) - a(0)| \rightarrow 0 \text{ as } \delta \rightarrow 0_+, \end{aligned}$$

since $h(t)$ is continuous. Hence $V(t, x)$ is continuous for $x \in \mathbb{R}^n$ and $t \neq \tau_k$.

Let $x \in \mathbb{R}^n$, $t \in \mathbb{R}_+$, $t \neq \tau_k$ and $\delta > 0$. Then

$$\begin{aligned} &D_{(2,1)}^+ V(t, x) \\ &= \lim_{\delta \rightarrow 0+} \sup \frac{1}{\delta} [V(t + \delta, x(t + \delta, t, x)) - V(t, x)] \\ &= \lim_{\delta \rightarrow 0+} \sup \frac{1}{\delta} [\sup_{\tau > 0} |x(t + \delta + \tau, t + \delta, x(t + \delta, t, x))|h(t + \delta + \tau)^{-1}h(t + \delta) \\ &\quad - \sup_{\tau > 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t)] \\ &\leq \lim_{\delta \rightarrow 0+} \sup \frac{1}{\delta} [\sup_{\tau > \delta} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t + \delta) \end{aligned}$$

$$\begin{aligned}
 & - \sup_{\tau > 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t)] \\
 \leq & \lim_{\delta \rightarrow 0_+} \sup_{\tau > 0} \frac{1}{\delta} [\sup_{\tau > 0} |x(t + \tau, t, x)|h(t + \tau)^{-1}h(t)(h(t + \delta)h(t)^{-1} - 1)] \\
 \leq & \lim_{\delta \rightarrow 0_+} [\sup_{\tau > 0} \frac{1}{\delta} [h(t + \delta)h(t)^{-1} - 1]V(t, x)] \\
 \leq & \frac{h'(t)}{h(t)}V(t, x), \quad t \neq \tau_k.
 \end{aligned}$$

Since, for small $\delta > 0$ and $t \neq \tau_k$,

$$\begin{aligned}
 & |V(t + \delta, x + \delta f(t, x)) - V(t, x)| \\
 \leq & |V(t + \delta, x + \delta f(t, x)) - V(t + \delta, x(t + \delta, t, x))| \\
 & + |V(t + \delta, x(t + \delta, t, x)) - V(t, x)| \\
 \leq & L|x + \delta f(t, x) - x(t + \delta, t, x)| + |V(t + \delta, x(t + \delta, t, x)) - V(t, x)|,
 \end{aligned}$$

it follows that

$$D_{(2.1)}^+ V(t, x) \leq \frac{h'(t)}{h(t)}V(t, x), \quad t \neq \tau_k.$$

Thus, the property (iii) is proved for $t \neq \tau_k$ for each $k \in \mathbb{N}$.

Let $\tau_k \in \mathbb{R}_+$ and $x \in \mathbb{R}^n$ be fixed and $t_i \in (\tau_k, \tau_{k+1})$, $x_i \in \mathbb{R}^n$ for $i = 1, 2$. Then

$$\begin{aligned}
 (3.10) \quad |V(t_1, x_1) - V(t_2, x_2)| \leq & |V(t_1, x_1) - V(t_1, x(t_1, \tau_k, x))| \\
 & + |V(t_2, x_2) - V(t_2, x(t_2, \tau_k, x))| \\
 & + |V(t_1, x(t_1, \tau_k, x)) - V(t_2, x(t_2, \tau_k, x))|.
 \end{aligned}$$

Since $V(t, x)$ and $f(t, x)$ are locally Lipschitz continuous, we obtain

$$\begin{aligned}
 |V(t_i, x_i) - V(t_i, x(t_i, \tau_k, x))| & \leq c|x_i - x(t_i, \tau_k, x)| \\
 & \leq c|x_i - x| + c|x - x(t_i, \tau_k, x)| \\
 & \rightarrow 0 \text{ as } x_i \rightarrow x, \quad t_i \rightarrow \tau_k, \quad i = 1, 2,
 \end{aligned}$$

where $\lim_{t_i \rightarrow \tau_k} x(t_i, \tau_k^+, x) = x$ for $i = 1, 2$. Also, we obtain

$$\begin{aligned}
 & |V(t_1, x(t_1, \tau_k^+, x)) - V(t_2, x(t_2, \tau_k^+, x))| \\
 = & \sup_{s > 0} |x(t_1 + s, t_1, x(t_1, \tau_k^+, x))|h(t_1 + s)^{-1}h(t_1) \\
 & - \sup_{s > 0} |x(t_2 + s, t_2, x(t_2, \tau_k^+, x))|h(t_2 + s)^{-1}h(t_2)| \\
 = & |a(t_1 - \tau_k)h(\tau_k^+)h(t_1)^{-1} - a(t_2 - \tau_k)h(\tau_k^+)h(t_2)^{-1}| \rightarrow 0 \text{ as } t_i \rightarrow \tau_k, \quad i = 1, 2,
 \end{aligned}$$

since $a(\delta) = \sup_{\tau > \delta} |x(t + \tau, t, y)|h(t + \tau)^{-1}h(t)$ is non-increasing for $\delta \geq 0$ and $\lim_{\delta \rightarrow 0_-} a(\delta) = a(0)$. This implies that the limit $V(\tau_k^+, x)$ exists. From the similar manner, we have the existence of limit $V(\tau_k^-, x)$. From the equality of $V(\tau^-, x) = V(\tau_k, x)$, we have $V \in v_0$.

Let $\eta(t, t_0, x_0)$ be the solution of the initial value problem

$$\frac{d\eta}{dt} = f(t, \eta), \quad \eta(t_0) = x_0.$$

In fact that the relation $x(s, \mu, \eta(\mu, \tau_k, x + I_k(x))) = x(s, \lambda, \eta(\lambda, \tau_k, x))$ holds for $\tau_{k-1} < \lambda < \tau_k < \mu < \tau_{k+1}$ and $s > \mu$, we obtain

$$V(\mu, \eta(\mu, \tau_k, x + I_k(x))) \leq V(\lambda, \eta(\lambda, \tau_k, x)).$$

Passing to the limit as $\mu \rightarrow \tau_k^+$ and $\lambda \rightarrow \tau_k^-$ yields

$$V(\tau_k^+, x + I_k(x)) \leq V(\tau_k^-, x) = V(\tau_k, x).$$

This completes the proof of theorem. □

- Remark 3.9.*
- (i) In the case where system (2.1) has no impulses, i.e., all I_k are identically zero, then Theorem 3.8 reduces to Theorem 2.4 in [7].
 - (ii) Note that the convergence of the series $\sum_{k=1}^{\infty} l_k$ is equivalent to the convergence of the infinite product $\prod_{k=1}^{\infty} l_k$.
 - (iii) The questions about known stability properties of the solutions of various classes of differential systems with impulse effect have been studied in [2, 10, 11, 12, 16, 19, 20]. We improved well-known results on the converse Lyapunov theorems on various types of stability of solutions for system (2.1).

We can obtain the following Massera type converse theorem for the uniformly exponential asymptotic stability of impulsive differential equations as a special case of Theorem 3.8.

Corollary 3.10. *Assume that $(f_x(t, 0), \frac{\partial I_k(0)}{\partial x}) \in \mathcal{M}$ is t_∞ -similar to*

$$(f_x(t, x(t, t_0, x_0)), \frac{\partial I_k(x(\tau_k))}{\partial x}) \in \mathcal{M}.$$

Suppose further that the zero solution of system (2.1) is globally h -stable with $h(t) = e^{-\lambda t}$ for a nonnegative constant λ . Then, there exist a positive constant c and a function $V : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that

- (i) $V \in v_0$ and $V(t, x)$ is Lipschitzian in x with Lipschitz constant L for each $t \in \mathbb{R}_+$;
- (ii) $|x| \leq V(t, x) \leq c|x|$, $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^n$;
- (iii) $D_{(2.1)}^+ V(t, x) \leq -\lambda V(t, x)$, $t \neq \tau_k$;
- (iv) $V(\tau_k^+, x + I_k(x)) \leq V(\tau_k, x)$, $k \in \mathbb{N}$, $x \in \mathbb{R}^n$.

4. Examples

In this section we give two examples which illustrate some results from the previous section.

Example 4.1 ([5, Example 2.10]). To illustrate Lemma 3.7, we consider the linear impulsive differential equation

$$(4.1) \quad \begin{cases} x' &= a(t)x, \quad t \neq \tau_k, \\ \Delta x &= a_k x, \quad t = \tau_k, \quad k \in \mathbb{N}, \end{cases}$$

where $a \in PC(\mathbb{R}_+, \mathbb{R})$, $a_k \in \mathbb{R}$, and $\det(1 + a_k) \neq 0, k \in \mathbb{N}$. Suppose that $\int_{t_0}^\infty |a(s)|ds < \infty$ and $\sum_{t_0 \leq \tau_k \leq \infty} |a_k| < \infty$ for each $t_0 \in \mathbb{R}_+$. Then the solution $x(t, t_0, x_0)$ of equation (4.1) given by

$$x(t, t_0, x_0) = \prod_{t_0 \leq \tau_k < t} (a_k + 1) \exp\left(\int_{t_0}^t a(s)ds\right)x_0$$

satisfies

$$\begin{aligned} |x(t, t_0, x_0)| &= \left| \prod_{t_0 \leq \tau_k < t} (a_k + 1) \exp\left(\int_{t_0}^t a(s)ds\right) \right| |x_0| \\ &\leq |x_0| \exp\left(\int_{t_0}^t |a(s)|ds + \sum_{t_0 \leq \tau_k < t} |a_k|\right) \\ &\leq c|x_0|h(t)h(t_0)^{-1}, \quad t \geq t_0, \end{aligned}$$

where $c = 1$ and $h : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a positive bounded left-continuous function given by

$$h(t) = \begin{cases} \exp(\int_0^t |a(s)|ds), & t_0 \leq t \leq \tau_1, \\ \dots, & \\ \exp(\int_0^t |a(s)|ds + \sum_{t_0 \leq \tau_k < t} |a_k|), & \tau_k < t \leq \tau_{k+1}, \quad k \in \mathbb{N}. \end{cases}$$

Hence the zero solution of equation (4.1) is h -stable.

Example 4.2 ([7, 11]). To illustrate Theorem 3.8, we consider the Riccati scalar equation with impulse effect

$$(4.2) \quad \begin{cases} x' = f(t, x) &= \lambda(t)(-x + x^2), \quad t \neq \tau_k, \\ \Delta x = I_k(x) &= d_k x, \quad t = \tau_k, \quad k = 1, 2, \dots, \\ x(t_0^+) &= x_0, \end{cases}$$

where $\lambda : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function on $[\tau_{k-1}, \tau_k]$, $k = 1, 2, \dots$, with points of discontinuity of first kind at $t = \tau_k$, and $\det(d_k + 1) \neq 0, k \in \mathbb{N}$. Assume $\int_{t_0}^\infty \lambda(s)ds < \infty$ for each $t_0 \in \mathbb{R}_+$ and $\sum_{k=1}^\infty |d_k| < \infty$. Then the solution $x(t, t_0, x_0)$ of equation (4.2) with $x_0 \in D \subset \mathbb{R}$ is given by

$$x(t) = \begin{cases} \frac{x_0}{x_0 + (1-x_0) \exp(\int_{t_0}^t \lambda(s)ds)}, & t_0 \leq t \leq \tau_1, \\ \frac{(d_1+1)x_0}{\exp(\int_{t_0}^t \lambda(s)ds) + x_0[(d_1+1) - \exp(\int_{t_0}^{\tau_1} \lambda(s)ds) - d_1 \exp(\int_{\tau_1}^t \lambda(s)ds)]}, & \tau_1 < t \leq \tau_2, \\ \dots, & \\ \frac{\prod_{i=1}^k (d_i+1)x_0}{e^{\int_{t_0}^t \lambda(s)ds} + x_0 g(t,k)}, & \tau_k < t \leq \tau_{k+1}, \quad k = 2, 3, \dots, \end{cases}$$

where

$$g(t, k) = \prod_{i=1}^k (d_i + 1) - e^{\int_{t_0}^t \lambda(s) ds} - d_1 e^{\int_{\tau_1}^t \lambda(s) ds} - \dots - d_k \prod_{i=1}^{k-1} (d_i + 1) e^{\int_{\tau_k}^t \lambda(s) ds}.$$

Then, we obtain the associated impulsive variational systems for equation (4.2) along the solution $x(t, t_0, x_0)$ of equation (4.2) as the following:

$$(4.3) \quad \begin{cases} v' = f_x(t, 0)v & = -\lambda(t)v, \quad t \neq \tau_k, \\ \Delta v = \frac{\partial I_k(0)}{\partial x} v & = d_k v, \quad t = \tau_k, \quad k \in \mathbb{N}, \\ v(t_0 + 0) & = v_0 \end{cases}$$

and

$$(4.4) \quad \begin{cases} z' = f_x(t, x(t, t_0, x_0))z & = \lambda(t)[-1 + 2x(t, t_0, x_0)]z, \quad t \neq \tau_k, \\ \Delta z = \frac{\partial I_k(x(\tau_k))}{\partial x} z & = d_k z, \quad t = \tau_k, \quad k \in \mathbb{N}, \\ z(t_0 + 0) & = z_0. \end{cases}$$

Then we have

$$\begin{aligned} |\Phi(t, t_0, 0) = \frac{\partial x(t, t_0, x_0)}{\partial x_0}| &= \left| \prod_{t_0 \leq \tau_k < t} (d_k + 1) \exp\left(-\int_{t_0}^t \lambda(s) ds\right) \right| \\ &\leq \exp\left(\int_{t_0}^t \lambda(s) ds + \sum_{t_0 \leq \tau_k < t} |d_k|\right) \\ &\leq ch(t)h(t_0)^{-1}, \quad t \geq t_0, \end{aligned}$$

where $h(t) = \exp(-\int_0^t \lambda(s) ds)$ and $c = \exp(\sum_{t_0 \leq \tau_k \leq \infty} |d_k|)$. Thus the zero solution of equation (4.3) is h -stable. Also, there exist $F_0 \in L_1$ with $t \neq \tau_k$ and $F_k \equiv 0 \in l_1, k \in \mathbb{N}$, such that

$$\begin{aligned} S'(t) - f_x(t, 0)S(t) + S(t)f_x(t, x(t, t_0, x_0)) &\equiv F_0 \in L_1, \quad t \neq \tau_k, \\ \Delta S(\tau_k) - d_k S(\tau_k) + S(\tau_k^+)d_k &\equiv F_k \in l_1, \quad t = \tau_k, \quad k \in \mathbb{N}, \end{aligned}$$

where $\Delta S(\tau_k) = S(\tau_k^+) - S(\tau_k)$ and $S(t) = 1$, since

$$\begin{aligned} \int_0^\infty |F_0(s)| ds &= \int_0^\infty |f_x(s, x(s, t_0, x_0)) - f_x(s, 0)| ds \\ &= \int_0^\infty |2\lambda(s)x(s, t_0, x_0)| ds \\ &\leq 2 \left| \prod_{i=1}^k (d_i + 1) \right| |x_0| \int_0^\infty \left| \frac{\lambda(s)}{e^{\int_{t_0}^s \lambda(\tau) d\tau} + x_0 g(s, k)} \right| ds \\ &\leq |x_0| 2 \exp\left(\sum_{k=1}^\infty |d_k|\right) \int_0^\infty \lambda(s) \exp\left(-\int_{t_0}^s \lambda(\tau) d\tau\right) ds < \infty, \end{aligned}$$

where the limit $\lim_{k \rightarrow \infty} x_0 g(t, k)$ exists and $0 < \lim_{k \rightarrow \infty} x_0 g(t, k) \equiv C(x_0)$. Furthermore, we have

$$\Delta S(\tau_k) - d_k S(\tau_k) + S(\tau_k^+) d_k = 0 \equiv F_k \in l_1, \quad t = \tau_k, \quad k \in \mathbb{N}.$$

Thus $(f_x(t, 0), d_k) \in \mathcal{M}$ is t_∞ -similar to $(f_x(t, x(t, t_0, x_0)), d_k) \in \mathcal{M}$ for each $k \in \mathbb{N}$. Therefore, the zero solution of equation (4.4) is also h -stable by Lemma 3.6.

Finally, all assumptions of Theorem 3.8 are satisfied, then there exist a positive constant c and a function $V : \mathbb{R}_+ \times I \rightarrow \mathbb{R}$ with $I \subset \mathbb{R}$ which satisfies conditions (i)-(iv) of Theorem 3.8.

Acknowledgments. The authors are thankful to the anonymous referee for his/her valuable comments to improve this paper.

References

- [1] D. D. Bainov, S. I. Kostadinov, and A. D. Myshkis, *Kinematical similarity and exponential dichotomy of linear abstract impulsive differential equations*, Internat. J. Theoret. Phys. **33** (1994), no. 2, 487–497.
- [2] D. D. Bainov and P. S. Simeonov, *Systems with Impulse Effect: Stability, Theory and Applications*, Ellis Horwood Limited, Chichester, 1989.
- [3] ———, *Impulsive Differential Equations: Asymptotic Properties of the Solutions*, World Scientific Publishing Co., Inc., River Edge, NJ, 1995.
- [4] S. K. Choi and N. Koo, *Variationally stable impulsive differential systems*, Dyn. Syst. **30** (2015), no. 4, 435–449.
- [5] S. K. Choi, N. Koo, and C. Ryu, *h-Stability of linear impulsive differential equations via similarity*, J. Chungcheong Math. Soc. **24** (2011), 393–400.
- [6] ———, *Stability of linear impulsive differential equations via t_∞ -similarity*, J. Chungcheong Math. Soc. **26** (2013), 811–819.
- [7] S. K. Choi, N. Koo, and H. S. Ryu, *h-Stability of differential systems via t_∞ -similarity*, Bull. Korean Math. Soc. **34** (1997), no. 3, 371–383.
- [8] R. Conti, *Sulla t_∞ -similitudine tra matrici e la stabilit  dei sistemi differenzialelineari*, Atti. Acc. Naz. Lincei, Rend. Cl. Fis. Mat. Nat. (8) **19** (1955), 247–250.
- [9] F. M. Dannan and S. Elaydi, *Lipschitz stability of nonlinear systems of differential equations*, J. Math. Anal. Appl. **113** (1986), no. 2, 562–577.
- [10] J. V. Devi, *A variation of the Lyapunov second method to impulsive differential equations*, J. Math. Anal. Appl. **177** (1993), no. 1, 190–200.
- [11] M. Dlala and M. A. Hammami, *Uniform exponential practical stability of impulsive perturbed systems*, J. Dyn. Control Syst. **13** (2007), no. 3, 373–386.
- [12] G. K. Kulev and D. D. Bainov, *Lipschitz stability of impulsive systems of differential equations*, Internat. J. Theoret. Phys. **30** (1991), no. 5, 737–756.
- [13] V. Lakshmikantham, D. D. Bainov, and P. S. Simeonov, *Theory of Impulsive Differential Equations*, World Scientific Publishing Co. Pte. Ltd., Teaneck, NJ, 1989.
- [14] V. Lakshmikantham and S. G. Deo, *Method of Variation of Parameters for Dynamical Systems*, Gordon and Breach Science Publishers, Amsterdam, 1998.
- [15] V. Lakshmikantham and S. Leela, *Differential and Integral Inequalities, Vol I and II*, Academic Press, New York, 1969.
- [16] R. Medina and M. Pinto, *Uniform asymptotic stability of solutions of impulsive differential equations*, Dynam. Systems Appl. **5** (1996), no. 1, 97–107.
- [17] M. Pinto, *Perturbations of asymptotically stable differential systems*, Analysis **4** (1984), no. 1-2, 161–175.

- [18] ———, *Stability of nonlinear differential systems*, Appl. Anal. **43** (1992), no. 1-2, 1–20.
- [19] P. S. Simeonov and D. D. Bainov, *Stability of the solutions of singularly perturbed systems with impulse effect*, J. Math. Anal. Appl. **136** (1988), no. 2, 575–588.
- [20] ———, *Exponential stability of the solutions of singularly perturbed systems with impulse effect*, J. Math. Anal. Appl. **151** (1990), no. 2, 462–487.
- [21] J. Shen and X. Liu, *Global existence results for impulsive differential equations*, J. Math. Anal. Appl. **314** (2006), no. 2, 546–557.
- [22] T. Yoshizawa, *Stability Theory by Liapunov's Second Method*, The Mathematical Society of Japan, Tokyo, 1966.

SUNG KYU CHOI
DEPARTMENT OF MATHEMATICS
CHUNGNAM NATIONAL UNIVERSITY
DAEJEON 305-764, KOREA
E-mail address: sgchoi@cnu.ac.kr

NAMJIP KOO
DEPARTMENT OF MATHEMATICS
CHUNGNAM NATIONAL UNIVERSITY
DAEJEON 305-764, KOREA
E-mail address: njkoo@cnu.ac.kr