

A dust-enshrouded tidal disruption event with a resolved radio jet in a galaxy merger

DOI:

[10.1126/science.aao4669](https://doi.org/10.1126/science.aao4669)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Mattila, S., Perez-Torres, M., Efstathiou, A., Mimica, P., Fraser, M., Kankare, E., Alberdi, A., Aloy, M. A., Heikkila, T., Jonker, P. G., Lundqvist, P., Marti-Vidal, I., Meikle, W. P. S., Romero-Canizales, C., Smartt, S. J., Tsygankov, S., Varenus, E., Alonso-Herrero, A., Bondi, M., ... Östlin, G. (2018). A dust-enshrouded tidal disruption event with a resolved radio jet in a galaxy merger. *Science*, *361*(6401), 482-485. <https://doi.org/10.1126/science.aao4669>

Published in:

Science

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



Supplementary Materials for

A dust-enshrouded tidal disruption event with a resolved radio jet in a galaxy merger

S. Mattila*[†], M. Pérez-Torres*[†], A. Efstathiou, P. Mimica, M. Fraser, E. Kankare,
A. Alberdi, M. Á. Aloy, T. Heikkilä, P. G. Jonker, P. Lundqvist, I. Martí-Vidal,
W. P. S. Meikle, C. Romero-Cañizales, S. J. Smartt, S. Tsygankov, E. Varenius,
A. Alonso-Herrero, M. Bondi, C. Fransson, R. Herrero-Illana, T. Kangas, R. Kotak,
N. Ramírez-Olivencia, P. Väisänen, R. J. Beswick, D. L. Clements, R. Greimel,
J. Harmanen, J. Kotilainen, K. Nandra, T. Reynolds,
S. Ryder, N. A. Walton, K. Wiik, G. Östlin

*These authors contributed equally to this work.

[†]Corresponding author. Email: sepma@utu.fi (S.M.); torres@iaa.es (M.P.-T.)

Published 14 June 2018 on *Science* First Release
DOI: 10.1126/science.aao4669

This PDF file includes:

Materials and Methods
Figs. S1 to S7
Tables S1 to S8
References

Materials and Methods

The host galaxy of the transient Arp 299-B AT1

The merging pair of galaxies Arp 299 (Fig. 1) at a 44.8 Mpc distance (33; corrected to the cosmic microwave background reference frame; $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$) is one of the brightest nearby LIRGs with an IR luminosity $L_{\text{IR}} \sim 7 \times 10^{11} L_{\odot}$ (34; $L_{\odot} = 3.833 \times 10^{26} \text{ W}$). The system consists of two galaxies, Arp 299-A and Arp 299-B, and the merger is thought to have begun around 750 Myr ago (35). Arp 299 has an estimated total star formation rate of $\sim 150\text{-}200 M_{\odot} \text{ yr}^{-1}$ and a total core-collapse SN rate of $\sim 1.5\text{-}2 \text{ SNe yr}^{-1}$ (9). The system has continued to experience star formation triggered over the last ~ 50 Myr (11). Arp 299-B hosts two prominent nuclei, B1 and B2. A strongly obscured AGN was discovered in the B1 nucleus using data in the hard X-ray band collected by the BeppoSAX observatory in 2001 (36).

Optical and infrared observations and data analysis

We monitored Arp 299 in the near-IR *Ks*-band from 2002 Jan. to 2005 Jan. as part of a program to search for obscured SNe in the nuclear regions of nearby starburst galaxies (37-38), using the Isaac Newton Group Red Imaging Device (INGRID; 39) and Long-slit Intermediate Resolution Infrared Spectrograph (LIRIS; 40) instruments on the 4.2-m William Herschel Telescope (WHT). We reduced the images and compared them to a reference image using standard Image Reduction and Analysis Facility (IRAF; 41) routines and the ISIS2.2 image subtraction package (42-43), to produce difference images. This procedure revealed a strong outburst in the B1 nucleus of Arp 299 in the *Ks*-band images obtained on 2005 Jan. 30 (13). The type II SN2005U in Arp 299 was also discovered in the same images (44).

To test whether the transient Arp 299-B AT1 was detectable in the optical, we obtained *r* and *i* band imaging with the Wide Field Camera (WFC) on the 2.5-m Isaac Newton Telescope (INT) in 2005 and 2006. For the photometric calibration of the images we used the *r* and *i* band magnitudes (in the AB system; 45) of six field stars available from the Sloan Digital Sky Survey (SDSS; 46). We compared these images with pre-outburst images obtained with the INT in 2002 using the same filters and instrument setup. The post-outburst images were aligned with the pre-outburst images and image subtraction was carried out using ISIS2.2. However, our *r* and *i*-band observations did not yield an optical detection of Arp 299-B AT1, and we estimate an *i*-band limiting magnitude of ~ 17.5 for a point source at the position of the *K*-band nucleus B1 in the 2005 Mar. 24 epoch.

Arp 299 has been observed on multiple occasions by the *HST*. We used pipeline-reduced and drizzled images from the Hubble Legacy Archive obtained using the *F814W* filter (close to *I*-band) and several different instruments: in 1998 Oct. 25.1 Wide-Field Planetary Camera 2 (WFPC2; 47) with Planetary Camera (PC), in 2000.0 July 11.2 WFPC2, in 2001 Mar. 16.4 WFPC2 with PC, in 2001 May 31.1 WFPC2 with PC, in 2006 Mar. 19.3 Advanced Camera for Surveys (ACS; 48) with Wide-Field Channel (WFC) and 2010 Jun. 24.9 Wide-Field Camera 3 (WFC3; 49) with Ultraviolet-Visible channel (UVIS). In cases where multiple consecutive exposures were taken using the same filter, we combined them into a single deep image. We derived a geometric transformation between each of the earlier images and the 2010 image, using 17 point sources and allowing for rotation, an independent scaling in *x* and *y*, and

a polynomial term. Once all the *F814W* frames were in the same pixel coordinates, we performed difference imaging between the 2010 frame and all other images. As the images were taken with a range of instruments, with different pixel scales and point spread functions (PSFs), we used the HOTPANTS package (50) to convolve the images and match their PSFs before scaling them to a common flux level and subtracting them.

The best subtraction, as judged by the absence of residuals and a clean uniform background across the difference image, was obtained when subtracting the 2006 ACS image from the 2010 WFC3 image, as shown in Fig. S1. In this subtraction, we see two sources close to Arp 299-B1. The first source corresponds to the nucleus, and is coincident (within 7 mas) with the source seen in *K*-band difference imaging. This source is brighter in 2010 than in 2006. The second source is offset by 0.2" to the East of the nucleus, and is brighter in 2006 compared to 2010. We performed aperture photometry on both sources in the difference image using a 0.1" aperture (Fig. S1), and set the background flux to zero. We corrected our measured fluxes to an infinite aperture but did not correct for charge transfer efficiency losses, as these should be small (~ 0.01 mag) given the date of the observations and the high background levels (51-52). We obtain a magnitude of $F814W = 19.76 \pm 0.01$ for the source coincident with the nucleus, and $F814W = 22.07 \pm 0.01$ for the offset source in 2006. The photometric uncertainties correspond solely to the Charge-Coupled Device (CCD) noise although there is likely a larger systematic error due to the difference imaging.

We obtained ground-based near-IR (*J*, *H*, *Ks*-band) imaging follow-up of Arp 299-B AT1 with LIRIS on the WHT, the 3.8-m United Kingdom Infrared Telescope (UKIRT) and UKIRT Fast-Track Imager (UFTI; 53), NOTCam (54) on the 2.6-m Nordic Optical Telescope (NOT), and The Instrument Formerly Known As Mosaic (TIFKAM) on the 2.4-m Hiltner telescope. Our near-IR imaging follow-up thus covers epochs between 2005 Jan. and 2016 Dec. The near-IR images were reduced and the quiescent host contribution subtracted following the procedures detailed above and in (55). A WHT/LIRIS image obtained on 2004 Jun. 5 was used as the reference image in *K*-band and images from National Science Foundation camera (NSFCAM; 56) on the 3.0-m NASA Infrared Telescope Facility (IRTF) obtained on 1998 Mar. 2 were used as the reference images in *J*- and *H*-bands (57).

For the astrometric calibration the LIRIS *Ks*-band image was registered with a Two Micron All-Sky Survey (2MASS; 58) *Ks*-band image with absolute astrometric information available. The centroid coordinates for Arp 299-B AT1 were then measured in the subtracted image at the epoch of discovery, yielding a right ascension (RA) of 11h28m30.97s and declination (Dec) of +58°33'40.9" (J2000.0 equinox) with an estimated uncertainty of 0.5" in each coordinate. For the photometric calibration of the reference images we used the *J*, *H*, and *K* band magnitudes (on the Vega system) of five field stars surrounding Arp 299 available from 2MASS (55) and transformed the calibration to the IRTF images using NOTCam images with a larger field of view. We carried our aperture photometry of Arp 299-B AT1 in the subtracted images using the Starlink package GAIA (59). An aperture radius of 3" was used. The photometric errors are dominated by the systematic uncertainties in the calibration and uncertainties arising from the image subtraction method. The former was estimated as the standard error of the mean of the zero point magnitudes obtained with the different field stars and the latter from the scatter between the consecutive light curve points during a phase of no apparent light curve evolution. We estimate a photometric uncertainty of 4%, including both of these effects. See Table S1 for the log of near-IR observations and the photometric results.

We observed Arp 299 using the Gemini-North Telescope (Gemini-N) for a total of ten epochs (between 2007 Mar. 6 and 2012 Jan. 29; PIs: S. Smartt and S. Ryder) with the Near-Infrared Imager (NIRI) and the ALTAIR laser guide star adaptive optics (AO) system

($0.022''/\text{pix}$, FWHM $\sim 0.1''$). The individual NIRI images were reduced with IRAF using the NIRI package in IRAF and registered and co-added using a bright star as a reference (for details see 55). *HST* Near Infrared Camera and Multi-Object Spectrometer (NICMOS, 60) observations (61) with the NIC2 camera ($0.076''/\text{pix}$, FWHM $\sim 0.1''$) and *F222M* filter on 1997 Nov. 5 provided a reference image with the nucleus B1 at the quiescent level.

We derived a geometric transformation between each of the Gemini-N images and the *HST* image using 9 point-like sources common between the images and a transformation including x and y shifts, a scaling factor and a rotation angle. We made use of this transformation to align the *HST* reference image to each Gemini-N image prior to subtracting between these two using the ISIS2.2 package. The centroid coordinates of the outburst were then measured in the subtracted images. The r.m.s. uncertainties of the geometric transformation and the calculated offsets between the coordinates of the quiescent nucleus and the outburst are listed in Table S2. We found the RA of the outburst in the nucleus B1 to be coincident (within the 1σ uncertainty) with the quiescent nucleus in all the epochs (also see Fig. 1). In Dec, there are offsets of up to 37 mas for the individual epochs between the outburst and the quiescent nucleus, although the offsets are not statistically significant (about 2σ). It is also likely that our method underestimates the astrometric uncertainties given the different instruments used for the pre- and post-outburst observations. Therefore, we conclude that Arp 299-B AT1 is coincident with the quiescent nucleus B1 and we see no evidence for any systematic shift in the nuclear position between the epochs.

Mid-IR follow-up observations of the outburst were serendipitously obtained from the *Spitzer* Space Telescope. Arp 299 was monitored at $4.5\ \mu\text{m}$ and $8.0\ \mu\text{m}$ with the Infrared Array Camera (IRAC, 62) instrument from 2004 May every ~ 6 months over 10 epochs as a part of a search for obscured SNe (PI: C. Lawrence). We continued the mid-IR monitoring of the outburst as a part of the *Spitzer* warm mission at $4.5\ \mu\text{m}$ (PI: S. Mattila) for 14 epochs in 2009-2016. We also included archival $3.6\ \mu\text{m}$ IRAC observations obtained in 2013-2015 (PI: O. Fox). IRAC observations at 3.6 , 4.5 , and $8.0\ \mu\text{m}$ obtained in 2003 (PI: G. Fazio) provided reference images of Arp 299-B1 at the quiescent level. For the log of mid-IR observations see Table S3.

The 4.5 and $8.0\ \mu\text{m}$ IRAC observations comprised four exposures per dithering position, the first with an exposure time of 0.6 sec, and the other three each with an exposure time of 12 sec. Although the nucleus B1 was badly saturated in all 12 sec frames, the 0.6 sec frames were useful for accurate flux measurements. We used the post-basic calibrated data (PBCD) product short exposure mosaics throughout (pipeline versions S18.25, 19.1 and 19.2). In the case of the warm mission $3.6\ \mu\text{m}$ observations only one short exposure had been obtained per epoch and the basic calibrated data (BCD) products were used. We carried out aperture photometry of Arp 299-B AT1 in the subtracted images using GAIA. A circular aperture of radius $3.66''$ was used and the photometry was corrected to an infinite aperture (63). The sky was measured using a clipped mean sky estimator and concentric annuli outside the galaxy. For nucleus B1, which is a very bright source, we assume that the photometric uncertainty is dominated by the *Spitzer* absolute flux calibration. We adopt 3% errors for all the cryogenic observations and post-cryogenic PBCD mosaics processed with a pipeline version earlier than S19.2, 2% errors for the post-cryogenic mosaics processed with the S19.2 pipeline and 10% errors for the $3.6\ \mu\text{m}$ measurements made on the individual BCD frames (63). The photometric results are presented in Table S3 (before subtracting the quiescent nuclear flux) and Fig. S2 (after subtracting the quiescent flux based on the first epoch of $3.6\ \mu\text{m}$ observation and the first three epochs of $4.5\ \mu\text{m}$ and $8.0\ \mu\text{m}$ observations). The epoch when the mid-IR light curves were observed to start rising, 2004 Dec. 21.6 corresponding to the Julian Date (JD) 2453361.1,

was adopted as day 0 (Julian Date is used by astronomers and corresponds to the number of days since year 4713 BC plus the fraction of a day since the preceding noon in Universal Time).

Radio observations and data analysis

We imaged the central few parsecs of Arp 299-B1 using radio interferometry data taken with the European VLBI Network (EVN) (PI: M. Pérez-Torres), the VLBA (PI: M. Pérez-Torres) and global VLBI, which included both the EVN and the VLBA (PI: M. Pérez-Torres) at 1.5, 1.7, 2.3, 5.0 and 8.4 GHz. These observations were part of a VLBI monitoring program aimed at characterizing the population of compact radio objects in the central starburst regions and estimating the (radio) SN rates therein (64, 65). We also used publicly available archival data taken with the Very Large Array (VLA) at multiple frequencies, as well as VLBA data at 2.3 and 8.4 GHz. (See Table S4 for a log of the observations.)

The EVN and global VLBI data were correlated at the EVN MkIV Data Processor at the Joint Institute for VLBI in Europe (JIVE) in The Netherlands, while the VLBA data were correlated at the Array Operations Center in Socorro (USA). We chose time and frequency intervals small enough to guarantee negligible time and bandwidth-smearing effects in the data. We phase-referenced our observations to the VLBI calibrator NVSS J112813+592515. We used the strong source 4C 39.25 as fringe finder and bandpass calibrator and, when this was not possible, we used J1128+5925 for those purposes. The data were later correlated at the different nuclei of Arp 299, which included our target source, the nucleus Arp 299-B1. We performed standard a-priori gain calibration using the measured gains and system temperatures of each antenna. This calibration, as well as the data inspection and flagging, were done within the National Radio Astronomy Observatory (NRAO) Astronomical Image Processing System (*AIPS*; 66) assisted with *ParselTongue* (67), which we used as an interface to *AIPS*. We also corrected for ionosphere effects (of particular relevance at 1.7 and 2.3 GHz) and source-structure effects of the phase-reference source at all three frequencies, following the same procedures described in (64-65).

We complemented our datasets with archival VLA and VLBA + Green Bank Telescope (GBT) data taken at 2.3 and 8.4 GHz, as well as with VLA data at 8.4 and 22.5 GHz. The angular resolution of the VLA is much poorer than that provided by VLBI arrays, so the VLA cannot resolve the compact radio emission. Fortunately the nuclear region of Arp 299-B1 is faint enough at high frequencies that the strong burst in 2005 could be detected with the VLA [(8) and Table S4]. Also, the VLA flux density measurements could contain contributions from surrounding sources, so we have assumed a 30% calibration error for those measurements. The archival NRAO VLBA+GBT observations at 2.3 and 8.4 GHz included a pre-discovery (Jan. 2005) and a post-discovery epoch (Jul. 2005). These data were published in (14). We independently re-calibrated and re-analyzed these data, following the same data reduction procedures described above. We also analyzed archival VLBA+GBT data prior to 2005. However, these data had severe calibration problems, which prevented us from detecting any source in the Arp 299 system, and therefore were discarded.

We carried out all imaging and deconvolution procedures within *AIPS*, using task *IMAGR* with Briggs weighting (68) and a *ROBUST* parameter of 0.5 (Figs. 1 and S4). This scheme was a good compromise to combine a very high angular resolution and a very low off-source r.m.s. value in our images. For the archival VLBA+GBT data, we used instead a *ROBUST* parameter of 0 for comparison purposes with the imaging in (14) (see also Fig. S3). The off-source r.m.s. value varied significantly from epoch to epoch, but was significantly

lower than the calibration uncertainty, which was typically 10%, depending on the array and observing frequency. We list in Table S4 the multi-frequency radio interferometric observations of Arp 299-B AT1, spanning more than 12 years of observations for a total of more than 30 epochs of milliarcsecond angular resolution radio observations, plus lower angular resolution observations taken with the VLA at 8.4 and 22.5 GHz.

X-ray observations and data analysis

In the energy range below ~ 10 keV, Arp 299 has been observed with the *Chandra* and *XMM-Newton* observatories several times during 2002-17 resulting in 5 publicly available datasets including our own observations obtained through DDT requests (PI: P. Nandra and PI: M. Pérez-Torres, respectively; see Table S5). Despite the strong absorption and existence of additional sources of emission in this energy band we can use these observations to constrain the AGN variability on \sim -year time scales. We reanalyzed all *Chandra* and *XMM-Newton* data in a consistent way.

We extracted the X-ray spectrum of Arp 299-B1 from the *Chandra* data for three epochs: 2001, 2005 and 2013 (where the last epoch consists of observation done on consecutive days in 2013). First, all the data were aligned, and all standard corrections were applied using CIAO 4.8 (69) and CALDB 4.7.0 (70). The spectra were extracted using CIAO's `specextract` tool from a circular source aperture at the Arp 299-B1 position with a radius of $1.5''$. The background was measured from an empty region outside the galaxy. The spectra from observations in 2013 were combined. All the extracted spectra were grouped so that each spectral bin had a minimum of one count. The spectra were analyzed using XSPEC 12.9.0i (71), using C-statistics.

Only channels in the energy range 0.3-8.0 keV were included for the purposes of fitting. LIRGs such as Arp 299 are well known X-ray emitters from several different sources (thermal emission of hot gas, X-ray binaries associated with the starburst, AGN). Furthermore, due to the very strong absorption inferred from the spectral analysis (corresponding to a hydrogen column density of $N_{\text{H}} = 3 \times 10^{24} \text{ cm}^{-2}$) direct emission from the AGN is detectable only at energies above 10 keV. We adopted the best fitting model from (72) as initial values, and fitted this model for the combined spectrum from 2013 allowing all parameters to vary freely. All the emission components were absorbed via photoelectric absorption (we used the XSPEC model `phabs(apec+pcfabs(po+gau))`). For the AGN, we adopted the spectral model (72) consisting of a power-law with additional Gaussian line components to represent the ~ 6.4 keV iron emission line seen through a partial covering absorber plus a thermal plasma component. The results of this fit were consistent within the errors with the values obtained in (72). We then took our best-fit values for this observation as starting values for the spectra extracted from the 2001 and 2005 observations. We fitted the same model for both of these spectra separately, but only allowing the power-law photon index and the normalization parameters of all components (thermal plasma, power-law and Gaussian) to vary freely. We also measured, for all three spectra, the flux of the fully absorbed model as well as the unabsorbed flux of the power-law component (both in the 0.3-8.0 keV energy range) and the equivalent width of the iron line (see Table S5). We find no evidence for statistically significant variability in the full model flux, power-law component flux, photon index or equivalent width of the iron line.

XMM-Newton observed Arp 299 twice, in May 2001 and Dec. 2011. However, the poorer angular resolution than *Chandra* did not permit us to isolate photons from the Arp 299-B1 nucleus alone. For these data the source aperture had a radius of $15''$ and therefore collected

almost all photons from the western component of Arp 299. The background region was located outside the galaxy and contained neither point sources nor a diffuse thermal component. The fitting model was taken to be exactly the same as for the *Chandra* data. However, taking into account the very different source apertures, the absolute values of the resulting fluxes obtained with *XMM-Newton* and *Chandra* observatories differ by a factor of a few owing to a contribution from the unresolved high-mass X-ray binaries residing in the central parts of the western component of Arp 299. Nevertheless, we can compare the fluxes derived from the two *XMM-Newton* observations well before and after the discovery of Arp 299-B AT1. We conclude that the soft X-ray flux from the B1 nucleus has remained constant within the uncertainties.

Proper motion of Arp 299-B AT1

Our VLBI observations allowed us to accurately determine the average proper motion of Arp 299-B AT1. We show in Fig. S5 model fits to the proper motion in both the RA and Dec directions, over a time span of more than 12 yr. We used only the highest angular resolution VLBI data (at 5.0 and 8.4 GHz) up to 4000 days for fitting the proper motion of the jet, as the lower frequency data have a much lower angular resolution, especially in the North-South direction. The fitted angular velocities are $\mu_{\text{RA}} \sim -(0.322 \pm 0.032)$ mas yr⁻¹ and $\mu_{\text{Dec}} \sim (0.151 \pm 0.015)$ mas yr⁻¹, which correspond to an average apparent transverse linear velocity of $\beta_{\text{app}} = (0.25 \pm 0.03)$.

The apparent change in the proper motion in the Dec coordinate after 4000 days occurs when observations switched from the EVN/Global VLBI observations to VLBA-only observations, so the change in proper motion beyond ~ 4000 days is likely an artifact caused by the poorer angular resolution of the VLBA synthesized beam. We emphasize that the proper motion estimates found in the paper are not affected by this potential change at late times.

The nature of Arp 299-B AT1

Pre-discovery VLBA+GBT observations of Arp 299-B AT1 showed four compact sources within the B1 nucleus at the frequency of 2.3 GHz, but no counterparts at 8.4 GHz [see (14), their figure 4]. We show in Fig. S3-4 the resulting images of our reanalysis of the 8.4 GHz VLBA+GBT observations in Jan. 2005 and Jul. 2005. The 8.4 GHz observations in Jan. 2005 do not show evidence for radio emission above the noise, while the observations in Jul. 2005 show clear evidence for a new, bright radio source (at RA = 11h28m30.9875529s, Dec = 58°33'40.783601"; J2000.0.0) with a peak luminosity at 8.4 GHz of $\nu L_{\nu} \sim 1.6 \times 10^{38}$ erg s⁻¹.

We monitored the evolution of the flux density and radio morphology of Arp 299-B AT1 using VLBI observations. The high flux density and compactness of the radio emission from Arp 299-B AT1 implies a non-thermal origin. Initially, the source appeared unresolved (Fig. S3, right), and therefore its compactness and high flux density could have been explained by a number of scenarios, either extrinsic to the SMBH (e.g., an extremely energetic SN, or a GRB), or intrinsic (e.g., accretion-induced SMBH variability, such as an AGN flare, or a TDE). Our VLBI monitoring shows that Arp 299-B AT1 developed a jet-like morphology, indicating an apparent expansion speed $\beta_{\text{app}} = 0.25 \pm 0.03$ (Figs. 1 and S4). Long GRBs occur preferentially in low-metallicity environments (e.g. 73) whereas Arp 299 has a roughly solar metallicity (74). While GRBs may exhibit jet-like features, the time-to-peak (a few days to a few tens of days),

8.4 GHz peak radio luminosity and flux density evolution of both short- and long-GRBs (15) are at odds with the observed radio behavior of Arp 299-B AT1. A TDE from an intermediate mass black hole (IMBH) is an unlikely scenario, given the very small number of known IMBH candidates and the fact that the location of the transient coincides with the dynamical mass center of Arp 299-B1 (as traced by the 2.2 μm light). In addition, the IMBH scenario would also require the disruption of a subsolar mass white dwarf (25) which is not consistent with our observations (see below). We therefore conclude that Arp 299-B AT1 is an event related to the SMBH.

The compact 2.3 GHz emission detected with VLBI in January 2005 very close to the VLBI position of Arp 299-B AT1 in July 2005 (Fig. S3), together with the long-term evolving radio morphology, suggests that this 2.3 GHz emission pinpoints the location of the AGN in Arp 299-B1. This persistent 2.3 GHz emission could in principle be explained by two alternative scenarios: (i) the AGN core itself; or (ii) a stationary component of the jet structure very close to the base of the AGN. In the AGN-core scenario, the initial 8.4 GHz compact emission observed in Jul. 2005 would correspond to the emission of a single plasmoid from the AGN core, which is shifted at 8.4 GHz with respect to the 2.3 GHz peak (Fig. S3) because of the well-known core-shift (75), whereby the peaks of emission at different frequencies are shifted due to opacity effects. The observed shift would correspond to a linear offset of (0.35 ± 0.11) pc, which is not unexpected for AGN (76). An alternative scenario would be that of the 2.3 GHz persistent radio emission being due to a stationary component of the pre-existing AGN jet in Arp 299-B1. However, since the AGN torus is seen almost edge-on (11) and the activity of a normal AGN jet in Arp 299-B1 would be along the polar direction, essentially no beamed emission from such a pre-existing jet would be detected. Therefore, the observed behavior of the compact VLBI radio emission appears to be at odds with the 2.3 GHz radio emission being from a stationary component, and we assume hereafter that the 2.3 GHz VLBI peak observed in 2005 corresponds to the core of the AGN in Arp 299-B1. We emphasize that in either case the 2.3 GHz peak would correspond to pre-existing emission of the AGN.

We conclude that Arp 299-B AT1 is the result of a sudden change in the accretion rate onto the SMBH induced by the tidal disruption of a star. Our high-angular resolution VLBI imaging at 8.4 GHz shows (Figs. 1 and S4) that the radio emission at the initial location has faded completely a few years later, while the radio jet has been evolving and expanding over about 10 years. Therefore, the pre-existing low-luminosity AGN, probably being fed at a very low accretion rate, returned to its steady state shortly after the TDE flare while the TDE-launched jet continued expanding into a dense surrounding medium, and therefore decelerated. The average expansion speed of Arp 299-B AT1 requires a substantial deceleration of the radio jet over a 12 year period, which is at odds with a jet launched by a low-luminosity AGN (32; see also Fig. 2B).

The 8.4 GHz VLBI burst in Arp 299-B AT1 and its observed radio jet morphology and evolution are most naturally explained by a TDE. We therefore modeled the data in Table S4 using the same techniques as outlined in (16, 77-78), which we briefly outline here. We first simulated the hydrodynamical evolution using the MRGENESIS code (79-80). The radio emission was then modeled using the SPEV code (77-78). The (radio) synchrotron emission arises from relativistic electrons that follow a power-law in energy, $N(E) \sim E^{-p}$, where p is the index of the injected power-law of relativistic electrons. The energy feeding the relativistic electrons and the magnetic field comes from the post-shock thermal energy: a fraction, ϵ_e , of the post-shock thermal energy goes to the relativistic electrons, and a fraction ϵ_B , to the magnetic field. The values of ϵ_e and ϵ_B are uncertain, but are unlikely to be much larger than ~ 0.1 (e.g. 16). Additionally, the code in (16, 77-78) allows the specification of a fraction ζ_e of

the available electrons to be accelerated at the shock. Here, we adopted $\zeta_e = 1$. Fig. 2A shows the best-fit modeled radio light curves, which are obtained for $p = 3.0$, i.e., a rather steep power-law, and for $\epsilon_e = 0.125$ and $\epsilon_B = 0.10$.

We reproduce the overall multi-frequency radio light behavior with a jet that has a half-opening angle $\theta_j \sim 6^\circ$ (Table S7). The jet has an isotropic equivalent kinetic energy, E_{iso} , of 8×10^{53} erg, corresponding to a beaming-corrected kinetic jet energy (16) $E_{\text{jet}} = E_{\text{iso}} (1 - \cos \theta_j)/2 \sim 2 \times 10^{51}$ erg. This energy is injected at a distance of $r_{\text{inj}} = 10^{16}$ cm from the central engine, and the jet is embedded in a circumnuclear ambient medium of a constant number density of $n_e \sim 3.7 \times 10^4 \text{ cm}^{-3}$ up to a fiducial distance within the AGN torus of $r_f = 6.75 \times 10^{17}$ cm (Fig. S6A). This dense medium causes the jet to rapidly decelerate. While densities orders of magnitude higher can be found in the clouds of the broad line regions (BLR) surrounding the AGNs, the filling factors within these regions are very low ($\sim 10^{-2}$).

Our modeling of the radio jet indicates that while it is initially relativistic ($\Gamma = 10$ and $\beta \approx 0.99$ at day zero, and $\Gamma = 1.8$ and $\beta \approx 0.87$ at ~ 1.1 days), as soon as ~ 16 days (in the observer frame) after the ejection the jet has already decelerated significantly ($\Gamma \approx 1.3$; $\beta \approx 0.64$). At a distance r_f , the jet reaches the edge of the torus and encounters an external medium with a 100 times lower density falling off then with distance as $(r/r_f)^{-2.5}$. The apparent kink seen in the radio light curves at all frequencies (Fig. 2A) happens also when the jet head crosses the outer boundary of the constant density medium, r_f . This distance corresponds to about 270 light days. Since the jet head crosses this boundary at about 1065 days after the ejection (as measured in the galaxy frame), the time at which this is observed in the light curves is roughly around day 795 (Figs. 2A and S6A). The sudden drop in density causes the jet to accelerate from $\beta \sim 0.09$ to $\beta \sim 0.22$ (Fig. S6A), converting a fraction of its internal energy into kinetic energy. The jet therefore reaches quasi-asymptotic values $\Gamma \approx 1.0$ and $\beta \approx 0.22$ by day ~ 800 . We note that the actual model simulations were done for a 1-dimensional jet (with a viewing angle of 0°). However, as demonstrated in (16), the results for a larger viewing angle of θ_{obs} up to $\sim 40^\circ$ are similar.

The model reproduces well both the early and late time radio emission from Arp 299-B AT1. The early rise at high frequencies implies the existence of significant external absorption close to the supermassive black hole of Arp 299-B1. This is due to free-free absorption of dense gas ($n_{e, \text{torus}} \sim 3.7 \times 10^4 \text{ cm}^{-3}$) made of thermal electrons within the torus, with $T_e \sim 10^4$ K. The two earliest observations at 8.4 GHz, together with the late time radio observations, provide a stringent constraint on the distribution of T_e within the torus. The early radio observations ($t < 300$ days) require that the electron temperature in the inner regions of the torus ($r \leq 6 \times 10^{17}$ cm) must be at least $T_e \sim 10^4$ K. On the other hand, T_e must be around or lower than 3000 K close to the outer edge of the torus, providing the larger absorption required by the radio observations. If the temperature at the outer edge of the torus was of $\sim 10^4$ K instead, the parameters of the jet required to explain the first two epochs of 8.4 GHz observations would yield a much lower flux density for the late time ($t \sim 400$ -500 days) radio emission than is actually observed.

Constraints on the viewing angle and intrinsic velocity of the radio jet

Our VLBI observations provide constraints on the viewing angle, θ_{obs} , and the average intrinsic speed, β , of the jet. We can constrain the allowed combinations using standard relations in the twin-relativistic jet model (81), which assumes that the ejecta are double-sided and intrinsically

the same (jet and the counter-jet have equal intrinsic emitting powers and equal velocities, β , in opposite directions), and that the interstellar medium which the jet and the counter-jet impact with is similar. The apparent jet speed, β_{app} , and the ratio between the flux densities of the jet and the counterjet, R , are then related with β and θ_{obs} as follows:

$$\beta_{\text{app}} = \beta \sin \theta_{\text{obs}} / (1 - \beta \cos \theta_{\text{obs}}) \quad (\text{S1})$$

and

$$R = [(1 + \beta \cos \theta_{\text{obs}})/(1 - \beta \cos \theta_{\text{obs}})]^{3-\alpha} \quad (\text{S2})$$

where α is the spectral index, defined as $S_\nu \sim \nu^\alpha$, with S_ν the flux density at frequency ν . Both the apparent speed of the jet, β_{app} , and the ratio of the flux density of the jet to the (undetected) counterjet, R , can thus be used to directly constrain θ_{obs} and β . We show in Fig. S7 the constraints on θ_{obs} and β , considering 3- σ uncertainties for the measured proper motion, and a realistic 5- σ upper limit for the counter-jet of Arp 299-B AT1 at late times (where σ is the off-source r.m.s. noise within the B1 nucleus, and is typically about 30 $\mu\text{Jy/b}$). The measured average value β_{app} and its uncertainties delimit a region of allowed values of θ_{obs} and β (solid lines in Fig. S7). In addition, the non-detection of the counterjet implies a curve corresponding to a constant value of R (dashed curve in Fig. S7).

For the estimate of R , we have accounted for the free-free absorption from both the AGN torus and the external medium (as per Table S8), as well as for relativistic boosting, deboosting and light-travel-time effects. For values of the average intrinsic jet speed, β , larger than about 0.3, free-free absorption from the AGN torus and the external medium can be neglected, and the observed ratio R coincides with the value predicted by Eq. (S2), which assumes no absorption. For smaller values of β , free-free absorption effects quickly become very significant, implying that a counter-jet at a relatively large angle ($\sim 25^\circ$ - 35°) would remain undetected. Note also that there is a minimum value for β . For values of β lower than about 0.10, the deboosting is not high enough to compensate for the free-free absorption and produce a brightness ratio as high as the one observed.

In summary, our VLBI observations and radio modeling yield an average intrinsic jet speed of $\beta = 0.22 \pm 0.02$, and constrain the jet viewing angle to be within the range $\theta_{\text{obs}} \sim 25^\circ$ - 35° . Viewing angles larger than $\sim 50^\circ$ are ruled out by the VLBI observations, which disfavors the radio jet being produced by normal, quiescent AGN accretion activity, where the jet would be expected to be launched nearly perpendicular to our line-of-sight to the AGN in Arp 299-B1, while it is seen nearly edge-on (11).

Origin of the IR emission

Inspection of the IR SED of Arp 299-B AT1 shows that it can be described well by a single blackbody component (see Fig. 3). We used the EMCEE python implementation (82) of the Markov chain Monte Carlo (MCMC) method to estimate the best-fit blackbody parameters with 1 σ confidence intervals for each epoch with 4.5 μm Spitzer photometry. For this we interpolated the ground-based *JHKs* and 3.6 μm Spitzer photometric measurements to the

epochs of the 4.5 μm observations. The blackbody parameters are listed in Table S6, and their evolution is shown in Fig. S2, together with the blackbody luminosity and total radiated energy, obtained by integrating the luminosity over time assuming it changes linearly between the epochs. We note that this luminosity does not include the contribution of any cooler dust, which would radiate at longer wavelengths, hence this yields a lower limit to the total radiated energy of 1.5×10^{52} erg.

We assume that the near- and mid-IR flux was due to thermal emission from dust heated by optical and UV emission from the transient event (e.g., 83). Here, the transient is surrounded by local dust that absorbs and re-radiates most of the UV and optical photons emitted and radiates initially as a $T \sim 1050$ K blackbody. The blackbody radius indicates a size for the IR emitting region of ~ 0.04 pc initially, and later increasing to ~ 0.13 pc. The fact that the IR SED is well described by a blackbody up to 4.5 and 8.0 μm means that the dust shell local to the transient must be optically thick at least up to these wavelengths, corresponding to a visual extinction of $A_V > 20$ mag for a Galactic extinction law (84-85).

Given the high core-collapse SN rate of ~ 0.3 SN yr^{-1} within the circumnuclear region of Arp 299-B1 (8), a SN is the most likely explanation for the source detected close to the nucleus B1 in the Mar. 2006 HST F814W filter observations (see Fig. S1). We emphasize that its contribution to the IR light curves is very small. Indeed, assuming a normal Type II SN (the most common core collapse SN type), we expect a total radiated energy less than a few times 10^{49} erg (e.g., 86) for this event. This is negligible compared to the total radiated energy of Arp 299-B AT1 event, if most of this energy was absorbed by dust and re-radiated in the IR. We do not see any evidence of this transient in our IR light curves of Arp 299-B AT1 (see Fig. S2). Furthermore, the optical HST images show a brightening of the nucleus from 2006 to 2010 corresponding to a new source of F814W = 19.8 (corresponding to 0.03 mJy) which is substantially brighter than extrapolated from our IR blackbody fits. For comparison the B1 nucleus brightened by ~ 0.8 mJy in J , ~ 5 mJy in H and ~ 16 mJy in K -band over the same period of time. Therefore the optical detection is not consistent with emission from hot dust. Instead, the I -band flux could be explained by optical emission of Arp 299-B AT1 scattered from free electrons above/below the broad line region (87) and/or from the polar dust.

For a transient event close to the SMBH in Arp 299-B1 we would expect a large extinction due to the AGN dusty torus (10-11), given the large column density of $\sim 3 \times 10^{24}$ cm^{-2} towards the accreting SMBH. The shape of the IR SED is very close to a single component blackbody in all the epochs, indicating a narrow range of dust temperatures. This is also not expected if there was substantial 'foreground' extinction due to the reddening altering the shape of the SED and the likely contribution of cooler dust. Therefore, absorption and re-radiation by dust in the polar regions of the AGN torus (24,88) appears to be the most plausible scenario for explaining the observed IR SED. In the surroundings of Arp 299-B1 we expect such polar dust clouds to suffer from a relatively low foreground extinction (89) in agreement with our single component blackbody fits to the IR SED of the transient. The presence of dust in the polar regions of at least some AGN, which is most likely concentrated in discrete optically thick clouds, is now a generally well accepted idea supported by mid-IR interferometry (88).

We have fitted the spectral energy distributions of the pre- and post-outburst components of Arp 299-B1 using the CYGNUS (90) radiative transfer code, which models the emission from a starburst and the AGN torus predicted by the standard unified model. The GYGNUS starburst model is described in (91-92) and has been used extensively for modeling the IR emission of both local and high- z starburst galaxies. The model treats a starburst as an ensemble of giant molecular clouds (GMCs) centrally illuminated by recently formed stars, incorporates the stellar population synthesis model (93), and considers the effect of *transiently*

heated grains/PAHs in the radiative transfer. An important feature of the model is that the GMCs are at different evolutionary stages. The key parameters of the model are the initial optical depth of the molecular clouds τ_v , the age of the starburst t_* and the e-folding time of the exponentially decaying star formation history.

The CYGNUS AGN model follows the prescription developed by (94) and can additionally include the effect of polar dust (24, 95). The model self-consistently solves the radiative transfer problem in the AGN torus to determine the temperature distribution of dust grains which have a range of sizes and composition. The model also self-consistently takes into account absorption of emission from the hot inner torus by dust in the cooler outer parts of the torus depending on the inclination. The model treats the AGN disc as a tapered one, i.e., the height of the disc increases with distance from the black hole, but tapers off to a constant height in its outer part. In this work we used a large grid of tapered discs (94). The parameters of the model are the ratio of outer to inner radius r_2/r_1 the equatorial UV optical depth τ_{UV} , the opening angle of the disc θ_0 and the inclination i .

The pre-outburst SED was adapted from (9) including also data points at near-IR wavelengths whereas the post-outburst SED was obtained by combining the former with the near- and mid-IR fluxes of the outburst at 734 days after its first mid-IR detection (corresponding to 54095.4 MJD) which is the last epoch with 8.0 μm photometry available. The spectral resolution of the Spitzer/IRS data was reduced so that they are better matched to the resolution of the radiative transfer models. For fitting the SEDs we used a grid of 6475 AGN torus templates, 360 starburst templates and a single polar dust template for each epoch. The SEDs are fitted with the MCMC SED fitting code SATMC (96) and after post-processing of the fitting data the covering factor of polar dust f_c and its uncertainty are computed. SATMC combines and interpolates these grids to generate the required number of models to reach convergence. The models presented in Fig. 4 (for model parameters see Table S7) demonstrate that both the pre- and post-outburst SEDs can be modeled successfully adopting plausible parameters for the starburst, the AGN dusty torus, and the polar dust (incl. the opening angle, inclination, polar dust covering factor). We further assume that these parameters do not change as a result of the outburst, except for the temperature and luminosity of the polar dust that are left as a free parameter. In the post-outburst case $f_c = 23\text{-}78\%$. This implies that the observed radiated energy from the outburst must be divided by f_c to give the total radiated energy i.e. $\sim 1.9\text{-}6.5 \times 10^{52}$ erg.

Origin of the X-ray emission

The absence of significant variability in the soft X-ray band (0.3-8.0 keV), as given by pre- and post-burst *Chandra* and *XMM-Newton* observations, can be easily explained by the large column density ($\sim 3 \times 10^{24}$ cm²) towards the AGN in the nucleus B1: low-energy X-ray photons are in this case scattered multiple times, removing any evidence of an X-ray flare due to the different light travel delays introduced by the scattering.

Therefore, the only possibility of clearly detecting a sign of TDE emission at X-rays towards the nucleus of B1 would have been hard (> 10 keV) X-ray observations (97). While no hard X-ray observations were made shortly after the burst, the hard X-ray emission decreased by a factor of two between 2001 and 2013 (10). Such X-ray variability is not uncommon in AGNs (98) but could be expected also in the case of a TDE, as the increase in the number of soft (optical) photons locally to the SMBH cools the corona, partially depleting the region from hard X-ray photons.

The mass of the disrupted star

Based on our IR observations and modeling, the total radiated energy of Arp 299-B AT1 is between 1.9×10^{52} erg and 6.5×10^{52} erg. Following (99) we can estimate the maximum radiated energy by disruption of a star with a mass M .

$$E_{\text{rad,max}} = \eta f_{\text{acc}} M c^2 / 2 \sim 10^{52} (\eta/0.1) (f_{\text{acc}}/0.1) (M/M_{\odot}) \text{ erg} \quad (\text{S3})$$

where f_{acc} is the accreted fraction and η is the radiative efficiency. Assuming $f_{\text{acc}} = 0.1$ and $\eta = 0.1$, which are consistent with observations of optical TDEs (99), we find that the disruption of a star in the range of 1.9 to $6.5 M_{\odot}$ can explain the energetics of Arp 299-B AT1. Stars in this mass range can be disrupted by the $\sim 2 \times 10^7 M_{\odot}$ (10) SMBH in Arp 299-B1 (24).

Emitting and absorbing regions surrounding Arp 299-B AT1

In Fig. 4 (right panel), we present a schematic diagram showing the assumed geometry of the emitting and absorbing regions surrounding Arp 299-B AT1. The disruption of a star by the tidal forces of the SMBH in the nucleus Arp 299-B1 launches a relativistic jet that propagates outwards from the SMBH. The jet crosses the AGN torus, which is about 100 times denser than the surrounding circumnuclear medium. Most of the emission from the transient in the direction of the observer is thus absorbed and scattered by the dense gas and dust within the torus. However, a fraction of the photons can escape the torus in the polar direction and are observable in the near- and mid-IR range as thermal re-radiation by the polar dust.

Once the jet breaks out from the high-density region within the AGN torus, it expands freely and can be observed at radio wavelengths. The main source of opacity for the jet is external absorption. Consequently, the emission of the radio jet is detected first at high frequencies (Fig. S3; bottom right panel) and later at lower frequencies. The drop in radio flux density results from the period in which the jet drills its way through the very dense regions of the obscuring torus. The early rise of the 8.4 GHz flux density of Arp 299-B AT1, together with its apparent proper motion as measured with VLBI, imply that the distance travelled by the jet inside the high-density region of the torus is $l_t \sim 6.8 \times 10^{17}$ cm, or ~ 0.22 pc.




Figure S1. HST images of the site of Arp 299-B AT1. HST F814W (close to *I*-band) images of Arp 299-B AT1 taken on 2006 Mar. 19 (A; 2453813.8 JD) and 2010 Jun. 24 (B; 2455372.4 JD). (C) Difference image. The source 0.2'' to the East from the nucleus which appears white in the difference image was present in 2006 but not in 2010, while the center of the nucleus, which was brighter in 2010 appears as a dark source. The dashed circles indicate the 0.1'' apertures used for photometry.




Figure S2. IR observations and derived properties for Arp 299-B AT1. (A) *K*-band (2.2 μm) and Spitzer 3.6, 4.5 and 8.0 μm light curves (after removing the quiescent nuclear flux). The epoch when the mid-IR light curves were observed to start rising (2004 Dec. 21.6 corresponding to 2453361.1 JD) has been adopted as day 0. (B) The evolution of the derived blackbody radius, temperature, luminosity and total energy over time. Our blackbody model fits (see Fig. 3) suggest that an expanding (from about 0.04 pc to 0.13 pc) and cooling (from about 1050 to 750K) source can account for the observed near- and mid-IR behavior. Integrating the blackbody luminosities to the current epoch yields a total radiated energy of about 1.5×10^{52} erg.




Figure S3. The appearance of the transient Arp 299-B AT1 in the nucleus B1 at radio wavelengths. Pre-discovery (2453372.5 JD) VLBI observations of Arp 299-B AT1 at 2.3 (A) and 8.4 GHz (C). Pre-existing radio emission is seen at 2.3 GHz, which we identify with the core of the AGN in Arp 299-B1. Post-discovery (2453568.5 JD) VLBI observations at 2.3 (B) and 8.4 GHz (D), showing the appearance of Arp 299-B AT1 at high radio frequencies (dashed vertical and horizontal lines are centered at the peak position of 8.4 GHz emission). In each map, the contours are drawn at (3,4,5,7,10,15) times the off-source r.m.s. All maps are centered at RA = 11h28m30.9875529s, Dec = 58°33' 40.783601" (J2000.0).




Figure S4. The radio jet associated with Arp 299-B AT1. VLBI observations of Arp 299-B AT1 at 5.0 GHz on 2011 Jun. 1 (**A**; 2455713.5, JD), 2013 Oct. 23 (**B**; 2456588.5 JD), and 2015 Jun. 5 (**C**; 2457178.5 JD) and at 8.4 GHz on 2012 May 27 (**D**; 2456074.5 JD), 2013 Nov. 5 (**E**; 2456601.5 JD), and 2015 Jun. 3 (**F**; 2457176.5 JD). There is evidence for resolved emission at 5.0 GHz in Jun. 2011 (**A**), which is confirmed by the subsequent 5.0 and 8.4 GHz observations (**B** through **F**). Data are displayed in the same way as in Fig. S3.




Fig. S5. Proper motion of Arp 299-B AT1 as measured from VLBI observations. Angular shift in right ascension (RA) (A) and declination (Dec) (B), as function of time. The epoch when the mid-IR light curves were observed to start rising (2004 Dec. 21.6 corresponding to 2453361.1 JD) has been adopted as day 0. The data show that the head of the jet consistently moved to the North-West for the first ~4000 days at an average apparent speed of $(0.25 \pm 0.03) c$.




Figure S6. (A) Model speed of the TDE jet in Arp 299-B AT1 vs. time (solid line). The dotted, dashed and dot-dashed lines highlight relevant times for the evolution of the jet speed: early time (at about 1 day in the observer frame, dotted line), Sedov time (the time at which the rest mass energy of the accumulated material equals the jet E_{iso} , dot-dashed line), and the epochs at which the jet reaches the lowest velocity (close to the outer edge of the AGN torus; thin dashed line in the inset figure) and the acceleration (very shortly after leaving the dense torus and entering the much less dense circumnuclear material; thick dashed line in the inset figure). **(B)** Modeled (beaming-corrected) jet kinetic energy versus speed of the jet (as in Fig. 2), highlighting the same moments as in panel A.




Figure S7. Viewing angle versus average intrinsic jet speed (θ_{obs} , β) parameter space for Arp 299-B AT1. The shaded area below the dashed line corresponds to the allowed range of viewing angles and average intrinsic speeds for the jet, as constrained by our VLBI observations. The solid lines correspond to the $3\text{-}\sigma$ uncertainties for the measured proper motion of the jet. The dashed line corresponds to the $5\text{-}\sigma$ upper limit for the counter-jet of Arp 299-B AT1 at late times. The dash-dotted lines correspond to the $3\text{-}\sigma$ range of values of the intrinsic jet speed found from our radio modeling ($\beta = 0.22 \pm 0.02$). Our VLBI observations and radio modeling yield an average intrinsic jet speed of $\beta \sim (0.2\text{-}0.3)$ corresponding to a viewing angle of $\theta_{\text{obs}} \sim (25^\circ\text{-}35^\circ)$.

Table S1. Log of ground-based near-IR photometry. Col. 1: Julian Date since day 2400000; col. 2: the telescope and instrument used; col. 3: the flux densities given after subtraction of the quiescent nuclear flux. The photometric errors are dominated by the systematic uncertainties and are assumed to be 4%.

Julian Date (2400000+)	Telescope/instrument	Flux density <i>J</i>-band (mJy)	Flux density <i>H</i>-band (mJy)	Flux density <i>K</i>-band (mJy)
50875.0	IRTF	-	-	-
53162.5	WHT/LIRIS	-	-	-
53400.8	WHT/LIRIS	-	-	7.36
53410.1	UKIRT/UFTI	1.22	3.43	8.33
53446.7	WHT/LIRIS	-	-	16.27
53455.6	NOT/NOTCam	-	-	21.72
53500.7	TIFCAM	3.58	-	29.33
53833.7	NOT/NOTCam	-	7.56	24.75
54156.6	NOT/NOTCam	2.59	8.61	26.95
54250.4	NOT/NOTCam	-	-	31.92
54608.4	NOT/NOTCam	-	12.15	39.82
54786.7	NOT/NOTCam	-	14.29	45.76
54905.5	NOT/NOTCam	-	-	43.76
55027.5	NOT/NOTCam	4.20	13.70	46.04
55234.8	NOT/NOTCam	-	-	43.11
55315.5	NOT/NOTCam	4.03	12.97	40.89
55543.7	NOT/NOTCam	3.46	11.07	37.95
55641.4	NOT/NOTCam	-	-	36.66
55697.5	NOT/NOTCam	2.52	8.35	30.25
55787.4	NOT/NOTCam	-	-	25.88
55934.7	NOT/NOTCam	-	6.46	24.30
56054.5	NOT/NOTCam	2.16	4.27	19.05
56221.8	NOT/NOTCam	-	-	19.41
56318.7	NOT/NOTCam	-	4.08	13.79
56408.5	NOT/NOTCam	-	2.96	13.39
56496.4	NOT/NOTCam	-	3.45	12.68
56621.7	NOT/NOTCam	-	-	11.59
56701.6	NOT/NOTCam	-	-	11.85
56818.6	NOT/NOTCam	1.88	3.01	10.91
57028.6	NOT/NOTCam	-	-	9.97
57206.4	NOT/NOTCam	-	-	9.33
57379.6	NOT/NOTCam	-	-	8.07
57502.5	NOT/NOTCam	1.42	2.28	8.45
57561.4	NOT/NOTCam	-	-	9.08

Table S2. Log of high-resolution near-IR imaging observations. Col. 1: Julian Date since day 2400000; col. 2-3: the telescope, instrument and filter used; col. 4-5: the root-mean-square (rms) uncertainty of the derived geometric transformation between each Gemini image and the reference HST image in right ascension (RA) and declination (Dec); cols. 6-7: the offset between the position of Arp 299-B AT1 (as measured in the difference images) and the B1 nucleus (as measured in the reference HST image) in RA and Dec. In RA Arp 299-B AT1 is found to be coincident with the quiescent nucleus in all epochs within 1σ . In Dec we find a maximum offset of 37 milliarcseconds (at about 2σ level).

Julian Date (2400000+)	Telescope/ Instrument	Filter	rms in RA (mas)	rms in Dec (mas)	Offset in RA (mas)	Offset in Dec (mas)
50757.5	HST/NICMOS2	<i>F222M</i>	-	-	-	-
54165.9	Gemini/NIRI	<i>K</i>	17	13	-7	-22
54639.8	Gemini/NIRI	<i>K</i>	17	14	-3	-23
54818.1	Gemini/NIRI	<i>K</i>	18	15	-10	-23
54928.8	Gemini/NIRI	<i>K</i>	19	16	-6	-23
54958.8	Gemini/NIRI	<i>K</i>	19	15	-6	-28
54859.1	Gemini/NIRI	<i>K</i>	17	16	-12	-37
55252.0	Gemini/NIRI	<i>K</i>	18	16	-20	-37
55317.8	Gemini/NIRI	<i>K</i>	19	15	-5	-27
55668.8	Gemini/NIRI	<i>K</i>	18	16	-14	-2
55956.1	Gemini/NIRI	<i>K</i>	14	12	-4	-11

Table S3. Log of Spitzer observations. Col. 1: Julian Date since day 2400000; col. 2-4: the flux densities are listed before removing the quiescent nuclear flux (96.4, 164.3 and 879 mJy at 3.6, 4.5 and 8.0 μm , respectively) and the 1σ uncertainties are listed in parentheses. The epoch when the mid-IR light curves were observed to start rising (2004 Dec. 21.6 corresponding to 2453361.1 JD) has been adopted as day 0 in Figures 2, 3 and S2.

Julian Date (2400000+)	Flux density 3.6μm (mJy)	Flux density 4.5μm (mJy)	Flux density 8.0μm (mJy)
52991.7	96.4(2.9)	165.1(5.0)	875(26)
53152.2	-	162.7(4.9)	882(26)
53328.3	-	165.2(5.0)	881(26)
53361.1	-	169.1(5.1)	886(27)
53497.2	-	232.5(7.0)	923(28)
53698.6	-	265.4(8.0)	961(29)
53740.3	-	272.0(8.2)	975(29)
53854.0	-	285.0(8.6)	990(30)
53889.9	-	290.0(8.7)	1000(30)
54064.6	-	314.7(9.4)	1039(31)
54095.4	-	318.6(9.6)	1047(31)
55183.5	-	419(8)	-
55346.4	-	421(8)	-
55569.5	-	423(8)	-
55732.2	-	408(8)	-
55940.9	-	391(8)	-
56078.1	-	374(8)	-
56302.7	193(19)	-	-
56324.0	-	347(7)	-
56360.3	195(20)	-	-
56450.7	190(19)	-	-
56460.1	-	334.3(7)	-
56505.7	191(19)	-	-
56677.9	-	321(6)	-
56682.7	185(19)	-	-
56737.4	179(18)	-	-
56820.7	181(18)	-	-
56822.1	-	312(6)	-
57052.0	168(17)	-	-
57056.7	-	299(6)	-
57195.4	-	291(6)	-
57433.4	-	282(6)	-
57568.3	-	276(6)	-

Table S4. Log of interferometric radio observations for the source Arp 299-B AT1. Col. 1: Julian Date since day 2400000; col. 2: the observing array (EVN=European VLBI Network, VLBA=Very Long Baseline Interferometry, VLA = Very Large Array, GBT = Green Bank Telescope); col. 3: observing frequency, in GHz; col. 4: Size of the radio interferometric synthesized beam at each epoch and for the given array, in (mas \times mas); col. 5: peak flux density, in $\mu\text{Jy beam}^{-1}$; col. 6: total flux density in μJy ; col. 7: flux density uncertainty, in μJy . The values in cols. 5 and 6 for the VLA array have been obtained after having subtracted a baseline flux density at each frequency, following (17). The quoted flux density uncertainties in col. 7 are 1σ when the source is detected, and include the sum in quadrature of the off-source r.m.s. noise in the image plus the calibration uncertainties. When the source was not detected, col. 7 gives the 3σ upper limit, where 1σ corresponds to the off-source r.m.s. noise of the image. 2004 Dec. 21.6 was adopted as day 0, corresponding to the Julian Date (JD) 2453361.1.

Julian Date (2400000+)	Array	Freq. (GHz)	Beam size (mas \times mas)	Peak flux density ($\mu\text{Jy beam}^{-1}$)	Total flux density (μJy)	Flux density uncertainty (μJy)
53311.5	VLA	8.4	780 \times 730	0	0	390
53372.5	VLBA+GBT	8.4	2.2 \times 1.2	0	0	99
53372.5	VLBA+GBT	2.3	6.6 \times 4.0	0	0	195
53409.5	VLA	22.5	296 \times 102	2704	2704	820
53539.5	VLA	22.5	850 \times 300	11395	11395	3450
53568.5	VLBA+GBT	8.4	1.8 \times 1.2	966	1150	132
53568.5	VLBA+GBT	2.3	6.5 \times 4.8	0	0	210
53840.5	VLA	8.4	780 \times 730	7670	7670	2320
54989.5	EVN	1.7	11.1 \times 8.2	10161	9574	963
54994.5	EVN	5.0	2.7 \times 1.1	5299	7645	767
55344.5	EVN	5.0	3.0 \times 1.0	2124	3967	416
55352.5	EVN	1.7	5.9 \times 2.3	13288	15143	1522
55713.5	EVN	5.0	2.6 \times 0.9	1791	3428	361
55724.5	EVN	1.7	8.7 \times 4.0	8195	8893	892
55855.5	EVN	5.0	3.2 \times 1.7	2931	3815	391
56074.5	EVN	8.4	2.1 \times 1.1	697	1557	171
56090.5	EVN	1.7	10.0 \times 3.8	9037	9547	959
56231.5	EVN	5.0	2.4 \times 0.8	4653	4653	489
56348.5	EVN	5.0	3.2 \times 1.4	2149	3187	337
56358.5	EVN	8.4	1.7 \times 0.8	767	2416	263
56364.5	EVN	1.7	6.4 \times 3.0	6483	9826	985
56438.5	EVN	5.0	2.8 \times 0.9	2150	3078	312
56449.5	EVN	1.7	6.9 \times 3.3	6161	9804	983
56452.5	EVN	8.4	1.4 \times 0.5	680	770	107
56588.5	EVN	5.0	2.6 \times 1.2	1306	3010	307
56601.5	EVN	8.4	1.6 \times 0.7	760	1349	146
56720.5	EVN	5.0	2.7 \times 1.0	1899	2204	224
57085.5	EVN	5.0	3.5 \times 1.4	738	1520	179
57176.5	EVN+VLBA	8.4	0.7 \times 0.4	95	878	153
57178.5	EVN+VLBA	5.0	1.3 \times 0.7	486	2721	316
57185.5	EVN+VLBA	1.7	5.1 \times 2.7	2443	5913	603
57605.5	VLBA	5.0	5.2 \times 1.5	1152	1841	201

57605.5	VLBA	1.5	13.7×4.3	4501	5813	597
57605.5	VLBA	8.4	2.3×0.7	274	562	114
57788.5	VLBA	5.0	5.1×1.5	774	1880	200

Table S5. Log of soft X-ray observations. Col. 1: Julian Date since day 2400000; col. 2: name of the observatory and identifier of the specific observation; col. 3: full observed flux in the 0.3-8 keV energy band, in 10^{-13} erg s^{-1} cm^{-2} ; col. 4: model flux of the power-law spectral component in the 0.3-8 keV energy band corrected for the photoelectric absorption, in 10^{-12} erg s^{-1} cm^{-2} ; col. 4: photon index of the power-law spectral component; col. 4: equivalent width of the iron 6.4-keV fluorescent emission line, keV.

Julian Date (2400000 +)	Instrument, ObsId	Full flux 0.3-8 keV (10^{-13} erg cm^2 s^{-1})	Power-law flux 0.3-8 keV, unabsorbed (10^{-12} erg cm^2 s^{-1})	Power-law photon index	Fe K Equivalent Width (keV)
52035.5	<i>XMM-Newton</i> , ID 0112810101	$7.16^{+0.22}_{-0.13}$	$14.90^{+0.70}_{-0.64}$	$1.87^{+0.08}_{-0.08}$	$-0.64^{+0.27}_{-0.28}$
55910.5	<i>XMM-Newton</i> , ID 0679381101	$8.26^{+0.38}_{-0.25}$	$16.35^{+0.90}_{-0.90}$	$1.82^{+0.09}_{-0.09}$	$-0.31^{+0.18}_{-0.28}$
52103.5	<i>Chandra</i> , ObsId 1641	$1.84^{+0.19}_{-0.18}$	$2.53^{+0.31}_{-0.28}$	$1.10^{+0.26}_{-0.27}$	$-0.31^{+0.23}_{-0.30}$
53415.5	<i>Chandra</i> , ObsId 6227	$1.48^{+0.27}_{-0.26}$	$1.76^{+0.76}_{-0.35}$	$0.99^{+0.56}_{-0.64}$	$-0.59^{+0.48}_{-0.67}$
56363-56364	<i>Chandra</i> , ObsId 15077+15619	$2.13^{+0.10}_{-0.10}$	$3.17^{+0.19}_{-0.17}$	$1.36^{+0.11}_{-0.11}$	$-0.75^{+0.16}_{-0.15}$

Table S6. Derived Infrared properties of Arp 299-B AT1. Col. 1: Julian Date since day 2400000; col. 2-3: blackbody parameters derived for the dates with 4.5 μ m photometry (the 1 σ uncertainties in the last digit are listed in parentheses); col. 4: the blackbody luminosity; col. 5: the total radiated energy obtained by integrating the luminosity over time assuming it changes linearly between the epochs.

Julian Date (2400000+)	Radius (pc)	Temperature (K)	Luminosity (10⁴³ erg s⁻¹)	Energy (10⁵¹ erg)
53497.2	0.0423(6)	1045(7)	1.54	0.091
53698.6	0.0596(7)	983(6)	2.35	0.434
53740.3	0.0646(7)	965(5)	2.53	0.522
53854.0	0.0699(7)	957(5)	2.85	0.786
53889.9	0.0738(8)	943(5)	2.85	0.874
54064.6	0.0883(9)	902(5)	3.56	1.36
54095.4	0.0914(9)	894(4)	3.66	1.45
55183.5	0.108(1)	919(5)	5.90	5.95
55346.4	0.112(1)	904(5)	5.90	6.78
55569.5	0.119(2)	877(5)	5.92	7.92
55732.2	0.126(2)	840(5)	5.57	8.73
55940.9	0.133(2)	803(6)	5.10	9.69
56078.1	0.135(2)	783(5)	4.72	10.3
56324.0	0.127(3)	771(6)	4.05	11.2
56460.1	0.124(3)	770(6)	3.76	11.7
56677.9	0.123(3)	759(7)	3.53	12.3
56822.1	0.123(3)	748(6)	3.11	12.8
57056.7	0.117(3)	752(7)	3.08	13.4
57195.4	0.113(5)	755(7)	2.87	13.7
57433.4	0.112(4)	747(7)	2.70	14.3
57568.3	0.103(3)	766(7)	2.53	14.6

Table S7. Pre- and post-outburst properties of the nucleus B in Arp 299. The luminosities of the AGN torus (L_{AGN}), starburst (L_{SB}), and polar dust (L_{polar}) components, the AGN fraction (f_{AGN}) and the polar dust temperature (T_{polar}) and covering factor (f_C) obtained from the infrared SED fits for Arp 299-B before and after (734 days after the first mid-IR detection corresponding to 2454095.4 JD) the outburst.

	Pre-outburst	Post-outburst
L_{AGN} (10 ¹⁰ L \odot)	3.18 ^{+0.77} _{-0.87}	3.02 ^{+1.31} _{-0.66}
L_{SB} (10 ¹⁰ L \odot)	9.84 ^{+1.15} _{-0.79}	9.91 ^{+0.07} _{-0.16}
L_{polar} (10 ⁹ L \odot)	0.52 ^{+4.68} _{-0.52}	7.78 ^{+4.05} _{-4.07}
f_{AGN}	0.24 ^{+0.04} _{-0.06}	0.23 ^{+0.07} _{-0.04}
T_{polar} (K)	500	900
f_C	0.03 ^{+0.40} _{-0.03}	0.52 ^{+0.26} _{-0.29}

Table S8. Best-fitting parameters of the TDE jet model for the radio light curves of Arp 299-B AT1. The jet (col. 1) has a beaming-corrected kinetic energy, E_{jet} , and a (constant) half-opening angle θ_j , and its bulk motion is described by the Lorentz factor, Γ ; p is the index of the injected power-law of relativistic electrons, ε_e and ε_B are the fractions of post-shock thermal energy going to relativistic electrons and magnetic fields, respectively, and ζ_e is the fraction of electrons being accelerated at the shock; r_{inj} is the distance from the central engine at which E_{jet} is injected; n_e and T_e are the thermal electron density and electron temperature inside the AGN torus (col. 2) and in the external medium (col. 3).

Jet	Torus	External medium
$E_{jet} = 1.8 \times 10^{51}$ erg $\theta_j = 6$ degrees Γ (initial) = 10 $p = 3.0$ $\varepsilon_e = 0.125$ $\varepsilon_B = 0.1$ $\zeta_e = 1$ $r_{inj} = 10^{16}$ cm	$n_e (r < r_f) = 3.7 \times 10^4 \text{ cm}^{-3}$ $T_e (r < r_i) = 10^4$ K $T_e (r_i < r < r_f) = 3 \times 10^3$ K $r_i = 6.0 \times 10^{17}$ cm $r_f = 6.75 \times 10^{17}$ cm	$n_e (r > r_f) = 370 (r/r_f)^{-2.5} \text{ cm}^{-3}$ $T_e (r > r_f) = 3 \times 10^3$ K $r_f = 6.75 \times 10^{17}$ cm

References

1. J. G. Hills, Possible power source of Seyfert galaxies and QSOs. *Nature* **254**, 295–298 (1975). doi:[10.1038/254295a0](https://doi.org/10.1038/254295a0)
2. M. J. Rees, Tidal disruption of stars by black holes of 10^6 - 10^8 solar masses in nearby galaxies. *Nature* **333**, 523–528 (1988). doi:[10.1038/333523a0](https://doi.org/10.1038/333523a0)
3. S. van Velzen, G. E. Anderson, N. C. Stone, M. Fraser, T. Wevers, B. D. Metzger, P. G. Jonker, A. J. van der Horst, T. D. Staley, A. J. Mendez, J. C. A. Miller-Jones, S. T. Hodgkin, H. C. Campbell, R. P. Fender, A radio jet from the optical and x-ray bright stellar tidal disruption flare ASASSN-14li. *Science* **351**, 62–65 (2016). doi:[10.1126/science.aad1182](https://doi.org/10.1126/science.aad1182) [Medline](#)
4. B. A. Zauderer, E. Berger, A. M. Soderberg, A. Loeb, R. Narayan, D. A. Frail, G. R. Petitpas, A. Brunthaler, R. Chornock, J. M. Carpenter, G. G. Pooley, K. Mooley, S. R. Kulkarni, R. Margutti, D. B. Fox, E. Nakar, N. A. Patel, N. H. Volgenau, T. L. Culverhouse, M. F. Bietenholz, M. P. Rupen, W. Max-Moerbeck, A. C. S. Readhead, J. Richards, M. Shepherd, S. Storm, C. L. H. Hull, Birth of a relativistic outflow in the unusual γ -ray transient Swift J164449.3+573451. *Nature* **476**, 425–428 (2011). doi:[10.1038/nature10366](https://doi.org/10.1038/nature10366) [Medline](#)
5. S. Komossa, Tidal disruption of stars by supermassive black holes: Status of observations. *J. High Energy Astrophys.* **7**, 148–157 (2015). doi:[10.1016/j.jheap.2015.04.006](https://doi.org/10.1016/j.jheap.2015.04.006)
6. D. Giannios, B. D. Metzger, Radio transients from stellar tidal disruption by massive black holes. *Mon. Not. R. Astron. Soc.* **416**, 2102–2107 (2011). doi:[10.1111/j.1365-2966.2011.19188.x](https://doi.org/10.1111/j.1365-2966.2011.19188.x)
7. Materials and methods are available as supplementary materials.
8. C. Romero-Cañizales, S. Mattila, A. Alberdi, M. A. Pérez-Torres, E. Kankare, S. D. Ryder, The core-collapse supernova rate in Arp 299 revisited. *Mon. Not. R. Astron. Soc.* **415**, 2688–2698 (2011). doi:[10.1111/j.1365-2966.2011.18886.x](https://doi.org/10.1111/j.1365-2966.2011.18886.x)
9. S. Mattila, T. Dahlen, A. Efstathiou, E. Kankare, J. Melinder, A. Alonso-Herrero, M. Á. Pérez-Torres, S. Ryder, P. Väisänen, G. Östlin, Core-collapse Supernovae Missed by Optical Surveys. *Astrophys. J.* **756**, 111 (2012). doi:[10.1088/0004-637X/756/2/111](https://doi.org/10.1088/0004-637X/756/2/111)
10. A. Ptak, A. Hornschemeier, A. Zezas, B. Lehmer, M. Yukita, D. Wik, V. Antoniou, M. K. Argo, L. Ballo, K. Bechtol, S. Boggs, R. Della Ceca, F. E. Christensen, W. W. Craig, C. J. Hailey, F. A. Harrison, R. Krivonos, T. J. Maccarone, D. Stern, M. Tatum, T. Venters, W. W. Zhang, A Focused, Hard X-Ray Look at Arp 299 with NuSTAR. *Astrophys. J.* **800**, 104 (2015). doi:[10.1088/0004-637X/800/2/104](https://doi.org/10.1088/0004-637X/800/2/104)
11. A. Alonso-Herrero, P. F. Roche, P. Esquej, O. González-Martín, M. Pereira-Santaella, C. Ramos Almeida, N. A. Levenson, C. Packham, A. Asensio Ramos, R. E. Mason, J. M. Rodríguez Espinosa, C. Alvarez, L. Colina, I. Aretxaga, T. Díaz-Santos, E. Perlman, C. M. Telesco, Uncovering the Deeply Embedded Active Galactic Nucleus Activity in the Nuclear Regions of the Interacting Galaxy Arp 299. *Astrophys. J.* **779**, L14 (2013). doi:[10.1088/2041-8205/779/1/L14](https://doi.org/10.1088/2041-8205/779/1/L14)

12. X. Chen, A. Sesana, P. Madau, F. K. Liu, Tidal Stellar Disruptions by Massive Black Hole Pairs. II. Decaying Binaries. *Astrophys. J.* **729**, 13 (2011). [doi:10.1088/0004-637X/729/1/13](https://doi.org/10.1088/0004-637X/729/1/13)
13. S. Mattila *et al.*, Arp 299. *IAU Circ.* **8477**, 2 (2005).
14. J. S. Ulvestad, Radio Emission from Young Supernovae and Supernova Remnants in Arp 299. *Astron. J.* **138**, 1529–1538 (2009). [doi:10.1088/0004-6256/138/5/1529](https://doi.org/10.1088/0004-6256/138/5/1529)
15. P. Chandra, D. A. Frail, A Radio-selected Sample of Gamma-Ray Burst Afterglows. *Astrophys. J.* **746**, 156 (2012). [doi:10.1088/0004-637X/746/2/156](https://doi.org/10.1088/0004-637X/746/2/156)
16. P. Mimica, D. Giannios, B. D. Metzger, M. A. Aloy, The radio afterglow of Swift J1644+57 reveals a powerful jet with fast core and slow sheath. *Mon. Not. R. Astron. Soc.* **450**, 2824–2841 (2015). [doi:10.1093/mnras/stv825](https://doi.org/10.1093/mnras/stv825)
17. S. Rosswog, E. Ramirez-Ruiz, W. R. Hix, Tidal Disruption and Ignition of White Dwarfs by Moderately Massive Black Holes. *Astrophys. J.* **695**, 404–419 (2009). [doi:10.1088/0004-637X/695/1/404](https://doi.org/10.1088/0004-637X/695/1/404)
18. C. Romero-Cañizales, J. L. Prieto, X. Chen, C. S. Kochanek, S. Dong, T. W.-S. Holoien, K. Z. Stanek, F. Liu, The TDE ASASSN-14li and Its Host Resolved at Parsec Scales with the EVN. *Astrophys. J.* **832**, L10 (2016). [doi:10.3847/2041-8205/832/1/L10](https://doi.org/10.3847/2041-8205/832/1/L10)
19. K. D. Alexander, E. Berger, J. Guillochon, B. A. Zauderer, P. K. G. Williams, Discovery of an Outflow from Radio Observations of the Tidal Disruption Event ASASSN-14li. *Astrophys. J.* **819**, L25 (2016). [doi:10.3847/2041-8205/819/2/L25](https://doi.org/10.3847/2041-8205/819/2/L25)
20. E. S. Perlman, E. T. Meyer, Q. D. Wang, Q. Yuan, R. Henriksen, J. Irwin, M. Krause, T. Wiegert, E. J. Murphy, G. Heald, R.-J. Dettmar, Compact Resolved Ejecta in the Nearest Tidal Disruption Event. *Astrophys. J.* **842**, 126 (2017). [doi:10.3847/1538-4357/aa71b1](https://doi.org/10.3847/1538-4357/aa71b1)
21. J. Yang, Z. Paragi, A. J. van der Horst, L. I. Gurvits, R. M. Campbell, D. Giannios, T. An, S. Komossa, No apparent superluminal motion in the first-known jetted tidal disruption event Swift J1644+5734. *Mon. Not. R. Astron. Soc.* **462**, L66 (2016). [doi:10.1093/mnrasl/slw107](https://doi.org/10.1093/mnrasl/slw107)
22. S. van Velzen, A. J. Mendez, J. H. Krolik, V. Gorjian, Emission from Dust Heated by Stellar Tidal Disruption Flares. *Astrophys. J.* **829**, 19 (2016). [doi:10.3847/0004-637X/829/1/19](https://doi.org/10.3847/0004-637X/829/1/19)
23. L. Dou, T. Wang, L. Yan, N. Jiang, C. Yang, R. M. Cutri, A. Mainzer, B. Peng, Discovery of a Mid-infrared Echo from the TDE Candidate in the Nucleus of ULIRG F01004-2237. *Astrophys. J.* **841**, L8 (2017). [doi:10.3847/2041-8213/aa7130](https://doi.org/10.3847/2041-8213/aa7130)
24. A. Efstathiou, A model for the infrared emission of FSC 10214+4724. *Mon. Not. R. Astron. Soc.* **371**, L70–L73 (2006). [doi:10.1111/j.1745-3933.2006.00210.x](https://doi.org/10.1111/j.1745-3933.2006.00210.x)
25. J. Law-Smith, M. MacLeod, J. Guillochon, P. Macias, E. Ramirez-Ruiz, Low-mass White Dwarfs with Hydrogen Envelopes as a Missing Link in the Tidal Disruption Menu. *Astrophys. J.* **841**, L132 (2017). [doi:10.3847/1538-4357/aa6ffb](https://doi.org/10.3847/1538-4357/aa6ffb)
26. S. Dong, B. J. Shappee, J. L. Prieto, S. W. Jha, K. Z. Stanek, T. W.-S. Holoien, C. S. Kochanek, T. A. Thompson, N. Morrell, I. B. Thompson, U. Basu, J. F. Beacom, D. Bersier, J. Brimacombe, J. S. Brown, F. Bufano, P. Chen, E. Conseil, A. B. Danilet, E. Falco, D. Grupe, S. Kiyota, G. Masi, B. Nicholls, F. Olivares E, G. Pignata, G.

- Pojmanski, G. V. Simonian, D. M. Szczygiel, P. R. Woźniak, Astronomy. ASASSN-15lh: A highly super-luminous supernova. *Science* **351**, 257–260 (2016). doi:10.1126/science.aac9613 [Medline](#)
27. G. Leloudas *et al.*, The Superluminous Transient ASASSN-15lh as a Tidal Disruption Event from a Kerr Black Hole. *New Astron.* **1**, 2 (2016).
 28. E. Kankare *et al.*, A population of highly energetic transient events in the centres of active galaxies. *New Astron.* **1**, 865 (2017).
 29. T. Piran, G. Svirski, J. Krolik, R. M. Cheng, H. Shiokawa, Disk Formation Versus Disk Accretion—What Powers Tidal Disruption Events? *Astrophys. J.* **806**, 164 (2015). doi:10.1088/0004-637X/806/2/164
 30. C. Tadhunter, R. Spence, M. Rose, J. Mullaney, P. Crowther, A tidal disruption event in the nearby ultra-luminous infrared galaxy F01004-2237. *Nat. Astron.* **1**, 0061 (2017). doi:10.1038/s41550-017-0061
 31. B. Magnelli, P. Popesso, S. Berta, F. Pozzi, D. Elbaz, D. Lutz, M. Dickinson, B. Altieri, P. Andreani, H. Aussel, M. Béthermin, A. Bongiovanni, J. Cepa, V. Charmandaris, R.-R. Chary, A. Cimatti, E. Daddi, N. M. Förster Schreiber, R. Genzel, C. Gruppioni, M. Harwit, H. S. Hwang, R. J. Ivison, G. Magdis, R. Maiolino, E. Murphy, R. Nordon, M. Pannella, A. Pérez García, A. Poglitsch, D. Rosario, M. Sanchez-Portal, P. Santini, D. Scott, E. Sturm, L. J. Tacconi, I. Valtchanov, The deepest Herschel-PACS far-infrared survey: Number counts and infrared luminosity functions from combined PEP/GOODS-H observations. *Astron. Astrophys.* **553** (Suppl.), A132 (2013). doi:10.1051/0004-6361/201321371
 32. R. Ma, F.-G. Xie, S. Hou, Relationship between the Kinetic Power and Bolometric Luminosity of Jets: Limitation from Black Hole X-Ray Binaries, Active Galactic Nuclei, and Gamma-Ray Bursts. *Astrophys. J.* **780L**, 14 (2017).
 33. Z. Y. Huo, X. Y. Xia, S. J. Xue, S. Mao, Z. G. Deng, Chandra Observations of Ultraluminous Infrared Galaxies: Extended Hot Gaseous Halos in Merging Galaxies. *Astrophys. J.* **611**, 208–219 (2004). doi:10.1086/422164
 34. D. B. Sanders, J. M. Mazzarella, D. C. Kim, J. A. Surace, B. T. Soifer, The IRAS Revised Bright Galaxy Sample. *Astron. J.* **126**, 1607–1664 (2003). doi:10.1086/376841
 35. J. E. Hibbard, M. S. Yun, A 180 Kiloparsec Tidal Tail in the Luminous Infrared Merger ARP 299. *Astron. J.* **118**, 162–185 (1999). doi:10.1086/300928
 36. R. Della Ceca, L. Ballo, F. Tavecchio, L. Maraschi, P. O. Petrucci, L. Bassani, M. Cappi, M. Dadina, A. Franceschini, G. Malaguti, G. G. C. Palumbo, M. Persic, An Enshrouded Active Galactic Nucleus in the Merging Starburst System Arp 299 Revealed by BeppoSAX. *Astrophys. J.* **581**, L9 (2002). doi:10.1086/345925
 37. S. Mattila, W. P. S. Meikle, Supernovae in the nuclear regions of starburst galaxies. *Mon. Not. R. Astron. Soc.* **324**, 325–342 (2001). doi:10.1046/j.1365-8711.2001.04255.x
 38. S. Mattila, W. P. S. Meikle, R. Greimel, Highly extinguished supernovae in the nuclear regions of starburst galaxies. *New Astron. Rev.* **48**, 595–600 (2004). doi:10.1016/j.newar.2003.12.033

39. C. Packham, K. L. Thompson, A. Zurita, J. H. Knapen, I. Smail, R. Greimel, D. F. M. Folha, C. Benn, A. Humphrey, R. Rutten, D. Ciardi, M. Bec, R. Bingham, S. Craig, K. Dee, D. Ives, P. Jolley, P. Moore, M. Pi i Puig, S. Rees, G. Talbot, S. Worswick, INGRID: A near-infrared camera for the William Herschel Telescope. *Mon. Not. R. Astron. Soc.* **345**, 395–405 (2003). [doi:10.1046/j.1365-8711.2003.06982.x](https://doi.org/10.1046/j.1365-8711.2003.06982.x)
40. A. Manchado *et al.*, LIRIS: a long-slit intermediate-resolution infrared spectrograph for the WHT. *Proc. SPIE*, ed. A.M. Fowler, **3354**, 448 (1998).
41. D. Tody, The *IRAF* Data Reduction and Analysis System. *Proc. SPIE*, ed. D.L. Crawford, **627**, 733 (1986).
42. C. Alard, R. H. Lupton, A method for Optical Image Subtraction. *Astrophys. J.* **503**, 325–331 (1998). [doi:10.1086/305984](https://doi.org/10.1086/305984)
43. C. Alard *et al.*, Image subtraction using a space-varying kernel. *Astron. Astrophys.* **144** (Suppl.), 363 (2000).
44. S. Mattila *et al.*, Supernovae 2005Q, 2005R, 2005S, 2005T, 2005U. *IAU Circ.* **8473**, 1 (2005).
45. J. B. Oke, J. E. Gunn, Secondary standard stars for absolute spectrophotometry. *Astrophys. J.* **266**, 713 (1983). [doi:10.1086/160817](https://doi.org/10.1086/160817)
46. B. Abolfathi *et al.*, The Fourteenth Data Release of the Sloan Digital Sky Survey: First Spectroscopic Data from the extended Baryon Oscillation Spectroscopic Survey and from the second phase of the Apache Point Observatory Galactic Evolution Experiment. *arXiv:1707.09322* (2017).
47. J. A. Holtzman, C. J. Burrows, S. Casertano, J. J. Hester, J. T. Trauger, A. M. Watson, G. Worthey, The Photometric Performance and Calibration of WFPC2. *Publ. Astron. Soc. Pac.* **107**, 1065 (1995). [doi:10.1086/133664](https://doi.org/10.1086/133664)
48. M. Sirianni, M. J. Jee, N. Benítez, J. P. Blakeslee, A. R. Martel, G. Meurer, M. Clampin, G. De Marchi, H. C. Ford, R. Gilliland, G. F. Hartig, G. D. Illingworth, J. Mack, W. J. McCann, The Photometric Performance and Calibration of the Hubble Space Telescope Advanced Camera for Surveys. *Publ. Astron. Soc. Pac.* **117**, 1049–1112 (2005). [doi:10.1086/444553](https://doi.org/10.1086/444553)
49. R. A. Kimble *et al.*, Wide Field Camera 3: a powerful new imager for the Hubble Space Telescope. *Proc. SPIE*. Vol. **7010**, 12 (2008).
50. A. Becker. HOTPANTS: High Order Transform of PSF ANd Template Subtraction. *Astrophysics Source Code Library*, ascl:1505.004 (2015).
51. S. Gonzaga *et al.*, ACS Data Handbook, v. 7.2 (STScI, Baltimore, 2014).
52. A. Rajan *et al.*, WFC3 Data Handbook, v. 2.1 (STScI, Baltimore, 2011).
53. P. F. Roche *et al.*, UFTI: the 0.8 - 2.5 μm fast track imager for the UK infrared telescope. *Proc. SPIE*, Vol. **4841**, 901-912 (2003).
54. T. M. Abbott *et al.*, SWIR at the Nordic Optical Telescope: NOTCam. *Proc. SPIE*, Vol. **4008**, 14-719 (2000).

55. E. Kankare *et al.*, The nature of supernovae 2010O and 2010P in Arp 299 - I. Near-infrared and optical evolution. *Mon. Not. R. Astron. Soc.* **415**, 2688–2698 (2014).
56. M. A. Shure, D. W. Toomey, J. T. Rayner, P. M. Onaka, A. J. Denault, NSFCAM: a new infrared array camera for the NASA Infrared Telescope Facility. *Proc. SPIE*, Vol. **2198**, 614-622 (1994).
57. W. Wu, G. C. Clayton, K. D. Gordon, K. A. Misselt, T. L. Smith, D. Calzetti, A New Database of Observed Spectral Energy Distributions of Nearby Starburst Galaxies from the Ultraviolet to the Far-Infrared. *Astrophys. J.* **143** (Suppl), 377–418 (2002).
[doi:10.1086/342834](https://doi.org/10.1086/342834)
58. M. F. Skrutskie, R. M. Cutri, R. Stiening, M. D. Weinberg, S. Schneider, J. M. Carpenter, C. Beichman, R. Capps, T. Chester, J. Elias, J. Huchra, J. Liebert, C. Lonsdale, D. G. Monet, S. Price, P. Seitzer, T. Jarrett, J. D. Kirkpatrick, J. E. Gizis, E. Howard, T. Evans, J. Fowler, L. Fullmer, R. Hurt, R. Light, E. L. Kopan, K. A. Marsh, H. L. McCallon, R. Tam, S. Van Dyk, S. Wheelock, The Two Micron All Sky Survey (2MASS). *Astron. J.* **131**, 1163–1183 (2006). [doi:10.1086/498708](https://doi.org/10.1086/498708)
59. P. W. Draper, N. Gray, D. Berry, M. Taylor, Starlink User Note 214.44. Starlink project (2014). <http://star-www.rl.ac.uk/docs/sun214.htx/sun214.html>
60. R. Thompson, NICMOS - Near-Infrared Camera and Multi-Object Spectrometer. *Space Science Rev.* (ISSN 0038-6308), vol. **61**, 1-2, 69-93 (1992).
61. A. Alonso-Herrero, G. H. Rieke, M. J. Rieke, N. Z. Scoville, Extreme Star Formation in the Interacting Galaxy Arp 299 (IC 694+NGC 3690). *Astrophys. J.* **532**, 845–866 (2000).
[doi:10.1086/308622](https://doi.org/10.1086/308622)
62. G. G. Fazio, J. L. Hora, L. E. Allen, M. L. N. Ashby, P. Barmby, L. K. Deutsch, J.-S. Huang, S. Kleiner, M. Marengo, S. T. Megeath, G. J. Melnick, M. A. Pahre, B. M. Patten, J. Polizotti, H. A. Smith, R. S. Taylor, Z. Wang, S. P. Willner, W. F. Hoffmann, J. L. Pipher, W. J. Forrest, C. W. McMurty, C. R. McCreight, M. E. McKelvey, R. E. McMurray, D. G. Koch, S. H. Moseley, R. G. Arendt, J. E. Mentzell, C. T. Marx, P. Losch, P. Mayman, W. Eichhorn, D. Krebs, M. Jhabvala, D. Y. Gezari, D. J. Fixsen, J. Flores, K. Shakoorzadeh, R. Jungo, C. Hakun, L. Workman, G. Karpati, R. Kichak, R. Whitley, S. Mann, E. V. Tollestrup, P. Eisenhardt, D. Stern, V. Gorjian, B. Bhattacharya, S. Carey, B. O. Nelson, W. J. Glaccum, M. Lacy, P. J. Lowrance, S. Laine, W. T. Reach, J. A. Stauffer, J. A. Surace, G. Wilson, E. L. Wright, A. Hoffman, G. Domingo, M. Cohen, The Infrared Array Camera (IRAC) for the Spitzer Space Telescope. *Astrophys. J.* **154** (Suppl), 10–17 (2004). [doi:10.1086/422843](https://doi.org/10.1086/422843)
63. S. Laine (Ed), IRAC Instrument Handbook v. 2.1.2 (2015).
64. M. A. Pérez-Torres, C. Romero-Cañizales, A. Alberdi, A. Polatidis, A. An extremely prolific supernova factory in the buried nucleus of the starburst galaxy IC 694. *Astron. Astrophys.* **507**, L17–L20 (2009). [doi:10.1051/0004-6361/200912964](https://doi.org/10.1051/0004-6361/200912964)
65. M. Bondi, M. A. Pérez-Torres, R. Herrero-Illana, A. Alberdi, A. The nuclear starburst in Arp 299-A: From the 5.0 GHz VLBI radio light-curves to its core-collapse supernova rate. *Astron. Astrophys.* **539**, A134 (2012). [doi:10.1051/0004-6361/201118446](https://doi.org/10.1051/0004-6361/201118446)

66. E. Greisen, AIPS, the VLA, and the VLBA. in *Information Handling in Astronomy - Historical Vistas*, André Heck, Ed. (Kluwer Academic Publishers, 2003) *Astroph. and Space Science Library*, vol. **285**, p. 109.
67. M. Kettenis, H.-J. van Langevelde, C. Reynolds, B. Cotton, in *Astronomical Data Analysis Software and Systems XV*, Carlos Gabriel et al., Ed. (Astronomical Society of the Pacific, San Francisco, 2006), vol. **351**, p. 497 (2006).
68. D. S. Briggs, High-fidelity deconvolution of moderately resolved sources. Ph.D. Thesis, New Mexico Institute of Mining and Technology, Socorro, New Mexico (1995).
69. A. Fruscione *et al.*, CIAO: Chandra's data analysis system. Proc. SPIE, Vol. **6270**, 62701V (2006).
70. The Calibration Database User's Guide, NASA;
https://heasarc.gsfc.nasa.gov/docs/heasarc/caldb/docs/memos/cal_gen_94_002/cal_gen_94_002.html
71. K. A. Arnaud, XSPEC: The First Ten Years. *Astronomical Data Analysis Software and Systems. Vol 101*, 17 (1996).
72. K. Anastasopoulou, A. Zezas, L. Ballo, R. Della Ceca, A deep Chandra observation of the interacting star-forming galaxy Arp 299. *Mon. Not. R. Astron. Soc.* **460**, 3570–3586 (2016). [doi:10.1093/mnras/stw1200](https://doi.org/10.1093/mnras/stw1200)
73. T. Krühler, D. Malesani, J. P. U. Fynbo, O. E. Hartoog, J. Hjorth, P. Jakobsson, D. A. Perley, A. Rossi, P. Schady, S. Schulze, N. R. Tanvir, S. D. Vergani, K. Wiersema, P. M. J. Afonso, J. Bolmer, Z. Cano, S. Covino, V. D'Elia, A. de Ugarte Postigo, R. Filgas, M. Friis, J. F. Graham, J. Greiner, P. Goldoni, A. Gomboc, F. Hammer, J. Japelj, D. A. Kann, L. Kaper, S. Klose, A. J. Levan, G. Leloudas, B. Milvang-Jensen, A. Nicuesa Guelbenzu, E. Palazzi, E. Pian, S. Piranomonte, R. Sánchez-Ramírez, S. Savaglio, J. Selsing, G. Tagliaferri, P. M. Vreeswijk, D. J. Watson, D. Xu, GRB hosts through cosmic time. VLT/X-Shooter emission-line spectroscopy of 96 γ -ray-burst-selected galaxies at $0.1 < z < 3.6$. *Astron. Astrophys.* **581**, A125 (2015). [doi:10.1051/0004-6361/201425561](https://doi.org/10.1051/0004-6361/201425561)
74. J. P. Anderson, S. M. Habergham, P. A. James, On the multiple supernova population of Arp 299: Constraints on progenitor properties and host galaxy star formation characteristics. *Mon. Not. R. Astron. Soc.* **416**, 567 (2011).
75. J. M. Marcaide, I. I. Shapiro, VLBI study of 1038 + 528 A and B - Discovery of wavelength dependence of peak brightness location. *Astrophys. J.* **276**, 56 (1984). [doi:10.1086/161592](https://doi.org/10.1086/161592)
76. A. B. Pushkarev, T. Hovatta, Y. Y. Kovalev, M. L. Lister, A. P. Lobanov, T. Savolainen, J. A. Zensus, MOJAVE: Monitoring of Jets in Active galactic nuclei with VLBA Experiments. IX. Nuclear opacity. *Astron. Astrophys.* **545**, A113 (2012). [doi:10.1051/0004-6361/201219173](https://doi.org/10.1051/0004-6361/201219173)
77. P. Mimica, M.-A. Aloy, I. Agudo, J. M. Martí, J. L. Gómez, J. A. Miralles, Spectral Evolution of Superluminal Components in Parsec-Scale Jets. *Astrophys. J.* **696**, 1142–1163 (2009). [doi:10.1088/0004-637X/696/2/1142](https://doi.org/10.1088/0004-637X/696/2/1142)

78. P. Mimica, M. A. Aloy, D. Giannios, B. D. Metzger, Numerical simulations of the jetted tidal disruption event Swift J1644+57. *J. Phys. Conf. Ser.* **719**, 012008 (2016). doi:[10.1088/1742-6596/719/1/012008](https://doi.org/10.1088/1742-6596/719/1/012008)
79. M. A. Aloy, J. M. Ibanez, J. M. Martí, E. Müller, GENESIS: A High-Resolution Code for Three-dimensional Relativistic Hydrodynamics. *Astroph. J. S. S.* **122**, 151–166 (1999). doi:[10.1086/313214](https://doi.org/10.1086/313214)
80. P. Mimica, D. Giannios, M. A. Aloy, Deceleration of arbitrarily magnetized GRB ejecta: The complete evolution. *Astron. Astrophys.* **494**, 879–890 (2009). doi:[10.1051/0004-6361:200810756](https://doi.org/10.1051/0004-6361/200810756)
81. C. M. Urry, P. Padovani, Unified Schemes for Radio-Loud Active Galactic Nuclei. *Publ. Astron. Soc. Pac.* **107**, 803 (1995). doi:[10.1086/133630](https://doi.org/10.1086/133630)
82. D. Foreman-Mackey, D. W. Hogg, D. Lang, J. Goodman, emcee: The MCMC Hammer. *Publ. Astron. Soc. Pac.* **125**, 306–312 (2013). doi:[10.1086/670067](https://doi.org/10.1086/670067)
83. S. Kozłowski, C. S. Kochanek, D. Stern, J. L. Prieto, K. Z. Stanek, T. A. Thompson, R. J. Assef, A. J. Drake, D. M. Szczygieł, P. R. Woźniak, P. Nugent, M. L. N. Ashby, E. Beshore, M. J. I. Brown, A. Dey, R. Griffith, F. Harrison, B. T. Jannuzi, S. Larson, K. Madsen, B. Pilecki, G. Pojmański, J. Skowron, W. T. Vestrand, J. A. Wren, SDWFS-MIT-1: A self-obscured luminous supernova at $z \sim 0.2$. *Astrophys. J.* **722**, 1624–1632 (2010). doi:[10.1088/0004-637X/722/2/1624](https://doi.org/10.1088/0004-637X/722/2/1624)
84. M. Xue, B. W. Jiang, J. Gao, J. Liu, S. Wang, A. Li, A Precise Determination of the Mid-infrared Interstellar Extinction Law Based on the APOGEE Spectroscopic Survey. *Astrophys. J.* **224** (Suppl), 23 (2016). doi:[10.3847/0067-0049/224/2/23](https://doi.org/10.3847/0067-0049/224/2/23)
85. J. A. Cardelli, G. C. Clayton, J. S. Mathis, The relationship between infrared, optical, and ultraviolet extinction. *Astrophys. J.* **345**, 245 (1989). doi:[10.1086/167900](https://doi.org/10.1086/167900)
86. V. P. Utrobin, An optimal hydrodynamic model for the normal type IIP supernova 1999em. *Astron. Astrophys.* **461**, 233–251 (2007). doi:[10.1051/0004-6361:20066078](https://doi.org/10.1051/0004-6361:20066078)
87. R. R. J. Antonucci, J. S. Miller, Spectropolarimetry and the nature of NGC 1068. *Astrophys. J.* **297**, 621 (1985). doi:[10.1086/163559](https://doi.org/10.1086/163559)
88. S. F. Hönig, M. Kishimoto, K. R. W. Tristram, M. A. Prieto, P. Gandhi, D. Asmus, R. Antonucci, L. Burtscher, W. J. Duschl, G. Weigelt, Dust in the Polar Region as a Major Contributor to the Infrared Emission of Active Galactic Nuclei. *Astrophys. J.* **771**, 87 (2013). doi:[10.1088/0004-637X/771/2/87](https://doi.org/10.1088/0004-637X/771/2/87)
89. M. García-Marín, L. Colina, S. Arribas, A. Alonso-Herrero, E. Mediavilla, Integral Field Spectroscopy of the Luminous Infrared Galaxy Arp 299 (IC 694 + NGC 3690). *Astrophys. J.* **650**, 850–871 (2006). doi:[10.1086/507411](https://doi.org/10.1086/507411)
90. CYGNUS - CYprus models for Galaxies and their NUClear Spectra; <http://ahpc.euc.ac.cy/index.php/resources/cygnus>
91. A. Efstathiou, R. Siebenmorgen, Starburst and cirrus models for submillimeter galaxies. *Astron. Astrophys.* **502**, 541–548 (2009). doi:[10.1051/0004-6361/200811205](https://doi.org/10.1051/0004-6361/200811205)

92. A. Efstathiou, M. Rowan-Robinson, R. Siebenmorgen, Massive star formation in galaxies: Radiative transfer models of the UV to millimetre emission of starburst galaxies. *Mon. Not. R. Astron. Soc.* **313**, 734–744 (2000). [doi:10.1046/j.1365-8711.2000.03269.x](https://doi.org/10.1046/j.1365-8711.2000.03269.x)
93. G. Bruzual, S. Charlot, Stellar population synthesis at the resolution of 2003. *Mon. Not. R. Astron. Soc.* **344**, 1000–1028 (2003). [doi:10.1046/j.1365-8711.2003.06897.x](https://doi.org/10.1046/j.1365-8711.2003.06897.x)
94. A. Efstathiou, M. Rowan-Robinson, Dusty discs in active galactic nuclei. *Mon. Not. R. Astron. Soc.* **273**, 649–661 (1995). [doi:10.1093/mnras/273.3.649](https://doi.org/10.1093/mnras/273.3.649)
95. A. Efstathiou, J. H. Hough, S. Young, A model for the infrared continuum spectrum of NGC 1068. *Mon. Not. R. Astron. Soc.* **277**, 1134–1144 (1995). [doi:10.1093/mnras/277.3.1134](https://doi.org/10.1093/mnras/277.3.1134)
96. S. P. Johnson, G. W. Wilson, Y. Tang, K. S. Scott, SATMC: Spectral energy distribution analysis through Markov Chains. *Mon. Not. R. Astron. Soc.* **436**, 2535–2549 (2013). [doi:10.1093/mnras/stt1758](https://doi.org/10.1093/mnras/stt1758)
97. K. Hryniewicz, R. Walter, Unbeamed tidal disruption events at hard x-rays. *Astron. Astrophys.* **586**, A9 (2016). [doi:10.1051/0004-6361/201526591](https://doi.org/10.1051/0004-6361/201526591)
98. F. Vagnetti, S. Turriziani, D. Trevese, Ensemble x-ray variability of active galactic nuclei from serendipitous source catalogues. *Astron. Astrophys.* **536**, A84 (2011). [doi:10.1051/0004-6361/201118072](https://doi.org/10.1051/0004-6361/201118072)
99. B. D. Metzger, N. C. Stone, A bright year for tidal disruptions. *Mon. Not. R. Astron. Soc.* **461**, 948–966 (2016). [doi:10.1093/mnras/stw1394](https://doi.org/10.1093/mnras/stw1394)