


RESEARCH

Open Access



# A fractional Fourier integral operator and its extension to classes of function spaces

Shrideh K. Al-Omari<sup>1\*</sup> 

\*Correspondence:  
s.k.q.alomari@fet.edu.jo  
<sup>1</sup>Department of Physics and Basic  
Sciences, Faculty of Engineering  
Technology, Al-Balqa Applied  
University, Amman, Jordan

## Abstract

In this paper, an attempt is being made to investigate a class of fractional Fourier integral operators on classes of function spaces known as ultraBoehmians. We introduce a convolution product and establish a convolution theorem as a product of different functions. By employing the convolution theorem and making use of an appropriate class of approximating identities, we provide necessary axioms and define function spaces where the fractional Fourier integral operator is an isomorphism connecting the different spaces. Further, we provide an inversion formula and obtain various properties of the cited integral in the generalized sense.

**Keywords:** Fractional Fourier integral; Shrinking support; UltraBoehmians

## 1 Introduction and preliminary

Generalized functions, which are continuous linear functionals over a space of smooth functions, are useful in making discontinuous functions more likely smooth and describe physical phenomena as point charges that consequently lead to an extensive use in applied physics and engineering problems. The space of recent generalized functions known as the space of Boehmians is defined by an algebraic construction which is similar to that of field of quotients. When constructions are applied to function spaces and multiplications are interpreted as convolutions, the constructions yield various spaces of generalized functions. Approximating identities or delta sequences with shrinking support to the origin are needful in constructing Boehmian spaces. This, indeed, led among other things to uniqueness theorems which are interpreted as an uncertainty principle for Boehmian spaces. However, as Boehmians have an abstract algebraic definition, they allow different interpretations of those extended operators to be isomorphisms between the different spaces of Boehmians; see [1–7] for further details.

The fractional Fourier integral operator is a generalization of the classical Fourier integral operator into the fractional domains. Although various definitions of the fractional Fourier integral operator were given in literature, the most intuitive way of defining the fractional Fourier operator has been given by generalizing the concept of rotations over an angle  $\pi/2$  in the classical Fourier integral operator. As the classical Fourier integral operator corresponds to a rotation in the time frequency plane, the fractional Fourier integral operator corresponds to a rotation over an angle  $\alpha = a\pi/2$ ,  $a \in \mathbb{R}$ .

Let  $K_\alpha(t, w)$  be defined by

$$K_\alpha(t, w) = \begin{cases} \frac{c(\alpha)}{\sqrt{2\pi}} \exp\{ja(\alpha)((t^2 + w^2) - 2b(\alpha)wt)\}, \alpha \neq n\pi \\ e^{-j\omega t} / \sqrt{2\pi}, \alpha = \frac{\pi}{2} \end{cases}, \tag{1}$$

then the formal and direct definition of the fractional Fourier integral operator with an angle  $\alpha$  of a signal  $\vartheta(t)$  is defined by the integral equation [8] (see also Pathak et al. [9] and Prasad and Kumar [10])

$$F_\alpha(\vartheta)(w) = \int_{-\infty}^{\infty} \vartheta(t)K_\alpha(t, w) dt, \tag{2}$$

where  $a(\alpha) = (\cot \alpha)/2$ ,  $b(\alpha) = \sec \alpha$ , and  $c(\alpha) = \sqrt{1 - \cot \alpha}$ . The inversion formula has been recovered from (2) and (1) as follows:

$$\vartheta(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F_\alpha(\vartheta)(w)K_{-\alpha}(t, w) dw. \tag{3}$$

However, for certain values of  $\alpha$ , the fractional Fourier integral operator satisfies  $F_0(\vartheta)(w) = \vartheta(w)$ ,  $F_\pi(\vartheta)(w) = \vartheta(-w)$ , and  $\frac{\pi}{2}(\vartheta)(w) = F(\vartheta)(w)$  where  $F$  denotes the ordinary Fourier integral of  $\vartheta$ . The fractional Fourier integral operator has various applications in literature. Indeed, it has applications in the solution of ordinary differential equations, quantum optics (Garcia et al. [11]), quantum mechanics (Andez [12]), optical systems (Narayanana and Prabhu [13]), time filtering (Narayanana and Prabhu [13]), and some pattern recognitions (see Zayed and Garcia [14] and Zayed [15, 16]) as well. Various properties of such a remarkable integral, such as linearity, index additivity  $F_{\alpha_1}F_{\alpha_2} = F_{\alpha_1+\alpha_2}$ , commutativity  $F_{\alpha_1}F_{\alpha_2} = F_{\alpha_2}F_{\alpha_1}$ , and associativity  $(F_{\alpha_1}F_{\alpha_2})F_{\alpha_3} = F_{\alpha_1}(F_{\alpha_2}F_{\alpha_3})$  of the fractional Fourier integral operator, have been obtained in the literature; see Barbu [17] for some details.

As part of long-term research, the purpose of this article is to give an extension of the fractional Fourier integral operator into appropriately defined spaces of Boehmians and define an isomorphism between the different spaces of Boehmians. We divide this article into four sections. In Sect. 2 we define convolution products and prove a convolution theorem. In Sect. 3 we establish a class of approximating identities and derive spaces of ultra-Boehmians and, further, we extend the fractional integral operator to the defined spaces. In Sect. 4 we prove linearity and discuss an inversion formula.

### 2 Convolution product and convolution theorem

To establish a convolution theorem for the fractional Fourier integral operator, we introduce a convolution product as follows:

$$(\vartheta *^\alpha \vartheta_0)(t) = \int_{-\infty}^{\infty} \vartheta(z)\vartheta_0(t - z)\psi(t, z) dz, \tag{4}$$

where  $\psi(t, z) = e^{2jz(z-t)a(\alpha)}$ .

The convolution theorem for the fractional Fourier integral operator can be drawn as follows; see [7, 15] for similar discussion.

**Theorem 1** Let  $a(\alpha) = (\cot \alpha)/2$ ,  $b(\alpha) = \sec \alpha$ ,  $c(\alpha) = \sqrt{1 - \cot \alpha}$ , and  $*^\alpha$  be defined by (4), then we have

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) = \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j\frac{\cot \alpha}{2} w^2\right) (F_\alpha \vartheta)(w) (F_\alpha \vartheta_0)(w). \tag{5}$$

*Proof* By aid of (2) and (4), we are led to writing

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) = \frac{c(\alpha)}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \vartheta(z) \vartheta_0(t-z) \exp\left(j\frac{\cot \alpha}{2} t^2\right) \times \exp\left(j\frac{\cot \alpha}{2} w^2\right) \exp(-j \csc \alpha wt) \exp(j(z-t) \cot \alpha) dt dz. \tag{6}$$

The change of variables  $t - z = v$  gives

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) = \frac{c(\alpha)}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \vartheta(z) \vartheta_0(v) \exp\left(j\frac{\cot \alpha}{2} (v+z)^2\right) \times \exp\left(\frac{\cot \alpha}{2} w^2\right) \exp(-j(w+v+z) \csc \alpha) \exp(-jzv \cot \alpha) dz dv. \tag{7}$$

Hence, by rearranging the products in (7), we obtain

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) = \frac{c(\alpha)}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \vartheta(z) \exp(j \cot \alpha z^2) \exp\left(j\frac{\cot \alpha}{2} w^2\right) \times \exp(-jwz \csc \alpha) dz \vartheta_0(v) \exp\left(j\frac{\cot \alpha}{2} v^2\right) \exp(-j \cot \alpha wv) dv. \tag{8}$$

Multiplying (8) by  $\frac{c(\alpha)}{\sqrt{2\pi}} \exp(j\frac{\cot \alpha}{2} w^2)$  yields

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) \frac{c(\alpha)}{\sqrt{2\pi}} \exp\left(j\frac{\cot \alpha}{2} w^2\right) = (F_\alpha \vartheta)(w) (F_\alpha \vartheta_0)(w).$$

That is,

$$F_\alpha(\vartheta *^\alpha \vartheta_0)(w) = \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j\frac{\cot \alpha}{2} w^2\right) (F_\alpha \vartheta)(w) (F_\alpha \vartheta_0)(w).$$

The proof is finished. □

### 3 UltraBoehmians

With the help of the convolution theorem, we introduce a generalized convolution product and generate a class of ultraBoehmians. The spaces under construction generalize those obtained by Prasad and Kumar [10] and Pathak [9]. Due to Prasad and Kumar [9, (5)] and Zemanian [18] as well, an infinitely differentiable complex-valued function  $\vartheta$  is said to be in  $S_{\beta}^{\tilde{\beta}}(R)$  if and only if

$$\gamma_{\beta, \tilde{\beta}}(\vartheta) = \sup_{t \in R} |t^{\beta} D^{\tilde{\beta}} \vartheta(t)| < \infty \tag{9}$$

for every choice of constants  $\beta, \tilde{\beta}$ . Alternatively, (9) can often be written as

$$J_{m,\tilde{\beta}}(\vartheta) = \sup_{t \in \mathbb{R}} |(1 + |t|^2)^{\frac{m}{2}} D^{\tilde{\beta}} \vartheta(t)| < \infty \quad (m, \tilde{\beta} \in \mathbb{N}).$$

The dense subspace of  $S_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$  of smooth ( $C^\infty$ ) functions of compact supports over  $\mathbb{R}$  is denoted by  $D_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$ .

The following theorem is very needful in the sequel; see [9].

**Theorem 2** *The fractional Fourier integral operator is a continuous linear mapping from  $S_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$  into  $S_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$ .*

*Parseval's identity of  $F_\alpha$  is given as*

$$\int_{-\infty}^{\infty} \vartheta(t) \overline{\psi(t)} dt = \int_{-\infty}^{\infty} \hat{\vartheta}_\alpha(\xi) \overline{\hat{\psi}_\alpha(\xi)} d\xi,$$

and hence we have

$$\int_{-\infty}^{\infty} |\vartheta(t)|^2 dt = \int_{-\infty}^{\infty} |\hat{\vartheta}_\alpha(\xi)|^2 d\xi,$$

where  $\hat{\psi}_\alpha, \hat{\vartheta}_\alpha$  are the fractional Fourier integral operators of  $\psi_\alpha$  and  $\vartheta_\alpha$ , and  $\bar{\psi}$  is the conjugate of  $\psi$ , see Pathak et al. [9, Theorem 3.1, p. 243] for some details.

We state without proof the following two theorems as the proofs are a straightforward consequence of (4).

**Theorem 3** *The product  $*^\alpha$  is commutative in  $S_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$ , i.e.,  $\vartheta *^\alpha \vartheta_0 = \vartheta_0 *^\alpha \vartheta$ .*

**Theorem 4** *Let  $\vartheta, \vartheta_0, h \in S_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$ , then  $\vartheta *^\alpha (\vartheta_0 *^\alpha h) = (\vartheta *^\alpha \vartheta_0) *^\alpha h$ .*

Now we introduce a class of approximating identities as follows.

**Definition 5** Let  $\Delta_{\tilde{\beta}}^{\tilde{\beta}}$  be the set of sequences  $(\delta_n)_1^\infty$  of  $D_{\tilde{\beta}}^{\tilde{\beta}}(\mathbb{R})$  where the following formulas hold:

$$\int_{-\infty}^{\infty} \delta_n(t) e^{ja(\alpha)t^2} dt = 1 \quad (\forall n \in \mathbb{N}), \tag{10}$$

$$\int_{-\infty}^{\infty} |\delta_n(t)| dt < M \quad (M \in \mathbb{R}, n \in \mathbb{N}), \tag{11}$$

$$\max_{|t| \geq \delta} |\delta_n(t)| \rightarrow 0 \tag{12}$$

as  $n \rightarrow \infty, \forall \delta > 0$ . We claim that  $\Delta_{\tilde{\beta}}^{\tilde{\beta}}$  with the convolution product  $*^\alpha$  represents a class of approximating identities.

We establish the following theorem.

**Theorem 6** *The class  $(\Delta_{\tilde{\beta}}^{\tilde{\beta}}, *^\alpha)$  forms a class of approximating identities.*

*Proof* We show  $(\delta_n *^\alpha \gamma_n) \in \Delta_{\beta}^{\tilde{\beta}}$  when  $(\delta_n), (\gamma_n) \in \Delta_{\beta}^{\tilde{\beta}}$ . As the proofs of (11) and (12) are straightforward, it suffices to show that (10) holds. By (4) we write

$$\begin{aligned} \int_{-\infty}^{\infty} (\delta_n *^\alpha \gamma_n)(t) e^{ja(\alpha)t^2} dt &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \delta_n(z) \gamma_n(t-z) \psi(t,z) dz e^{ja(\alpha)t^2} dt \\ &= \int_{-\infty}^{\infty} \delta_n(z) \int_{-\infty}^{\infty} \gamma_n(t-z) \psi(t,z) e^{ja(\alpha)t^2} dt dz. \end{aligned}$$

Let  $t = z + w$ , then the preceding equation gives

$$\begin{aligned} \int_{-\infty}^{\infty} \delta_n(z) \int_{-\infty}^{\infty} \gamma_n(w) \psi(z+w, z) e^{ja(\alpha)(z+w)^2} dw dz &= \int_{-\infty}^{\infty} \delta_n(z) e^{ja(\alpha)z^2} dz \\ &\quad \times \int_{-\infty}^{\infty} \gamma_n(w) e^{ja(\alpha)w^2} dw. \end{aligned}$$

Hence the following is obtained by simple computations:

$$\int_{-\infty}^{\infty} (\delta_n *^\alpha \gamma_n)(t) e^{ja(\alpha)t^2} dt = 1.$$

The proof is completely finished. □

**Theorem 7** *If  $\psi \in D_{\beta}^{\tilde{\beta}}(R)$  and  $\vartheta, \vartheta_n, \vartheta_0 \in S_{\beta}^{\tilde{\beta}}(R)$ ,  $\vartheta_n \rightarrow \vartheta$ , as  $n \rightarrow \infty$ , then we have*

- (i)  $(\vartheta + \vartheta_0) *^\alpha \psi = \vartheta *^\alpha \psi + \vartheta_0 *^\alpha \psi$ .
- (ii)  $\vartheta_n *^\alpha \psi \rightarrow \vartheta *^\alpha \psi$  as  $n \rightarrow \infty$ .
- (iii)  $\lambda(\vartheta *^\alpha \psi) = (\lambda\vartheta *^\alpha \psi)$  for some  $\lambda \in C$ .

The proof of this theorem can be obtained by easy computations.

**Theorem 8** *If  $\vartheta \in S_{\beta}^{\tilde{\beta}}(R)$  and  $\vartheta_0 \in D_{\beta}^{\tilde{\beta}}(R)$ , then  $\vartheta *^\alpha \vartheta_0 \in S_{\beta}^{\tilde{\beta}}(R)$ .*

*Proof* Indeed, by (9), we get

$$\gamma_{\alpha,\beta}(\vartheta *^\alpha \vartheta_0) \leq \sup_{t \in R} |t^\alpha D_t^\beta (\vartheta * \vartheta_0)(t)| < \infty, \tag{13}$$

where  $*$  denotes the usual Fourier convolution product. The last inequality follows from the fact that  $\vartheta * \vartheta_0 \in S_{\beta}^{\tilde{\beta}}(R)$  and that  $|\psi| < 1$ , see Zemanian [18].

The proof is finished. □

**Theorem 9** *Let  $(\delta_n) \in \Delta_{\beta}^{\tilde{\beta}}$ ,  $\vartheta \in S_{\beta}^{\tilde{\beta}}(R)$ , then  $\vartheta *^\alpha \delta_n \rightarrow \vartheta$  as  $n \rightarrow \infty$ .*

*Proof* By aid of (10), we get

$$\begin{aligned} |t^\beta D_t^{\tilde{\beta}} (\vartheta *^\alpha \delta_n - \vartheta)(t)| &= \left| t^\beta D_t^{\tilde{\beta}} \left( \int_{-\infty}^{\infty} \vartheta_z(t) - \vartheta(t) \right) \delta_n(z) dz \right| \\ &\leq \int_K M |t^\beta D_t^{\tilde{\beta}} (\vartheta_z(t) - \vartheta(t))| dz \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ , where  $\vartheta(t-z) = \vartheta_z(t)$ ,  $M$  is a constant, and  $K$  is a compact subset of  $R$  satisfying  $|\delta_n| \leq M$  and  $K \supseteq \text{supp } \delta_n, n \in N$ . The proof is finished. □

The space  $\beta_1$  ( $\beta_1 \equiv \beta_1(S_{\beta}^{\tilde{\beta}}(R), (D_{\beta}^{\tilde{\beta}}(R), *^{\alpha}), *^{\alpha}, \Delta_{\beta}^{\tilde{\beta}})$ ) of Boehmians is defined.  $(\vartheta_n, \delta_n)$  and  $(\theta_n, t_n)$  in  $\beta_1$  are equivalent,  $(\vartheta_n, \delta_n) \sim (\theta_m, t_m)$ , if  $\vartheta_n *^{\alpha} t_m = \theta_m *^{\alpha} \delta_n$  ( $\forall m, n \in N$ ). Indeed,  $\sim$  defines an equivalence relation on  $\beta_1$ . The equivalence class in  $\beta_1$  containing  $(\vartheta_n, \delta_n)$  is denoted as  $((\vartheta_n)/(\delta_n))$  and is called Boehmian. An embedding between  $S_{\beta}^{\tilde{\beta}}(R)$  and  $\beta_1$  is expressed as  $x \rightarrow x *^{\alpha} \delta_n / \delta_n$  ( $\forall m, n \in N$ ). If  $((\vartheta_n)/(\delta_n)) \in \beta_1$  and  $z \in \beta_1$ , then

$$((\vartheta_n)/(\delta_n)) *^{\alpha} \varrho = (\vartheta_n *^{\alpha} \varrho) / (\delta_n).$$

To define the space of ultraBoehmians, let  $S_{\alpha}$  and  $D_{\alpha}$  be the fraction spaces of  $F_{\alpha}$  of all members of  $S_{\beta}^{\tilde{\beta}}(R)$  and  $D_{\beta}^{\tilde{\beta}}(R)$ , respectively, and, similarly, let  $\Delta_{\alpha}$  be the fractional set of  $F_{\alpha}$  of all sequences in  $\Delta^{\alpha}$ . Then, we introduce a product on  $S_{\alpha}$  as follows:

$$(F \times G)(w) = \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) F(w)G(w). \tag{14}$$

Then we have the following theorem.

**Theorem 10** *Let  $F, F_n, H, G \in S_{\alpha}, F_n \rightarrow F$  as  $n \rightarrow \infty$  and  $\Psi \in D_{\alpha}$ , then the following identities hold:*

- (i)  $(F + G) \times \Psi = F \times \Psi + G \times \Psi$ .
- (ii)  $F_n \times \Psi \rightarrow F \times \Psi$  as  $F_n \rightarrow F$  as  $n \rightarrow \infty$ .
- (iii)  $F \times G = G \times F$ .
- (iv)  $F \times (G \times H) = (F \times G) \times H$ .
- (v)  $\eta(F \times G) = (\eta F \times G), \eta \in C$ .

*Proof* Proofs of (i) and (ii) are straightforward being similar to the proofs assigned to the space  $\beta(S_{\beta}^{\tilde{\beta}}(R), (D_{\beta}^{\tilde{\beta}}(R), *^{\alpha}), *^{\alpha}, \Delta_{\beta}^{\tilde{\beta}})$ .

Proof of (iii) Let  $\vartheta, \vartheta_0 \in S_{\beta}^{\tilde{\beta}}(R)$  be such that  $F = F_{\alpha} \vartheta$  and  $G = F_{\alpha} \vartheta_0$ , then by (14) we have

$$\begin{aligned} (F \times G)(w) &= \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) F(w)G(w) \\ &= \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) (F_{\alpha} \vartheta)(w)(F_{\alpha} \vartheta_0)(w) \\ &= F_{\alpha}(\vartheta *^{\alpha} \vartheta_0)(w) \in S_{\alpha}. \end{aligned}$$

Since  $\vartheta *^{\alpha} \vartheta_0 = \vartheta_0 *^{\alpha} \vartheta$ , it follows from (14) that

$$(F \times G)(w) = F_{\alpha}(\vartheta_0 *^{\alpha} \vartheta)(w) = (G \times F)(w) \in S_{\alpha}. \tag{15}$$

Proof of (iv) is similar to that of (iii), whereas the proof of (v) is straightforward.

The proof is finished. □

**Theorem 11** *Let  $(\theta_n), (\varphi_n) \in \Delta_{\alpha}$  and  $F \in S_{\alpha}$ , then  $(\theta_n \times \varphi_n) \in \Delta_{\alpha}$  and  $\lim_{n \rightarrow \infty} F \times \theta_n = F$ .*

*Proof* Let  $(\delta_n), (\psi_n) \in \Delta_{\beta}^{\tilde{\beta}}$  be such that  $F_{\alpha} \delta_n = \theta_n$  and  $F_{\alpha} \psi_n = \varphi_n, \forall n \in N$ . Then by (14) we have

$$(\theta_n \times \varphi_n)(w) = \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) \theta_n(w) \varphi_n(w) = F_{\alpha}(\delta_n *^{\alpha} \psi_n)(w).$$

Hence  $(\theta_n \times \varphi_n)$  belongs to  $\Delta_\alpha$  since  $(\delta_n *^\alpha \psi_n)$  belongs to  $\Delta^\alpha$ . The proof of the second part of the theorem can similarly be drawn.

The proof of the theorem is therefore finished. □

The space  $\beta_2$  ( $\beta_2 \equiv \beta_2(S_\alpha, (D_\alpha, \times), \times, \Delta_\alpha)$ ) of ultraBoehmians is obtained. An ultra-Boehman in  $\beta_2$  is written as

$$(F_\alpha \vartheta_n)/(F_\alpha \theta_n),$$

where  $(F_\alpha \vartheta_n) \in S_\alpha$  and  $(F_\alpha \theta_n) \in \Delta_\alpha$ . For similar concepts of addition, convergence, and scalar multiplication in  $\beta_1$  and  $\beta_2$ , we refer to Omari [19, 20].

**Definition 12** Let  $(\delta_n) \in \Delta_\beta^{\tilde{\beta}}$  and  $(\vartheta_n) \in S_\beta^{\tilde{\beta}}(R)$ , then the fractional Fourier operator of a Boehman in  $\beta_1$  can be given as

$$\hat{F}_\alpha((\vartheta_n)/(\delta_n)) = (F_\alpha \vartheta_n)/(F_\alpha \delta_n), \tag{16}$$

which indeed is a member of  $\beta_2$ .

**4 Characteristics and an inversion formula**

To show that  $\hat{F}_\alpha$  is well defined, we assume  $(\vartheta_n)/(\delta_n) = (\theta_n)/(\varepsilon_n) \in \beta_1$ , then the idea of quotients of sequences in  $\beta_1$  suggests to have  $\vartheta_n *^\alpha \varepsilon_m = \theta_m *^\alpha \delta_n$  ( $m, n \in N$ ). Hence, applying  $F_\alpha$  gives

$$F_\alpha(\vartheta_n *^\alpha \varepsilon_m) = F_\alpha(\theta_m *^\alpha \delta_n) \quad (m, n \in N).$$

Therefore, the convolution theorem reads as follows:

$$\frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) (F_\alpha \vartheta_n)(F_\alpha \varepsilon_m) = \frac{\sqrt{2\pi}}{c(\alpha)} \exp\left(-j \frac{\cot \alpha}{2} w^2\right) (F_\alpha \theta_m)(F_\alpha \delta_n).$$

This equivalently can be written as

$$(F_\alpha \vartheta_n) \times (F_\alpha \varepsilon_m) = (F_\alpha \theta_m) \times (F_\alpha \delta_n).$$

Hence, the idea of quotients and equivalent classes of  $\beta_2$  and the preceding equation suggest to have

$$(F_\alpha \vartheta_n)/(F_\alpha \delta_n) = (F_\alpha \theta_m)/(F_\alpha \varepsilon_n) \quad (m, n \in N).$$

Thus, we obtain that

$$\hat{F}_\alpha((\vartheta_n)/(\delta_n)) = \hat{F}_\alpha((\theta_n)/(\varepsilon_n)) \quad (m, n \in N).$$

The proof is completely finished.

**Theorem 13** *The integral operator  $\hat{F}_\alpha : \beta_1 \rightarrow \beta_2$  is linear.*

Proof of this theorem follows from the concept of addition of Boehmian spaces. Hence it has been deleted.

**Theorem 14** *Let  $\varrho \in \beta_1, \varrho = 0$ , then  $\hat{F}_\alpha(\varrho) = 0$ .*

Proof of this theorem is straightforward. Details are, therefore, omitted.

**Theorem 15** *Let  $\varrho_0, \varrho \in \beta_1$ , then we have*

$$\hat{F}_\alpha(\varrho_0 *^\alpha \varrho)(w) = \frac{c(\alpha)}{\sqrt{2\pi}} \exp\left(j \frac{\cot \alpha}{2} w^2\right) \hat{F}_\alpha(\varrho_0) \hat{F}_\alpha(\varrho).$$

*Proof* Let  $\varrho_0 = (\vartheta_n)/(\delta_n), \varrho = (\theta_n)/(\varepsilon_n) \in \beta_1$  be given. Then, by employing  $*^\alpha$ , we get

$$\hat{F}_\alpha(\varrho_0 *^\alpha \varrho) = \hat{F}_\alpha\left(\left((\vartheta_n) *^\alpha (\theta_n)\right) / \left((\delta_n) *^\alpha (\varepsilon_n)\right)\right).$$

Hence, the convolution theorem reveals

$$\hat{F}_\alpha(\varrho_0 *^\alpha \varrho)(w) = \frac{c(\alpha)}{\sqrt{2\pi}} \exp\left(j \frac{\cot \alpha}{2} w^2\right) \hat{F}_\alpha(\varrho_0) \hat{F}_\alpha(\varrho).$$

The proof is completely finished. □

**Definition 16** Let  $\varrho_0 \in \beta_2, \varrho_0 = (F_\alpha \vartheta_n)/(F_\alpha \delta_n)$ , then we introduce the inverse operator of  $\hat{F}_\alpha$  as the mapping  $(\hat{F}_\alpha)^{-1} : \beta_2 \rightarrow \beta_1$  given by

$$(\hat{F}_\alpha)^{-1}(\varrho_0) = (\vartheta_n)/(\delta_n)$$

for each  $(\delta_n) \in \Delta_{\tilde{\beta}}$ .

**Theorem 17** *The inverse operator  $(\hat{F}_\alpha)^{-1} : \beta_2 \rightarrow \beta_1$  is well defined and linear.*

*Proof* Let  $\varrho_0 = \varrho$  in  $\beta_2, \varrho_0 = (F_\alpha \vartheta_n)/(F_\alpha \delta_n), \varrho = (F_\alpha \theta_n)/(F_\alpha \varepsilon_n)$ . Then

$$F_\alpha \vartheta_n \times F_\alpha \varepsilon_m = F_\alpha \theta_m \times F_\alpha \delta_n$$

for some  $(\theta_n), (\vartheta_n)$  in  $S_{\tilde{\beta}}(R)$ . By using the convolution theorem, we obtain

$$\hat{F}_\alpha(\vartheta_n *^\alpha \varepsilon_m) = \hat{F}_\alpha(\theta_m *^\alpha \delta_n) \quad (m, n \in N).$$

Therefore,  $\vartheta_n *^\alpha \varepsilon_m = \theta_m *^\alpha \delta_n \quad (m, n \in N)$ . Hence, we have

$$(\vartheta_n)/(\delta_n) = (\theta_n)/(\varepsilon_n).$$

To show that  $(\hat{F}_\alpha)^{-1}$  is linear, let  $\varrho_0 = (F_\alpha \vartheta_n)/(F_\alpha \delta_n), \varrho = (F_\alpha \theta_n)/(F_\alpha \varepsilon_n)$  be members in  $\beta_2$ , then by addition of  $\beta_2$  and the convolution theorem, we write

$$(\hat{F}_\alpha)^{-1}(\varrho_0 + \varrho) = (\hat{F}_\alpha)^{-1}\left(F_\alpha\left(\left(\vartheta_n\right) *^\alpha \left(\varepsilon_n\right)\right) + \left(\left(\theta_n\right) *^\alpha \left(\delta_n\right)\right)\right) / \left(F_\alpha\left(\delta_n *^\alpha \varepsilon_n\right)\right).$$



Therefore,

$$(\hat{F}_\alpha)^{-1}(\varrho_0 + \varrho) = ((\vartheta_n) *^\alpha (\varepsilon_n) + (\theta_n) *^\alpha (\delta_n)) / ((\delta_n) *^\alpha (\varepsilon_n)).$$

Hence addition in  $\beta_1$  finishes the proof of the theorem.  $\square$

#### Acknowledgements

The authors would like to express many thanks to the anonymous referees for their corrections and comments on this manuscript.

#### Funding

Not applicable.

#### Availability of data and materials

Not applicable.

#### Competing interests

The author declares that there is no conflict of interest regarding the publication of this paper. The author declares that he has no competing interests.

#### Authors' contributions

The author read and proved the final manuscript.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 11 January 2018 Accepted: 8 May 2018 Published online: 22 May 2018

#### References

- Nemzer, D.: A note on the convergence of a series in the space of Boehmians. *Bull. Pure Appl. Math.* **2**, 63–69 (2008)
- Al-Omari, S.K.Q., Kilicman, A.: An estimate of Sumudu transform for Boehmians. *Adv. Differ. Equ.* **2013**, 77 (2013)
- Mikusinski, P.: Tempered Boehmians and ultra distributions. *Proc. Am. Math. Soc.* **123**, 813–817 (1995)
- Nemzer, D.: Quasi-asymptotic behavior of Boehmians. *Novi Sad J. Math.* **46**, 87–102 (2016)
- Nemzer, D.: Boehmians of  $L_p$ -growth. *Integral Transforms Spec. Funct.* **27**, 653–666 (2016)
- Al-Omari, S.K.Q., Kilicman, A.: On generalized Hartley–Hilbert and Fourier–Hilbert transforms. *Adv. Differ. Equ.* **2012**, 232 (2012)
- Almeida, L.B.: Product and convolution theorems for the fractional Fourier transform. *IEEE Signal Process. Lett.* **4**(1), 15–17 (1997)
- Namias, V.: The fractional Fourier transform and its application to quantum mechanics. *J. Inst. Math. Appl.* **25**, 241–265 (1980)
- Pathak, R.S., Prasad, A., Kumar, M.: Fractional Fourier transform of tempered distributions and generalized pseudo-differential operator. *J. Pseud.-Differ. Oper. Appl.* **3**, 239–254 (2012)
- Prasad, A., Kumar, M.: Product of two generalized pseudo-differential operators involving fractional Fourier transform. *J. Pseud.-Differ. Oper. Appl.* **2**, 355–365 (2011)
- García, J., Mas, D., Dorsch, R.G.: Fractional-Fourier-transform calculation through the fast-Fourier-transform algorithm. *Appl. Opt.* **35**(35), 7013–7018 (1996)
- Andez, V.J.: Application of the fractional Fourier transform to image reconstruction in MRI. Thesis, Pontificia Universidad Católica De Chile (2009)
- Narayanan, V.A., Prabhu, K.: The fractional Fourier transform: theory, implementation and error analysis. *Microprocess. Microsyst.* **27**, 511–521 (2003)
- Zayed, A.I., García, A.G.: New sampling formulae for the fractional Fourier transform. *Signal Process.* **77**, 111–114 (1999)
- Zayed, A.I.: A convolution and product theorem for the fractional Fourier transform. *IEEE Signal Process. Lett.* **5**(4), 101–103 (1998)
- Zayed, A.I.: Fractional Fourier transform of generalized function. *Integral Transforms Spec. Funct.* **7**(3–4), 299–312 (1998)
- Barbu, M., Kaminsky, E.J., Trahan, R.E.: Sonar signal enhancement using fractional Fourier transform. In: *Proceed. SPIE*, vol. 5807, pp. 170–177. SPIE, Bellingham (2005)
- Zemanian, A.H.: *Distribution Theory and Transform Analysis: An Introduction to Generalized Functions, with Applications*. McGraw-Hill, New York (1965)
- Al-Omari, S.K.Q.: Hartley transforms on certain space of generalized functions. *Georgian Math. J.* **20**(3), 415–426 (2013)
- Al-Omari, S.K.Q., Kilicman, A.: On the generalized Hartley and Hartley–Hilbert transformations. *Adv. Differ. Equ.* **2013**, 222 (2013). <https://doi.org/10.1186/1687-1847-2013-222>