

 Open access • Journal Article • DOI:10.1111/J.1469-8137.2008.02619.X

## **A functional genomics resource for *Brassica napus*: development of an EMS mutagenized population and discovery of FAE1 point mutations by TILLING.**

— [Source link](#) 

Nian Wang, Yajie Wang, Fang Tian, Graham J.W. King ...+4 more authors

**Institutions:** Huazhong Agricultural University, Rothamsted Research

**Published on:** 01 Dec 2008 - New Phytologist (Wiley/Blackwell (10.1111))

**Topics:** TILLING, Paralogous Gene, Population, Erucic acid and Functional genomics

Related papers:

- [A rich TILLING resource for studying gene function in \*Brassica rapa\*.](#)
- [Discovery of chemically induced mutations in rice by TILLING](#)
- [Spectrum of Chemically Induced Mutations From a Large-Scale Reverse-Genetic Screen in \*Arabidopsis\*](#)
- [TILLING to detect induced mutations in soybean](#)
- [Discovery of induced point mutations in maize genes by TILLING.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/a-functional-genomics-resource-for-brassica-napus-3l1fdzq66>

# A functional genomics resource for *Brassica napus*: development of an EMS mutagenized population and discovery of *FAE1* point mutations by TILLING

Nian Wang<sup>1</sup>, Yajie Wang<sup>1</sup>, Fang Tian<sup>1</sup>, Graham J. King<sup>2</sup>, Chunyu Zhang<sup>1</sup>, Yan Long<sup>1</sup>, Lei Shi<sup>1</sup> and Jinling Meng<sup>1</sup>

<sup>1</sup>National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University, Wuhan, 430070, China; <sup>2</sup>Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK

## Summary

Author for correspondence:

Jinling Meng

Tel: +86 27 87282457

Fax: +86 27 87280016

Email: jmeng@mail.hzau.edu.cn

Received: 15 June 2008

Accepted: 18 July 2008

- Two ethylmethanesulfonate (EMS) mutant populations of the semi-winter rapeseed cv. Ningyou7 were constructed with high mutant load, to provide a TILLING platform for functional genomics in *Brassica napus*, and for introduction of novel allelic variation in rapeseed breeding.
- Forward genetic screening of mutants from the M2 populations resulted in identification of a large number of novel phenotypes. Reverse genetic screening focused on the potentially multi-paralogous gene *FAE1* (fatty acid elongase1), which controls seed erucic acid synthesis in rapeseed. A *B. napus* BAC library was screened, and loci in a reference mapping population (TNDH) were mapped to conclude that there are two paralogous copies of *FAE1*, one on each of the *B. napus* A and C genomes.
- A new procedure is demonstrated to identify novel mutations in situations where two or more very similar paralogous gene copies exist in a genome. The procedure involves TILLING of single plants, using existing SNPs as a positive control, and is able to distinguish novel mutations based on primer pairs designed to amplify both *FAE1* paralogues simultaneously.
- The procedure was applied to 1344 M2 plants, with 19 mutations identified, of which three were functionally compromised with reduced seed erucic acid content.

**Key words:** *Brassica napus*, EMS mutagenesis, *FAE1*, mutant population, point mutations, seed erucic acid content, TILLING.

*New Phytologist* (2008) **180**: 751–765

© National Key Laboratory of Crop Genetic Improvement (2008).

Journal compilation © *New Phytologist* (2008)

doi: 10.1111/j.1469-8137.2008.02619.x

## Introduction

Rapeseed is a staple oilseed crop of enormous economic importance worldwide. The oil derived from crushing harvested seed is a major provider of calorific value to the human food chain, with variations in fatty acid profile including combinations of erucic acid, oleic and linolenic acid that are of industrial value as oils, lubricants, surfactants and high-value plastics (Metzger & Bornscheuer, 2006). There is an increasing need to focus rapeseed crop improvement on optimizing performance characteristics such as fatty acid composition, seed oil content and yield. More generally, crop

genetic improvement needs to expand the available variation present within the existing gene pool in order to make significant advances in key traits. For many years, mutant populations generated with chemical mutagens such as ethylmethanesulfonate (EMS) and N-ethyl-N-nitrosourea (ENU) have been used to induce valuable variations in crop genomes (Konzak *et al.*, 1976). However, whilst precursory construction of an EMS population for *B. napus* has recently been reported (Wu *et al.*, 2007a), the resultant mutant populations have not been large enough to generate sufficient variation, nor has an efficient method for screening such variation been developed. There remains a requirement for

development of large mutant populations, along with suitable high-efficiency screening methods suitable for rapeseed improvement.

Functional characterization of genes is an important component of plant genomic research and is increasingly informing crop improvement. Loss of function mutants are of particular interest since they provide valuable evidence for the role of specific genes in regulatory, developmental, biochemical and metabolic networks. Several general methods such as insertional mutagenesis and RNA interference have been used to obtain reduction-of-function or knockout mutations, and used successfully in model plants such as *Arabidopsis* (*Arabidopsis thaliana*) (Long & Coupland, 1998) and crops such as rice (Waterhouse *et al.*, 1998). However, there are particular challenges in applying these approaches to rapeseed research and crop improvement, primarily because of either reliance on *Agrobacterium* T-DNA vectors for transmission or lack of availability of endogenous tagging systems. Transgenesis is inefficient in *B. napus*, and endogenous tagging is not accepted by regulatory bodies and consumers in most countries. TILLING (targeting-induced local lesions in genomes) is a more recent reverse genetic approach that has successfully been applied to both model plants such as *Arabidopsis* (McCallum *et al.*, 2000) and crops such as rice (Till *et al.*, 2007), maize (Till *et al.*, 2004b) and wheat (Slade *et al.*, 2005). TILLING is based on screening populations of plants that have been treated with traditional chemical mutagens that cause point mutations, followed by discovery in genes of interest using a very sensitive detection method. The main advantage is the ability to accumulate an allelic series of mutants with a range of modified functions, from wild-type to almost loss of function (Slade *et al.*, 2005). The original TILLING method involves seeds that are mutagenized by treatment with EMS, which primarily introduces G/C to A/T transitions. The resulting M1 plants are self-fertilized, and M2 individuals are used to prepare DNA samples for mutational screening, whilst an inventory of their seeds is established for future and downstream research (Henikoff *et al.*, 2004). The mutational screening relies on an enzyme which cleaves singleton DNA mismatches, as described by Oleykowski *et al.* (1998). Since TILLING relies on use of chemical mutagens that induce genome lesions randomly, it is relatively easy to construct a sufficiently large mutant population that has a high probability of containing mutations in all gene loci. Moreover, TILLING is applicable to targeting induction of mutations in genes of interest within the genome of almost any plant species (Till *et al.*, 2006). Species with duplicated genomes are more tolerant of induced mutation as a result of the functional redundancy conferred by paralogous sets of genes (Slade *et al.*, 2005). Since the rapeseed genome is particularly complex, it is likely to be quite tolerant of mutations induced by EMS. These considerations suggest that TILLING is better suited to rapeseed genomic research and trait improvement than other methods reported to date.

Although TILLING is potentially very powerful, there are some difficulties in applying the method to rapeseed. The most challenging issues arise from the need to develop gene locus-specific primer pairs within the complex rapeseed genome. *B. napus* (AACC,  $2n = 4x = 38$ ; c. 1150 Mbp) is an amphidiploid species originating from a spontaneous hybridization of *Brassica rapa* (AA,  $2n = 2x = 20$ ) and *Brassica oleracea* (CC,  $2n = 2x = 18$ ) (U, 1935), where the originating diploids have been retained essentially intact (Li *et al.*, 2005). Moreover, the *Arabidopsis*, *B. rapa* and *B. oleracea* genomes appear to have arisen from a common hexaploid ancestor c. 14.5–20.4 million yr ago (Lysak *et al.*, 2005). Sequential duplication events within ancestral genomes have resulted in regions that are represented as single copies within *Arabidopsis*, being present as three segmental copies within the *B. rapa* and *B. oleracea* genomes. This compounds the complexity of the *B. napus* A and C genomes (Mayer *et al.*, 2001; Park *et al.*, 2005; Yang *et al.*, 2006). From recent comparative genomic studies (Park *et al.*, 2005; Town *et al.*, 2006), it is possible to infer the presence of six paralogous regions in *B. napus*, with random gene loss resulting in, on average, four intact gene copies, with additional genomic rearrangements contributing further pseudogene fragments. Among the intact paralogous genes, some are able to be distinguished based on differences such as sequence lengths and intron composition. In such cases, it is relatively trivial to obtain gene-specific primer pairs based on sequence polymorphisms. Slade *et al.* (2005) have reported successful screening using a TILLING approach for this class of multi-paralogous gene, in both hexaploid and tetraploid wheat. For multi-paralogous genes that differ by only a few nucleotides among different paralogues, it is very difficult to design gene-specific primer pairs to screen for mutations, and also difficult to identify mutations reliably in distinct paralogues.

We were motivated to overcome the difficulties associated with application of TILLING in polyploid species, owing to the potential benefits for functional gene analysis in the closest crop relatives to the model *Arabidopsis*, and specifically for potential application in genetic crop improvement of *B. napus*. We therefore tested our system by focusing on screening using TILLING for mutations in the gene *FAE1* (*fatty acid elongase1*), which had previously been shown by Southern blotting analysis to be present with at least two very similar paralogous copies within the *B. napus* genome (Barret *et al.*, 1998). *FAE1* is the key gene in seed erucic acid biosynthesis in rapeseed (Roscoe *et al.*, 2001), and was originally cloned in *Arabidopsis* by directed transposon tagging with the maize element Activator (*Ac*) (James *et al.*, 1995). The product of the gene is a condensing enzyme that extends the chain length of fatty acids from C18 to C20 and C22 (Lassner *et al.*, 1996). More recently, the restriction enzyme *AvrII* has been used to discriminate between the two genes present in the A and C genomes of the two *B. napus* cultivars Zhongyou 821 and Zhongshuang No. 9, which possess high (HEA) and low seed

**Table 1** Concentrations of aqueous ethylmethanesulfonate (EMS) solution used to treat seeds of *Brassica napus* Ningyou7

	Treatment					
	0.0% <sup>a</sup>	0.4%	0.6%	0.8%	1.0%	1.2%
Concentration (w/v)						
Germination rate (%)	100	95	82.3	81.9	76.7	75.9
Germination time (d) <sup>b</sup>	2	7	12	12	19	19
Seed set (%) <sup>c</sup>	100	45.3	25	8.3	6.7	4.7

<sup>a</sup>Batches of 300 mature Ningyou7 seeds were imbibed at each of five different concentrations of EMS solution, with a control treatment (0%) of distilled water, for 18 h at 20°C.

<sup>b</sup>Germination time per treatment was recorded as the number of days from observation of the first germinated seed to the day when the last seed germinated.

<sup>c</sup>Seed set is expressed as the percentage of M1 plants from which viable seeds were set.

erucic acid (LEA) content respectively. The *B. napus* *FAE1* gene copies are intronless, with 98% sequence similarity between the paralogues (Wu *et al.*, 2007b). A reference doubled haploid population (TNDH) developed by our laboratory from Ningyou7 (a Chinese semi-winter cultivar) and Tapidor (a European winter cultivar) was used to detect functional QTL for seed erucic acid content, with evidence from four independent field environments (Qiu *et al.*, 2006). Two major seed erucic acid content QTL were detected, one located on linkage group/chromosome A8 (A genome) and the other on C3 (C genome), accounting for *c.* 71% of the genetic variation. Taken together, such evidence indicates there are likely to be two intact highly identical functional *FAE1* gene paralogues in *B. napus*, one in each of the A and C genomes.

Here we report development of a TILLING platform for *B. napus*, involving generation of two structured EMS mutagenized populations derived from homozygous cv. Ningyou7 (one parent of the TNDH mapping population). In order to demonstrate the utility of these populations for gene discovery, a forward genetic screen was carried out, resulting in a large number of novel phenotypes. Reverse genetics was tested within a potentially multi-paralogous gene system. Comparative genomic sequence analysis involved screening of a *B. napus* cv. Tapidor (the other parent of the TNDH population) BAC library to obtain a more precise estimate of paralogous *FAE1* gene copy. BAC-derived locus-specific sequences were then mapped to the A8 and C3 chromosomes within the TNDH mapping population (Qiu *et al.*, 2006). Reverse genetic screening for *FAE1* mutants by TILLING resulted in identification of 19 mutations from 1344 M2 plants, of which three had altered function as revealed by marked changes in the accumulation of erucic acid.

## Materials and Methods

### Construction of EMS mutant population

Self-pollinated seeds of the *Brassica napus* L. DH line cv. Ningyou7 were prepared in summer 2004. For the pilot

dosage-response experiment (autumn 2004), dry high-quality mature seeds of Ningyou7 were soaked in water for 8 h at room temperature and then sets of 300 seeds were imbibed in different concentrations of EMS (Sigma-Aldrich Biotech, St Louis, MO, USA) aqueous solution ranging from 0.0% (w/v) to 1.2% at 0.2% intervals in a 50 ml flask at room temperature in the dark for 18h. Seed germination rate and germination time of M0-treated seed was recorded, as well as percentage self-pollinated seed set of M1 plants derived from the M0 seed (Table 1). In subsequent large-scale population generation experiments (autumn 2005), seeds of Ningyou7 (generated in 2004 and stored under optimum conditions) were treated in either 0.6% or 0.3% EMS. Large-scale DNA isolation, and storage of M1 and M2 seed followed the methods described for the Arabidopsis TILLING project (McCallum *et al.*, 2000; Colbert *et al.*, 2001; Greene *et al.*, 2003).

Investigation of novel phenotypes amongst the two mutant populations was carried out on M2 plants grown in field conditions (Wuhan, China at 30°35'N 108°54'E). Morphological characters scored included plant architecture, leaf, floral and reproductive morphology, with mutant phenotypes being compared to untreated wild type plants. Phenotypic trait categories are outlined in Table 2. Seed oil content of M3 seed was estimated using near infrared spectroscopy (NIR) with a standard method (Mika *et al.*, 2003).

### BAC library screening and BAC clones analysis

The JBnB BAC library was developed from DNA isolated from a *B. napus* doubled haploid line derived from cv. Tapidor (Rana *et al.*, 2004) and screened by hybridization of a <sup>32</sup>P-labeled 426 bp Tapidor *FAE1* genomic DNA fragment (amplified by primer pair FAE426). Primer pairs FAE426 and Fae1primer1 to 5 (Table 3, Fig. 1) were designed from *B. napus* *FAE1* cDNA sequence AF274750 (GenBank accession) and *B. napus* *FAE1* promoter sequence AF275254. BAC DNA was isolated using a modified mini-alkaline lysis extraction procedure (Bendahmane, 1999). PCR walking on three representational BAC clones (BAC1, BAC7 and BAC11) followed standard procedures (Gibbons *et al.*, 1991). PCR products were

Morphological characters	No. of plants		Percentage (%)	
	0.3% (2604 plants)	0.6% (7110 plants)	0.3%	0.6%
<b>Architecture</b>				
Dwarf stature	3	9	0.12	0.13
Increased stature	23	112	0.88	1.58
others	34	189	1.31	2.66
<b>Flower</b>				
<b>Colors</b>				
Pure white	7	13	0.27	0.18
Light-yellow/creamy	25	98	0.96	1.38
Abnormal petal	21	123	0.81	1.73
Abnormal stamen	36	209	1.38	2.94
Flowers absent	5	27	0.19	0.38
others	21	57	0.81	0.80
<b>Leaves</b>				
Abnormal appearance	35	213	1.34	3.00
Abnormal color	37	145	1.42	2.04
<b>Reproductive</b>				
infertile	91	457	3.49	6.43

Mutants were categorized according to architectural, floral, leaf and reproductive traits. The numbers recorded represent characters scored on individual plants and thus may represent sibling aggregates. Furthermore, since the scores are nonexclusive, a single plant may contain more than one variation.

**Table 2** Summary of phenotypes observed in M2 *Brassica napus* plants from the two ethylmethanesulfonate (EMS)-treated (0.3 and 0.6%) populations


Primer name	Forward primer sequence (5'→3')	Reverse primer sequence (5'→3')
FAE426	TACTCATGCTACCTTCCAC	ACTACAACCCATGCCACCAA
Fae1primer1	CCCTGTTCAATTAATTAAGTAGCG	ACGTTCTTGAATCTTCCTCA
Fae1primer2	CCTCATAACCCATCGCTCC	ATGATAACTTGCTCCGTCTCT
Fae1primer3	GCTTGTCCGTGTTGGTG	AATCCGGGACGTAGTAATGTT
Fae1primer4	CCCGGATTTCAAACCTTGCTA	ATGCAGTGTCCCAAGGACTA
Fae1primer5	CCCGGATTTCAAACCTTGCTA	AAATACTAGAACCACCATTTG
Fae1p1-1	TTCTCTTCTTCCGCTTCCA	TGGGTGTGATGAATGCTTCT
Fae1p1-2	GGAGAGGAAAAGAGGAGAGGA	GACGACTCTGTCCGTGTTGA
Fae1p1-3	CATGAATTGAAACGAGAAGG	TTTCAAACCCACGTGTCCAC
Fae1p2-1	GGACCCATGGCAATAGCCAAG	ACGTTCTTGAATCTTCCTCA
Fae1E1	GGGTCAGGCTTTAAGTGTAACAG	CCAATCAATTCGGGAGCCAC
Fae1E2	GGGTCAGGCTTTAAGTGTAACAG	CCGTGCGAGAGCTATTAGGAC
M-FAE1-A8	ATCTCCGTGAGGAGCCAGAGAGAC	ACGTTCTTGAATCTTCCTCA
M-FAE1-C3	CCCGGATTTCAAACCTTGCTA	TTAACAGAGATCCTTAACCCC
S-FAE1-A8	TACTCATGCTACCTTCCAC	CCGTGCGAGAGCTATTAGGAC
S-FAE1-C3	TACTCATGCTACCTTCCAC	CCAATCAATTCGGGAGCCAC

**Table 3** Primer sequences used for PCR amplification

sequenced and six primer pairs (Fae1p1-1, Fae1p1-2, Fae1p1-3, Fae1E1, Fae1p2-1 and Fae1E2) (Table 3, Fig. 1) were designed based on these sequences. Gene-specific primer pairs, S-FAE1-A8 and S-FAE1-C3, were designed to provide specific amplification products that distinguished between the two groups of BAC clones.

Based on the *FAE1* sequence obtained from the *B. napus* cv. Ningyou7 and Tapidor DH parents of the TNDH mapping population, two polymorphic markers, FAE1-A8 and FAE1-C3 (using primer pairs M-Fae1-A8 and M-Fae1-C3, Table 3,

Fig. 1), were developed. This enabled us to identify and map loci in the context of the TNDH linkage map constructed by Long *et al.* (2007) using JoinMap 3.0 software (<http://www.kyazma.nl/index.php/mc.JoinMap>). Seed erucic acid content data were obtained from field trials of the TNDH population conducted over four environments (Qiu *et al.*, 2006). Seed erucic acid content QTLs were detected by the composite interval method (CIM) using Windows QTL cartographer 2.5 software (<http://statgen.ncsu.edu/qtlcart/WQTLCart.htm>).



**Fig. 1** Regions of the *FAE1* BAC clones covered by primer pairs on the two groups (the names of the primer pairs are marked above their corresponding amplicons). Ten BAC clones in group 1 correspond to *BnC3.FAE1* and three BAC clones in group 2 correspond to *BnA8.FAE1*. Lines represent 5' and 3' genomic sequence of *FAE1* (c. 1 kb for 5' and 3' of *BnA8.FAE1*, c. 3.5 kb for 5' of *BnC3.FAE1* and c. 1 kb for 3' sequence); rectangles represent coding region (1521 bp). Lines with triangles at the end indicate specific primers of *BnA8.FAE1*; lines with circles at the end indicate that primer pair can amplify both *BnA8.FAE1* and *BnC3.FAE1*; lines with diamonds at the end indicate specific primers of *BnC3.FAE1*. Fae1primer1 and Fae1primer5 were used in the PCR assay in Fig. 4.

### *FAE1* mutation screening

The primer pair FAE426 was used to screen the EMS mutant populations. The forward primer was labeled at the 5' end with 700 nm IRD fluorescence and the reverse primer was labeled with 800 nm IRD fluorescence (MWG Biotech, Inc., Ebersberg, Germany). DNA samples isolated from M2 plants of the 0.6 and 0.3% EMS populations were screened individually, with the screening protocol following that used in rice and Arabidopsis TILLING projects (Till *et al.*, 2004b; Till *et al.*, 2006). CEL 1 was extracted from celery and stored in  $-80^{\circ}\text{C}$  (Till *et al.*, 2004a). FAE426 with unlabeled primer was used to amplify products from genomic DNA of M2 plants which had yielded novel cleaved bands at 700 and 800 nm in the TILLING gel. The PCR products were electrophoresed on agarose gel so as to recover the 426 bp bands and then these 426 bp PCR products were sequenced.

### Mutation sequence analysis

Sequence derived from mutant and wild-type (WT) plants was analyzed to determine the likelihood of changes affecting protein function. The PARSESNP (Project Aligned Related Sequences and Evaluate SNPs; <http://www.proweb.org/parsesnp/>) and SIFT (Sorting Intolerant from Tolerant; <http://blocks.fhcrc.org/sift/SIFT>) software packages were used to predict the severity of each mutation (Ng & Henikoff, 2003; Taylor & Greene, 2003). SIFT values  $< 0.05$  and PSSM scores  $> 10$  indicated that the nucleotide change was likely to affect gene function.

The 426 bp sequences obtained following TILLING screening were used in BLAST comparison against the comprehensive nucleotide database of available *Brassica* sequences served at BrassicaDB (<http://brassica.bbsrc.ac.uk/BrassicaDB/>). All related *B. napus* DNA sequence accessions that were detected with an  $E$ -value  $< 10^{-15}$  were batch-downloaded from the NCBI GenBank nucleotide database. The sequences were aligned using a local implementation of the NCBI standalone Blast software (packageblast-2.2.18-ia32-win32.exe, <http://www.ncbi.nlm.nih.gov/BLAST/download.shtml>). The 426 bp

sequence was used as query and those downloaded sequences that appeared to contain a sequence error (a different base present in just one downloaded sequence) were deleted. Default parameters were used in Blast, except for the output format (its format was 'm 4' so that SNPs can be easily visualized). SNPs were assigned where different bases at a given nucleotide position appeared in two or more sequences.

### Phenotypic and genotypic analysis

The erucic acid content of seeds harvested from WT Ningyou7 and M2 plants which contained *FAE1* mutations was measured by gas chromatography (GC) using standard methods (Kaushik & Agnihotri, 2000) with multiple seeds tested from each plant. Wild-type Ningyou7 plants were grown in the same field and seeds were harvested at the same time as the mutant population. Single seed of the M3 generation of the two mutants L080-1 and N004-1 was used to establish the erucic acid content, using a half-seed GC analysis method (Conte *et al.*, 1989).

To detect the location of mutations in *fae1* paralogues in the A8 or C3 chromosomes, M2 DNA from mutants L080-1 and N004-1 was used to amplify products using the gene locus-specific primer pairs S-FAE1-A8 and S-FAE1-C3. PCR reactions were carried out with proof-reading *pfu taq* polymerase (Tiangen Biotech, Inc., Beijing, China), and the purified PCR products sequenced.

Genetic analysis of the two mutants L080-1 and N004-1 involved ANOVA and multi-comparison analysis of phenotypic and genotypic data, using SAS 8.2 software.

## Results

### Mutant population development

We first established an optimal concentration of EMS for treatment of *B. napus* seeds required to generate a mutant population. Three hundred self-pollinated seeds from a Ningyou7 DH line were treated with EMS at each of seven different concentrations, from 0.0 to 1.2% (w/v), at 0.2%

**Table 4** Summary characteristics of the two *Brassica napus* cv. Ningyou7 ethylmethanesulfonate (EMS) populations


	0.3% EMS	0.6% EMS
<i>M1 generation</i>		
Number of plants	~ 1080	~ 13 000
<i>M2 generation</i>		
Number of lines	933	3926
Number of plants	2604	7110
Mutations detected from x M2 plants	4	15
Mutation load within plant	614	730
Mutation load within plant	9174	28 941
Mutation density per plant	130.8 kb	41.5 kb

intervals. The M1 germination rate, germination time and self-pollinated percentage seed set were recorded (Table 1). At EMS concentrations of 0.8% or above, the germination rate was < 85%, and germination time was prolonged. Almost all the surviving M1 plants were unable to set seed through self-pollination, with M2 seed set < 10%. At 0.6% EMS, 25% of M1 plants had sufficient self-pollinated seeds, with a germination rate of *c.* 85%. An EMS concentration of 0.6% was therefore selected as an optimal treatment for Ningyou7 seeds in developing the mutant population. To determine whether a lower EMS concentration, generating a higher rate

of M1 self-pollinating seed set, would be suitable for TILLING, a 0.3% EMS concentration was also selected to develop a secondary mutant population. Between one and four M2 plants were generated by random selection from each M1 plant. In total, 3926 lines were developed from the 0.6% EMS-treated population with 7110 M2 plants, and a further 933 lines from the 0.3% EMS-treated population with 2604 M2 plants (Table 4). DNA was extracted from all M2 plants and diluted to the same concentration suitable for PCR reactions. Self-pollinated seeds were also harvested from the individual M2 plants and stored in a seed bank.


#### Forward genetic screening of mutants in the M2 population

To evaluate and compare the effect of mutagenesis in the two populations through forward genetics, all M2 plants were scrutinized for visible variant phenotypes. These included key developmental characters throughout the growing season, such as plant architecture, leaf, floral and reproductive morphology (summarized in Table 2). As expected, the population treated with 0.6% EMS yielded a higher percentage of mutant phenotypes, with examples of visible phenotypic variants shown in Fig. 2. In order to assess variation in total seed oil content, 652 M3 seeds from the



**Fig. 2** Examples of phenotypic mutants observed in the ethylmethanesulfonate (EMS)-treated M2 populations of *Brassica napus*. (a) Mutant plant with stocky (reduced stature) architecture; (b) curly leaf phenotype; (c–f) floral mutants: antherous inflorescence (c); white flowers (d); absent and distorted stamens (e); aborted buds (f). Both (e) and (f) were also recorded as infertile mutants.





**Fig. 3** Distribution of *Brassica napus* seed oil content within 1168 M3 plants (652 samples were from the 0.6% ethylmethanesulfonate (EMS)-treated population, and 516 were from the 0.3% EMS population). ANOVA did not detect any significant difference in the oil content between the two populations. However, the transgressive distribution appeared greater in the 0.6% EMS population, which contained both the maximum (30.62%) and minimum (47.93%) oil contents. The oil content of wild-type (WT) Ningyou7 was  $38.36 \pm 1.10\%$  ( $n = 34$ ). The maximum oil content of WT was 40.89% and the minimum was 36.36%. The values recorded were from M3 seed samples scored on an individual plant basis, and thus may represent sibling aggregates.



**Fig. 4** PCR assay to analyze *Brassica napus* cv. Tapidor BAC clones. Thirteen BAC clones were amplified with Fae1primer1 (a) and Fae1primer5 (b). Lane numbers are indicated on the figures, with M representing the DNA ladder DL2000 (Promega, Inc). (a) Fae1primer1 was not able to amplify a product from BAC11. (b) Fae1primer5 was able to classify the 13 BAC clones into two groups. Group 2 included BAC4, BAC7 and BAC10, with the remainder in group 1. Subsequent investigation revealed BAC11 to be in group 1.


0.6% EMS population and 516 from the 0.3% EMS population were compared with untreated seed (WT). ANOVA of these three groups indicated no significant difference in oil content between the two EMS populations, although a significant difference ( $P < 0.01$ ) was observed between the 0.6% EMS population and WT ( $n = 34$ ). A significant difference ( $P < 0.01$ ) was also detected between the 0.3% EMS populations and WT. The frequency distribution of oil content amongst a total of 1168 M3 seeds (Fig. 3) shows a maximum value of 47.8%, considerably higher than WT ( $38.36 \pm 1.10\%$ ).

#### *FAE1* homologous in *B. napus* cv. Tapidor and Ningyou7

To estimate the copy number of paralogous *FAE1* genes in the *B. napus* AC genome, and to identify sequence differences amongst these copies, a 426 bp probe generated from Tapidor genomic DNA was used to screen the JBnB BAC library, which was developed from the doubled haploid *B. napus* cv. Tapidor (Rana *et al.*, 2004). 13 BAC clones (JBnB060D12, JBnB015C15, JBnB060D12, JBnB154K14, JBnB170L17, JBnB160P09, JBnB079L09, JBnB009C03, JBnB065M23, JBnB149N05, JBnB032N18, JBnB178F05, and JBnB108J15),

identified here as BAC1 to BAC13, were identified with positive hybridization signals. Five primer pairs were designed to amplify each of these BAC clones (Fae1primer1 to Fae1primer5, see Table 3, Fig. 1). Fae1primer5 allowed classification of the BAC clones into two groups, with BAC4, BAC7 and BAC10 in group 2 generating single amplified bands of the same size when visualized on agarose gels, and no products obtained from the remaining BACs of group 1 (Fig. 4b). PCR with Fae1primer1 resulted in products of predicted length from all BAC clones apart from BAC11 (Fig. 4a). PCR-walking was then used to obtain the 5' and 3' sequences of the *FAE1* gene in each of three clones (BAC1, BAC4, BAC11) representing the different classes. The end of the BAC11 insert was located near the *FAE1* promoter region, which explained why the Fae1primer1 was unable to amplify a product from this BAC, and suggested that this clone may in fact belong to the same locus-class as BAC1. We then obtained sequences flanking the *FAE1* gene from BAC1, including *c.* 3.5 kb from 5' and 1 kb from 3' regions, together with *c.* 1 kb each of 5' and 3' sequence from BAC4. Using these sequences, six primer pairs were designed (Fae1p1-1, Fae1p1-2, Fae1p1-3, Fae1p2-1 amplifying the 5' end region, Fae1E1, and Fae1E2 amplifying the 3' end region) (Table 3,





**Fig. 5** Mapping of the two *FAE1* paralogues in the context of the reference TNDH linkage groups, and detection of QTLs for seed erucic acid content. The two linkage groups were constructed based on data presented in Qiu *et al.* (2006) and Long *et al.* (2007). *BnA8.FAE1* was located on *Brassica napus* chromosome A8, and *BnC3.FAE1* was located on chromosome C3. Two hundred and two TNDH lines were grown in four environments, two in the south of China (Wuhan, at 30°35'N 108°54'E) and two in the north (Dali, at 34°52'N 109°56'E). Linkage map construction and QTL detection followed the methods outlined by Qiu *et al.* (2006), with the length of the vertical lines representing 95% confidence intervals of these QTLs. All QTLs shown were significant at  $P < 0.01$ .

Fig. 1). In combination, these primers were able to be used in PCR reactions to generate products that enabled classification of the 13 BAC clones into two groups, corresponding to two distinct loci. The results were consistent with those obtained with *Fae1* primer5 (data not shown).


To further characterize the paralogues within the *B. napus* AC genomes, the *FAE1* gene and its flanking sequence were obtained from each of the 13 BAC clones using the six specific primer pairs. The 10 BAC clones in group 1 were identical throughout *c.* 6 kb of sequence (3.5 kb was 5' of the *FAE1* start codon, 1 kb was 3' of the *FAE1* stop codon, with a coding region of 1521 bp) (Fig. 1). The three BAC clones in group 2 also showed no sequence difference within the *c.* 3.5 kb region (with 1 kb 5' end of the *FAE1* start codon, and 1 kb 3' end of the *FAE1* stop codon) (Fig. 1).

Two markers, *FAE1-A8* and *FAE1-C3*, corresponding to each group of BAC clones, were found to be polymorphic between the parents of the TNDH mapping population. These enabled us to locate the existing *FAE1* BAC clones in the context of the TNDH genetic linkage map, on chromosomes A8 and C3. Both markers co-located with the peaks of the QTLs for seed erucic acid content (Fig. 5), indicating that these corresponded to the two paralogous copies of *FAE1* in the *B. napus* genome. The two genes were designated as *BnA.FAE1.a* and *BnC.FAE1.a*, according to the standard nomenclature of Ostergaard & King (2008). However, to simplify subsequent reference within this paper, *BnA8.FAE1* and *BnC3.FAE1* are used to represent paralogues in the A8

and C3 chromosomes of *B. napus*, respectively. Two gene locus-specific primer pairs, S-*FAE1-A8* and S-*FAE1-C3*, corresponding to *BnA8.FAE1* and *BnC3.FAE1*, respectively, were designed from the sequence flanking *FAE1*. These enabled us to obtain complete genomic sequences for both paralogues from the WT Ningyou7 DH parent of the EMS mutant populations that included 5' and 3' flanking regions. Alignment of these sequences (GenBank accessions: EU543282 and EU543283) with cDNA sequences present in the GenBank nucleotide database confirmed the lack of introns within this gene, as reported elsewhere (Wu *et al.*, 2007b). The coding region of both paralogues was 1521 bp in Ningyou7, with a sequence similarity of 98.6%. Twenty-one SNPs were identified between the two paralogues. There was also very little (*c.* 6%) variation in the *c.* 400 bp upstream region of the start codon.

#### Screening for *FAE1* mutations in the EMS mutant populations

To screen for mutations in EMS-treated populations using TILLING, it is essential to utilize gene-specific primer pairs (Colbert *et al.*, 2001). The maximum length of fragment that can be screened using the Li-Cor 4300 polyacrylamide gel system is just over 1 kb. Although other authors have reported detection of mutants by TILLING in PCR products in the range 300 bp–1.6 kb (Slade *et al.*, 2005), our experience was that we observed very diffuse bands with PCR fragments of



**Fig. 6** Mutation within the *FAE1* gene fragment from L209-1 identified by TILLING. 700 nm and 800 nm Li-Cor 4300 scanning gel photos are shown in (a) and (b). Two novel bands were found in the same lane, respectively, in both the 700 and 800 nm images, and these are magnified in (e) and (f). (c, d) Partial sequence trace corresponding region in L209-1 (c) and in wild type (d). The primer pair FAE426 was designed to amplify the two *FAE1* paralogues simultaneously – this is demonstrated at position 480 where double peaks are observed in the sequence trace (which indicates an induced nucleotide change G to A in one of the two paralogous copies at this position), whilst a single peak is observed in the wild type at the same position (indicates both of the two paralogous copies *FAE1* in this position are G).

> 1 kb on the Li-Cor 4300 gels. Moreover, although we were able to use gene-specific primer pairs for each of the paralogous copies of *FAE1*, the forward and reverse primers matched positions remote from the start and stop codons. As a result, it was not possible to obtain gene-specific primers that corresponded to the coding region suitable for TILLING of mutations within each of the two highly similar paralogues of *FAE1*. We therefore designed the primer pair FAE426, so that it simultaneously amplified 426 bp from both paralogues, for which five SNPs already existed to distinguish the WT alleles, and used this in TILLING for novel mutations in our EMS populations.

Co-amplification of two similar genes in a pooled sample can lead to an increased error (Cooper *et al.*, 2008). We therefore carried out direct screening for novel mutations from DNA isolated from individual M2 plants, using the standard TILLING protocol (Till *et al.*, 2006). TILLING gels were scanned using a Li-Cor 4300. Putative mutant lines and location of nucleotide changes were validated by sequencing of the PCR product generated from unlabeled FAE426 primers. Double peaks within the sequence trace not associated with the five known paralogous SNPs positions indicated an induced nucleotide change, which also corresponded with the cleaved band position in the Li-Cor TILLING gel. Fig. 6 shows one

**Table 5** *FAE1* mutations identified by TILLING

M2 plant	Nucleotide change	Amino acid change	PSSM difference <sup>c</sup>	SIFT score <sup>d</sup>	Seed erucic acid content (%)
L324-1	C266T <sup>a</sup>	P89L <sup>b</sup>			47.77
L457-1	C347T	T116M			50.27
N050-1	G384A	K128=			48.77
N057-1	G391A	E131K	<b>11.2</b>	0.22	43.56
N004-1	G400A	G134S	<b>21.3</b>	<b>0.01</b>	38.33
K026-1 <sup>e</sup>	C403T	L135=			49.34
L297-1	G412A	E138K	<b>10.8</b>	<b>0.05</b>	33.19
N128-1	C420T	H140=			52.53
L518-1	C446T	P149L			48.77
N142-1	C449T	P150L			49.34
N184-1	C450T	P150=			45.58
K024-1 <sup>e</sup>	G453A	R151=			48.59
L421-1	G478A	E160K			50.08
L209-1 <sup>e</sup>	G480A	E160=			51.26
L496-1	G508A	E170K			50.78
L495-1	A542C	K181T	<b>11.9</b>	0.17	38.26
L080-1 <sup>e</sup>	C599T	S200F	<b>19.4</b>	<b>0.00</b>	19.46
N377-1	C627T	L209=			48.56
L286-1	G629A	R210Q	<b>19.9</b>	<b>0.01</b>	no seed

<sup>a</sup>The code for nucleotide change refers to mutations at specific positions in the genomic sequence of the *FAE1* gene, referenced from the from the first base (A) of the start codon. Thus, C266T indicates that the 266th nucleotide had a transition from C to T.

<sup>b</sup>The consequence of the nucleotide mutation in the *FAE1* protein chain is referred to by a code whereby the initial and resulting amino acid is shown, referenced to the amino acid position of the start codon. Thus, P89L indicates that the 89th amino acid is predicted to be modified from P to L. =, indicates no change (synonymous mutation).

<sup>c</sup>SIFT scores < 0.05 and PSSM value > 10 are predicted to be deleterious to gene function.

<sup>e</sup>Plants indicated arose from the 0.3% EMS treated population, with the remainder from the 0.6% ethylmethanesulfonate (EMS)-treated population. Bold type indicates where SIFT scores or PSSM value were greater than the threshold for likelihood of causing a deleterious effect on protein function.

example of a mutation detected within M2 plant L209-1 (M2 plant field accession NO). From the Li-Cor gel system, 700 and 800 nm images are obtained (Fig. 6a,b). The cleaved bands corresponding to the induced mutation were located at *c.* 230 bp in the 700 nm image, and at *c.* 190 bp in the 800 nm image, thus totaling 426 bp. The sequence traces indicate the mutated region of L209-1 (Fig. 6c) and the corresponding region of nontreated WT Ningyou7 (Fig. 6d). In total, 19 mutations were discovered amongst 1344 M2 plants screened (one plant from each M2 line) (Table 5).

#### Relationship between EMS concentration and mutation frequency

Of the 19 mutations detected, 15 arose from the 0.6% EMS mutagenized population, following screening of 730 M2 plants, and four from the 0.3% population following screening of 614 M2 plants (Table 5). Given that *FAE426* was designed to amplify two *FAE1* paralogous copies simultaneously, this corresponds to a single screen of 852 bp within the *B. napus* genome. Within the 0.6% EMS population, the mutation frequency was calculated at one mutation per 41.5 kb in each plant ((0.426 kb × 2 × 730 plants)/15 mutations = one mutation every 41.5 kb) and 28 941 mutations per plant

(1 200 000 kb per plant/one mutation every 41.5 kb = 28 941 mutations per plant), and within the 0.3% EMS population at one mutation per 130.8 kb ((0.426 kb × 2 × 614 plants)/four mutations = one mutation every 130.8 kb) and 9174 mutations per plant (1 200 000 kb per plant/one mutation every 130.8 kb = 9174 mutations per plant) (Table 4).

Of the 19 *FAE1* mutations, 18 represented G/C to A/T transitions and one represented an A to C transversion (Table 5), which is a close fit to previous reports that EMS mutagenesis primarily induces G/C to A/T transitions.

#### A comparative analysis of natural *FAE1* SNPs in rapeseed cultivars and in the mutant population

Ecotilling is a new approach for detecting SNPs in natural populations (Comai *et al.*, 2004). To determine whether the 19 induced mutations were present within existing lines representing selected varieties, breeding or experimental material, an *in silico* Ecotilling analysis was carried out. A total of 112 *B. napus* *FAE1* sequences that corresponded to the 426 bp region screened in our TILLING experiments were aligned. This included 101 ESTs (note that this gene is intronless) and 11 genomic sequences. Once sequence errors were deleted (i.e. a different base present in just one downloaded

**Table 7** Genetic analysis of the *Brassica napus* M3 generation from the heterozygous mutant *c3fae1-400*

	Plant number	Seed erucic acid content (%)	LSD <sup>a</sup> all-pairwise comparisons test
<i>C3FAE1-400/C3FAE1-400</i>	9	52.01 ± 1.14	A <sup>b</sup>
<i>C3FAE1-400/c3fae1-400</i>	16	45.71 ± 0.85	B
<i>c3fae1-400/c3fae1-400</i>	6	31.83 ± 1.39	C

<sup>a</sup> Least significant difference multiple comparisons.<sup>b</sup> Significance at  $P < 0.01$ .**Table 6** *In silico* Ecotilling with *Brassica napus* FAE1

Positions	264 <sup>a</sup>	284	311	312	363	369
SNPs	G/A <sup>b</sup>	G/C	A/T	C/T	G/C	C/T
Positions	393	417	462	531	591	606
SNPs	G/A	C/T	T/G	T/C	A/T	A/T/G
Positions	623	635	648	657	666	667
SNPs	A/D <sup>c</sup>	A/D	T/D	G/T/A	G/T	G/T/D <sup>d</sup>

<sup>a</sup> Position from the FAE1 start codon ATG, to the 264th nucleotide.<sup>b</sup> Indicates that two alternative nucleotides were detected at this position, with G/A denoting that G was prevalent in some accessions, and A in others.<sup>c</sup> Indicates the presence of an indel – with A/D denoting that A was present in some accession, whereas a deletion was present at the corresponding position in others.<sup>d</sup> Indicates the presence of either two alternative nucleotides (G/T) or a deletion at this position. Comparison of the 18 SNPs to the 19 ethylmethanesulfonate (EMS)-induced mutations (Table 5) indicates that there was no overlap in the distribution of nucleotide changes.

sequence), nucleotide polymorphisms were detected at 18 positions, each of which was distinct from the 19 induced point mutations (Table 6).

### Functional characterization of *fae1* mutants with reduced seed erucic acid content


In total, seven (37%) synonymous mutations and 12 (63%) mis-sense mutations were identified from TILLING screening of 1344 M2 plants. Both SIFT score and PSSM values were used to predict whether the base change for individual mutations would be deleterious to FAE1 gene function (Ng & Henikoff, 2003; Taylor & Greene, 2003). Four of the 19 mutations were predicted to affect the FAE1 protein function since both the SIFT score and PSSM values for these events were at least at the threshold of severity (Table 5). Seed erucic acid content was determined by GC for M3 seeds of 18 available plants (multiple seeds tested from each plant). The M3 seed erucic acid content of plants N057-1, N004-1, L297-1 and L080-1 was lower than that of the wild-type. This corresponded well with the predicted probabilities of altered gene function derived from the four mutations (Table 5). Two

individual M2 plants, L080-1 and N004-1, were selected to identify the mutated locus within the genome using gene-specific primer pairs S-FAE1-A8 and S-FAE1-C3 (Table 3), which amplified *BnA8.FAE1* and *BnC3.FAE1*, respectively. The M2 mutant plant L080-1 was recessive homozygote *bnA8.fae1* at position 599bp (assigned mutant *a8fae1-599*), while the M2 plant N004-1 was heterozygous *BnC3.FAE1/bnC3.fae1* at 400 bp (assigned mutant *c3fae1-400*) (Fig. 7). Alignment of these sequences to the corresponding paralogous copies in WT Ningyou7 showed that *BnC3.FAE1* in L080-1 corresponded to WT *BnC3.FAE1*, and that *BnA8.FAE1* in N004-1 corresponded to WT *BnA8.FAE1*. We therefore deduced that in each mutant line the paralogous copy contained no other induced mutations.

Thirty-one M3 seeds were harvested from the M2 mutant plant N004-1, and the seed erucic acid content was determined using half of each seed. This involved dissection and analysis of one cotyledon. The remaining half seed included the viable shoot apical meristem and radicle, which was germinated and grown to yield M3 plants, allowing subsequent genetic analysis. The genotypic segregation in 31 M3 plants displayed Mendelian inheritance with 9 *C3FAE1-400/C3FAE1-400* : 16 *C3FAE1-400/c3fae1-400* : 6 *c3fae1-400/c3fae1-400*, corresponding to a 1 : 2 : 1 ratio supported by a chi-squared test ( $\chi^2 = 0.6130$ ). The erucic acid content in the 31 seeds was significantly ( $P < 0.01$ ) associated with their genotypes tested by LSD (least significant difference) all-pairwise comparison test multiple comparison (Table 7). We conclude that the point mutation in *c3fae1-400* appears to be causal in reducing the seed erucic acid of N004-1. Moreover, no significant difference was observed between the seed erucic acid content ( $19.21 \pm 0.53\%$ ) among 29 M3 seeds of L080-1 with ANOVA analysis, consistent with a recessive homozygote genotype at *bnA8.fae1*.

### Discussion

We have successfully constructed two rapeseed mutant populations suitable for both forward and reverse genetic detection methods. In the forward genetic screening of mutants, a large number of novel mutated phenotypes were identified, and many of these represent a valuable genetic resource affecting



**Fig. 7** Location of *FAE1* mutations in lines L080-1 and N004-1. DNA from single M2 plants (L080-01 and N004-1) was amplified with the primer pair FAE426. (a, d) The amplified PCR products were sequenced and partial sequence traces are shown with double peaks at positions of nucleotide transition. Two gene-specific primer pairs, S-FAE1-A8 and S-FAE1-C3, amplified DNA from M2 plant of the two mutants L080-1 and N004-1, and the PCR products were sequenced. (b) The partial *FAE1* sequence trace of L080-1 from chromosome A8. (c) Corresponding sequence from the C3 gene copy. (e) Partial *FAE1* sequence trace from N004-1 in A8 chromosome. (f) Sequence from N004-1 corresponding to the chromosome C3 paralogue. From these four trace files, it was possible to deduce that a C to T transition mutation had occurred in the L080-1 *FAE1* genomic sequence corresponding to the gene copy located on chromosome A8. Similarly, it was easy to deduce that a G to A transition mutation within the N004-1 *FAE1* genomic sequence corresponded to the paralogue located on chromosome C3. Moreover, it was possible to predict that the *FAE1* mutation within the L080-1 M2 plant was homozygous, whilst the *FAE1* mutation within the N004-1 M2 plant was heterozygous.

crop traits such as male sterility, reduced stature and plants with increased oil content. In the reverse genetic TILLING of mutants, we screened 1344 M2 plants representing 1344 distinct lines (i.e. siblings were not screened), and detected a total of 19 *FAE1* mutations, three of which had a change of function. In addition, *in silico* Ecotilling analysis demonstrated a clear distinction between sites of natural and induced SNPs. This is very encouraging for the application of TILLING within rapeseed crop genetic improvement as a highly efficient method for generating new genetic resources for breeding. We are also able to calculate that if a 1kb gene fragment were to be screened within the 0.6% EMS population, with a single plant selected from each line, then *c.* 100 mutations would be detected following a complete screening of the 0.6% EMS population (a total of 3926 lines). This suggests that the population carries a high mutation load, and is a valuable resource for breeding and research. However, for practical application of induced mutation within crop improvement, it is necessary to adopt a process that reduces the background caused by the mutations at unlinked loci. Such a schema should involve backcrossing to the WT or target elite cultivar,

using molecular marker and phenotypic tracking for recurrent selection of the mutant locus. It is also important to eliminate epistatic effects and carry out controlled self-pollination of the candidate mutant plant before segregation analysis of the mutation identified and modified phenotype. Such information can be substantiated by analysis of transcript profile and *in vitro* protein expression for the target gene (Heckmann *et al.*, 2006). We conclude that the populations generated are efficient for exploring novel sources of variation using both forward and reverse genetics approaches to screening mutants.

Although we were able to demonstrate development of an efficient new research platform through discovery of three novel induced functional low erucic acid mutant alleles, it is important to determine the optimal mutagen dosage for *B. napus*. The EMS concentration is a key factor in developing such populations, with higher concentrations leading to higher mutation frequencies. However, any higher mutation load achieved is offset by the effect of EMS on reducing seed set, viability and germination (Greene *et al.*, 2003; Kim *et al.*, 2006). For *B. napus* there have been no prior reports of an optimal mutagen dosage to guide development of a forward



genetics or reverse genetics TILLING population. Here, we found that at EMS concentration of 0.6% yielded a calculated mutation density of one per 41.5 kb in each plant, compared with almost a third of that in the population treated at 0.3% EMS. These estimates may not accurately reflect the true mutation density of the two EMS populations, as it was not possible to screen the first 50 bp in a TILLING gel. However, the mutation frequencies within our two *B. napus* mutants were within the range of those reported elsewhere for monocot and dicot species, with one mutation per 24 kb reported in hexaploid wheat, one per 40 kb in tetraploid wheat (Slade *et al.*, 2005), 300 kb in rice (Till *et al.*, 2007), 100 kb in barley (*Hordeum vulgare* L.) (Caldwell *et al.*, 2004), and 170 kb in *Arabidopsis* (Greene *et al.*, 2003). In the forward genetic investigation, 23.25% M2 plants displayed observable mutant phenotypes in the 0.6% EMS population, compared with 12.98% in the 0.3% EMS population. These estimates may be low and not reflect the true mutant load, as some plants may carry more than one mutant phenotype. Moreover, when M2 plants were grown in field conditions, we found that the performance within the 0.3% EMS population was stronger than the 0.6% population. At concentrations of EMS above 0.6% we found that M1-treated Ningyou7 plants set almost no seed. Therefore, taking into account the mutation frequency and the performance of mutagenized plants in the field, we consider an EMS concentration between 0.3 and 0.6% to be optimal for construction of mutant populations in *B. napus* and probably also in other amphidiploids of the *Brassica* triangle of U, for example *B. juncea* ( $2n = 36$ ; AB genomes) and *B. carinata* ( $2n = 34$ ; BC genomes).

Since the original report of TILLING in 2000, the technique has successfully been applied to many crops, notably in rice and wheat (Slade *et al.*, 2005; Till *et al.*, 2007). However, we are not aware of other reports to date of its application within any *Brassica* species. The main challenges arise from the complex genome of *B. napus* in which most of genes are multi-paralogous. This reduces the ease with which gene locus-specific primers can be designed, and also limits the ability to locate the mutations unequivocally to specific paralogous copies. Here we have successfully demonstrated that TILLING technology can also be applied in polyploid crops such as *B. napus* that comprise multiple closely related complex segmentally duplicated genomes. We have been able to allocate target genes within the *B. napus* genome into two classes. The first class comprises those genes which occur as multi-paralogous copies distinguished by clear sequence differences; and the second class includes multi-paralogous genes that possess only minor sequence differences between different copies. It is a relatively trivial exercise to design gene-specific primers for the first class. However, for the second class we have developed a novel strategy that is based on using a primer pair that is designed to amplify all paralogous copies of the gene simultaneously, albeit from individual rather than pooled DNA samples. The procedure is based on single plant

TILLING, to identify mutations in situations where two or more very similar paralogous gene copies exist in a genome. We made use of existing SNPs as a positive control, and demonstrated the ability to distinguish novel mutations, based on use of primer pairs designed to amplify both *FAE1* paralogues simultaneously. Slade *et al.* (2005) reported that using a primer pair that could amplify at least two gene copies would be less efficient. However, in our experiments we found that these types of primers provide a powerful approach to mutant detection. The precondition for successful screening of mutations using this approach is that the target fragment should be highly conserved amongst the different paralogues. In the case of *FAE1*, we found just five SNPs between the two paralogues in the 426 bp target region. These five SNPs were also detected as background signals in the Li-Cor gel images, but did not detract from our ability to detect novel induced mutations (Fig. 5). Within the TILLING gel, the cleaved bands that were present in all lanes with positions differing between 700 and 800 nm are predicted to arise from the SNPs that exist between the two *FAE1* paralogues. However, the cleaved bands that were detected in only one lane within the 700 nm image, together with the corresponding cleaved band in the 800 nm image, indicated discovery of a novel mutation. Therefore, we can conclude that with a modified TILLING primer design we are able to screen for mutations in any gene in the EMS population of *B. napus*, even though this consists of very complex duplicated genomes. This modified method may improve and widen the application of TILLING technology for crop genomic research and improvement.

For rapeseed, the ability to achieve zero or low seed erucic acid (LEA) content has long been a major breeding objective, since erucic acid has been regarded as an antinutritional component in seed oil (Vles *et al.*, 1978). In the 1960s, the first variant with LEA content was found in the animal feed rape cultivar 'Liho', with the first LEA rapeseed 'Oro' being derived following introduction of the LEA variation from 'Liho' (Downey & Craig, 1964). Subsequently, almost all LEA rapeseed cultivars have carried the LEA gene source from 'Liho' or 'Oro', and so this single genetic resource has probably contributed considerably to an inbreeding effect and associated genetic erosion through linkage drag in rapeseed breeding programmes (Sharpe & Lydiate, 2003). Although breeders have successfully developed many LEA rapeseed cultivars, new sources of LEA are still required to modulate the relevant pathways within the seed oil fatty acid synthesis and modification pathways (Barker *et al.*, 2007). Wu *et al.* (2008) reported discovery of a four base deletion in the *FAE1* gene that resulted in LEA content within six Chinese rapeseed cultivars. However, this new source of LEA may still contain 'Liho' or 'Oro' genetic background, as so many LEA rapeseed cultivars have derived from material including the parent 'Liho' or 'Oro'.

In the work reported here, the WT genotype of our EMS mutant population was Ningyou7, which is a high erucic acid

(HEA) content cultivar in China, distinct from many other modern rapeseed cultivars (Meng *et al.*, 1996). This cultivar has little or no allelic genetic background in common with 'Lih0' or 'Oro', and so the three novel LEA content rapeseed alleles induced by TILLING in this research represent a unique genetic resource. The gene locus-specific primer analysis confirmed that the mutant *a8fae1-599* and *c3fae1-400* were separately associated with *bn8.fae1* and *bnC3.fae1*, respectively. Seed erucic acid content in rapeseed has been shown to be primarily controlled by these two genes (Barret *et al.*, 1998; Qiu *et al.*, 2006; Wu, Y *et al.*, 2007; Wu *et al.*, 2008). Thus a cross between these two mutants would be expected to generate a new LEA cultivar within what is otherwise a HEA background. In conclusion, the three new LEA mutants, L080-1, N004-1 and L297-1, that were discovered by TILLING could be used as a new genetic resource for LEA rapeseed breeding without collateral inbreeding effect and genetic erosion.

In conclusion, we have reported the application of TILLING to the generation and identification of a novel LEA genetic resource for rapeseed improvement. Although TILLING was reported initially as a reverse genetic approach for Arabidopsis genomic research (McCallum *et al.*, 2000), it will increasingly be very powerful to combine these two approaches to crop functional genomics and genetic improvement. Alongside this work focused on TILLING, QTL for rapeseed seed oil and erucic acid content have been detected in the TNDH genetic mapping population, followed by development of near-isogenic lines (NILs) to resolve these two regions (F. Tian & J. Meng, unpublished). Fine mapping of the two QTLs has resulted in *FAE1* and several other genes being proposed as major candidates contributing to seed oil content (unpublished data). To validate gene function for these candidates, TILLING will now be the method of choice in *B. napus*, rather than conventional transgene knock-down approaches. Knockout of candidate gene function via TILLING overcomes complications associated with RNAi or anti-sense in a multi-paralogous system, and has the advantage of greatly reducing the time required to carry out such research. Taken together, we propose that the combination of QTL resolution making use of comparative genomic information and TILLING of candidates provides a powerful new approach to functional crop genomics, with the advantages of both forward and reverse genetics, and that this will accelerate genetic crop improvement.

## Acknowledgements

The authors thank to Dr Andy Phillips, Dr Hai-Chun Jing and Mr Carlos Bayon (Rothamsted Research, Harpenden, Herts AL5 2JQ, UK) for their support in TILLING technique training to NW. The authors thank Dr Ian Bancroft (Department of Crop Genetics John Innes Centre, Norwich Research Park Colney Lane, Norwich NR4 7UH, UK) for his con-

structive suggestion to this work. The authors thank Mr Dianrong Li (Hybrid Rapeseed Centre of Shaanxi, Dali 715105, China) for his contribution to the field work and NW thanks graduate students within his laboratory for their contribution to the work of mutant population DNA extraction. Financial support for this work was provided by the National Basic Research and Development Program (2006CB101600), the Key Project of Chinese Ministry of Education (103175), and the National Nature Science Foundation of China (30600373).

## References

- Barker GC, Larson TR, Graham IA, Lynn JR, King GJ. 2007. Novel insights into seed fatty acid synthesis and modification pathways from genetic diversity and quantitative trait loci analysis of the *Brassica* C genome. *Plant Physiology* 144: 1827–1842.
- Barret P, Delourme R, Renard M, Domergue F, Lessire R, Delseny M, Roscoe TJ. 1998. A rapeseed *FAE1* gene is linked to the E1 locus associated with variation in the content of erucic acid. *Theoretical and Applied Genetics* 96: 177–186.
- Bendahmane A. 1999. Zero-background plasmid vector for BAC library construction. *Biotechniques* 26: 228–230, 232.
- Caldwell DG, McCallum N, Shaw P, Muehlbauer GJ, Marshall DF, Waugh R. 2004. A structured mutant population for forward and reverse genetics in Barley (*Hordeum vulgare* L.). *Plant Journal* 40: 143–150.
- Colbert T, Till BJ, Tompa R, Reynolds S, Steine MN, Yeung AT, McCallum CM, Comai L, Henikoff S. 2001. High-throughput screening for induced point mutations. *Plant Physiology* 126: 480–484.
- Comai L, Young K, Till BJ, Reynolds SH, Greene EA, Codomo CA, Enns LC, Johnson JE, Burtner C, Odden AR *et al.* 2004. Efficient discovery of DNA polymorphisms in natural populations by Ecotilling. *Plant Journal* 37: 778–786.
- Conte LS, Leoni O, Palmieri S, Capella P, Lercker G. 1989. Half-seed analysis: rapid chromatographic determination of the main fatty acids of sunflower seed. *Plant Breeding* 102: 158.
- Cooper JL, Till BJ, Laport RG, Darlow MC, Kleffner JM, Jamai A, El-Mellouki T, Liu S, Ritchie R, Nielsen N *et al.* 2008. TILLING to detect induced mutations in soybean. *BMC Plant Biology* 8: 9.
- Downey RK, Craig BM. 1964. Genetic control of fatty acid biosynthesis in rapeseed (*Brassica napus* L.). *Journal of the American Oil Chemists' Society* 41: 475–478.
- Gibbons IR, Asai DJ, Ching NS, Dolecki GJ, Mocz G, Phillipson CA, Ren H, Tang WJ, Gibbons BH. 1991. A PCR procedure to determine the sequence of large polypeptides by rapid walking through a cDNA library. *Proceedings of the National Academy of Sciences, USA* 88: 8563–8567.
- Greene EA, Codomo CA, Taylor NE, Henikoff JG, Till BJ, Reynolds SH, Enns LC, Burtner C, Johnson JE, Odden AR *et al.* 2003. Spectrum of chemically induced mutations from a large-scale reverse-genetic screen in Arabidopsis. *Genetics* 164: 731–740.
- Heckmann AB, Lombardo F, Miwa H, Perry JA, Bunnewell S, Parniske M, Wang TL, Downie JA. 2006. *Lotus japonicus* nodulation requires two GRAS domain regulators, one of which is functionally conserved in a nonlegume. *Plant Physiology* 142: 1739–1750.
- Henikoff S, Till BJ, Comai L. 2004. TILLING. Traditional mutagenesis meets functional genomics. *Plant Physiology* 135: 630–636.
- James DW Jr, Lim E, Keller J, Plooy I, Ralston E, Dooner HK. 1995. Directed tagging of the Arabidopsis *FATTY ACID ELONGATION1* (*FAE1*) gene with the maize transposon activator. *Plant Cell* 7: 309–319.
- Kaushik N, Agnihotri A. 2000. GLC analysis of Indian rapeseed-mustard to study the variability of fatty acid composition. *Biochemical Society Transactions* 28: 581–583.



- Kim Y, Schumaker KS, Zhu JK. 2006. EMS mutagenesis of Arabidopsis. *Methods in Molecular Biology* 323: 101–103.
- Konzak CF, Nilan RA, Kleinhofs A. 1976. Artificial mutagenesis as an aid in overcoming genetic vulnerability of crop plants. *Basic Life Sciences* 8: 163–177.
- Lassner MW, Lardizabal K, Metz JG. 1996. A jojoba beta-Ketoacyl-CoA synthase cDNA complements the canola fatty acid elongation mutation in transgenic plants. *Plant Cell* 8: 281–292.
- Li M, Zhang C, Liu L, Yu L. 2005. Development of relationship between A, B and C genomes in *Brassica* genera. *Hereditas (Beijing)* 27: 671–676.
- Long D, Coupland G. 1998. Transposon tagging with Ac/Ds in Arabidopsis. *Methods in Molecular Biology* 82: 315–328.
- Long Y, Shi J, Qiu D, Li R, Zhang C, Wang J, Hou J, Zhao J, Shi L, Park B-S *et al.* 2007. Flowering time quantitative trait loci analysis of oilseed *Brassica* in multiple environments and genome-wide alignment with Arabidopsis. *Genetics* 177: 2433–2444.
- Lysak MA, Koch MA, Pecinka A, Schubert I. 2005. Chromosome triplication found across the tribe *Brassicaceae*. *Genome Research* 15: 516–525.
- Mayer K, Murphy G, Tarchini R, Wambutt R, Volckaert G, Pohl T, Dusterhoft A, Stiekema W, Entian K-D, Terryn N *et al.* 2001. Conservation of microstructure between a sequenced region of the genome of rice and multiple segments of the genome of *Arabidopsis thaliana*. *Genome Research* 11: 1167–1174.
- McCallum CM, Comai L, Greene EA, Henikoff S. 2000. Targeted screening for induced mutations. *Nature Biotechnology* 18: 455–457.
- Meng J, Tian Z, Fu T, Qian X. 1996. Analysis genetics diversity of *Brassica napus* L. with RFLP markers. *Acta Genetica Sinica* 23: 293–306.
- Metzger JO, Bornscheuer U. 2006. Lipids as renewable resources: current state of chemical and biotechnological conversion and diversification. *Applied Microbiology and Biotechnology* 71: 13–22.
- Mika V, Tillmann P, Koprna R, Nerusil P, Kucera V. 2003. Fast prediction of quality parameters in whole seeds of oilseed rape (*Brassica napus* L.). *Plant, Soil and Environment* 49: 141–145.
- Ng PC, Henikoff S. 2003. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Research* 31: 3812–3814.
- Oleykowski CA, Bronson Mullins CR, Godwin AK, Yeung AT. 1998. Mutation detection using a novel plant endonuclease. *Nucleic Acids Research* 26: 4597–4602.
- Ostergaard L, King GJ. 2008. Standardized gene nomenclature for the *Brassica* genus. *Plant Methods* 4: 10.
- Park JY, Koo DH, Hong CP, Lee SJ, Jeon JW, Lee SH, Yun PY, Park BS, Kim HR, Bang JW *et al.* 2005. Physical mapping and microsynteny of *Brassica rapa* ssp. *Pekinensis* genome corresponding to a 222 kbp gene-rich region of Arabidopsis chromosome 4 and partially duplicated on chromosome 5. *Molecular Genetics and Genomics* 274: 579–588.
- Qiu D, Morgan C, Shi J, Long Y, Liu J, Li R, Zhuang X, Wang Y, Tan X, Dietrich E *et al.* 2006. A comparative linkage map of oilseed rape and its use for QTL analysis of seed oil and erucic acid content. *Theoretical and Applied Genetics* 114: 67–80.
- Rana D, van den Boogaart T, O'Neill CM, Hynes L, Bent E, Macpherson L, Park JY, Lim YP, Bancroft I. 2004. Conservation of the microstructure of genome segments in *Brassica napus* and its diploid relatives. *Plant Journal* 40: 725–733.
- Roscoe TJ, Lessire R, Puyaubert J, Renard M, Delseny M. 2001. Mutations in the *fatty acid elongation 1* gene are associated with a loss of beta-ketoacyl-CoA synthase activity in low erucic acid rapeseed. *FEBS Letters* 492: 107–111.
- Sharpe AG, Lydiat DJ. 2003. Mapping the mosaic of ancestral genotypes in a cultivar of oilseed rape (*Brassica napus*) selected via pedigree breeding. *Genome* 46: 461–468.
- Slade AJ, Fuerstenberg SI, Loeffler D, Steine MN, Facciotti D. 2005. A reverse genetic, nontransgenic approach to wheat crop improvement by TILLING. *Nature Biotechnology* 23: 75–81.
- Taylor NE, Greene EA. 2003. PARSESNP: a tool for the analysis of nucleotide polymorphisms. *Nucleic Acids Research* 31: 3808–3811.
- Till BJ, Burtner C, Comai L, Henikoff S. 2004a. Mismatch cleavage by single-strand specific nucleases. *Nucleic Acids Research* 32: 2632–2641.
- Till BJ, Cooper J, Tai TH, Colowit P, Greene EA, Henikoff S, Comai L. 2007. Discovery of chemically induced mutations in rice by TILLING. *BMC Plant Biology* 7: 19.
- Till BJ, Reynolds SH, Weil C, Springer N, Burtner C, Young K, Bowers E, Codomo CA, Enns LC, Odden AR *et al.* 2004b. Discovery of induced point mutations in maize genes by TILLING. *BMC Plant Biology* 4: 12.
- Till BJ, Zerr T, Comai L, Henikoff S. 2006. A protocol for TILLING and Ecotilling in plants and animals. *Nature Protocols* 1: 2465–2477.
- Town CD, Cheung F, Maiti R, Crabtree J, Haas BJ, Wortman JR, Hine EE, Althoff R, Arbogast TS, Tallon LJ *et al.* 2006. Comparative genomics of *Brassica oleracea* and *Arabidopsis thaliana* reveal gene loss, fragmentation, and dispersal after polyploidy. *Plant Cell* 18: 1348–1359.
- U N. 1935. Genome-analysis in *Bassica* with special reference to the experimental formation of *B. napus* and its peculiar mode of fertilization. *Japanese Journal of Botany* 7: 389–452.
- Vles RO, Bijster GM, Timmer WG. 1978. Nutritional evaluation of low-erucic-acid rapeseed oils. *Archives of Toxicology Suppl* 1: 23–32.
- Waterhouse PM, Graham MW, Wang MB. 1998. Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA. *Proceedings of the National Academy of Sciences, USA* 95: 13959–13964.
- Wu G, Wu Y, Xiao L, Li X, Lu C. 2008. Zero erucic acid trait of rapeseed (*Brassica napus* L.) results from a deletion of four base pairs in the *fatty acid elongase 1* gene. *Theoretical and Applied Genetics* 116: 491–499.
- Wu GZ, Shi QM, Niu Y, Xing MQ, Xue HW. 2007a. Shanghai RAPESEED Database: a resource for functional genomics studies of seed development and fatty acid metabolism of *Brassica*. *Nucleic Acids Research* 36: 1044–1047.
- Wu Y, Xiao L, Wu G, Lu C. 2007b. Cloning of *fatty acid elongase1* gene and molecular identification of A and C genome in *Brassica* species. *Science in China. Series C, Life Sciences* 50: 343–349.
- Yang T-J, Kim JS, Kwon S-J, Lim K-B, Choi B-S, Kim J-A, Jin M, Park JY, Lim M-H, Kim H-I *et al.* 2006. Sequence-level analysis of the diploidization process in the triplicated flowering locus C region of *Brassica rapa*. *Plant Cell* 18: 1339–1347.