

# A Gene Cluster Responsible for Alkylaldoxime Metabolism Coexisting with Nitrile Hydratase and Amidase in *Rhodococcus globerulus* A-4<sup>†,‡</sup>

Sheng-Xue Xie,<sup>§</sup> Yasuo Kato, Hidenobu Komeda, Satoshi Yoshida, and Yasuhisa Asano\*

Biotechnology Research Center, Faculty of Engineering, Toyama Prefectural University,  
5180 Kurokawa, Kosugi, Toyama 939-0398, Japan

Received June 25, 2003; Revised Manuscript Received August 26, 2003

**ABSTRACT:** An enzyme “alkylaldoxime dehydratase (OxdRG)” was purified and characterized from *Rhodococcus globerulus* A-4, in which nitrile hydratase (NHase) and amidase coexisted with the enzyme. The enzyme contains heme *b* as a prosthetic group, requires reducing reagents for the reaction, and is most active at a neutral pH and at around 30 °C, similar to the phenylacetaldoxime dehydratase from *Bacillus* sp. OxB-1 (OxdB). However, some differences were seen in subunit structure, substrate specificity, and effects of activators and inhibitors. The corresponding gene, *oxd*, encoding a 1059-base pair ORF consisting of 353 codons, was cloned, sequenced, and overexpressed in *Escherichia coli*. The predicted polypeptide showed 30.3% identity to OxdB. The gene is mapped just upstream of the gene cluster encoding the enzymes involved in the metabolism of aliphatic nitriles, i.e., NHase and amidase, and their regulatory and activator proteins. We report here the existence of an aldoxime dehydratase genetically linked with NHase and amidase, and responsible for the metabolism of alkylaldoxime in *R. globerulus*.

Aldoximes derived from amino acids are considered to be intermediates in the biosynthesis of cyanogenic glucosides and glucosinolates in plants (1). However, information on aldoxime metabolism is quite limited and the genetics and enzymology have not been well characterized. One oxime-metabolizing enzyme (cytochrome P450 CYP71E1) has been reported to catalyze the conversion of aldoxime to  $\alpha$ -hydroxynitrile in the pathway for biosynthesis of cyanogenic glucoside dhurrin in *Sorghum bicolor* (2–4). Other cytochrome P450s, namely, CYP83 homologues (A1 and B1), have also been identified as oxime-metabolizing enzymes, which catalyze the conversion of indoleacetaldoxime to the corresponding *aci*-nitro compound, the first step in the biosynthesis of indole glucosinolates in *Arabidopsis thaliana*. However, the level of activity is quite low, and the mechanisms involved have not been studied. Indoleacetaldoxime is known to be a metabolic branch point between the production of indoleacetic acid and indole glucosinolates in *A. thaliana* (5–7), but the enzymes responsible for the metabolism have yet to be purified and characterized.

Asano et al. have isolated various nitrile-degrading microorganisms, e.g., *Rhodococcus rhodochrous* (formerly *Arthrobacter* sp.) strains J-1 and I-9 (K22, AKU 629) (8)

and *Pseudomonas chlororaphis* B23 (9). They first purified, characterized, and named nitrile hydratase (NHase, EC. 4.2.1.84) from *R. rhodochrous* J-1 (10–12). They also found that *P. chlororaphis* B23 accumulates large quantities of amides from nitriles and is suitable for the industrial production of acrylamide from acrylonitrile (9, 12). Moreover, nicotinamide and 5-cyanovaleramide are also industrially produced by NHase (13, 14). Despite its important uses, the physiological function of NHase in nature remains unclear.

We have studied the metabolism of aldoximes from a physiological as well as an applicative perspective, and have isolated *Bacillus* sp. OxB-1 and *Rhodococcus* sp. YH3-3, which can convert aldoximes to the corresponding carboxylic acids, from soil samples. The reaction occurs via intermediary nitriles and involves a combination of enzymes including aldoxime dehydratase and nitrile-hydrolyzing enzymes such as nitrilase (Nit),<sup>1</sup> and NHase, and/or amidase (15–17) (Figure 1). We previously reported the purification and characterization of phenylacetaldoxime dehydratase (OxdB) from *Bacillus* sp. OxB-1 and molecular cloning of the gene (*oxd*) (18). The enzyme was overexpressed in *E. coli* JM109 (19) and in a recombinant *E. coli* JM109/pOxD-90F strain employed in the enzymatic synthesis of various arylalkyl- and alkyl-nitriles from the corresponding aldoximes (20). The gene coding OxdB was linked with that for Nit in the genome of the *Bacillus* strain. We have also shown that aldoxime dehydratase activity coexists not only with Nit but also with

<sup>†</sup> This work was supported in part by a Grant-in-Aid from the Japan Society for the Promotion of Science (JSPS) for Foreign Researchers (P00319 to S.-X. Xie) and a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan.

<sup>‡</sup> The nucleotide sequence reported in this paper has been submitted to the GenBank/EMBL/DBJ Data Bank with accession number AB105912.

\* To whom correspondence should be addressed. Phone: +81-766-56-7500. Fax: +81-766-56-2498. E-mail: asano@pu-toyama.ac.jp.

<sup>§</sup> Postdoctoral fellow supported by the Japan Society for the Promotion of Science (JSPS). Present address: The High Throughput Screening Laboratory, University of Kansas, 1501 Wakarusa Dr., Lawrence, KS 66047.

<sup>1</sup> Abbreviations: OxdRG, alkylaldoxime dehydratase from *Rhodococcus globerulus* A-4; OxdB, phenylacetaldoxime dehydratase from *Bacillus* sp. OxB-1; Nit, nitrilase; NHase, nitrile hydratase; PAOx, phenylacetaldoxime; PyOx, pyridine-3-aldoxime; PPOx, phenylpropionaldoxime; VOx, *n*-valeraldoxime; IPTG, isopropyl- $\beta$ -D-thiogalactopyranoside; DMSO, dimethyl sulfoxide; ORF, open reading frame.




FIGURE 1: Microbial metabolism of aldoximes via the "aldoxime-nitrile" pathway. *Bacillus* sp. OxB-1 metabolizes Z-phenylacetaldoxime (Z-PAOx, R = PhCH<sub>2</sub>) to form phenylacetnitrile using PAOx dehydratase (OxdB), which is subsequently converted to phenylacetic acid by nitrilase. *Rhodococcus globerulus* A-4 metabolizes aldoximes such as *E/Z*-valeraldoxime (VOx, R = CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>-) and Z-PAOx to the corresponding nitrile by the action of alkylaldoxime dehydratase, which is converted to the corresponding amide by nitrile hydratase, and the amide is then hydrolyzed to the corresponding acid by amidase.

NHase and amidase in various bacteria, yeasts, and fungi (20). Until now, however, aldoxime dehydratase has not been purified from a strain having NHase and amidase. It is of interest to show the possible enzymological and genetic relationships between aldoxime dehydratase and NHase/amidase.

In this study, we examined the specificity of enzymes responsible for the metabolism of aldoximes and nitriles among newly isolated strains and stock cultures. Strain A-4, identified as *R. globerulus* exhibiting strong aldoxime dehydratase activity with a wide range of substrates together with NHase/amidase activity, was selected for the purification and characterization of an alkylaldoxime dehydratase. Its gene was cloned and overexpressed in *E. coli*, and its flanking region was sequenced. We found that the gene exists together with the NHase and amidase genes in the genome of *R. globerulus* A-4, providing the first evidence of a genetic relationship between aldoxime dehydratase and NHase/amidase and the metabolism of alkylaldoximes.

## MATERIALS AND METHODS

**Materials.** DEAE-Toyopearl 650 M, Butyl-Toyopearl 650 M, and HPLC columns G-3000SW and ODS-80TS were purchased from Tosoh Corp. (Tokyo, Japan). Gigapite was obtained from Seikagaku Kogyo (Tokyo, Japan). The FPLC columns Superdex 200 HR 10/30, MonoQ HR10/10, and Phenyl-Superose HR 5/5 were from Amersham Biosciences (Uppsala, Sweden). Aldoximes were synthesized from the corresponding aldehydes using the method demonstrated previously (15, 16, 22). Other chemicals were from commercial sources and were used without further purification.

**Analytical Methods.** The molecular weight ( $M_r$ ) of the enzyme was measured by HPLC on a TSK G3000SW column (7.5 × 600 mm) or by SDS-PAGE as described previously (18, 23). SDS- and native-PAGE and protein-concentration measurements were performed as described (24, 25). Heme staining was carried out using the method described by Goodhew C. F. et al (26). The absorption spectra were recorded on a HITACHI U-3210 spectrophotometer (Tokyo, Japan). The heme concentration was determined by the pyridine hemochromogen method (27) using the molecular extinction coefficient of the  $\alpha$ -peak at 557 nm of the hemochromogen of protoheme IX ( $\epsilon = 34.4 \text{ mM}^{-1} \text{ cm}^{-1}$ ) (28). To determine the N-terminal amino acid sequence, purified enzyme was blotted from the gel after SDS-PAGE (12.5% acrylamide) on a poly(vinylidene difluoride) membrane (PVDF, Bio-Rad, Hercules, CA) with the trans-blot semi-dry electrophoresis transfer cell (Nippon Eido,

Tokyo, Japan), analyzed with HP G1005A Protein Sequencing Systems (Hewlett-Packard Co., Palo Alto, CA).

**Enzyme Assay.** The standard assay mixture comprised, in 200  $\mu\text{L}$ , 0.1 M potassium phosphate buffer (KPB, pH 7.0), 5 mM of aldoxime, 10% (v/v) dimethyl sulfoxide (DMSO), and enzyme solution with 1 mM Na<sub>2</sub>S. After the enzyme reaction was run at 30 °C for 10–30 min, during which the enzyme reaction proceeded linearly, the reaction was stopped by adding an equal volume of 0.5 M H<sub>3</sub>PO<sub>4</sub> followed by vortexing. The mixture was analyzed by GLC or HPLC as described previously (18, 20, 21). One unit of enzyme activity is defined as the amount of enzyme that catalyzes the dehydration of aldoxime into nitrile at a rate of 1  $\mu\text{mol}$  per min under the assay conditions.

**Screening for Aldoxime-Degrading Microorganisms.** Microorganisms stocked in our laboratory and isolates from soil such as aldoxime or nitrile degraders were used for screening. TGY medium, which comprised 0.5% Bacto Tryptone (Difco, Detroit, MI), 0.5% yeast extract (Difco), 0.1% glucose, and 0.1% K<sub>2</sub>HPO<sub>4</sub>, pH 7.0, was used to culture microorganisms. Each microorganism was inoculated in a test tube (16.5 × 165 mm) containing 5 mL of TGY medium with 0.05% of various aldoximes as inducers, and then incubated at 30 °C for 12–36 h with reciprocal shaking (200 strokes/min). Five milliliters of seed culture was transferred to a 2 L Sakaguchi flask containing 500 mL of the same medium, followed by incubation at 30 °C for 12–24 h with reciprocal shaking (96 strokes/min). The cells were harvested by centrifugation at 15200g for 20 min at 4 °C, and washed with 0.85% saline and 50 mM KPB (pH 7.0). Each reaction mixture comprised 5% (w/v) of wet cells, 0.1 M KPB (pH 7.0), and 50–100 mM of substrate such as *E*-pyridine-3-aldoxime (*E*-PyOx), Z-PAOx, Z-3-phenylpropionaldoxime (Z-3-PPOx), *E/Z*-2-phenylpropionaldoxime (*E/Z*-2-PPOx), or *E/Z*-isobutyraldoxime in a total volume of 400  $\mu\text{L}$ . The reaction was run for 2 h at 30 °C, and then two volumes of ethyl acetate or one volume of 0.5 M H<sub>3</sub>PO<sub>4</sub> was added to stop it. The product formation was analyzed by GC or HPLC.

**Identification of Strain A-4.** The taxonomic characteristics of the strain were examined by the method described in *Bergey's Manual of Systematic Bacteriology* (29). Fatty acid composition of the cells was analyzed by GLC and evaluated by Sherlock Microbial Identification System (MIDI, Inc., Newark, DE) (30). Sequencing of the PCR-amplified 16S rDNA-500 was done by ABI PRISM377 DNA sequencer (Applied Biosystems, Foster City, CA) and compared with a database of MicroSeq Bacterial 500 Library V.0023 (Applied Biosystems).

*Purification of an Aldoxime Dehydratase from R. globerulus A-4.* All purification procedures were performed at temperatures lower than 5 °C. Twenty millimolar KPb (pH 7.0) containing 0.2 mM dithiothreitol and 5 mM 2-mercaptoethanol was used throughout the purification. Centrifugation was done at 19700g for 20 min unless otherwise specified. Washed cells (294 g of wet weight) of *R. globerulus* A-4 obtained from 100 L of TGY culture containing 0.05% E-PyOx were suspended in 2.5 L of the buffer, and then disrupted for 3 × 20 min at 4 °C by an Insonator model 201M (9 kHz, Kubota-Shoji, Tokyo, Japan). After centrifugation, the resulting supernatant was fractionated with solid (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. The precipitate obtained at 30–60% saturation was collected, dissolved, and dialyzed against four changes of 20 L of the buffer. The dialysate was put on a DEAE-Toyopearl column (6.3 × 40 cm) and eluted with a linear gradient of 0–1 M NaCl in 8 L of buffer and the active fractions were combined. After the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> concentration had been adjusted to 20% saturation, the enzyme solution was loaded onto a Butyl-Toyopearl column (5.7 × 20 cm) and eluted by lowering the ionic strength of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> linearly from 20% saturation to 0% in 5 L of buffer. The active fractions were collected and dialyzed against four changes of 10 L of buffer. The dialysate was collected and concentrated with a Centriprep YM-10 membrane (Millipore, MA), and then added to a Gigapite column (5.6 × 25 cm). The enzyme was eluted with a linear gradient of 0–5 M KPb (pH 7.0) in 4 L of the buffer. The active fractions were dialyzed, concentrated, and put on a second DEAE-Toyopearl column (2.8 × 45 cm). The enzyme was eluted with a linear gradient of 0.1–0.5 M NaCl in 1.5 L of buffer. The enzyme solution was again purified with a second Butyl-Toyopearl chromatography column (2.8 × 28 cm) using a linear gradient of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (15 to 0% saturation in 1 L of buffer). The dialyzed active fraction was purified with a Superdex 200 HR 10/30 column using buffer containing 0.15 M NaCl. The active fractions were dialyzed, concentrated, then loaded onto a MonoQ HR10/10 column and eluted with a linear gradient of 0–1 M NaCl in buffer. After adding NaCl to a concentration of 4 M, the active enzyme solution was applied to a Phenyl-Superose HR 5/5 column and eluted with a linear gradient of NaCl (4 to 0 M). The active fractions were collected, dialyzed, and concentrated for enzyme characterization.

*Bacterial Strains, Plasmids, and Culture Conditions.* *R. globerulus* A-4 was used as a source of chromosomal DNA throughout this study. *E. coli* JM109 (recA1, endA1, gyrA96, thi, hsdR 17, supE44, relA1, Δ(lac-proAB)/F' [traD36, proAB<sup>+</sup>, lacI<sup>q</sup>, lacZΔM15]) was used as a host strain for gene cloning and in DNA manipulations. Plasmids pBluescriptII SK- and pUC19 were employed as a cloning vector and for sequencing analysis. Plasmid DNA was isolated with an automatic plasmid isolation system PI-100 (Kurabo, Osaka, Japan) or by using a QIAGEN plasmid purification kit (Valencia, CA). Recombinant *E. coli* cells were cultivated at 25, 30, or 37 °C in Luria-Bertani (LB) medium (1% Bacto Tryptone, 0.5% Bacto yeast extract, and 1% NaCl, pH 7.0) containing 100 μg/mL of ampicillin. Terrific medium (1.2% Bacto Tryptone, 2.4% Bacto yeast extract, 0.4% glycerol, 17 mM KH<sub>2</sub>PO<sub>4</sub>, and 72 mM K<sub>2</sub>HPO<sub>4</sub>, pH 7.3) was used for overexpressing the enzyme. To induce the *lac*-promoter,

1 mM of isopropyl-β-D-thiogalactopyranoside (IPTG) was added to the medium.

*Cloning of oxd and Its Flanking Sequences.* The sense primer ATGGARWSSGCNATCGGNGAR and antisense primer CTG GTGSARSGTRTCRCNGC were synthesized on the basis of the sequences of MESAIGE and ADDTLHQ, respectively, which were part of the *N*-terminal amino acid sequence of the purified aldoxime dehydratase. Degenerate positions are indicated by R for A or G, W for A or T, S for C or G, and N for all bases. The genomic DNA of *R. globerulus* A-4 was isolated as described previously (31) and used as a template for PCR amplification with a PTC-200 thermal cycler (MJ Research Inc., Watertown, MA). Reaction mixtures contained 2.5 ng of chromosomal DNA, 100 pmol of each primer, and 1 unit of *Thermus aquaticus* DNA polymerase (EX Taq, Takara, Japan) in a volume of 50 μL. Thirty-five thermal cycles were employed, each consisting of 95 °C for 0.5 min, 55 °C for 1.5 min, and 72 °C for 2.5 min. The gel-purified PCR product (~120-bp) was cloned into pT7Blue vector (Novagen, Darmstadt, Germany) to construct a plasmid pOxdRGNS, and then introduced into *E. coli* JM109. The PCR product was purified from the plasmid and used as a radiolabeled probe for Southern hybridization to clone the full-length aldoxime dehydratase gene as described previously (18). The 4.5–6.0 kb fragment of the genomic DNA digested with *Bam*HI was purified and inserted into the same site of pBluescript II KS- to construct a genomic library of *R. globerulus* A-4. While colony hybridization was performed, one positive clone, harboring pOxdRGB97, which contains about a 4.7-kb *Bam*HI fragment, was selected for further analysis. The positive fragment was inserted into the *Bam*HI site of pUC19 for sequencing with the Kilo-sequence Deletion kit (Takara). Sequence analysis of the deletion templates was performed by the dideoxy chain termination procedure (32) with an automatic DNA sequencer 4000L (LiCor, Lincoln, NB) or a model 310 (Applied Biosystems). The positive 3.9-kb *Fba*I and 6.3-kb *Pst*I fragments were also obtained in the manner described above. Another 3.7-kb *Hind*III-*Sph*I fragment was cloned by the DNA-probing method with a 0.7-kb *Hind*III-*Pst*I fragment as a probe that was obtained by *Hind*III digestion of the 6.3-kb *Pst*I fragment. The entire 11.6-kb DNA containing *oxd* and its flanking region was sequenced. A homology search was performed with the sequence similarity searching programs FASTA (33) and BLAST (34) and the ClustalW method was used to align the sequences (35). The GENE-TYX software system (Software Development Co., Tokyo, Japan) was used for computer analysis of nucleotide sequences and deduced amino acid sequences.

*Overexpression of oxd Gene Product of R. globerulus A-4 in E. coli.* Restriction mapping of pOxdRGB97, subcloning into the vector pBluescriptII SK-, and subsequent Southern blot hybridization analysis yielded a 1.5-kb *Hinc*II fragment containing *oxd* of *R. globerulus* A-4, which was subcloned into the vector pUC19 to make pOxdRG HII. The 5'-flanking region of the ATG start codon of the *oxd* gene was modified by PCR with pOxdRGHII as a template. PCR was carried out in a reaction mixture (50 μL) containing 25 ng of template DNA, 100 pmol of each primer: for the *N*-terminal, 5'-TCGACCAAGCTTTAAGGAGGAATAGCTCATGGAATCTGCAATCGGCGAACACC-3' (the translation start codon is underlined; ribosome-binding site is in bold;

Table 1: Purification of the Aldoxime Dehydratase from *Rhodococcus globerulus* A-4

step	total protein (mg)	total activity (units)	specific activity <sup>a</sup> (units/mg)	yield (%)	specific activity ratio <sup>b</sup>
cell-free extract	16900	14.3	0.000845	100	0.956
ammonium sulfate	5970	7.96	0.00133	55.7	0.858
DEAE-Toyopearl (I)	2560	4.43	0.00173	31.0	0.924
Butyl-Toyopearl (I)	567	3.25	0.00574	22.7	1.81
Gigapite	347	2.86	0.00826	20.0	3.49
DEAE-Toyopearl (II)	236	2.54	0.0107	17.8	6.73
Butyl-Toyopearl (II)	176	2.37	0.0134	16.6	8.24
Superdex 200 HR 10/30	40.4	1.92	0.0475	13.4	16.3
MonoQ HR 10/10	26.2	1.67	0.0636	11.7	17.7
Phenyl-Superose HR 5/5	8.77	0.772	0.0879	5.41	8.93

<sup>a</sup> The reaction was carried out with 5 mM Z-PAOx as the substrate in the presence of 1 mM Na<sub>2</sub>S. <sup>b</sup> Ratio of specific activities determined with and without Na<sub>2</sub>S.

restriction site is in italics), and for C-terminal, 5'-AA-CATATCTAGAGTTGCCTCCGAGGG-3' (restriction site is in italics), and 0.5 U of *Pwo* polymerase (Roche, Mannheim, Germany). Thirty thermal cycles were employed, each consisting of 94 °C for 0.5 min, 55 °C for 0.5 min, and 72 °C for 2 min. The PCR product was digested with *Hind*III and *Xba*I, and the gel-purified fragment was inserted into the same site of pUC19. The plasmid thus obtained was used to transform *E. coli* JM109 afforded the overexpression plasmid pOxdRGP9. Base sequencing confirmed the nucleotide sequence of the complete amplified gene and showed that no error had been incorporated during PCR.

**Purification of the Recombinant Enzyme.** *E. coli* JM109/pOxdRGP9 was cultured in 3 mL of LB medium containing 100 µg/mL of ampicillin at 37 °C for 12 h. The seed culture was transferred to a 2-L Erlenmeyer flask containing 1.5 L of Terrific medium with 100 µg/mL of ampicillin and 1 mM IPTG, and was incubated at 25 °C for 158 h with a 200-rpm shaking rate. The cells were harvested by centrifugation at 15200g for 20 min, and washed with 0.85% NaCl and 50 mM KPB, pH 7.0. The cells (20 g of wet weight) were sonicated and fractionated with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (20–60% saturation). The dialyzed enzyme was subsequently purified with DEAE-Toyopearl (4.3 × 25 cm), Butyl-Toyopearl (2.8 × 30 cm), MonoQ HR10/10, Phenyl-Superose HR 5/5, Superdex 200 HR 10/30, Gigapite, and MonoQ HR5/5 columns. The active fractions were collected, dialyzed, and concentrated for further analysis.

## RESULTS

**Screening for Aldoxime-Degrading Microorganisms.** Various bacterial strains in TPU stock cultures and isolates from soil were screened for aldoxime- and nitrile-degrading activities. Among them, the unidentified strain A-4, *Rhodococcus* sp. YH3-3 (16), *Corynebacterium* sp. C5 TPU 6015 (21), and *Rhodococcus* sp. N-774 (21) converted not only Z-PAOx, but also substituted alkylaldoximes, such as *E/Z*-2-PPOx and *E/Z*-isobutyraldoxime, and arylaldoxime, *E*-PyOx, to the corresponding nitriles. The nitriles were further degraded to the corresponding carboxylic acids through amides, suggesting that the strains metabolize the aldoximes via the actions of aldoxime dehydratase, NHase, and amidase. The active strains were different from *Bacillus* sp. OxB-1 which did not act on the aryl- and substituted aldoximes and degraded nitriles by the action of Nit. Strain A-4 was selected for further study because it showed the highest levels of activity for aldoxime- and nitrile-degradation. Strain A-4 was

identified as *R. globerulus* based on the following taxonomical characteristics according to *Bergey's Manual of Systematic Bacteriology* (29). The cells were singular, non-spore forming, and gram-positive irregular rods. A typical rod-coccus growth cycle was observed. The colonies were orange on a nutrient agar. Catalase was present and oxidase was not. No acid or gas was produced from typical sugars, such as glucose, ribose, xylose, mannitol, maltose, lactose, and glycogen. Alkaline phosphatase and α-glucosidase were detected. β-Glucuronidase, β-galactosidase, and *N*-acetyl-β-glucosaminidase were absent. Major fatty acids present were tetradecanoic (5.3%), pentadecanoic (5.6%), hexadecanoic (19.1%), 8-*cis*-heptadecenoic (5.7%), 10-methyl-octadecanoic (9.5%), 9-*cis*-octadecenoic (17.6%), and nonadecenoic (6.7%) acids. The partial sequencing of the 16S rDNA showed 99.8% similarity with that of *R. globerulus*.

**Purification of the Wild-Type Aldoxime Dehydratase.** The wild-type enzyme was purified to homogeneity through a 10-step purification procedure as shown in Table 1. The enzyme was purified 104-fold with a yield of 5.4% from the cell-free extract of *R. globerulus* A-4, by measuring the enzyme activity with Z-PAOx as the substrate. The purified enzyme (OxdRG) gave a single band on SDS-PAGE (Figure 2) and a single peak by HPLC on a TSK G-3000 SW column. The *M<sub>r</sub>* of the subunit was estimated to be 42 000 by comparing the mobility on SDS-PAGE to that of standard proteins and the *M<sub>r</sub>* of the native enzyme was 76 200 according to gel filtration chromatography. These results suggest that the enzyme has a dimeric structure. The *N*-terminal 40 amino acid sequence of the native OxdRG protein was determined to be MESAIGEHLQCPRTLTR-RVPDITYTPPFPMWVGRADDTLHQ. It did not show significant similarity to any known proteins including OxdB.

**Optical Spectrum and Prosthetic Group.** The absorption maxima of the purified OxdRG enzyme were at 273, 359, 421, and 549 nm, indicating that the enzyme contains heme. Furthermore, the enzyme was positively stained by heme staining on native-PAGE gel (26). The pyridine hemochrome was prepared by mixing the enzyme with alkaline-pyridine for examining the properties of the heme in the enzyme. The absorption peaks of the hemochromogen occurred at 417 (Soret), 524 (α-band), and 556 (β-band) (Figure 3A, dotted line), which are characteristic of the reduced pyridine hemochrome of protoheme IX (28). The addition of a few crystals of Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> caused the increase of each band of the hemochromogen without any changes in their absorbance wavelength (Figure 3A, solid line). These results indicate




FIGURE 2: SDS-PAGE of the aldoxime dehydratase purified from *R. globerulus* A-4. Forty micrograms of protein were loaded onto the lane. Lane A, purified enzyme. Lane B, marker proteins: phosphorylase ( $M_r = 97\,400$ ), bovine serum albumin (66 267), ovalbumin (45 000), carbonic anhydrase (30 000), soybean trypsin inhibitor (20 100), and  $\alpha$ -lactalbumin (14 400).

that the heme iron was present in a reduced form. The heme prosthetic group could be extracted from the enzyme by HCl/acetone-treatment (28) and the extract then concentrated in vacuo. The pyridine hemochrome of the extracted heme exhibited the oxidized-type spectrum of protoheme IX (Figure 3B, dotted line) and the same spectrum as that in Figure 3A was obtained on adding  $\text{Na}_2\text{S}_2\text{O}_4$  (Figure 3B, solid line). The heme content of the enzyme was calculated to be 0.37 mol heme/mol enzyme determined from the spectrum of its pyridine hemochromogen. The value is quite close to that of OxdB, 0.33 (18). The absorption spectra of the hemochromogen were measured with several types of compounds. In the presence of CO, the peaks of Soret, the  $\alpha$ -band, and the  $\beta$ -band decreased and a new shoulder at 430 nm appeared. On the other hand, no significant change in the spectrum was observed with KCN and various aldoximes and nitriles.

**Effects of Reducing Reagents on the Enzyme Activity.** The purified enzyme activity was remarkably enhanced by the addition of reducing reagents to the assay. The activity for Z-PAOx dehydration was enhanced 8.35-, 7.51-, 8.13-, 5.84-, 2.13-, 4.06-, 2.39-, and 3.93-fold in the presence of 1 mM of  $\text{Na}_2\text{S}$ ,  $\text{Na}_2\text{SO}_3$ ,  $\text{Na}_2\text{S}_2\text{O}_4$ ,  $\text{Na}_2\text{S}_2\text{O}_5$ , 2-mercaptoethanol, thioglycerol, L-cysteine, and cysteamine, respectively. The activity was not increased by addition of  $\text{NaHSO}_3$ ,  $\text{Na}_2\text{SO}_4$ ,  $\text{NaHSO}_4$ , or  $\text{Na}_2\text{S}_2\text{O}_7$ . Due to ease of handling, 1 mM of  $\text{Na}_2\text{S}$  was added to the assay mixtures. The apparent  $K_m$  value for  $\text{Na}_2\text{S}$  was estimated to be 40.8  $\mu\text{M}$ . No further enhancement of the activity was seen when reducing reagents coexisted with  $\text{Na}_2\text{S}$  in the assay mixture. The activity was not enhanced when the assay was carried out anaerobically, conditions under which OxdB activity was increased over 5-fold (18).




FIGURE 3: Absorption spectra of the pyridine hemochromogen of (A) native- and (B) extracted-heme of the aldoxime dehydratase from *R. globerulus* A-4. (A) 150  $\mu\text{L}$  of the enzyme solution (4.0 mg/mL) in 20 mM potassium phosphate buffer (pH 7.0) was mixed with 450  $\mu\text{L}$  of pyridine containing 0.1 N KOH. (B) The heme prosthetic group was extracted from 150  $\mu\text{L}$  of the enzyme (4.0 mg/mL) by HCl/acetone-treatment then concentrated in vacuo, and the extracted heme was mixed with 600  $\mu\text{L}$  of 50% pyridine containing 0.1 N KOH. Solid and dotted lines represent results obtained in the presence and absence of  $\text{Na}_2\text{S}_2\text{O}_4$ , respectively.

The activity was increased 1.7-, 1.2-, and 1.1-fold by 1 mM of the flavins, FMN, FAD, and riboflavin, respectively. Furthermore, the activity was increased 2.4-, 4.5-, and 4.2-fold, respectively, when  $\text{Na}_2\text{S}$  was omitted from the assay. The following coenzymes did not affect the reaction: pantothenate, phosphoenolpyruvate, NaF, pyridoxal, pyridoxal-5'-phosphate (PLP), ascorbate, dehydroascorbate, glu-

Table 2: Kinetic Parameters of the Enzyme for Various Aldoximes

aldoxime	$K_m$ (mM)	$V_{max}$ (units/mg)	$V_{max}/K_m$ (units/mg/mM)	relative act (%) <sup>a</sup>
Alkylalldoxime				
<i>E/Z</i> -cyclohexanecarboxaldehyde oxime	1.13	0.386	0.342	100
<i>E/Z</i> -propionaldoxime	5.13	0.430	0.084	7.76
<i>E/Z</i> - <i>n</i> -butyraldoxime	1.73	0.689	0.398	48.9
<i>E/Z</i> -isobutyraldoxime	5.54	0.041	0.007	26.5
<i>E/Z</i> -VOx	1.13	1.64	1.45	39.3
<i>E/Z</i> -isovaleraldoxime	3.97	0.239	0.060	58.6
<i>E/Z</i> - <i>n</i> -capronaldoxime	2.94	1.66	0.565	64.2
<i>E/Z</i> -isocapronaldoxime	6.76	1.32	0.195	88.3
Arylalkylalldoxime				
Z-PAOx	1.40	0.140	0.100	26.6
<i>E/Z</i> -2-PPOx	11.9	0.810	0.068	5.23
Z-3-PPOx	2.31	0.392	0.170	67.3
<i>E/Z</i> -4-phenylbutyraldoxime	ND <sup>b</sup>	ND <sup>b</sup>	ND <sup>b</sup>	5.23
<i>E/Z</i> -indoleacetaldoxime	3.91	0.281	0.072	7.29
<i>E/Z</i> -mandelalldoxime	3.23	0.572	0.177	3.09
Arylalldoxime				
<i>E</i> -PyOx	20.0	0.065	0.003	0.781

<sup>a</sup> Substrate concentration was 5 mM. Low levels of activity (0.04–0.8%) were seen for *E/Z*-cinnamalldoxime, *Z*-*p*-chlorophenylacetalldoxime, *Z*-naphthoacetalldoxime, *E*-thiophene-2-carboxalldoxime, *E*-thiophene-2-acetalldoxime, *E*-benzalldoxime, *E*-*p*-chlorobenzalldoxime, *E*-furfurylalldoxime, and *E*-*p*-tolualldoxime. <sup>b</sup> ND: not determined.

tathione, glutathione disulfide, NAD(P)(H), CoA-SH, AMP, ADP, ATP, GMP, GDP, GTP, IMP, IDP, and ITP.

**Effects of pH and Temperature on the Enzyme Activity and Stability.** The effects of pH and temperature on the enzyme activity and stability were measured in 0.1 M buffers at various pHs: AcOH/AcONa, pH 3.5–6.0; KPB, pH 6.0–8.5; Tris/HCl, pH 7.5–9.0; ethanolamine/HCl, pH 8.5–11.0; NH<sub>4</sub>Cl/NH<sub>4</sub>OH, pH 8.0–10.5; and glycine/NaCl/NaOH, pH 8.5–13.0, or at various temperatures between 20 and 80 °C in 0.1 M KPB, pH 8.0. The enzyme exhibited the highest activity in KPB, pH 8.0, at 30 °C. About 65% of the activity remained after incubation at 40 °C for 30 min in 0.1 M KPB, pH 8.0. The activity was maintained at approximately 75–99% between pH 6.0 and 9.5 after incubation at 30 °C for 30 min.

**Substrate Specificity and Kinetic Properties.** The substrate specificity of the purified enzyme was examined with various “synthetic” alldoximes as shown in Table 2. The enzyme was active toward various arylalkyl- and alkyl-alldoximes, and to a lesser extent to arylalldoximes, converting them to the corresponding nitriles. Since all the tested arylalldoximes were of the *E*-form, we synthesized *Z*-PyOx as described previously (22) and tested it as a substrate. However, we could not detect any activity with this alldoxime, suggesting that the enzyme does not utilize arylalldoximes to any great extent regardless of the geometry about the CN double bond. The enzyme also acted on *E/Z*-2-PPOx, *E/Z*-mandelalldoxime, and *E/Z*-isobutyralldoxime, all of which have a substituent group at the alldoxime  $\alpha$ -site although relative activity levels were low. The following compounds were inert as substrates: *E/Z*-*p*-hydroxyphenylacetalldoxime, *E/Z*-diphenylacetalldoxime, *E*-1-naphthoalldoxime, *E*-anisalldoxime, *E*-quinoline-2-carboxaldehyde oxime, *E*-terephthalaldehyde oxime, *E*-isophthalaldehyde oxime, *E*-pyrazinecarboxalldoxime, *Z*-crotonalldoxime, *E/Z*-methacrylalldoxime, *E/Z*-*O*-benzyl-PAOx, *E*-PAOx-hydrazone, *E/Z*-*O*-acetyl-PAOx, *E/Z*-phenylacetone oxime, and *E/Z*-acetophenone oxime. Values for  $V_{max}$  and  $K_m$  were determined using Lineweaver–Burk double reciprocal plots (Table 2). The  $K_m$  values for arylalkyl- and alkyl-alldoximes were nearly the same and much lower than the

value for arylalldoxime, whereas the  $V_{max}$  values for the former except for *E/Z*-isobutyralldoxime were greater than that of the latter. Arylalldoxime was not a good substrate for the enzyme. On the basis of the results, we tentatively name the enzyme “alkylalldoxime dehydratase (EC 4.3.1.-)”.

**Effects of Various Compounds on the Enzyme Activity.** The enzyme activity was assayed under standard conditions in the presence of various metal ions and compounds using *E/Z*-VOx and Z-PAOx as alkyl- and arylalkyl-type substrate, respectively, and compared with that of a control (Table 3). *E/Z*-VOx and Z-PAOx dehydration activity was enhanced by 1 mM of Fe<sup>2+</sup> 2.92- and 5.34-fold, respectively, while 1 mM Fe<sup>3+</sup> also increased the activity, 8.35- and 15.8-fold, respectively. In the absence of Na<sub>2</sub>S, the enhancement by Fe<sup>3+</sup> was 5.29- and 6.32-fold, respectively. The metal ions, Cu<sup>+</sup>, Cu<sup>2+</sup>, Co<sup>2+</sup>, Zn<sup>2+</sup>, and Cd<sup>2+</sup>, strongly inhibited the reaction for both substrates at 1 mM; however, they enhanced the activity when their concentration was low (0.1 mM). The enzyme activity was also inhibited by 1 mM concentrations of heavy metal ions and sulfhydryl reagents. Phenylhydrazine inhibited Z-PAOx dehydration, but not *E/Z*-VOx dehydration, as did Tiron (4,5-dihydroxy-1,3-benzene-disulfonic acid) whereas typical carbonyl reagents, such as hydrazine, KCN, NaN<sub>3</sub>, and NH<sub>2</sub>OH, did not reduce activity for either substrate. The enzyme activity was slightly inhibited by electron donors and carriers such as trimethylhydroquinone, *p*-phenylenediamine, dimethylphenylenediamine, tetramethylphenylenediamine, guaiacol, pyrogallol, miconazole, nitroblue tetrazolium, and phenazine methosulfate, while the electron acceptors, such as vitamin K<sub>3</sub> and duroquinone, were found to enhance the activity.

**Cloning and Sequence Analysis of the *oxd*.** To clone the gene encoding the enzyme, PCR was performed using primers designed on the basis of its *N*-terminal protein sequence with the total genomic DNA of *R. globerulus* A-4 as the template. A single 120-bp fragment was consistently amplified and the putative protein sequence derived from this fragment was identical to the *N*-terminal sequence of the purified enzyme. Using this 120-bp fragment as a probe, genomic Southern hybridization was performed to obtain the

Table 3: Effect of Various Compounds on the Enzyme Activity Measured with *E/Z*-VOx and *Z*-PAOx as Substrates

compound	conc (mM)	relative activity (%) <sup>a</sup>	
		<i>E/Z</i> -VOx	<i>Z</i> -PAOx
none		100	100
Cu <sup>+</sup> , Cu <sup>2+</sup> , Co <sup>2+</sup> , Zn <sup>2+</sup> , Cd <sup>2+</sup>	1	11.5–72.6	13.6–63.8
	0.1	204–368	172–986
Fe <sup>2+</sup>	1	292	534
Fe <sup>3+</sup>	1	835	1,580
	1	529 <sup>b</sup>	632 <sup>b</sup>
Hg <sup>2+</sup>	0.1	71.6	69.7
Sn <sup>2+</sup>	1	39.9	29.0
Tiron	1	80.6	5.31
	0.1	103	64.0
phenylhydrazine	1	72.1	1.60
	0.1	100	26.7
K <sub>3</sub> [Fe(CN) <sub>6</sub> ]	0.1	196	291
	0.1	136 <sup>b</sup>	163 <sup>b</sup>
	1	263	378
duroquinone	1	135	167

<sup>a</sup> The enzyme activity for both substrates was partially (30–80%) inhibited by 1 mM of metal ions such as Al<sup>3+</sup>, V<sup>+</sup>, As<sup>2+</sup>, Nb<sup>5+</sup>, Ag<sup>+</sup>, Hg<sup>2+</sup>, Tl<sup>2+</sup>, Pb<sup>2+</sup>, and penicillamine, thioglycerol, vitamin K<sub>1</sub>, vitamin K<sub>12</sub>, 8-hydroxyquinoline, *o*-phenanthroline, iodoacetate, dithiobis (4-nitrobenzoic acid), *p*-chloromercuribenzoate, diphenylhydantoin, glutathione, glutathione disulfide, phenazinemetosulfate, trimethylhydroquinone, phenylenediamine, tetramethylphenylenediamine, dimethylphenylenediamine, guaiacol, pyrogallol, miconazole, and nitroblue tetrazolium. No inhibitory effect was seen in the presence of 0.01–1 mM of other metal ions, EDTA, EGTA, bipyridyl, *N*-ethylmaleimide, phenylmethanesulfonyl fluoride, KCN, NH<sub>2</sub>OH, Na<sub>3</sub>, hydrazine, D-cycloserine, barbital, or avidine. <sup>b</sup> Activity measurements were done in the absence of Na<sub>2</sub>S.

entire gene with its flanking sequence. The nucleotide sequence analysis of the total 11.6-kb DNA revealed an ORF (*orf2*) for the enzyme with a start codon ATG, a stop codon TGA, and a potential ribosome-binding site (AGGGAG) just before the start codon. The predicted protein was translated as 353 amino acids, with an amino acid sequence identical to that obtained by *N*-terminal sequencing (Figure 4). The *M<sub>r</sub>* of the protein was estimated to be 39 892, which agree with the calculated *M<sub>r</sub>* of the subunit (42 000). The ORF was thus named *oxd* as the structural gene for the enzyme. The predicted polypeptide (OxdRG) encoded by *oxd* showed 30.3% identity with OxdB (Figure 5) and 97.8% identity to the predicted polypeptide containing 137 amino acids encoded by an unidentified part of the gene that was located just upstream of the regulator 2 gene in the NHase-amidase operon in the iron-type NHase producer *Rhodococcus* sp. N-771 (35). No significant similarity was found between OxdRG and the gene products for other known proteins in the databases.

**Analysis of the Flanking Region of the *oxd* Gene.** Further sequencing of the flanking 5'-regions of *oxd* revealed another ORF (*orfa*), which consisted of 978 bp starting with an ATG initiation codon located 1182 bp upstream from the *oxd* start codon. The deduced amino acid sequence of *orfa* showed significant similarity to the following transcriptional regulatory (DNA-binding) proteins: transcription factor *nitR* (26.1%), which regulates nitrilase in *R. rhodochrous* J-1; a transcriptional regulator (25.5%) from *Bradyrhizobium japonicum*; a probable DNA-binding regulatory protein (24.6%) from *Streptomyces coelicolor*; and a putative *araC* family regulatory protein (23.9%) from *S. coelicolor* A3(2). The protein encoded by *orfa* could be a regulatory protein,

although its function is not yet clear. Upstream of *orfa*, an incomplete ORF (*orfC*) without a stop codon was present in a reverse orientation. The deduced amino acid sequence of the ORF showed remarkable similarity to acyl-CoA ligase (synthase), such as fatty acid CoA ligase of *Bacillus cereus* (AAP07877, 33% identity), feruloyl CoA synthase of *Amycolatopsis* sp. (Q9EY88, 34%), and acyl-CoA synthase of *Streptomyces avermitilis* (BAC68086, 35%). As shown in Figure 6 and Table 4, the 3'-flanking region of *oxd* demonstrated the presence of genes for NHase and its regulatory proteins and amidase having high identity as in *Rhodococcus* sp. N-771 (36) and *Rhodococcus* sp. AJ270. The deduced amino acid sequences of *orf3*, 4, 5, 6, 7, and 8 displayed 82.8, 85.9, 96.4, 98.6, 98.6, and 73.2% identity to those of NHase regulator 2, NHase regulator 1, amidase, NHase  $\alpha$ -subunit, NHase  $\beta$ -subunit, and NHase activator of *Rhodococcus* sp. N-771, respectively. The deduced amino acid sequences of *orf4*, 5, 6, 7, and 8 displayed 98.0, 96.4, 98.6, 98.6, and 98.0% identity with those of *Rhodococcus* sp. AJ270, respectively. We therefore changed the names of *orf3*, 4, 5, 6, 7, and 8 to *nhr2*, *nhr1*, *ami*, *nha1*, *nha2*, and *nhr3*, respectively. Immediately downstream of the activator gene *nhr3* there is a 912-bp *orfB*. It shared more than 91% sequence identity to a part of a hypothetical protein gene (302-bp) of *Rhodococcus* sp. AJ270. Promoter-like sequences were found upstream of *orfa*, *nhr2*, and *ami* (data not shown).

**Overexpression and Characterization of Recombinant OxdRG.** The complete *oxd* amplified by PCR from pOxdRGHII, with a modified Shine-Dalgarno sequence (AGGAGG), was inserted into pUC19 so that the enzyme was under the control of the *lac* promoter. The resulting plasmid (pOxdRGP9) was introduced into *E. coli* JM109 cells. The activity shown by *E. coli* cells containing pOxdRGP9 was about 300 units/liter culture in Terrific medium using *Z*-PAOx as the substrate; it was 15 times higher than that of *R. globerulus* A-4 cells cultured in TGY medium with *E*-3-PyOx as the inducer.

The recombinant OxdRG was purified through a 9-step purification procedure (Supporting Information, Table S1). It was purified 14.9-fold with a yield of 4.75% from the cell-free extract by measuring activity with *Z*-PAOx as the substrate. The purified recombinant OxdRG showed a single band on SDS-PAGE and a single peak on the G-3000 SW column in agreement with the *M<sub>r</sub>* of wild-type OxdRG. The heme content of the recombinant enzyme determined from the spectrum of its pyridine hemochromogen was calculated to be 0.35 mol of heme/mol of enzyme, the same as wild-type OxdRG. Recombinant OxdRG was activated by several reducing reagents similar to wild-type OxdRG. The specific activity of recombinant OxdRG for *Z*-PAOx was 2.18 units/mg which was about 24.8-fold higher than that of wild-type (0.088 unit/mg of protein) in the presence of 1 mM Na<sub>2</sub>S. The effect of Na<sub>2</sub>S at the various stages of purification of the recombinant enzyme was similar to its effect on wild-type OxdRG (data not shown).

## DISCUSSION

To understand the relationship between aldoxime dehydratase and NHase/amidase, we purified and characterized

*HincII*

```

10      20      30      40      50      60      70      80      90      100     110     120
GTCGACTCGACGGTTGCCGTGAGATCGTGCAACGACACGGAGTAGAGCGGTGCGTGACCCAGATTTCTCACCGCTGGGGATTCCCGACCTGGCCGCTTCAGTCGAGCCTTCAAGG

130     140     150     160     170     180     190     200     210     220     230     240
CGGAATTCGGGATCAGCCCCGCGGGTACCGGTGCAAGCAGCGGCCCAAGTACTACCCGCTGATCTGTAACATCTTCGGACACTTACCCATTGAACATGGTGTCAAATTTTCT

250     260     270     280     290     300     310     320     330     340     350     360
GTCGCCGAAGCGTATCCCAATTTCTGGCACGGTAGGCCAAGCATCGGGCGCTGACGGCCAAAGTTTGCTGGGCTGAGCGGGCTAGATTTTCGGCACGGCGTTGCCAAGTGCAGGCACC

370     380     390     400     410     420     430     440     450     460     470     480
CGATTGCAACCTACATCACAGGGGACTCATGGAATCTGCAATCGGGCAACCTTCAATGCCCGCGACTCTGACCAGGCGGGTTCGGGATACCTATACGCCACCGTTTCCCATGT

          SD      M E S A I G E H L Q C P R T L T R R V P D T Y T P P F P M W
490     500     510     520     530     540     550     560     570     580     590     600
GGGTGGGCGCGCAGACGACACATTGCACCAAGTCTGATGGGCTATCTCGGCGTGAGTTCCGCGGCGAGGATCAGCGCCCGCAGCACTGCGGGCGATGCGGGATATCGTCGCGGGCT

V G R A D D T L H Q V V M G Y L G V Q F R G E D Q R P A A L R A M R D I V A G F

610     620     630     640     650     660     670     680     690     700     710     720
TCGACTTGCCTGGACCGGACCGGCACACCAAGTCTCACCCATCACATCGAACAACAGGGTACGAGAACCTGATCGTGGTACTCGAAAGATGTTTCTCCCAACATCGTTGGGGCA

D L P D G P A H H D L T H H I D N Q G Y E N L I V V G Y W K D V S S Q H R W S T

730     740     750     760     770     780     790     800     810     820     830     840
CATCACCTCCAGTGTCTCTGCTGGGGAGTCCGAAGACCGCTCTGCGGACGGATTGGGGTTCTTCGCGAGATCGTGGCACCGAGAGCGCAACCAATTCGAACCGCTCTACCGTTCCCAAG

S P P V S S W W E S E D R L S D G L G F F R E I V A P R A E Q F E T L Y A F Q D

850     860     870     880     890     900     910     920     930     940     950     960
ACGACCTCCCGGGTGGAGCTGTATGGACGGTGTGACGGCGAAATCAACGAGCAGCGGTACTGGGGTTCGATGCGCGAGCGGTTCCGATCTCCCAACCGACTGGATGACGGGCT

D L P G V G A V M D G V S G E I N E H G Y W G S M R E R F P I S Q T D W M Q A S

970     980     990     1000    1010    1020    1030    1040    1050    1060    1070    1080
CGGGCGAACTGCGGGTCTGCGCGGTGACCCGCGTGGTGGACGGCTAGTAGTGGCGGGACACGACAACATCGCACTGATCAGGTCCGGGCGAGACTGGGCGGACCGGAAGCAGACG

G E L R V V A G D P A V G G R V V V R G H D N I A L I R S G Q D W A D A E A D E

1090    1100    1110    1120    1130    1140    1150    1160    1170    1180    1190    1200
AGCGCAGCCTTACTTGGACGAAATCCTGCCACTCTTCAATCGGGCATGGACTTCCTCCGCGACAACGGCCCGCGCTGGGTGCTACAGCAACCGATTCTGACGCAATATCGACATCG

R S L Y L D E I L P T L Q S G M D F L R D N G P A V G C Y S N R F V R N I D I D

1210    1220    1230    1240    1250    1260    1270    1280    1290    1300    1310    1320
ACGGAACTTCTCGACTTGAGTCTACAACATCGGCCACTGGGCTCCCTCGACCAACTCGAACGCTGGGAAATCCCAACCGACCCATCTACGGATCTTACAGCGTTCCTCCGGGTGG

G N F L D L S Y N I G H W A S L D Q L E R W S E S H P T H L R I F T T F F R V A

1330    1340    1350    1360    1370    1380    1390    1400    1410    1420    1430    1440
CCGAGGCGCTGCGAAATTGCGTCTTACCATGAGGTCTCGGTTATTCGATGCCCGCATCAGCTCTACGAGTACATCAATTGCCATCCCGGGACCGGAATGCTGCGCGACGCGGTGATCA

E G L S K L R L Y H E V S V F D A A D Q L Y E Y I N C H P G T G M L R D A V I T

1450    1460    1470    1480    1490    1500    1510
CCGCCGAGCACTGACCGAAGTCGCCGAAATCTTCGGCCACATTCTTCCCTCGGAGGCAACTCGAGATATGTT

A E H *                               HincII
```

FIGURE 4: Nucleotide and predicted amino acid sequences of the aldoxime dehydratase gene (*oxd*) from *R. globerulus* A-4. The underlined amino acid sequence corresponds with that of the N-terminal peptide of OxdRG. The putative ribosome-binding sequence is shown in italics. An asterisk indicates the stop codon.

an aldoxime dehydratase OxdRG from *R. globerulus* A-4 isolated from soil, in which NHase/amidase were present together with the aldoxime dehydratase. Table 5 shows a comparison of the properties of OxdRG and OxdB. Although both enzymes contain heme *b* as a prosthetic group and have a similar optimum pH and temperature, some differences are apparent. OxdRG has a dimeric structure, but OxdB is a monomer. Wild-type OxdRG exhibits a lower *Z*-PAOx dehydration rate (0.088 unit/mg) than OxdB (8.5 units/mg) (*18*); however, the rate for the recombinant OxdRG is similar (2.2 units/mg). The substrate specificity of OxdRG is broader than that of OxdB. Aliphatic aldoximes are more effective

substrates than arylalkyl aldoximes for OxdRG, and aromatic aldoximes are also dehydrated. Moreover, alkyl- and arylalkyl-aldoximes substituted at their  $\alpha$ -position, such as *E/Z*-isobutyraldoxime and *E/Z*-2-PPOx, are acceptable to OxdRG, although the activity is significantly diminished compared with the corresponding unsubstituted aldoximes (Table 2). Due to a wide substrate specificity, direct use of whole recombinant *E. coli* cells overexpressing OxdRG may be a useful biocatalyst for the production of various nitriles from the corresponding aldoximes. Since OxdRG can act on 2-PPOx or mandelaldoxime, the possibility exists for the production of optically active nitriles from aldoximes.






FIGURE 5: Alignment of the deduced amino acid sequences of the aldoxime dehydratases from *R. globerulus* A-4 (OxdRG) with *Bacillus* sp. OxB-1 (OxdB) (GenBank/EBI, AB028892). The alignment was performed with the ClustalW method. Gaps denoted by dashes were inserted to obtain maximum homology. Gray and open boxes show identical and similar residues between the sequences, respectively.




FIGURE 6: Genetic organization of the sequenced 11.6-kb DNA fragment and comparison of the gene cluster from *R. globerulus* A-4 with that of *Bacillus* sp. OxB-1, *Rhodococcus* sp. N-771, and *Rhodococcus* sp. AJ270. Four fragments (3.9-kb *Fba*I, 4.7-kb *Bam*HI, 6.3-kb *Pst*I, and 3.7-kb *Hind*III-*Sph*I fragments) are also indicated. The positions and orientations of the different ORFs detected within the locus are shown by large arrows. F, *Fba*I; E, *Eco*RI; B, *Bam*HI; P, *Pst*I; H, *Hind*III; S, *Sph*I.

The purified OxdRG activity was greatly enhanced by reducing reagents, such as  $\text{Na}_2\text{S}$ ,  $\text{Na}_2\text{S}_2\text{O}_4$ , and some sulfhydryl compounds. The activity was further enhanced by  $\text{Fe}^{2+}$  as well as OxdB (18). On the basis of these observations, it is concluded that OxdRG and OxdB both require reduction to stimulate the reaction process. Distinct from OxdB activity, however, OxdRG activity was not enhanced under anaerobic conditions. Oxidized irons ( $\text{Fe}^{3+}$  and  $\text{K}_3\text{-[Fe(CN)}_6\text{]}$ ) also enhanced activity both in the presence and absence of  $\text{Na}_2\text{S}$ , but their effect on the activity is unclear. The mechanisms for the effect of some metal ions,  $\text{Cu}^+$ ,  $\text{Cu}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Zn}^{2+}$ , and  $\text{Cd}^{2+}$ , which act as both activators and inhibitors depending on their concentrations, are also not understood. The fact that phenylhydrazine and Tiron inhibited Z-PAOx dehydration but not E/Z-VOx dehydration suggests that they might inhibit the reaction by interfering

with the binding of substrate by their aromatic ring. The enhancement of the enzyme activity by flavins, vitamin  $\text{K}_3$ , duroquinone, and sulfite ion suggests that the enzyme requires an electron acceptor during the reaction progress. Partially purified indoleacetaldoxime hydrolyase (EC 4.2.1.29) is activated by PLP, ascorbic acid, and dehydroascorbic acid (37), but OxdRE is not activated by the compounds, suggesting that OxdRG is different from indoleacetaldoxime hydrolyase.

The heme content of the purified wild-type and recombinant OxdRG was low as it was for OxdB. It is still unclear whether the heme was lost during the purification or the enzyme was originally produced with a low heme content. Also, the actual role of the reducing reagents and electron acceptors in the enzyme reaction is little understood. The crystal structures of the active site of both enzymes may help

Table 4: ORFs Deduced by Sequencing Analysis of the 11.6-kb Fragment from *R. globerulus* A-4 and Their Corresponding Gene Products and Related Gene Products

gene <i>or orf</i>	function	gene product (aa) <sup>a</sup>	related gene and its product				
			gene	function	organism	identity i% j	accession no.
1 <i>orfa</i>	putative regulatory protein	326	<i>nitR</i>	transcription factor <i>nitR</i>	<i>R. rhodochrous</i> J1	26.1	JC6117
				transcriptional regulator probable DNA-binding regulatory protein	<i>Bradyrhizobium japonicum</i> <i>Streptomyces coelicolor</i>	25.5 24.6	AP005958 T36475
				putative <i>araC</i> family regulatory protein	<i>S. coelicolor</i> A3(2)	23.9	AL939112
2 <i>oxd</i>	aldoxime dehydratase	353	<i>oxd</i>	phenylacetaldoxime dehydratase	<i>Bacillus</i> sp. OxB-1	30.3	AB028892
3 <i>nhr2</i>	NHase regulator2	122	<i>nhr2</i> <i>nh1D</i>	undefined	<i>Rhodococcus</i> sp. N-771	97.8 (137) <sup>b</sup>	AB016078
				NHase regulator2	<i>Rhodococcus</i> sp. N-771	82.8	AB016078
				NHase regulator2 transposase	<i>R. rhodochrous</i> J1 <i>R. erythropolis</i> AJ270	41.8 95.4 (65)	D67028 RER490527
4 <i>nhr1</i>	NHase regulator1	410	<i>nhr1</i> <i>nhr1</i>	NHase regulator1	<i>R. erythropolis</i> AJ270	98.0	RER490527
				NHase regulator1	<i>Rhodococcus</i> sp. N-771	85.9	AB016078
5 <i>ami</i>	amidase	521	<i>ami</i> <i>ami</i> <i>ami</i>	amidase	<i>Rhodococcus</i> sp. N-771	96.4	AB016078
				amidase	<i>Rhodococcus</i> sp. N-774	96.4	X54074
				amidase	<i>R. erythropolis</i> AJ270	96.4	RER490527
6 <i>nha1</i>	NHase $\alpha$ -subunit	207	<i>nha1</i> <i>nha1</i> <i>nha</i>	NHase $\alpha$ -subunit	<i>Rhodococcus</i> sp. N-771	98.6	AB016078
				NHase $\alpha$ -subunit	<i>Rhodococcus</i> sp. N-774	98.6	X54074
				NHase $\alpha$ -subunit	<i>R. erythropolis</i> AJ270	98.6	RER490527
7 <i>nha2</i>	NHase $\beta$ -subunit	212	<i>nha2</i> <i>nha2</i> <i>nhb</i>	NHase $\beta$ -subunit	<i>Rhodococcus</i> sp. N-771	98.6	AB016078
				NHase $\beta$ -subunit	<i>Rhodococcus</i> sp. N-774	98.6	X54074
				NHase $\beta$ -subunit	<i>R. erythropolis</i> AJ270	98.6	RER490527
8 <i>nhr3</i>	NHase activator	399	<i>nha3</i> <i>nha3</i>	NHase activator	<i>R. erythropolis</i> AJ270	98.0	RER490527
				NHase activator	<i>Rhodococcus</i> sp. N-771	73.2	AB016078
				NHase activator	<i>R. erythropolis</i> AJ270	43.6 (101)	RER490527
9 <i>orfb</i>	hypothetical protein	304	<i>orfe</i>	hypothetical protein	<i>R. erythropolis</i> AJ270	43.6 (101)	RER490527

<sup>a</sup> aa, amino acid. <sup>b</sup> The number in parentheses indicates the partial deduced amino acid sequence of the gene product that has been used for comparison.

Table 5: Comparison of the Properties of Aldoxime Dehydratases from *R. globerulus* A-4 and *Bacillus* sp. OxB-1

property	value with enzyme from			
	<i>R. globerulus</i> A-4		<i>Bacillus</i> sp. OxB-1	
molecular weight (sequencing)	39,892		40,018	
(subunit)	42,000		40,000	
(native)	76,200		42,000	
no. of subunits	2		1	
optimum pH	8.0 (KPB)		7.0 (KPB)	
temp	30 °C		30 °C	
stable pH	6.0–9.5		6.5–11.5	
temp	<40 °C		<45 °C	
substrate specificity	aryl-, arylalkyl-, alkylaldoxime		arylalkyl-, alkylaldoxime	
prosthetic group	heme <i>b</i>		heme <i>b</i>	
activator	FMN, Fe <sup>2+</sup> , Sn <sup>2+</sup> , Na <sub>2</sub> SO <sub>3</sub> , Na <sub>2</sub> S, vitamin K <sub>3</sub>		FMN, Fe <sup>2+</sup> , Sn <sup>2+</sup> , Na <sub>2</sub> SO <sub>3</sub>	
inhibitor (cations)	Co <sup>2+</sup> , Zn <sup>2+</sup> , Cd <sup>2+</sup> , Cu <sup>+</sup> , Cu <sup>2+</sup> , Ag <sup>+</sup> , Hg <sup>2+</sup> , Ti <sup>+</sup> , As <sup>3+</sup>		Cu <sup>+</sup> , Cu <sup>2+</sup> , Ag <sup>+</sup> , Hg <sup>2+</sup>	
(chemical compounds)	electron donors and carriers		EDTA, 8-hydroxyquinoline tetramethylphenylenediamine	
<i>K<sub>m</sub></i> (mM) and <i>V<sub>max</sub></i> (units/mg) value for	<i>K<sub>m</sub></i>	<i>V<sub>max</sub></i>	<i>K<sub>m</sub></i>	<i>V<sub>max</sub></i>
Z-PAOx	1.40	0.140	0.872	19.5
Z-3-PPOx	2.31	0.392	1.36	14.3
<i>E/Z</i> -indoleacetaldoxime	3.91	0.281	2.40	5.42
<i>E/Z</i> -2-PPOx	11.9	0.810		no reaction
<i>E</i> -PyOx	20.0	0.065		no reaction
<i>E/Z</i> -cyclohexanecarboxaldehyde oxime	1.13	0.386		no reaction
<i>E/Z</i> - <i>n</i> -butyraldoxime	1.73	0.689	11.1	9.49
<i>E/Z</i> -isobutyraldoxime	5.54	0.041		no reaction
<i>E/Z</i> - <i>n</i> -capronaldoxime	2.94	1.66	6.12	32.3
<i>E/Z</i> -isocapronaldoxime	6.76	1.32	2.98	10.1

to explain the mechanism of the reaction. Interestingly, the specific activity of the recombinant OxdRG for PAOx was 25 times that of the wild-type, although the heme contents

of the two were nearly the same. Since the effect of Na<sub>2</sub>S on enzyme activity was similar in the wild type and recombinant OxdRGs, it is possible to consider that an

unknown cofactor was lost from wild-type OxdRG during the purification and that the cofactor property could not be restored by the addition of Na<sub>2</sub>S. However, the actual reasons for the observations are not clear at the present time.

Sequence analysis of the 11.6-kb DNA fragment revealed the presence of at least nine complete open reading frames (*orfa*, *oxd*, *nh2*, *nh1*, *ami*, *nha1*, *nha2*, *nh3*, and *orfb*). These genes encode the proteins involved in aldoxime metabolism, a combination of aldoxime dehydratase, NHase, and amidase, together with the regulatory proteins. An aldoxime dehydratase gene homologue in *Rhodococcus* sp. N-771 is likely to be located just upstream of the regulator 2 gene, given its high sequence similarity to a part of *oxd* (unpublished data). It is anticipated that the gene cluster regulating aldoxime metabolism in *Rhodococcus* sp. N-771 might be similar to that of *R. globerulus* A-4. The coexistence of the structural genes for OxdRG, amidase, and NHase, together with regulatory proteins, favors the expression of three enzymes in *R. globerulus* A-4, suggesting that the enzymes may be related and coexist in a single metabolic pathway for aldoxime catabolism. *Rhodococcus* sp. AJ270 has a similar gene assembly and shows high homology to *R. globerulus* A-4 and *Rhodococcus* sp. N-771, except for a transposase gene which is located just upstream of the NHase regulator 1 gene. The deduced 65-amino acid sequence of *nh2* of *R. globerulus* A-4 displayed 95.4% identity with that of transposase. Since *R. globerulus* A-4, *Rhodococcus* sp. N-771, and *Rhodococcus* sp. AJ270 probably evolved from the same ancestor, an evolutionary change in *Rhodococcus* sp. AJ270 was the insertion of the transposon into the aldoxime/nitrile metabolic operons.

Even though it has not been purified or characterized yet, the NHase encoded by the genes present in the 3'-flanking region of *oxd* may preferentially act on aliphatic nitriles because NHase of *Rhodococcus* sp. N-771 encoded by *nha1* and *nha2*, which showed high identity with the NHase in *R. globerulus* A-4, prefers aliphatic nitriles to aromatic nitriles (38). We conclude that OxdRG, in combination with NHase and amidase, plays a significant role in the metabolism of alkylaldoxime, according to the enzymological and genetic evidence.

Studies are under way to clarify the existence of biosynthetic pathways of aldoximes from amino acids in microorganisms similar to those in plants and their evolutionary relationships.

## SUPPORTING INFORMATION AVAILABLE

Table of purification of the recombinant aldoxime dehydratase from *E. coli* JM109/pOxDRGP9. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## REFERENCES

- Sibbesen, O., Koch, B., Rouz , P., M ller, B. L., and Halkier, B. A. (1995) in *Amino Acids and Their Derivatives in Higher Plants* (Wallsgrove, R. M., Ed.) pp 227–241, Cambridge University Press, Cambridge.
- Andersen, M. D., Busk, P. K., Svendsen, I., and M ller, B. L. (2000) *J. Biol. Chem.* 275, 1966–1975.
- Nielsen, J. S., and M ller, B. L. (2000) *Plant Physiol.* 122, 1311–1321.
- Bak, S., Tax, F. E., Feldmann, K. A., Galbraith, D. A., and Feyereisen, R. (2001) *Plant Cell.* 13, 101–111.
- Bak, S., Olsen, C. E., Halkier, B. A., and M ller, B. L. (2001) *Plant Physiol.* 123, 1437–1448.
- Bak, S., and Feyereisen, R. (2001) *Plant Physiol.* 127, 108–118.
- Hansen, C. H., Du, L.-C., Naur, P., Olsen, C. E., Axelsen, K. B., Hick, A. J., Pickett, J. A., and Halkier, B. A. (2001) *Biol. Chem.* 276, 24790–24796.
- Yamada, H., Asano, Y., Hino, T., and Tani, Y. (1979) *J. Ferment. Technol.* 57, 8–14.
- Asano, Y., Yasuda, T., Tani, Y., and Yamada, H. (1982) *Agric. Biol. Chem.* 46, 1183–1189.
- Asano, Y., Tani, Y., and Yamada, H. (1980) *Agric. Biol. Chem.* 44, 2251–2252.
- Asano, Y., Fujishiro, K., Tani, Y., and Yamada, H. (1982) *Agric. Biol. Chem.* 46, 1165–1174.
- Asano, Y. (1991) *Nippon Nogekagaku Kaishi* 65, 1617–1626.
- Yamada, H., and M. Kobayashi. (1996) *Biosci. Biotechnol. Biochem.* 60, 1391–1400.
- Hann, E. C., Eisenberg, A., Fager, S. K., Perkins, N. E., Gallagher, F. G., Cooper, S. M., Gavagan, J. E., Stieglitz, B., Hennessey, S. M., and DiCosimo, R. (1999) *Bioorg. Med. Chem.* 7, 2239–2245.
- Asano, Y., and Kato, Y. (1998) *FEMS Microbiol. Lett.* 158, 185–190.
- Kato, Y., Ooi, R., and Asano, Y. (1998) *Arch. Microbiol.* 170, 85–90.
- Asano, Y. (2002) *J. Biotechnol.* 94, 65–72.
- Kato, Y., Nakamura, K., Sakiyama, H., Mayhew, S. G., and Asano, Y. (2000) *Biochemistry*, 39, 800–809.
- Kato, Y., and Asano, Y. (2003) *Protein Express. Purif.* 28, 131–139.
- Xie, S.-X., Kato, Y., and Asano, Y. (2001) *Biosci. Biotechnol. Biochem.* 65, 2666–2672.
- Kato, Y., Ooi, R., and Asano, Y. (2000) *Appl. Environ. Microbiol.* 66, 2290–2296.
- Kato, Y., Ooi, R., and Asano, Y. (1999) *J. Mol. Catal. B: Enzymatic* 6, 249–256.
- Asano, Y., Itoh, H., Dairi, T., and Kato, Y. (1996) *J. Biol. Chem.* 271, 30256–30262.
- Davis, B. J. (1964) *Ann. New York Acad. Sci.* 121 (Part 2), 404–427.
- Laemmli, U. K. (1970) *Nature* 227, 680–685.
- Goodhew, C. F., Brown, K. R., and Pettigrew, G. W. (1986) *Biochim. Biophys. Acta.* 852, 288–294.
- Appleby, C. A., and Morton, R. K. (1959) *Biochem. J.* 73, 539–550.
- Siegel, L. M., Murphy, M. J., and Kamin, H. (1973) *J. Biol. Chem.* 248, 251–264.
- Lechevalier, H. A. (1986) Nocardioforms in *Bergey's Manual of Systematic Bacteriology* (Sneath, P. H. A., Ed.) Vol. 2, pp 1458–1506, Williams and Wilkins, Baltimore and London.
- Miller, L. T. (1982) *J. Clin. Microbiol.* 16, 54–586.
- Nagamune, T., Kurata, H., Hirata, M., Honda, J., Koike, H., Ikeuchi, M., Inoue, Y., Hirata, A., and Endo, I. (1990) *Biochem. Biophys. Res. Commun.* 168, 437–442.
- Sanger, F. S., Nicklen, S., and Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. U.S.A.* 74, 5463–5467.
- Pearson, W. R., and Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. U.S.A.* 85, 2444–2448.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403–410.
- Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994) *Nucleic Acid Res.* 22, 4673–4680.
- Nojiri, M., Yohda, M., Odaka, M., Matsushita, Y., Tsujimura, M., Yoshida, T., Dohmae, N., Takio, K., and Endo, I. (1999) *J. Biochem.* 125, 696–704.
- Shukla, P. S., and Mahadevan, S. (1970) *Arch. Biochem. Biophys.* 137, 166–174.
- Banerjee, A., Sharma, R., and Banerjee, U. C. (2002) *Appl. Microbiol. Biotechnol.* 60, 33–44.

BI035092U