

A GENERALIZATION OF THE AZ IDENTITY

RUDOLF AHLSEWEDE and NING CAI

Received July 2, 1990

The identity discovered in [1] can be viewed as a sharpening of the LYM inequality ([3], [4], [5]). It was extended in [2] so that it covers also Bollobás' inequality [6]. Here we present a further generalization and demonstrate that it shares with its predecessors the usefulness for uniqueness proofs in extremal set theory.

1. Introduction

A few years ago Ahlswede and Zhang [1] found the following identity.

Theorem AZ₁. For every family $\mathcal{A} \subset 2^\Omega$ of non-empty subsets of $\Omega = \{1, 2, \dots, n\}$

$$\sum_{X \subset \Omega} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 1, \text{ where } W_{\mathcal{A}}(X) = \left| \bigcap_{X \supset A \in \mathcal{A}} A \right|.$$

We associate with every $\mathcal{E} \subset 2^\Omega$ the upset $\mathcal{U}(\mathcal{E}) = \{U \subset \Omega : U \supset E \text{ for some } E \in \mathcal{E}\}$ and the downset $\mathcal{D}(\mathcal{E}) = \{D \subset \Omega : D \subset E \text{ for some } E \in \mathcal{E}\}$.

When \mathcal{A} is an antichain in the poset $(2^\Omega, \supset)$, then the identity becomes

$$(1) \quad \sum_{X \in \mathcal{A}} \frac{1}{\binom{n}{|X|}} + \sum_{X \in \mathcal{U}(\mathcal{A}) \setminus \mathcal{A}} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 1.$$

The LYM inequality is obtained by omission of the second summand, which by definition of $W_{\mathcal{A}}$ can also be written in the form $\sum_{X \notin \mathcal{D}(\mathcal{A})} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}}$. We call this the deficiency of the inequality.

More generally, in [2] the Bollobás inequality was lifted to an identity.

Theorem AZ₂. For two families $\mathcal{A} = \{A_1, \dots, A_N\}$ and $\mathcal{B} = \{B_1, \dots, B_N\}$ of subsets of Ω with the properties

- (a) $A_i \subset B_i$ for $i = 1, 2, \dots, N$
- (b) $A_i \not\subset B_j$ for $i \neq j$

$$(2) \quad \sum_{i=1}^N \frac{1}{\binom{n-|B_i \setminus A_i|}{|A_i|}} + \sum_{X \notin \mathcal{D}(\mathcal{B})} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 1.$$

In [1] it was explained that Theorem AZ₁ gives immediately, what LYM does not, namely the uniqueness part in Sperner’s Theorem. In [2] the uniqueness of an optimal configuration of unrelated chains of subsets due to Griggs, Stahl and Trotter [7] was proved with the help of Theorem AZ₂.

Recently, Körner and Simonyi [10] observed the LYM-type inequality:

For $\mathcal{A} = \{A_1, \dots, A_N\}, \mathcal{B} = \{B_1, \dots, B_N\} \subset 2^\Omega$ with

$$A_i \cap B_i = \emptyset, A_i \not\subset A_j \cup B_j, B_i \not\subset A_j \cup B_j \text{ for } i \neq j$$

$$(3) \quad \sum_{i=1}^N \binom{n-|A_i|}{|B_i|}^{-1} + \binom{n-|B_i|}{|A_i|}^{-1} - \binom{n}{|A_i|+|B_i|}^{-1} \leq 1$$

and they asked (Problem 2) “Is this inequality ever tight?”.

This rather modest question was a challenging test of the power of the identities in [1], [2] or, more precisely, of the procedure to produce new identities described in [1].

The outcome is an Ahlswede–Zhang type identity (Theorem 1) which goes considerably beyond Theorem AZ₂. From a special case of this identity we derive a full characterization of the cases with equality (Theorem 2) even for a generalized version of (3). In other words we characterize the cases with deficiency zero.

2. The identity

Theorem 1. Suppose that for a family $\mathcal{B} = \{B_1, \dots, B_N\}$ of subsets of Ω and a family $\mathcal{A}^* = \{\mathcal{A}_1, \dots, \mathcal{A}_N\}$ of subsets of 2^Ω , where $\mathcal{A}_i = \{A_i^t : t \in T_i\}$ for a finite index set T_i , we have the properties

- (a) $A_i^t \subset B_i$ for $t \in T_i$ and $i = 1, 2, \dots, N$
- (b) $A_i^t \not\subset B_j$ for $t \in T_i$ and $i \neq j$.

Then with $\mathcal{A} = \bigcup_{i=1}^N \mathcal{A}_i$

$$(4) \quad \sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \sum_{S \subset T_i, |S|=k} \binom{n-|B_i - \bigcup_{t \in S} A_i^t|}{|\bigcup_{t \in S} A_i^t|}^{-1} + \sum_{X \notin \mathcal{D}(\mathcal{B})} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 1.$$

The specialisation $|T_i| = 1$ for $i = 1, \dots, N$ gives Theorem AZ₂. The proof goes again by counting chains. A key tool in [2] was

Lemma 1. For two sets $A, B \subset \Omega$ with $A \subset B$ exactly $\frac{n!}{\binom{n-|B \setminus A|}{|A|}}$ maximal chains in $(2^\Omega, \subset)$ meet $\{X : A \subset X \subset B\}$. ■

Using the principle of inclusion-exclusion this generalizes to

Lemma 2. For $B \subset \Omega$ and $\mathcal{C} \subset 2^\Omega$ with $C \subset B$ for all $C \in \mathcal{C}$ exactly

$$n! \sum_{k=1}^{|\mathcal{C}|} (-1)^{k-1} \sum_{\mathcal{C}' \subset \mathcal{C}, |\mathcal{C}'|=k} \binom{n - |B \setminus \cup_{C \in \mathcal{C}'} C|}{|\cup_{C \in \mathcal{C}'} C|}^{-1}$$

maximal chains in $(2^\Omega, \subset)$ meet $\{X : C \subset X \subset B \text{ for some } C \in \mathcal{C}\}$. ■

Proof of Theorem 1. The number of maximal chains leaving $\mathcal{U}(\mathcal{A})$ at U is

$$(n - |U|)! W_{\mathcal{A}}(U) (|U| - 1)!$$

Since the sets $\mathcal{X}_i = \{X : A_i^t \subset X \subset B_i \text{ for some } t \in T_i\}$ ($i = 1, 2, \dots, N$) are disjoint we have

$$\sum_{i=1}^N \sum_{X \in \mathcal{X}_i} (n - |X|)! W_{\mathcal{A}}(X) (|X| - 1)! + \sum_{X \in \mathcal{U}(\mathcal{A}) - \cup \mathcal{X}_i} (n - |X|)! W_{\mathcal{A}}(X) (|X| - 1)! = n!$$

By the definition of $W_{\mathcal{A}}$ the last summand can be written in the form

$$\sum_{X \notin \mathcal{D}(\mathcal{B})} \frac{W_{\mathcal{A}}(X) n!}{|X| \binom{n}{|X|}} \text{ and by Lemma 2}$$

$$\sum_{X \in \mathcal{X}_i} (n - |X|)! W_{\mathcal{A}}(X) (|X| - 1)! = n! \sum_{k=1}^{|T_i|} (-1)^{k-1} \sum_{S \subset T_i, |S|=k} \binom{n - |B_i \setminus \cup_{t \in S} A_i^t|}{|\cup_{t \in S} A_i^t|}^{-1}. \quad \blacksquare$$

3. On zero deficiency

We characterize here a case of zero deficiency, that is, the property

$$(5) \quad \sum_{X \notin \mathcal{D}(\mathcal{B})} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 0.$$

Theorem 2. Under the assumptions of Theorem 1 and the additional conditions

- (c) $A_i^t \cap A_i^{t'} = \emptyset$ for all i and $t, t' \in T_i$ with $t \neq t'$
- (d) $|T_i| \geq 2$ and $N \geq 2$

we have, that the identity

$$(6) \quad \sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \sum_{S \subset T_i, |S|=k} \binom{n - |B_i - \cup_{t \in S} A_i^t|}{|\cup_{t \in S} A_i^t|}^{-1} = 1$$

holds exactly if

- (i) $|A_i^t| = 1$ for all $t \in T_i$ and all i .
- (ii) $B_1 \setminus \cup_{t \in T_1} A_1^t = B_2 \setminus \cup_{t \in T_2} A_2^t = \dots = B_N \setminus \cup_{t \in T_N} A_N^t = B$, say.
- (iii) $\sum_{i=1}^N |T_i| = n - |B|$.

In words, the B_i have a common part B and each B_i has a rest of singletons A_i^t . The B_i 's exhaust Ω .

In the proof we use a well-known identity, which follows by iterative application of Pascal's identity.

Lemma 3. $\sum_{k=1}^m (-1)^{k-1} \binom{M}{m-k} = \binom{M-1}{m-1}$. ■

Proof of Theorem 2. From (i) and (ii) we derive in terms of $\beta = |B|$

$$\begin{aligned} & \sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \sum_{S \subset T_i, |S|=k} \binom{n - |B_i \setminus \cup_{t \in S} A_i^t|}{|\cup_{t \in S} A_i^t|}^{-1} \\ &= \sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \binom{|T_i|}{k} \binom{n - (\beta + |T_i|) + k}{k}^{-1} \\ &= \sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \frac{|T_i|!(n - (\beta + |T_i|))!}{(|T_i| - k)!(n - (\beta + |T_i|) + k)!} \\ &= \sum_{i=1}^N \frac{|T_i|!(n - (\beta + |T_i|))!}{(n - \beta)!} \sum_{k=1}^{|T_i|} (-1)^{k-1} \binom{n - \beta}{|T_i| - k} \\ &= \sum_{i=1}^N \frac{|T_i|!(n - (\beta + |T_i|))!}{(n - \beta)!} \frac{(n - \beta - 1)!}{(|T_i| - 1)!(n - (\beta + |T_i|))!}, \end{aligned}$$

by Lemma 3, and now by (iii)

$$(7) \quad = \sum_{i=1}^N \frac{|T_i|}{n - \beta} = 1.$$

We assume now that (6) holds and derive (i), (ii), and (iii). By Theorem 1 we have deficiency zero, that is,

$$(8) \quad W_{\mathcal{A}}(X) = 0 \text{ for all } X \notin \mathcal{D}(\mathcal{B}).$$

For the quantity

$$(9) \quad m = \min\{|A_i^t| : 1 \leq i \leq N, t \in T_i\}$$

we show first that it equals 1, then we establish (i) and (ii), and finally (iii).

Step 1. W.l.o.g. we can assume $|A_1^1| = m$. For any $y \in \Omega \setminus B_1$ consider $A_1^1 \cup \{y\}$. Thus clearly $A_1^1 \cup \{y\} \not\subset B_1$ and by condition (b) also $A_1^1 \cup \{y\} \not\subset B_j$ for $j \neq 1$. Therefore $A_1^1 \cup \{y\} \notin \mathcal{D}(\mathcal{B})$ and by (8) $W_{\mathcal{A}}(A_1^1 \cup \{y\}) = 0$. By the minimality of A_1^1 in \mathcal{A} and the definition of $W_{\mathcal{A}}$ every m -subset of $A_1^1 \cup \{y\}$ must be in \mathcal{A} . In particular for any $a \in A_1^1$ the set $(A_1^1 \setminus \{a\}) \cup \{y\}$ is in \mathcal{A} . Since it is not in \mathcal{A}_1 it must be in some \mathcal{A}_j with $j \neq 1$.

W.l.o.g. we can assume it to be A_2^1 . Furthermore, since $A_1^1 \neq A_1^2$ we can require the a chosen above to be from $A_1^1 \setminus A_1^2$. Also, since by (b) $A_1^2 \not\subset B_2$ there is $z \in A_1^2 \setminus B_2, z \neq a$. As previously we conclude that $A_2^1 \cup \{z\} \notin \mathcal{D}(\mathcal{B})$ and that the m -set

$$(A_2^1 \cup \{z\}) \setminus \{y\} = (A_1^1 \setminus \{a\}) \cup \{z\} \in \mathcal{A}.$$

However, we also have $(A_1^1 \setminus \{a\}) \cup \{z\} \in \mathcal{A}_1$ and by (c) $A_1^1 \cap ((A_1^1 \setminus \{a\}) \cup \{z\}) = \emptyset$. This implies $A_1^1 = \{a\}$ and $m = 1$.

Step 2. After relabelling we can assume now $A_1^1 = \{1\}$ and $B_1 = \{1, 2, \dots, \ell\}$. By the arguments in Step 1 we get $\{1, k\} \notin \mathcal{D}(\mathcal{B})$ and $\{1, k\} \supset \{k\} \in \mathcal{A}$ whenever $k > \ell$. By (b) for all $t \in T_i$ and $i \geq 2$ A_i^t has an element, say e , with $e > \ell$. However, since $\{e\} \in \mathcal{A}$ by (a), (b) and (c) actually A_i^t must equal $\{e\}$. We thus know that A_i^t is a singleton for all $i \geq 2$ and $t \in T_i$. Now we can let any $i \geq 2$ take the role of 1 in the previous argument and get that all A_1^t are also singletons. We have proved (i).

Also we have arrived at the following configuration: $B_i \supset A_i = \cup_{t \in T_i} A_i^t$ and $B_i \cap A_j = \emptyset$ for $i \neq j$. We claim now that $B_i = A_i \cup C$, where $C = \Omega \setminus \cup_{i=1}^N A_i$. To see this, suppose that $c \in C$ and $c \notin B_i$. Then for any $a \in A_i$ $\{a, c\} \notin \mathcal{D}(\mathcal{B})$ and thus $W_{\mathcal{A}}(\{a, c\}) = 0$. This, however, contradicts $W_{\mathcal{A}}(\{a, c\}) = |\{a\}| = 1$.

We have established (ii) with $B = C$. (6), together with the equations leading to (7), give now also (iii). ■

Finally we present a consequence of Theorem 2, which in particular gives a positive answer to the question of Körner and Simonyi mentioned in the Introduction.

Corollary. *If we are given for $t = 1, 2$ and $i = 1, 2, \dots, N$ sets $A_i^t \subset \Omega$ with $A_i^1 \cap A_i^2 = \emptyset$ and $A_i^t \not\subset A_j^1 \cup A_j^2$ for $t = 1, 2$ and $i \neq j$ then*

$$(10) \quad \sum_{i=1}^N \binom{n - |A_i^1|}{|A_i^1|}^{-1} + \binom{n - |A_i^2|}{|A_i^2|}^{-1} - \binom{n}{|A_i^1| + |A_i^2|}^{-1} = 1$$

exactly if

$$(i') \quad |A_i^t| = 1 \text{ for } t = 1, 2 \text{ and } i = 1, 2, \dots, N$$

(ii') n is even and $N = \frac{n}{2}$.

There is a direct proof of this Corollary which is shorter than the one via Theorem 2.

Proof. With the choice $\mathcal{B}_i = \bigcup_{t \in T_i} A_i^t$ formula (6) takes the form

$$\sum_{i=1}^N \sum_{k=1}^{|T_i|} (-1)^{k-1} \sum_{S \subset T_i, |S|=k} \binom{n - \sum_{t \notin S} |A_i^t|}{\sum_{t \in S} |A_i^t|}^{-1} = 1$$

and if $T_i = \{1, 2\}$ for all i this becomes (10). (i) specializes to (i'), (ii) is true by definition of B_i with $B = \emptyset$. (iii) specializes to $N \cdot 2 = n$ and thus (ii'). ■

4. On general cloud antichains

A family $\mathcal{A}^* = \{\mathcal{A}_1, \dots, \mathcal{A}_N\}$ of subsets of 2^Ω is a cloud-antichain, if

$$(1') \quad A_i \not\subset A_j \text{ for } A_i \in \mathcal{A}_i, A_j \in \mathcal{A}_j \text{ with } i \neq j.$$

They have been analyzed in [2] for $N = 2$. In particular, in the case $|\mathcal{A}_i| = M$ for $i = 1, \dots, N$ we are interested in the maximal length $N(n, M)$ of these antichains.

Clearly, for $\mathcal{A} = \bigcup_{i=1}^N \mathcal{A}_i$

$$(12) \quad W_{\mathcal{A}}(X) = W_{\mathcal{A}_i}(X) \text{ for } X \in \mathcal{A}_i$$

and therefore by Theorem AZ₁

$$(13) \quad \sum_{i=1}^N \sum_{X \in \mathcal{A}_i} \frac{W_{\mathcal{A}_i}(X)}{|X| \binom{n}{|X|}} + \sum_{X \in \mathcal{U}(\mathcal{A}) \setminus \mathcal{A}} \frac{W_{\mathcal{A}}(X)}{|X| \binom{n}{|X|}} = 1.$$

Notice that $n! \sum_{X \in \mathcal{B}} \frac{W_{\mathcal{B}}(X)}{|X| \binom{n}{|X|}}$ counts the number, say $\sigma(\mathcal{B})$, of saturated chains meeting a member of \mathcal{B} .

We can derive from (13) a bound on $N(n, M)$, if we have a bound or even exact result for the following seemingly basic quantity:

$$(14) \quad s(M, n) = \min\{\sigma(\mathcal{B}) : \mathcal{B} \subset 2^\Omega, |\mathcal{B}| = M\}.$$

References

[1] R. AHLWEDE, and Z. ZHANG: An identity in combinatorial extremal theory, *Advances in Mathematics* **80** (2) (1990), 137–151.
 [2] R. AHLWEDE, and Z. ZHANG: On cloud-antichains and related configurations, *Discrete Mathematics* **85** (1990), 225–245.

- [3] K. YAMAMOTO: Logarithmic order of free distributive lattices, *J. Math. Soc. Japan* **6** (1954), 343–353.
- [4] L.D. MESHALKIN: A generalization of Sperner's theorem on the number of subsets of a finite set, *Theor. Probability Appl.* **8** (1963), 203–204.
- [5] D. LUBELL: A short proof of Sperner's theorem, *J. Combinatorial Theory* **1** (1966), 299.
- [6] B. BOLLOBÁS: On generalized graphs, *Acta Math. Acad. Sci. Hungar.* **16** (1965), 447–452.
- [7] J.R. GRIGGS, J. STAHL, and W.T. TROTTER: A Sperner theorem on unrelated chains of subsets, *J. Comb. Theory, Ser. A* **36** (1984), 124–127.
- [8] K. ENGEL, and H.D.O.F. GRONAU: *Sperner Theory in Partially Ordered Sets*, Texte zur Mathematik Bd. 78, Teubner, Leipzig, 1985.
- [9] E. SPERNER: *Ein Satz über Untermengen einer endlichen Menge*, *Math. Z.* **27** (1928), 544–548.
- [10] J. KÖRNER, and G. SIMONYI: A Sperner-type theorem and qualitative independence, *J. Comb. Theory, Ser. A* **59** (1992), 90–103.

Rudolf Ahlswede, Ning Cai

Universität Bielefeld,

Fakultät für Mathematik,

D-4800 Bielefeld 1, Germany

`hollmann@math.10.mathematik.`

`uni-bielefeld.de`