



REVIEW

A global perspective of fragmentation on a declining taxon — the sturgeon (Acipenseriformes)

Tim J. Haxton^{1,*}, Tim M. Cano²

¹Aquatic Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry, 1600 West Bank Dr., Peterborough, ON K9L 0G2, Canada

²Northwest Regional Operations Division, Ontario Ministry of Natural Resources and Forestry, 173 25th Side Road, Rosslyn, ON P7K 0B9, Canada

ABSTRACT: Acipenseriformes (sturgeons and paddlefishes) are considered to be one of the most globally imperiled taxon, with 25 of the 27 species listed by the International Union for the Conservation of Nature (IUCN). Overharvest, habitat degradation, fragmentation and water quality issues have contributed to their decline worldwide. These stressors have been ameliorated in some areas, but in others they remain a limiting factor to sturgeon. Barriers impeding upstream migrations to natural spawning areas and manifesting alterations to natural flows continue to compromise sturgeon recruitment and limit natural recovery. Watersheds in the Northern Hemisphere have been categorized as being strongly affected, moderately affected or unaffected based on the degree of fragmentation and water flow regulation. An overlay (i.e. intersect) of the sturgeons' status with this watershed categorization revealed that a small area remains in which sturgeon are not considered at risk and where rivers are unaffected in northern Canada. These relatively unperturbed populations provide a much needed opportunity to learn about sturgeon biology, habitat needs and reproductive potential in a natural riverine environment, which may facilitate conservation and recovery efforts in affected watersheds.

KEY WORDS: Acipenseriformes · Fragmentation · Imperiled

INTRODUCTION

Acipenseriformes (an order of basal ray-finned fishes that includes the sturgeons and paddlefishes) are considered to be one of the most globally imperiled vertebrate groups (Birstein et al. 1997, Pikitch et al. 2005). Sturgeon are unique and have been described as 'living fossils', dating back to the Lower Jurassic some 200 million years ago (Bemis & Kynard 1997, Bemis et al. 1997b, Pikitch et al. 2005) and have undergone few morphological changes since that time (Choudhury & Dick 1998). Rivers dominated the landscape during the period when sturgeon purportedly evolved; fossil records excavated strictly from these historical drainages support an ancestral af-

finity for flowing waters (Choudhury & Dick 1998). As such, extant species are highly migratory (Auer 1996, Welsh & McLeod 2010), spawning only in freshwater—including the many marine, anadromous sturgeon species (Rochard et al. 1990, Bemis & Kynard 1997). There are currently 27 recognized species of Acipenseriformes (25 species of sturgeon and 2 species of paddlefish) and they are found only in the Northern Hemisphere (Fig. 1; Bemis & Kynard 1997, Bemis et al. 1997a, Birstein et al. 1997), which is consistent with all sturgeon fossil locations (Bemis et al. 1997a). Eurasia supports the greatest diversity with 17 species of sturgeon, while North America hosts the remaining 10 species (Birstein 1993, Birstein et al. 1997, Billard & Lecointre 2001).

*Corresponding author: tim.haxton@ontario.ca




Fig. 1. Global distribution of the 27 Acipenseriformes species (depicted in gray)

GLOBAL DECLINE OF ACIPENSERIFORMES

Sturgeon are a resilient group, having adapted to large-scale global changes (Bemis & Kynard 1997) and survived mass global species extinctions (Jackson & Johnson 2000, Wang et al. 2014). However, many populations have become decimated despite their historical abundance and widespread distribution across the Northern Hemisphere (Birstein 1993, Bemis & Findeis 1994, DeVore et al. 1995). Non-regulated harvests for the highly sought caviar and meat of many sturgeon species initiated their decline in the mid-1800s in North America (Birstein 1993, Bemis & Findeis 1994, Pikitch et al. 2005), exacerbated stock declines from the Caspian Sea in the early 1960s (Khodorevskaya et al. 1997, Ivanov et al. 1999, Khodorevskaya & Krasikov 1999) and continue to depress or limit the recovery of some populations (Khodorevskaya et al. 1997, Williamson 2003, Peterson et al. 2007, Johnson & Iyengar 2015). Major alterations to many rivers were concurrent with, or subsequent, to overharvests. Dams were constructed on large rivers for mechanical or hydroelectric power,

for navigation purposes or to facilitate transport of material downstream, directly blocking access to and/or altering sturgeon spawning habitat. Dramatic changes to water quality due to use of rivers for disposal of industrial or municipal effluent, deforestation and urbanization of watersheds were all prevalent practices in the period post over-harvesting concurrent with dam construction (Rochard et al. 1990, Birstein et al. 1997, Lenhardt et al. 2006). Consequently, riverine fish (i.e. fluvial-dependent species) including Acipenseriformes are among the most endangered species globally (Cooke et al. 2012).

RECENT IMPROVEMENTS AND CONTINUED THREATS

More recently, water quality conditions in many regions have improved (Hensel & Holcik 1997, Schram et al. 1999, Kampa et al. 2014) and stricter harvest controls have been implemented (Khodorevskaya et al. 1997, Baker & Borgeson 1999, Peterson et al. 2007) in addition to the enforcement of import




Fig. 2. Large river systems in the Northern Hemisphere as categorized by Dynesius & Nilsson (1994). Red: areas in which the large river systems are strongly affected by river regulation, amber: moderately affected areas, green: unaffected area

restrictions on many sturgeon species through the Convention on International Trade in Endangered Species (Williamson 2003, Lenhardt et al. 2006, Ludwig 2006). However, poaching purportedly remains a conservation concern for all sturgeon species, given the exorbitant prices of caviar (Cohen 1997, Pikitch et al. 2005); this seems to be especially relevant for European stocks (Khodorevskaya et al. 1997, Ivanov et al. 1999, Ludwig 2006). In addition, hydroelectric dams are a long-term fixture on the landscape. In fact, hydro development has recently undergone a renaissance as a source of renewable energy to meet growing demands (Nilsson et al. 2005, Grill et al. 2015, Zarfl et al. 2015), with over 3700 large dams either planned or currently under construction globally (Zarfl et al. 2015). River fragmentation has been extensive, with in excess of 45 000 large dams (i.e. >15 m height) in existence globally (Nilsson et al. 2005). Fragmentation has contributed to local extirpation of sturgeon in numerous rivers (Bemis & Findley 1994, Ferguson & Duckworth 1997, Hensel & Holcik 1997). Dams act as barriers to upstream movement

to historic spawning areas (Birstein 1993, DeVore et al. 1995, Wei et al. 1997), congregating migrating sturgeon and exposing them to harvest and the perils of poaching (Cohen 1997, Fernández-Pasquier 1999), and altering flows, affecting the success of natural sturgeon recruitment within rivers (Khoroshko 1972, Ivanov et al. 1999, Jager et al. 2002, Goto et al. 2015, Haxton et al. 2015). Moreover, variation in sturgeon abundance across the landscape has been attributed to the presence of hydroelectric generating stations (DeVore et al. 1995, Haxton & Findley 2008, Haxton et al. 2014), demonstrating that dams can play a significant role in limiting the recovery processes for sturgeon.

The degree of fragmentation and water flow regulation has been categorized on a watershed basis for large river systems (those with mean annual discharge of at least $350 \text{ m}^3 \text{ s}^{-1}$ in a river channel within a catchment) on a global scale (Dynesius & Nilsson 1994, Nilsson et al. 2005). River watersheds have been classified, based on level of fragmentation and flow alteration, as strongly affected, moderately

Table 1. IUCN Red List status of sturgeon and paddlefish by continent in 2016. CR: Critically Endangered; EN: Endangered; VU: Vulnerable; NT: Near Threatened; LC: Least Concern

Species	Authority	Continent	Red List status	Year assessed
<i>Acipenser baerii</i>	Brandt, 1869	Eurasia	EN	2010
<i>Acipenser brevirostrum</i>	Lesueur, 1818	North America	VU	2004
<i>Acipenser dabryanus</i>	Duméril, 1869	Eurasia	CR	2010
<i>Acipenser fulvescens</i>	Rafinesque, 1817	North America	LC	2004
<i>Acipenser gueldenstaedtii</i>	Brandt & Ratzeburg, 1833	Eurasia	CR	2010
<i>Acipenser medirostris</i>	Ayres, 1854	North America	NT	2006
<i>Acipenser mikadoi</i>	Hilgendorf, 1892	Eurasia	CR	2010
<i>Acipenser naccarii</i>	Bonaparte, 1836	Eurasia	CR	2013
<i>Acipenser nudiventris</i>	Lovetsky, 1828	Eurasia	CR	2010
<i>Acipenser oxyrinchus</i>	Mitchill, 1815	North America	NT	2006
<i>Acipenser persicus</i>	Borodin, 1897	Eurasia	CR	2010
<i>Acipenser ruthenus</i>	Linnaeus, 1758	Eurasia	VU	2010
<i>Acipenser schrenckii</i>	Brandt, 1869	Eurasia	CR	2010
<i>Acipenser sinensis</i>	Gray, 1835	Eurasia	CR	2010
<i>Acipenser stellatus</i>	Pallas, 1771	Eurasia	CR	2010
<i>Acipenser sturio</i>	Linnaeus, 1758	North America	CR	2010
<i>Acipenser transmontanus</i>	Richardson, 1836	North America	LC	2004
<i>Huso dauricus</i>	Georgi, 1775	Eurasia	CR	2010
<i>Huso huso</i>	Linnaeus, 1758	Eurasia	CR	2010
<i>Pseudoscaphirhynchus fedtschenkoi</i>	Kessler, 1872	Eurasia	CR	2010
<i>Pseudoscaphirhynchus hermanni</i>	Kessler, 1877	Eurasia	CR	2010
<i>Pseudoscaphirhynchus kaufmanni</i>	Kessler, 1877	Eurasia	CR	2010
<i>Scaphirhynchus albus</i>	Forbes & Richardson, 1905	North America	EN	2004
<i>Scaphirhynchus platyrhynchus</i>	Rafinesque, 1820	North America	VU	2004
<i>Scaphirhynchus suttkusi</i>	Williams & Clemmer, 1991	North America	CR	2004
<i>Polyodon spathula</i>	Walbaum, 1792	North America	VU	2004
<i>Psephurus gladius</i>	Martens, 1862	Eurasia	CR	2010

affected or unaffected (Fig. 2; Dynesius & Nilsson 1994, Nilsson et al. 2005). Rivers in which less than one-quarter of the main channel was left without dams were considered strongly affected. Unaffected rivers contained no dams in their main channel, although dams in tributaries were permitted if the mean annual discharge was changed by less than 2%. Rivers that fell between these 2 categories were considered moderately affected (Dynesius & Nilsson 1994). In the Northern Hemisphere, within the sturgeon's historical range, approximately 75% of river discharge was considered to have been moderately to strongly affected by river regulation (Dynesius & Nilsson 1994).

Impacts of habitat fragmentation resulting from dam construction are widespread and well documented (Wozney et al. 2011, McDermid et al. 2011, Haxton et al. 2014). Barriers can present significant obstacles to the continued presence of sturgeon in historically occupied waters given that sturgeon often utilize entire river systems (Welsh & McLeod 2010, Gerig et al. 2011) to find the habitat necessary to fulfill the requirements of particular stages of the life cycle (Smith & King 2005). While safe, unimpeded movement up and downstream at barriers could

possibly ameliorate the situation, passage efficiency in the limited instances where it is known to occur has been minimal (Cooke et al. 2002, Parsley et al. 2007, Thiem et al. 2011, 2016) and at a high energetic cost to the fish (Thiem et al. 2016). Recovery of sturgeon populations in highly fragmented rivers could therefore be limited or reliant on artificial propagation given the negative effects dams could have on recruitment (e.g. Khodorevskaya et al. 1997, Khodorevskaya & Krasikov 1999, Schram et al. 1999). Furthermore, the effects of fragmentation and adult mortality can be confounding. Fragmentation can affect recruitment, which impedes recovery, whereas poaching generally targets adults. Species or populations that are vulnerable to poachers may never naturally recover when they inhabit a fragmented river.

INTERSECT: RIVER FRAGMENTATION AND STURGEON STATUS

These multiple stressors have led the International Union on the Conservation of Nature (IUCN; www.iucnredlist.org) to list (i.e. categorize based on the risk of extinction) 25 of the 27 Acipenseriformes (Table 1,




Fig. 3. IUCN's Red List status of the 27 sturgeon and paddlefish species globally. Status is indicated by colour — red: Critically Endangered or Endangered; peach: Near Threatened or Vulnerable; green: Least Concern

Fig. 3) worldwide. Only 2 sturgeon species, lake sturgeon *Acipenser fulvescens* Rafinesque, 1817 and white sturgeon *Acipenser transmontanus* Richardson, 1836, are considered by the IUCN to be not at risk (i.e. categorized as Least Concern) globally (Table 1); however, their status varies widely across their natural range at national or state/province scales.

On a global scale, few areas remain in which sturgeon populations are not considered to be at risk and where large rivers remain intact. An intersect (a geometric intersection; see the Appendix) of river fragmentation categorization (Fig. 2; Dynesius & Nilsson 1994, Nilsson et al. 2005) on sturgeon distribution and status (Fig. 3) identified those areas where sturgeon populations were not at risk and where rivers were not fragmented (Fig. 4). White sturgeon, indigenous to the west coast of North America, are anadromous, spawning in only 3 rivers: the Sacramento, Columbia and Fraser Rivers (Billard & Lecointre 2001). These rivers are all highly fragmented and as such, extensive management has been required to sustain or facilitate recovery of white sturgeon popu-

lations (e.g. Ireland et al. 2002, Parsley et al. 2002, 2007, Crossman & Hildebrand 2014). Lake sturgeon, a potamodromous species, is the only species of sturgeon not considered at risk by the IUCN that inhabits an area categorized as unaffected by fragmentation and barriers (Fig. 4). Northern Canada supports populations of lake sturgeon that purportedly avoided decimation in the late 1800s and early 1900s given the remoteness of the area (Bemis & Findeis 1994, Ferguson & Duckworth 1997). While robust lake sturgeon populations exist in areas categorized as strongly affected (e.g. Lake Winnebago, Wisconsin, Bruch 1999; the St. Lawrence River, Quebec, Mailhot et al. 2011; Lake of the Woods, Ontario, Rusak & Mosindy 1997), they are generally associated with large lakes accompanied with altered tributaries.

SUMMARY AND RECOMMENDATIONS

Sturgeon recovery efforts have historically been extensive and expensive (Bacalbasa-Dobrovici &




Fig. 4. Intersect of large river system categorization (Dynesius & Nilsson 1994) with sturgeon status and distribution (see Appendix). Dark green: areas where the species is not listed and where watersheds are unaffected. Light green: areas where species are listed but watersheds are unaffected. Amber: areas where species are not listed or listed as Near Threatened or Vulnerable with moderate fragmentation. Red: areas that are highly fragmented and species are listed

Patriche 1999, Lenhardt et al. 2006, 2008, Guti 2014) in areas where populations have drastically declined, such as the Danube River (Hensel & Holcik 1997) or Yangtze River (Wei et al. 1997, Dudgeon 2010). The relatively unperturbed populations in northern Canada provide an opportunity to learn about sturgeon biology, and to identify habitat needs and reproductive potential in a natural riverine environment. There is an identified need to learn more about endangered riverine fish, especially in unperturbed natural ecosystems (Cooke et al. 2012). Such research could help ascertain the life-history requirements of a fluvial-dependent species, which may facilitate conservation and recovery efforts in perturbed systems, and potentially unaltered systems in northern regions.

Acknowledgements. We thank Ron Bruch (WI, USA), Paul Sampson, Sarah Hogg, Josh Annett, Glenn Desy and Beth Wright (Ontario Ministry of Natural Resources and Forestry)

for their insightful comments on the manuscript. And a special thanks to Kevin Middel (Ontario Ministry of Natural Resources and Forestry) for GIS advice.

LITERATURE CITED

- Auer NA (1996) Importance of habitat and migration to sturgeons with emphasis on lake sturgeon. *Can J Fish Aquat Sci* 53(Suppl 1):152–160
- Bacalbasa-Dobrovici N, Patriche N (1999) Environmental studies and recovery actions for sturgeon in the lower Danube River system. *J Appl Ichthyol* 15:114–115
- Baker EA, Borgeson DJ (1999) Lake sturgeon abundance and harvest in Black Lake, Michigan, 1975–1999. *N Am J Fish Manage* 19:1080–1088
- Bemis WE, Findeis EK (1994) The sturgeons' plight. *Nature* 370:602
- Bemis WE, Kynard B (1997) Sturgeon rivers: an introduction to acipenseriform biogeography and life. *Environ Ecol Stat* 48:167–183
- Bemis WE, Birstein VJ, Waldman JR (1997a) Sturgeon biodiversity and conservation: an introduction. *Environ Biol Fishes* 48:13–14

- Bemis WE, Findeis EK, Grande L (1997b) An overview of Acipenseriformes. *Environ Biol Fishes* 48:25–71
- Billard R, Lecointre G (2001) Biology and conservation of sturgeon and paddlefish. *Rev Fish Biol Fish* 10:355–392
- Birstein VJ (1993) Sturgeons and paddlefishes: threatened fishes in need of conservation. *Conserv Biol* 7:773–787
- Birstein VJ, Bemis WE, Waldman JR (1997) The threatened status of acipenseriform species: a summary. *Environ Biol Fishes* 48:427–435
- Bruch RM (1999) Management of lake sturgeon on the Winnebago System - long term impacts of harvest and regulations on population structure. *J Appl Ichthyol* 15:142–152
- Choudhury A, Dick TA (1998) The historical biogeography of sturgeons (Osteichthyes: Acipenseridae): a synthesis of phylogenetics, palaeontology and palaeogeography. *J Biogeogr* 25:623–640
- Cohen A (1997) Sturgeon poaching and black market caviar: a case study. *Environ Biol Fishes* 48:423–426
- Cooke DW, Leach SD, Isely JJ (2002) Behaviour and lack of upstream passage of shortnose sturgeon at a hydroelectric facility and navigation lock complex. In: Van Winkle W, Anders PJ, Secor DH, Dixon D (eds) *Biology, management, and protection of North American sturgeon*. *Am Fish Soc Symp Ser* 28:101–110
- Cooke SJ, Paukert C, Hogan Z (2012) Endangered river fish: factors hindering conservation and restoration. *Endang Species Res* 17:179–191
- Crossman JA, Hildebrand LR (2014) Evaluation of spawning substrate enhancement for white sturgeon in a regulated river: effects on larval retention and dispersal. *River Res Appl* 30:1–10
- DeVore JD, James BW, Tracy CA, Hale DA (1995) Dynamics and potential production of white sturgeon in the unimpounded Lower Columbia River. *Trans Am Fish Soc* 124:845–856
- Dudgeon D (2010) Editorial requiem for a river: extinctions, climate change and the last of the Yangtze. *Aquat Conserv* 20:127–131
- Dynesius M, Nilsson C (1994) Fragmentation and flow regulation of river systems in the northern third of the world. *Science* 266:753–762
- Ferguson MM, Duckworth GA (1997) The status and distribution of lake sturgeon, *Acipenser fulvescens*, in the Canadian provinces of Manitoba, Ontario and Quebec: a genetic perspective. *Environ Biol Fishes* 48:299–309
- Fernández-Pasquier V (1999) *Acipenser sturio* L. in the Guadaquivir river, Spain. Water regulation and fishery as factors in stock decline from 1932 to 1967. *J Appl Ichthyol* 15:133–135
- Gerig B, Moerke A, Greil R, Koproski S (2011) Movement patterns and habitat characteristics of lake sturgeon (*Acipenser fulvescens*) in the St. Marys River, Michigan, 2007–2008. *J Gt Lakes Res* 37:54–60
- Goto D, Hamel MJ, Hammen JJ, Rugg ML, Pegg MA, Forbes VE (2015) Spatiotemporal variation in flow-dependent recruitment of long-lived riverine fish: model development and evaluation. *Ecol Modell* 296:79–92
- Grill G, Lehner B, Lumsdon AE, MacDonald GK, Zar C, Liermann CR (2015) An index-based framework for assessing patterns and trends in river fragmentation and flow regulation by global dams at multiple scales. *Environ Res Lett* 10:015001
- Guti G (2014) Can anadromous sturgeon populations be restored in the middle Danube River? *Acta Zool Bulg* 66: 63–67
- Haxton TJ, Findlay CS (2008) Variation in lake sturgeon (*Acipenser fulvescens*) abundance and growth among river reaches in a large regulated river. *Can J Fish Aquat Sci* 65:645–657
- Haxton TJ, Friday M, Cano T, Hendry C (2014) Variation in lake sturgeon (*Acipenser fulvescens* Rafinesque, 1817) abundance in rivers across Ontario, Canada. *J Appl Ichthyol* 30:1335–1341
- Haxton TJ, Friday M, Cano T, Hendry C (2015) Assessing the magnitude of effect of hydroelectric production on lake sturgeon abundance in Ontario. *N Am J Fish Manage* 35:930–941
- Hensel K, Holcik J (1997) Past and current status of sturgeons in the upper and middle Danube River. *Environ Biol Fishes* 48:185–200
- Ireland SC, Beamesderfer RCP, Paragamian VL, Wakkinen VD, Siple JT (2002) Success of hatchery-reared juvenile white sturgeon (*Acipenser transmontanus*) following release in the Kootenai River, Idaho, USA. *J Appl Ichthyol* 18:642–650
- Ivanov VP, Vlasenko AD, Khodorevskaya RP, Raspopov VM (1999) Contemporary status of Caspian sturgeon (Acipenseridae) stock and its conservation. *J Appl Ichthyol* 15:103–105
- Jackson JBC, Johnson KG (2000) Life in the last few million years. *Paleobiology* 26:221–235
- Jager HI, Van Winkle W, Chandler JA, Lepla KB, Bates P, Counihan TD (2002) A simulation study of factors controlling white sturgeon recruitment in the Snake River. *Am Fish Soc Spec Publ* 28:127–150
- Johnson TA, Iyengar A (2015) Phylogenetic evidence for a case of misleading rather than mislabeling in caviar in the United Kingdom. *J Forensic Sci* 60:S248–S253
- Kampa J, Hatzenbeler G, Jennings M (2014) Status and management of lake sturgeon (*Acipenser fulvescens* Rafinesque, 1817) in the upper St. Croix River and Namekagon River, Wisconsin, USA. *J Appl Ichthyol* 30: 1387–1392
- Khodorevskaya RP, Krasikov YV (1999) Sturgeon abundance and distribution in the Caspian Sea. *J Appl Ichthyol* 15:106–113
- Khodorevskaya RP, Dovgopol GF, Zhuravleva OL, Vlasenko AD (1997) Present status of commercial stocks of sturgeons in the Caspian Sea basin. *Environ Biol Fishes* 48: 209–219
- Khoroshko PN (1972) The amount of water in the Volga basin and its effect on the reproduction of sturgeon (Acipenseridae) under conditions of normal and regulated discharge. *J Ichthyol* 12:608–616
- Lenhardt M, Jaric I, Kalauzi A, Cvijanovic G (2006) Assessment of extinction risk and reasons for decline in sturgeon. *Biodivers Conserv* 15:1967–1976
- Lenhardt M, Jari I, Cvijanovi G, Smederevac-Lali M (2008) The key threats to sturgeons and measures for their protection in the lower Danube Region. In: Lagutov V (ed) *Rescue of sturgeon species in the Ural River basin*. NATO Science for Peace and Security Series C: Environmental Security. Springer, Dordrecht, 87–96
- Ludwig A (2006) A sturgeon view on conservation genetics. *Eur J Wildl Res* 52:3–8
- Mailhot Y, Dumont P, Vachon N (2011) Management of the lake sturgeon *Acipenser fulvescens* population in the lower St Lawrence River (Québec, Canada) from the 1910s to the present. *J Appl Ichthyol* 27:405–410
- McDermid JL, Wozney KM, Kjartanson SL, Wilson CC

- (2011) Quantifying historical, contemporary, and anthropogenic influences on the genetic structure and diversity of lake sturgeon (*Acipenser fulvescens*) populations in northern Ontario. *J Appl Ichthyol* 27:12–23
- Nilsson C, Reidy CA, Dynesius M, Revenga C (2005) Fragmentation and flow regulation of the world's large river systems. *Science* 308:405–408
- Parsley MJ, Anders PJ, Miller AI, Beckman LG, McCabe GT Jr (2002) Recovery of white sturgeon populations through natural production: understanding the influence of abiotic and biotic factors on spawning and subsequent recruitment. In: Van Winkle W, Anders PJ, Secor DH, Dixon D (eds) *Biology, management, and protection of North American sturgeon*. *Am Fish Soc Symp Ser* 28: 55–66
- Parsley MJ, Wright CD, van der Leeuw BK, Kofoot EE, Peery CA, Moser ML (2007) White sturgeon (*Acipenser transmontanus*) passage at the Dalles Dam, Columbia River, USA. *J Appl Ichthyol* 23:627–635
- Peterson DL, Vecsei P, Jennings CA (2007) Ecology and biology of the lake sturgeon: a synthesis of current knowledge of a threatened North American Acipenseridae. *Rev Fish Biol Fish* 17:59–76
- Pikitch EK, Doukakis P, Lauck L, Chakrabarty P, Erickson DL (2005) Status, trends and management of sturgeon and paddlefish fisheries. *Fish Fish* 6:233–265
- Rochard E, Castelnaud G, LePage M (1990) Sturgeons (Pisces: Acipenseridae); threats and prospects. *J Fish Biol* 37:123–132
- Rusak JA, Mosindy T (1997) Seasonal movements of lake sturgeon in Lake of the Woods and the Rainy River, Ontario. *Can J Zool* 75:383–395
- Schram ST, Lindgren J, Evrard LM (1999) Reintroduction of lake sturgeon in the St. Lois River, western Lake Superior. *N Am J Fish Manage* 19:815–823
- Smith KM, King DK (2005) Movement and habitat use of yearling and juvenile lake sturgeon in Black Lake, Michigan. *Trans Am Fish Soc* 134:1159–1172
- Thiem JD, Binder TR, Dawson JW, Dumont P and others (2011) Behaviour and passage success of upriver-migrating lake sturgeon *Acipenser fulvescens* in a vertical slot fishway on the Richelieu River, Quebec, Canada. *Endang Species Res* 15:1–11
- Thiem JD, Dawson JW, Hatin D, Danylchuk AJ and others (2016) Swimming activity and energetic costs of adult lake sturgeon during fishway passage. *J Exp Biol* 219: 2534–2544
- Wang Y, Sadler PM, Shen S, Erwin DH and others (2014) Quantifying the process and abruptness of the end-Permian mass extinction. *Paleobiology* 40:113–129
- Wei Q, Ke F, Zhang J, Zhuang P, Luo J, Zhou R, Yang W (1997) Biology, fisheries, and conservation of sturgeons and paddlefish in China. *Environ Biol Fishes* 48:241–255
- Welsh AB, McLeod DT (2010) Detection of natural barriers to movement of lake sturgeon (*Acipenser fulvescens*) within the Namakan River, Ontario. *Can J Zool* 88:390–397
- Williamson DF (2003) *Caviar and conservation*. TRAFFIC North America, World Wildlife Fund, Washington, DC
- Wozney KM, Haxton TJ, Kjartanson SL, Wilson CC (2011) Genetic assessment of lake sturgeon (*Acipenser fulvescens*) population structure in the Ottawa River. *Environ Biol Fishes* 90:183–195
- Zarfl C, Lumsdon AE, Tockner K (2015) A global boom in hydropower dam construction. *Aquat Sci* 77:161–170

Appendix.

To determine areas of low fragmentation and non-threatened sturgeon populations, an intersect was conducted in ArcMap v.10.1 (ESRI) of the categorization of fragmentation in large river systems (Fig. 2) (Dynesius & Nilsson 1994, Nilsson et al. 2005) and IUCN's Red List status (Fig. 3). A numerical value was assigned to the categorization for both fragmentation and status (fragmentation: unaffected = 1, moderately affected = 5, strongly affected = 10; status: Least Concern = 1; Near Threatened = 2, Vulnerable = 3, Endangered = 4, Critically Endangered = 5). The product of the 2 values provided an indication of Red List status and fragmentation (Fig. 4). A value of 1 represents areas where the species is not listed and watersheds are unaffected. Values between 2 and 9 represents areas where species are listed but watersheds are unaffected. Values between 10 and 19 represent areas where species are not listed or listed as Near Threatened or Vulnerable with moderate fragmentation. Values greater than 20 represent areas that are highly fragmented and species are listed.