arXiv:2012.02365v2 [math.AP] 10 Jan 2022

A HELE-SHAW LIMIT WITHOUT MONOTONICITY

NESTOR GUILLEN, INWON KIM, AND ANTOINE MELLET

ABSTRACT. We study the incompressible limit of the porous medium equation with a right hand side representing either a source or a sink term, and an injection boundary condition. This model can be seen as a simplified description of non-monotone motions in tumor growth and crowd motion, generalizing the congestion-only motions studied in recent literature ([AKY14], [PQV14], [KP18], [MPQ17]). We characterize the limit density, which solves a free boundary problem of Hele-Shaw type in terms of the limit pressure. The novel feature of our result lies in the characterization of the limit pressure, which solves an obstacle problem at each time in the evolution.

1. INTRODUCTION

The porous media equation is a nonlinear evolution equation which is commonly used to model many natural phenomena involving diffusion or heat propagation. In its simplest form, it consists of a continuity equation with a flux given by Darcy's law:

(1.1)
$$\partial_t \rho - \operatorname{div}\left(\rho \nabla p\right) = 0, \quad p = \frac{m}{m-1} \rho^{m-1}, \qquad m > 1.$$

The exponent m > 1 describes the anti-crowd tendency of the density motion, where the diffusion is larger at higher density ([BGHP84], [BH86], [Mur07], [Wit97]). Due to the degeneracy of the diffusion at lower densities, it is well-known that the density stays compactly supported if initially so (see for example [V07, Chapter 1]).

In this paper, we consider the porous media equation with a source term:

(1.2)
$$\partial_t \rho - \operatorname{div}(\rho \nabla p) = \lambda \rho \quad \text{in } \Omega \times \mathbb{R}^+,$$

set in the (exterior) domain $\Omega = \mathbb{R}^n \setminus K$, where K is a bounded subset of \mathbb{R}^n with smooth boundary and supplemented with the "injection" boundary condition,

(1.3)
$$\rho(x,t) = f(x,t)^{\frac{1}{m-1}} > 0 \quad \text{on } \partial K.$$

as well as the initial data $\rho(t=0) = \rho_m^0 \ge 0$. Importantly we will assume that the function $\lambda(x,t)$ is bounded but can take both positive or negative values. When $\lambda < 0$, the term $\lambda \rho$ is an absorption term which is competing with the injection at ∂K . Assumptions on the initial and boundary data are given in Section 2.1. Classically, (1.2) can also be written as the following equation for the pressure p(x,t):

(1.4)
$$\partial_t p = (m-1)p(\Delta p + \lambda) + |\nabla p|^2.$$

Our interest is with the *incompressible limit* of this equation, that is the limit $m \to \infty$. Heuristically speaking, if (ρ_m, p_m) denotes a sequence of solution of (1.2), then – provided there is an actual limit in a good enough sense – the limits ρ_{∞} and p_{∞} should satisfy

(1.5)
$$\partial_t \rho_\infty - \operatorname{div}(\rho_\infty \nabla p_\infty) = \lambda \rho_\infty \quad \text{in } \Omega \times \mathbb{R}^+, \qquad p_\infty = f \quad \text{on } \partial K, \qquad \rho_\infty(\cdot, 0) = \rho^0,$$

and taking the limit in the relation $p_m = \frac{m}{m-1}\rho_m^{m-1}$, we may guess that in the limit ρ_{∞} and p_{∞} are connected by what is known as the Hele-Shaw graph

(1.6)
$$p_{\infty} \in P_{\infty}(\rho_{\infty}) := \begin{cases} 0 & \text{if } 0 \le \rho_{\infty} < 1\\ [0,\infty) & \text{if } \rho_{\infty} = 1\\ \infty & \text{if } \rho_{\infty} > 1. \end{cases}$$

In particular, the pressure can be viewed as a Lagrange multiplier for the constraint $\rho_{\infty} \leq 1$ ([MRCS10]). In our framework, as in many of the related works discussed below, a priori estimates (under appropriate assumptions on f and on the initial data) will allow us to make the derivations of (1.5)-(1.6) rigorous.

Equations (1.5)-(1.6) fully characterize the evolution of ρ_{∞} (see the uniqueness result, Proposition 2.5). However, one would like to give a more geometrical description of the evolution of ρ_{∞} , and in particular of the evolution of the "saturated region"

$$\Sigma(t) := \{\rho_{\infty}(t) = 1\}.$$

Classically, such a description is provided by a Hele-Shaw type free boundary problem. Indeed, formally at least, we can pass to the limit in (1.4) to get the so-called *complementarity condition*:

(1.7)
$$p_{\infty}(\Delta p_{\infty} + \lambda) = 0 \text{ in } \Omega \times \mathbb{R}^{+}$$

which implies that $p_{\infty}(\cdot, t)$ solves $-\Delta p_{\infty} = \lambda$ in the set $\{p_{\infty}(\cdot, t) > 0\}$, and equation (1.5) implies (in a weak form) that the normal velocity of the interface $\partial \Sigma(t)$ is proportional to $|\nabla p_{\infty}|$. However, the derivation of (1.7) is less straightforward than that of (1.5)-(1.6) in general (see for instance [DP21]), and it is not obvious that we should always have $\{p_{\infty}(\cdot, t) > 0\} = \Sigma(t)$.

Incompressible limits were first studied for equation (1.1) (that is when $\lambda = 0$). In the absence of K, there are classical works starting by Bénilan and Crandall [BC81], followed by results with more general initial data by Caffarelli and Friedman [CF87] and numerical studies describing the shape of the limit by Elliot et al [EHKO86] (also see [GQ03] for its rigorous justification). In [BKM09] a similar weak formulation for the Hele-Shaw problem (still without right hand side) is derived as a "mesa" limit from the Stefan problem. The last decade has seen significant advances in the study of these asymptotics when the right hand side is monotone increasing in ρ – corresponding to the case $\lambda > 0$ in our framework. The convergence as $m \to \infty$ and characterization of the limit as a Hele-Shaw type flow has been achieved for models of congested crowd motion [AKY14, KPW19] and of tumor growth [PQV14, MPQ17, KP18]. It is important to note that monotonicity properties are present in the systems studied in these papers and are essential for proving that $\{p_{\infty}(\cdot, t) > 0\} = \Sigma(t)$. For instance, the monotonicity of the density was a key feature in characterizing the limiting problem in [MPQ17, KP18]. In [AKY14, KPW19] which features a drift field, the monotonicity of ρ along the streamline was crucial to characterize the limiting problem in terms of viscosity solutions.

In our work, the function $\lambda(x, t)$ is not necessarily positive so that one no longer expects ρ_{∞} to be monotone in time thus complicating the analysis. Moreover, a Hele-Shaw type problem with a single phase is typically monotone in time, suggesting that the lack of monotonicity should be reflected by having some modification of the one-phase Hele-Shaw model in the limit. One of the main contribution of this paper is to identify the pressure $p_{\infty}(\cdot, t)$ for all time t > 0 by showing that it solves an obstacle problem in the set $\Sigma(t)$ and might thus be such that $\{p_{\infty}(\cdot, t) > 0\} \subseteq \Sigma(t)$ (see Theorem 2.7), causing the saturated set $\Sigma(t)$ to shrink. Though our result appears to be new, its proof is relatively simple and can be generalized to problems with nonlinear source terms. As an illustration of this latter point, we apply these ideas to a tumor growth model which involves nonlinear terms (see Appendix A). Even in the monotone cases mentioned above, our result provides a new approach to the derivation of the complementarity condition (1.7). Equation (1.2) is simple but it allows us to study a very general and important behavior. Indeed, the monotonicity in the aforementioned works is characteristic of systems with only congestive effects. However, it is clear that de-congestion effects are important for applications. In [PQV14], a model for tumor growth which takes into account the evolution of the density of nutrients is introduced and studied. In that case, the tumor cells decrease their density in the event of insufficient nutrient, which yields to "de-congestion" or recession of the tumor cells. The consequent lack of monotonicity significantly complicates the analysis: The derivation of the complementarity condition was only achieved recently [DP21] and the geometric description of the tumor growth still remains to be understood. Similarly, the study of congested crowd motion that involve decongestion phenomena is of great interest (see [MRCS10],[San18]).

Our interest in studying the toy problem (1.2) is thus to better understand such behavior. By allowing λ to take both positive and negative value, we generate a motion that consists of both congestion and de-congestion. The presence of a fixed boundary condition on ∂K is by no mean necessary for our analysis (there is no K in the tumor growth model studied in Appendix A), but such injection boundary conditions are a classical feature of Hele-Shaw problems. In the context of crowd motion, our model describes a congested crowd coming out of the door (∂K) to the outdoors ($\mathbb{R}^n \setminus K$). In the context of the classical Hele-Shaw flow (with $\lambda = 0$) the boundary condition describe the injection of the fluids.

In our setting it seems natural to expect that p_{∞} , which acts against congestion, may vanish even when the density is fully saturated. Indeed we will see that when λ is not necessarily positive the support of the pressure $p_{\infty}(t)$ may be a strict subset of $\Sigma(t)$. In general, $p_{\infty}(t)$ must be found by solving an obstacle problem in $\Sigma(t)$. As a result, while $\Sigma(t)$ will expand according to a Hele-Shaw type law when $|\nabla p_{\infty}| > 0$ along $\partial \Sigma(t)$, it might recede when $|\nabla p_{\infty}| = 0$. Formally, the motion law of $\Sigma(t)$ can be written as

(1.8)
$$|\nabla p_{\infty}| = (1 - \rho^E) V \text{ on } \partial \Sigma(t),$$

where V denotes the outer normal velocity of $\partial \Sigma(t)$ and ρ^E is the trace of the "external density", namely the trace of ρ_{∞} on $\partial \Sigma(t)$ from $\{\rho_{\infty} < 1\}$ (this is well defined if $\partial \Sigma(t)$ smooth since ρ_{∞} is in $BV_{loc}(\mathbb{R}^n \setminus K)$).

The velocity law (1.8) can be formally justified from the weak equation (1.5) as follows (where ν denotes the inward normal unit vector on ∂K):

$$\begin{split} \int_{\partial K} \rho \nabla p \cdot \nu \, dS + \int_{\Omega} \lambda \rho \, dx &= \frac{d}{dt} \int_{\Omega} \rho \, dx = \frac{d}{dt} \left[\int_{\Sigma(t)} \rho \, dx + \int_{\Omega \setminus \Sigma(t)} \rho^E \, dx \right] \\ &= \int_{\Sigma(t)} \partial_t \rho \, dx + \int_{\Omega \setminus \Sigma(t)} \partial_t \rho^E \, dx + \int_{\partial \Sigma(t)} V(1 - \rho^E) \, dS \\ &= \int_{\Sigma(t)} \operatorname{div} \left(\rho \nabla p \right) + \lambda \rho \, dx + \int_{\Omega \setminus \Sigma(t)} \lambda \rho^E \, dx + \int_{\partial \Sigma(t)} V(1 - \rho^E) \, dS \\ &= \int_{\partial K} \rho \nabla p \cdot \nu \, dS + \int_{\partial \Sigma(t)} \rho \nabla p \cdot \nu \, dS + \int_{\Omega} \lambda \rho \, dx + \int_{\partial \Sigma(t)} V(1 - \rho^E) \, dS, \end{split}$$
from which we deduce (since $\rho = 1$ in $\Sigma(t)$) that $\int_{\Omega} [\nabla p \cdot \nu + V(1 - \rho^E)] \, dS = 0$

from which we deduce (since $\rho = 1$ in $\Sigma(t)$) that $\int_{\partial \Sigma(t)} [\nabla p \cdot \nu + V(1 - \rho^E)] dS = 0.$

We note that our motion law is different from [KPW19] where the free boundary can move back and forth under the action of a force field. Here the receding and advancing behavior of the free boundary takes place via completely different mechanisms. The motion law (1.8) is closer to the one obtained in [KM14] in the context of liquid drops sliding down on inclined plane. In this context, at the receding end of the drop, the contact angle between the liquid drop and the plane may vanish. In that moment the nature of the velocity law suddenly changes: it is no longer dictated by the local value of the pressure, but rather by the bulk behavior of the liquid via an obstacle problem.

Finally, we believe that our approach developed for the model problem (1.2) is quite general and is of independent interest. To illustrate this point we prove in Appendix A that it can be applied to the tumor growth problem with nutrient, considered in [PQV14, DP21].

Here is a brief outline of the paper. In Section 2 we collect and discuss implications of our results. In Section 3 we show convergence of the density and pressure variables. In section 4 we derive the novel characterization of the pressure via an obstacle problem. Section 5 introduces the comparison principle, as well as the uniqueness, of the limit problem, which will be used in the rest of the paper. Section 6 - 8 describes the motion law of the saturated region, starting with the measure theoretic representation in Section 6. An alternative characterization, in the flavor of viscosity solutions, is given in Sections 7-8.

1.1. Acknowledgements. The authors would like to acknowledge the generous support of the National Science Foundation. Inwon Kim was partially supported by National Science Foundation grant DMS-1900804 and Antoine Mellet was partially supported by National Science Foundation grant DMS-2009236.

2. NOTATIONS AND MAIN RESULTS

2.1. Assumptions. Throughout the paper, we denote by (ρ_m, p_m) the solution of the following initial boundary value problem:

(2.1)
$$\begin{cases} \partial_t \rho_m - \operatorname{div}\left(\rho_m \nabla p_m\right) = \lambda \rho_m & \text{in } Q, \qquad p_m = \frac{m}{m-1} \rho_m^{m-1}, \\ \rho_m(x,t) = f(x,t)^{\frac{1}{m-1}} & \text{on } \partial K \times \mathbb{R}_+ \\ \rho_m(0,x) = \rho_m^0(x) & \text{in } \Omega \end{cases}$$

where we denote

$$\Omega := \mathbb{R}^n \setminus K, \quad Q := \Omega \times \mathbb{R}_+, \quad Q_T := \Omega \times (0, T].$$

Below are the main assumptions to be used throughout our analysis:

Assumption 2.1. There is a constant $\Lambda > 0$ such that (i) The function $\lambda(x,t)$ satisfies

(2.2) $|\lambda(x,t)| \le \Lambda \quad \forall (x,t) \in Q,$

(2.3)
$$\lambda \in BV_{loc}(\Omega \times \mathbb{R}_+).$$

(ii) The boundary data f(x,t) satisfies

(2.4)
$$0 < \Lambda^{-1} \le f \le \Lambda, \quad |\nabla f| \le C, \quad |\partial_t f| \le C \quad on \; \partial K \times \mathbb{R}^+.$$

In order to write the assumptions on the initial condition ρ_m^0 , we first introduce appropriate barriers. Given $0 \leq \underline{R} < \overline{R}$, we consider $\overline{\varphi}(x)$ and $\varphi(x)$ solutions of

(2.5)
$$-\Delta\overline{\varphi} = \Lambda + 1 \text{ in } B_{\overline{R}} \setminus K, \quad \overline{\varphi} = f^{\frac{m}{m-1}} \text{ on } \partial K, \quad \overline{\varphi} = 0 \text{ on } \partial B_{\overline{R}}$$

and

(2.6)
$$-\Delta \underline{\varphi} = -\Lambda \text{ in } B_{\underline{R}} \setminus K, \quad \underline{\varphi} = f^{\frac{m}{m-1}} \text{ on } \partial K, \quad \underline{\varphi} = 0 \text{ on } \partial B_{\underline{R}},$$

where we assume that $\underline{\varphi} > 0$ in $B_{\underline{R}} \setminus K$ (if necessary we can replace $B_{\underline{R}}$ by a smaller set sufficiently close to K).

Assumption 2.2.

(i) The initial condition $\rho_m^0(x)$ satisfies

(2.7)
$$\underline{\varphi}(x)^{\frac{1}{m}} \le \rho_m^0(x) \le \overline{\varphi}^{\frac{1}{m}}(x) \qquad \forall x \in \Omega$$

(2.8)
$$\|\Delta(\rho_m^{0\ m}) + \lambda \rho_m^0\|_{L^1(\Omega)} + \|\nabla \rho_m^0\|_{L^1(\Omega)} \le C$$

(ii) The sequence $\{\rho_m^0\}_{m>1}$ converges in $L^1(\Omega)$ to ρ^0 .

Condition (2.8) may seem restrictive, but the following result shows that a wide range of initial condition ρ^0 can fit into this framework:

Lemma 2.3. Let $\Sigma \supset K$ be a bounded open set with C^2 boundary in \mathbb{R}^n and let $\rho^0(x)$ be given by $\rho^0 := \chi_{\Sigma} + \rho^E \chi_{\Sigma C}$ in Ω ,

where $\rho^E \in C_c^{1,1}(\Omega)$ satisfies $0 \le \rho^E < 1$. Then there exists a sequence ρ_m^0 satisfying Assumption 2.2.

The construction is simple, so we give it here: First, we define the pressure p_0 by

$$-\Delta p_0 = 0$$
 in $\Sigma \setminus K$ with $p_0 = 0$ on $\partial \Sigma$ and $p_0 = f$ on K.

We clearly have $p_0 \ge 0$ in K and $|\nabla p_0| \ne 0$ on $\partial \Sigma$. We can then define

$$\rho_m^0 := \max\{p_0^{1/m}, (\rho^E - a_m)_+\},\$$

where a_m is a nonnegative sequence such that $a_m \to 0$ and $(1 - a_m)^m \to 0$ as $m \to \infty$ (for instance $a_m = (\ln m)^{-1}$). Note that with this definition (2.7) holds for sufficiently large m. To check (2.8), first note that $p_0^{1/m}$ is in BV, since

$$\|Dp^{1/m}\|_{L^1} = \left\|\frac{p^{\frac{1}{m}-1}}{m}Dp\right\|_{L^1} \le C \sup_{p\ge 1/m} |Dp| + o(1),$$

for sufficiently large m, where we have used the fact that p grows at most linearly near the regular boundary $\partial \Sigma$. Lastly, note that

$$(\rho_m^0)^m = \max\{p_0, (\rho^E - a_m)^m\},\$$

which is a maximum of two C^2 functions. Moreover for large m we have $\nabla(p_0 - (\rho^E - a_m)^m) \neq 0$ where they coincide, since $\nabla p_0 \neq 0$ due to the regularity of $\partial \Sigma$ and $\nabla(\rho^E - a_m)^m$ uniformly vanishes as m grows. This nondegeneracy yields the regularity of the set $\Gamma := \{p_0 = (\rho^E - a_m)^m\}$. Collecting the facts we conclude (2.8), where $\Delta \rho_m^{0 m}$ is interpreted as a measure.

2.2. Limit and weak formulation of the limiting problem. By generalizing classical a priori estimates to our equation, we will first establish the convergence of ρ_m and p_m and prove the following result:

Theorem 2.4. Under Assumptions 2.1 and 2.2 and up to a subsequence, the density ρ_m and pressure p_m solution of (2.1) converge strongly in $L^1(Q_T)$ for all T > 0 to limits ρ_∞ and p_∞ which satisfy

$$\rho_{\infty}, \ p_{\infty} \in BV(Q_T),$$

$$\rho_{\infty} \in C^s([0,\infty); H^{-1}(\Omega)) \quad \forall s < 1/2, \quad p_{\infty} \in L^2(0,T; H^1(\Omega)),$$

$$0 \le p_{\infty}(x,t) \le C \quad a.e. \ (x,t) \in Q, \quad 0 \le \rho_{\infty}(x,t) \le 1 \quad a.e. \ x \in \Omega, \ \forall t > 0$$

and

(2.9)
$$\begin{cases} \partial_t \rho_{\infty} = \Delta p_{\infty} + \lambda \rho_{\infty} & \text{ in } \mathcal{D}'(\Omega \times \mathbb{R}^+), \qquad p_{\infty} \in P_{\infty}(\rho_{\infty}); \\ p_{\infty}(x,t) = f(x,t) & \text{ on } \partial K \times \mathbb{R}^+; \\ \rho_{\infty}(x,0) = \rho^0(x) & \text{ in } \Omega, \end{cases}$$

where P_{∞} is the Hele-Shaw graph (1.6).

Following [PQV14], we can prove the following result which shows that the result above fully characterizes the function ρ_{∞} :

Proposition 2.5. Suppose $\lambda \in L^2([0,T]; H^1(\Omega))$, then equation (2.9) has at most one solution $(\rho, p) \in X := L^{\infty}(\Omega \times (0,T]) \times L^2(0,T; H^1(\Omega)).$

Furthermore, if (ρ_1, p_1) and (ρ_2, p_2) are respectively sub and super-solutions of (2.9) in X satisfying $\rho_1(\cdot, 0) \leq \rho_2(\cdot, 0)$ and $p_1|_{\partial K} \leq p_2|_{\partial K}$, then $\rho_1 \leq \rho_2$ in $\Omega \times \mathbb{R}^+$.

Remark 2.6. This uniqueness result implies in particular that any subsequence of (ρ_m, p_m) converges to the same limit, and thus the entire sequence converges to $(\rho_{\infty}, p_{\infty})$.

When $\lambda = 0$, equation (2.9) implies that the saturated region $\Sigma(t) = \{\rho_{\infty}(t) = 1\}$ coincides with the set $\{p_{\infty}(t) > 0\}$, and $\Sigma(t)$ evolves according to the classical Hele-Shaw free boundary problem:

$$\begin{cases} \Delta p_{\infty} = 0 \text{ in } \Sigma(t), \quad p_{\infty} = f \text{ on } \partial K, \quad p_{\infty} = 0 \text{ on } \partial \Sigma(t); \\ V = |\nabla p_{\infty}| \text{ on } \partial \Sigma(t), \end{cases}$$

where V denotes the outer normal velocity of the interface $\partial \Sigma(t)$ ([Kim03], [QV99]). This provides a simple geometric description of the evolution of the set { $\rho_{\infty} = 1$ }. As explained in the introduction, our goal in this paper is to provide a similar characterization when $\lambda \neq 0$.

2.3. The pressure $p_{\infty}(t)$. Our first task is to determine how the pressure $p_{\infty}(t)$ depends on the set $\{\rho_{\infty} = 1\}$. An important and new feature in our framework is that that we may have $\{p_{\infty}(t) > 0\} \subsetneq \{\rho_{\infty}(t) = 1\}$. Indeed we prove that $p_{\infty}(t)$ is determined by solving an obstacle problem in the set $\{\rho_{\infty} = 1\}$.

First, we note that for all $t_0 \ge 0$ we have $p_{\infty} \in BV(\Omega \times (t_0, T))$ and so we can define the trace of the function p_{∞} on $\{t = t_0\}$. The interested reader might consult Giusti's book [Giu84, Chapter 2] for a thorough discussion on traces of BV functions (it is worth emphasizing that p_{∞} is of bounded variation in space and time). We denote this trace $p^+(x, t_0)$ since it is defined as a limit as $t \to t_0^+$. It satisfies in particular

(2.10)
$$\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} |p_{\infty}(x,t) - p^+(x,t_0)| \, dx \, dt \le \int_{t_0}^{t_0+\delta} \int_{\Omega} |\partial_t p_{\infty}| \, dx \to 0 \qquad \text{as } \delta \to 0.$$

and (by Lebesgue differentiation theorem) $p_{\infty}(x,t) = p^+(x,t)$ almost everywhere. Since $\lambda \in BV$, we can similarly define the trace $\lambda^+(\cdot,t)$ for all t > 0. We then prove:

Theorem 2.7. Under the conditions of Theorem 2.4 and for all $t \ge 0$, $p^+(\cdot, t)$ is the unique solution of the minimization problem

(2.11)
$$\min_{v \in E_t} \int_{\Omega} \frac{1}{2} |\nabla v|^2 - \lambda^+ (\cdot, t) v \, dx$$

where E_t denotes the functional space

$$E_t = \left\{ v \in H^1(\Omega) \cap L^1(\Omega) \, ; \, v = f \text{ on } \partial K, \, v \ge 0 \text{ in } \Omega, \, \langle v, 1 - \rho_\infty(t) \rangle_{H^1, H^{-1}} = 0 \right\}.$$

Equivalently, $p^+(\cdot, t)$ is the unique solution of the variational inequality

(2.12)
$$\begin{cases} p \in E_t \\ \int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda^+ (\cdot, t) (p-u) \, dx \le 0 \qquad \forall u \in E_t. \end{cases}$$

If the set $\Sigma(t) = \{\rho_{\infty}(\cdot, t) = 1\}$ is a smooth enough subset of Ω , then (2.11) is a classical obstacle problem in $\Sigma(t)$ with Dirichlet boundary conditions p = f on ∂K , p = 0 on $\partial \Sigma(t)$. The proof of this result is surprisingly simple and quite flexible (see Section 4). It does not require any additional a priori estimates besides the ones already used to prove Theorem 2.4. It can easily be adapted to more complicated models, such as the tumor growth model with nutrient, as we show in Appendix A (see Proposition A.2).

By using the approach developed in [MRCS10], it is also possible to show that for any weak solutions of (2.9), the pressure $p_{\infty}(\cdot, t)$ satisfies, for a.e. t > 0

$$\int_{\Omega} \nabla p \cdot \nabla u - \lambda u \, dx = 0, \qquad \forall u \in E_t.$$

So the pressure $p_{\infty}(\cdot, t)$ solves the equation $\Delta p + \lambda = 0$ in the set $\{\rho_{\infty}(t) = 1\}$ for almost every time. As explained in the introduction, we cannot expect this to hold for all time, since either λ or the set $\{\rho_{\infty}(t) = 1\}$ may evolve discontinuously over time. In the event where the solution of the obstacle problem (2.12) has its support strictly smaller that $\{\rho_{\infty}(t) = 1\}$, the set $\{\rho_{\infty}(t) = 1\}$ will shrink instantaneously. The result of [MRCS10] does not see these instantaneous collapses (which can happen over a large set of time, albeit one of measure zero). Our characterization of p_{∞} , which holds for all time t > 0, identifies how such collapses take place.

When $\lambda = 0$, Theorem 2.7 provides a simple proof of the harmonicity of p_{∞} in $\{\rho_{\infty}(\cdot, t) = 1\}$. In the general case, it implies in particular the so-called complementarity condition:

$$p_{\infty}(\Delta p_{\infty} + \lambda) = 0 \text{ in } \mathcal{D}'(\Omega \times (0, \infty))$$

which is readily obtained by taking $u = p(1 \pm \epsilon \varphi)$ in (2.12) with $\varphi \in \mathcal{D}(\Omega \times (0, \infty))$ and ϵ small enough so that $1 \pm \epsilon \varphi \geq 0$.

This complementarity condition is proved for the tumor growth model in [PQV14] (model without nutrient) and in [DP21] (model with nutrient). In both cases, the derivation relies on further estimates on the pressure (in particular the Aronson-Bénilan estimate or some variant of it). Our result thus provides an alternative derivation of this condition that does not require any of these additional estimates.

Given the interest for the complementarity condition in the literature, it is worth noting that it is equivalent to the obstacle problem formulation in the following sense:

Proposition 2.8. Let $(\rho, p) \in L^{\infty}(0, T; L^{1}(\Omega) \cap L^{\infty}(\Omega)) \times L^{2}(0, T; H^{1}(\Omega))$ be a solution of (2.9) with $p \in BV_{loc}(\Omega \times \mathbb{R}_{+})$. If p satisfies the complementarity condition

$$p(\Delta p + \lambda) = 0 \text{ in } \mathcal{D}'(\Omega \times (0, \infty))$$

then for every t > 0 the trace $p^+(\cdot, t)$ (as defined in (2.10)) is the unique solution of problem (2.11).

Note that given a weak solution of (2.9), we are not able to prove directly that it satisfies the obstacle problem formulation of Theorem 2.7) or the complementarity condition, but this proposition shows that these two properties are equivalent.

In general little is known on the boundary regularity of the set $\{\rho_{\infty}(\cdot, t) = 1\}$, including whether its boundary has measure zero. Thus for pointwise characterization of the pressure p_{∞} , we define the support of the measure $1 - \rho_{\infty}$ by

Supp
$$(1 - \rho_{\infty}(t)) := \left\{ x_0 \in \Omega; \int_{B_r(x_0)} (1 - \rho_{\infty})(\cdot, t) \, dx > 0 \text{ for all } r > 0 \right\}.$$

While it may differ from the set $\{\rho_{\infty} < 1\}$ by a measure zero set, this set has the advantage of being closed by its definition. Then the solution of the obstacle problem (2.11) has the usual properties in the open set

(2.13)
$$\mathcal{O}(t) := \Omega \setminus \operatorname{Supp}\left(1 - \rho_{\infty}(t)\right) = \left\{ x_0 \in \Omega : \int_{B_r(x_0)} (1 - \rho_{\infty})(\cdot, t) = 0 \text{ for some } r > 0 \right\}$$

which can be seen as the "interior" of the set $\{\rho_{\infty}(\cdot, t) = 1\}$. More precisely, we have:

Proposition 2.9. The function p^* , solution of the minimization problem (2.11), is in $C^{1,1}_{loc}(\mathcal{O}(t))$ and satisfies

(2.14)
$$-\Delta p^* = \lambda \chi_{\{p^* > 0\}} \text{ in } \mathcal{O}(t).$$

2.4. Velocity law: Measure theoretic results. In view of Theorem 2.7, we can redefine p_{∞} a.e. so that for each time t > 0 the function $p_{\infty}(\cdot, t)$ is the unique solution of the obstacle problem (2.11). We would now like to characterize the evolution of the saturated region. We start with the following proposition:

Proposition 2.10. For all t > 0, $\mathcal{P}(t) := \{p_{\infty}(\cdot, t) > 0\}$ the positivity set of the solution of the obstacle problem (2.11). Then the density equation in (2.9) can be rewritten as

(2.15)
$$\partial_t \rho_{\infty} = \mu_t + \lambda \rho_{\infty} (1 - \chi_{\mathcal{P}})$$

here $\mu_t := \Delta p_{\infty}(\cdot, t) + \lambda(\cdot, t) \chi_{\mathcal{P}(t)}$, which is a non-negative Radon measure supported in $\partial \mathcal{P}(t) \setminus \mathcal{O}(t)$.

When $\lambda \leq 0$, Equation (2.15) shows that the growth of ρ_{∞} can only occur when the measure μ is non zero (thus only on $\partial \mathcal{P}(t) \setminus \mathcal{O}(t)$) while the density can only decay when $\rho_{\infty}(1 - \chi_{\mathcal{P}(t)}) > 0$. Growth and decay thus take place according to different mechanisms. One is dictated by a singular measure, the other by an L^{∞} function. Note that $\mathcal{P}(t)$ is almost the saturated set $\Sigma(t)$, in the sense that their parabolic closures coincide (see Theorem 2.11.)

Heuristically, (2.15) can have a geometric interpretation as follows. Since $\rho_{\infty} = 1$ in $\mathcal{P}(t)$, we can always write

$$\rho_{\infty}(x,t) = \chi_{\mathcal{P}(t)}(x) + \rho^{E}(x,t)(1-\chi_{\mathcal{P}(t)}(x))$$

for some function ρ^E . Splitting the singular and regular part of (2.15), we get the following:

(2.16)
$$\begin{cases} (1 - \chi_{\mathcal{P}(t)}(x))(\partial_t \rho^E - \lambda \rho^E) = 0; \\ (1 - \rho^E(x, t))\partial_t \chi_{\mathcal{P}(t)} = \mu. \end{cases}$$

The first equation determines the value of ρ_{∞} outside of the congested set $(\partial_t \rho^E = \lambda \rho^E$ when p(x,t) = 0, supplemented by the condition that $\rho^E = 1$ when p(x,t) > 0).

Formally, we have $\mu = |\nabla p| dS$, where S is the surface measure on $\partial \mathcal{P}(t)$, so if $|\nabla p(x_0, t_0)| \neq 0$, the second equation in (2.16) gives $(1 - \rho^E)V(x_0, t_0) = |\nabla p(x_0, t_0)|$ (expansion of the congested region), while if $|\nabla p(x_0, t_0)| = 0$, then either $\partial_t \chi_{\mathcal{P}(t)} = 0$ or $\rho^E(x_0, t_0) = 1$. The later can only happen if $\chi_{\mathcal{P}(t)}(x_0) = 1$ as $t \to t_0^-$ and so $\partial_t \chi_{\mathcal{P}(t)} \leq 0$ (retraction of the congested region). Altogether, this gives the free boundary condition (1.8), assuming that the boundary of $\mathcal{P}(t)$ coincides with $\Sigma(t)$.




FIGURE 1. The external density ρ^E has boundary values on red and blue parts of the boundary: $\rho^E = 1$ on the red parts, and $\rho^E = \rho^0$ on the blue part.

Making these statement rigorous in the classical framework would require the development of a regularity theory which is not the topic of the present paper. Instead, in what follows, we will use the comparison principle to make sense of this in the spirit of viscosity solutions.

2.5. Velocity law: barrier approach. We define the external density ρ^E in the set $\{p = 0\}$ by solving the first-order equation $\partial_t \rho = \lambda \rho$ with appropriate boundary condition. More precisely, given x, the open set $\text{Int}(\{t; p(x,t) = 0\})$ can be written as $\bigcup_{i \in I} (a_i, b_i)$ and $\rho^E(x, t)$ for $t \in (a_i, b_i)$ is the solution of the first order ODE $\partial_t \rho = \lambda \rho$ with initial condition

$$\rho^{E}(x, a_{i}) = \begin{cases} \rho_{0}(x) & \text{if } a_{i} = 0\\ 1 & \text{if } a_{i} > 0 \end{cases}$$

See Figure 1 above for an illustration.

With this definition of ρ^E , and using the comparison principle for the limiting problem (Proposition 2.5), we obtain the following description on the motion of the congested zone $\{\rho_{\infty} = 1\}$:

Theorem 2.11. Let $(\rho, p) \in L^{\infty}(Q) \times L^{2}_{loc}(0, \infty; H^{1}(\Omega))$ be a weak solution of (2.9) with initial data $0 \leq \rho^{0}(x) \leq 1$. Then the following holds:

(a) If $\lambda \in C(Q_T) \cap L^2(0,T; H^1(\Omega))$, then we have, in the sense of comparison with barriers,

(2.17)
$$(1-\rho^E)V = |\nabla p| \quad on \ \partial\{\rho=1\}.$$

(b) ρ^E coincides with ρ a.e. outside of $\overline{\{\rho=1\}}$.

(c) If λ is negative, then for any T > 0

$$\overline{\{p>0\}\cap Q_T} = \overline{\{p>0\}}\cap Q_T = \overline{\{\rho=1\}}\cap Q_T.$$

The barriers used to make sense of (2.17) are local smooth sub- and super-solutions of (2.9). Their description can be found in in section 7. Such comparison property is akin to the viscosity solutions approach taken by [KP18] for $\lambda > 0$. We do not touch upon the issue of whether the barrier properties are enough for a complete characterization of the limit solution: see [KPW19] and [KP18] for analysis in this direction.

Part (c) in above theorem says that when λ is negative, the closure of the pressure support coincides with that of $\{\rho = 1\}$, and that the congested zone $\{\rho = 1\}$ cannot all of a sudden expand. This is not true when λ is positive, due to the nucleation of the congested zone generated by the

growth of the external density. The set $\{\rho = 1\}$ certainly can discontinuously shrink. For instance if λ decreases over time, the pressure decreases and the set $\{\rho = 1\}$ may start shrinking. While shrinking, if a component of the set gets disconnected at $t = t_0$ from K, the pressure in this region will drop to zero and ρ will immediately decrease below one after t_0 . Such scenario makes it difficult to describe ρ^E in an explicit way, except when λ only increases over time.

Theorem 2.12. Suppose that $\lambda \in C(\Omega \times [0,T]) \cap L^2(0,T; H^1(\Omega))$ is non-decreasing over time, and let (ρ, p) be the weak solution of (2.9) with initial data $\rho^0 \in BV$. Then the set $\{p(\cdot,t) > 0\}$ is monotone increasing in time. Moreover for all $t \ge 0$

$$\rho(\cdot,t) = \chi_{\Sigma(t)} + \rho^E \chi_{\mathbb{R}^n \setminus \Sigma(t)}, \text{ where } \rho^E(x,t) := \rho^0(x) \exp^{\int_0^t \lambda(x,s) ds}.$$

In particular $\overline{\Sigma(t)} = \overline{\{\rho(\cdot,t)=1\}}$ for all t > 0.

If ρ^0 is a characteristic function and $\Sigma_0 = \{\rho^0 = 1\} = \{\rho^0 > 0\}$, then ρ remains a characteristic function for all positive times.

Note that we may initially have $\{p(\cdot, 0) > 0\}$ as a strict subset of $\{\rho^0 = 1\}$. In this case this last theorem states that $\{\rho = 1\}$ experiences an initial discontinuous shrinkage.

2.6. Numerical examples. Figure 2.6 shows the evolution of the density and pressure in a simple framework to illustrate the receding and expanding motion of the free boundary. We consider the one dimension porous media equation

$$\partial_t \rho - \partial_x (\rho \partial_x p) = \lambda(t) \rho$$
 in $(0, \infty) \times (0, T)$, $p = \frac{m}{m-1} \rho^{m-1}$

with the boundary condition $\rho(0) = 1$ and m = 40 (so we are close to the limiting problem. In particular, the density is close to, but not equal to 1 when p > 0). The coefficient $\lambda(t)$ is independent of x but changes value discontinuously in time:

(2.18)
$$\lambda(t) = \begin{cases} -1 & \text{if } t \in [0, .75) \\ -5 & \text{if } t \in [.75, 1) \\ -1 & \text{if } t \ge 1. \end{cases}$$

The set $\{p(t) > 0\}$ is expanding with finite speed for $t \in (0, .75)$ (first row) and receding instantaneously at $t = .75^+$. The density is then decreasing for $t \in [.75, 1)$ in the region where p = 0 since $\partial_t \rho = -5\rho$ in that region (second row). Finally, for t > 1 (third row) the set $\{p(t) > 0\}$ is again expanding with finite speed.

3. Proof of Theorem 2.4

The proof of this theorem uses many classical techniques (see in particular [PQV14]), though we have to be careful with the two main differences between our framework and that of [PQV14]: the lack of sign of λ and the presence of the fixed boundary ∂K .

3.1. Notion of solutions for (2.1). First, we recall some well known facts about the porous media equation (2.1) (we refer the interested reader to [V07], Chapters 5 (Definition 5.5 and Theorem 5.14).

Definition 3.1. For $\rho^0 \in L^1(\Omega)$, $g \in L^2(0,T; H^1(\Omega))$ and $\lambda \in L^1(Q_T)$, we say that a non-negative function $\rho \in L^1(Q_T)$ is a weak solution of (2.1) with $\rho_m^0 = \rho^0$ and $f := g^{1-1/m}$ if

- (i) $\rho^m \in L^2(0,T; H^1(\Omega))$ with its trace on $\partial K \times [0,T]$ equal to g;
- (ii) $\rho \in L^2(Q_T);$




FIGURE 2. Graph of the density (upper curve) and pressure (lower curve) when m = 40 and λ given by (2.18). The functions are shown at time $t = 0^+$, t = 0.35, $t = 0.75^-$, $t = 0.75^+$, t = 0.9, $t = 1^-$, $t = 1^+$, t = 1.2 and t = 1.8

(iii) ρ satisfies the identity

$$\int \int_{Q_T} (\rho \partial_t \psi - \nabla \rho^m \cdot \nabla \psi + \lambda \psi) dx dt = -\int_{\Omega} \rho_m^0(x) \psi(x,0) dx$$

for any function $\psi \in C^1(Q_T)$ which vanishes on $\partial K \times [0,T]$ and for t = T.

Existence of a weak solution can be established by approximation with smooth functions, which either solves the porous media equation with strictly positive initial data or solves a regularized equation with strictly positive diffusion (see Theorem 5.14 of [V07]). Uniqueness of the weak solution is a consequence of the following comparison principle, which we will use often in our analysis.

Lemma 3.2. Let ρ and $\tilde{\rho}$ be two weak solutions of (2.1) with initial data $\rho_m^0, \tilde{\rho}_m^0$ and fixed boundary data f and \tilde{f} . If $\rho_m^0 \leq \tilde{\rho}^0$ a.e. and $f \leq \tilde{f}$ a.e., then $\rho \leq \tilde{\rho}$ a.e.

3.2. Maximum principle: L^{∞} bounds for ρ_m and p_m and $\nabla p_m \cdot \nu|_{\partial K}$.

Lemma 3.3. Under conditions (2.2), (2.4) and (2.7), and for all T > 0, there exists a constant C = C(T) > 0 independent of m such that the following holds:

For sufficiently large m (depending on T) the pressure p_m satisfies:

(3.1) $0 \le p_m(x,t) \le C \qquad \text{for all } (x,t) \in Q_T,$

and

(3.2)
$$-C \le \nu(x) \cdot \nabla p_m(x,t) \le C \quad \text{for all } x \in \partial K, \quad 0 \le t \le T.$$

Moreover,

(3.3)
$$\rho_m(x,t) \to 1$$
, locally uniformly in $U \times \mathbb{R}_+$

for some neighborhood U of K and

(3.4)
$$\operatorname{supp} \rho_m(\cdot, t) \subset B_{\overline{R}+C(T)} \quad \text{for all } t \in (0, T).$$

Remark 3.4. Note that by (3.3), ρ_m stays uniformly positive and solves a uniformly parabolic equation in U. It is thus smooth, a fact we will use repeatedly when dealing with the boundary data on ∂K .

Proof. We fix T > 0. This lemma follows from the maximum principle for the pressure p_m , which, we recall solves

$$\partial_t p = (m-1)p(\Delta p + \lambda) + |\nabla p|^2.$$

In view of (2.6), $\underline{\varphi}(x)$ satisfies $\Delta \underline{\varphi} + \lambda = \Lambda + \lambda \ge 0$ and is therefore a subsolution for this equation. Assumption (2.7) thus implies

$$p_m(x,t) \ge \varphi(x) \qquad \forall (x,t) \in \Omega \times \mathbb{R}^+.$$

For the upper barrier, we define the function $\bar{u}(x,t)$ as follows: For all t > 0, the function $x \mapsto \bar{u}(x,t)$ solves

$$-\Delta v = \Lambda + 1$$
 in $B_{R(t)} \setminus K$, $v = f^{\frac{m}{m-1}}$ on ∂K , $v = 0$ on $\partial B_{R(t)}$

where $R(t) := \overline{R} + \int_0^t M(s)ds$, with $M(s) := 2 \sup_{x \in \partial B_{R(t)}} |\nabla \overline{u}(\cdot, t)|$. The function \overline{u} is extended by 0 outside $B_{R(t)}$. Since R(t) depends on $\overline{u}(x,t)$, the function \overline{u} can be constructed for instance by discrete-time approximation. We note that (2.5) implies in particular that $\overline{u}(x,0) = \overline{\varphi}(x)$

We claim then \bar{u} is a supersolution for the pressure equation for sufficiently large m. To see this, note first that when $\bar{u} \ge (m-1)^{-1/2}$ we have

$$\partial_t \bar{u} \ge 0 \ge (m-1)\bar{u}(\Delta \bar{u} + \lambda) + |\nabla \bar{u}|^2$$
 if $m \ge \sup_{0 \le t \le T} |\nabla \bar{u}|^4 (\cdot, t)$.

On the other hand, since $\partial_t \bar{u} \geq 2|\nabla \bar{u}|^2 > 0$ on its zero level set $\partial B_{R(t)}$, it is clear that for small enough $\epsilon = \epsilon(T)$ we have $\partial_t \bar{u} \geq |\nabla \bar{u}|^2$ in $0 \leq \bar{u} \leq \epsilon$. Our claim follows if m is large enough (depending on T).

The comparison principle for the pressure equation now yields:

$$\varphi(x) \le p_m(x,t) \le \bar{u}(x,t) \qquad \forall (x,t) \in Q_T.$$

The results now follow: (3.1) follows from upper bound, while the lower bound together with the fact that $\rho_m \sim p_m^{\frac{1}{m-1}}$ implies (3.3). The fact that \bar{u} is supported in $B_{R(t)}$ implies (3.4) and since $\varphi(x) = p_m(x,t) = \bar{u}(x,t)$ on ∂K , we get

$$-C \le \nu(x) \cdot \nabla \underline{\varphi}(x) \le \nu(x) \cdot \nabla p_m(t,x) \le \nu(x) \cdot \nabla \overline{u}(x,t) \le C \qquad \forall x \in \partial K, \ t \in (0,T].$$

3.3. L^1 bounds for ρ_m and p_m .

Lemma 3.5. For all T > 0, there exists a constant C(T) depending on Λ and T such that

(3.5)
$$\|\rho_m(t)\|_{L^1(\Omega)} \le \|\rho_m^0\|_{L^1(\Omega)} e^{\Lambda T} + C(T)$$

and

(3.6)
$$\|p_m(t)\|_{L^1(\Omega)} \le C \|\rho_m^0\|_{L^1(\Omega)} e^{\Lambda T} + C(T)$$

for $t \in [0, T]$ and $m \ge 2$.

Proof. Integrating (2.1) on Ω yields

$$\frac{d}{dt} \int_{\Omega} \rho_m(t) \, dx = \int_{\Omega} \lambda(t) \rho_m(t) \, dx + \int_{\partial K} \rho_m \nabla p_m \cdot \nu \, dS$$
$$\leq \Lambda \int_{\Omega} \rho_m(t) \, dx + C,$$

where we used (3.1), (3.2). The bound (3.5) follows by a Gronwall argument. The bound (3.6) then follows from (3.5) and (3.1) since $p_m = \frac{m}{m-1}\rho_m^{m-1} \leq \frac{m}{m-1}C\rho_m$.

3.4. Bounds on the derivatives of ρ_m and p_m . For $\delta > 0$, we define

$$\Omega^{\delta} := \{ x \in \mathbb{R}^n \, ; \, \operatorname{dist}(x, K) > \delta \}.$$

Lemma 3.6. For any $\delta > 0$, there exists a constant C_{δ} independent on m such that

(3.7)
$$\|\partial_t \rho_m(t)\|_{L^1(\Omega^{\delta})} \le C_{\delta} \qquad \forall t > 0$$

(3.8)
$$\|\partial_{x_i}\rho_m(t)\|_{L^1(\Omega^{\delta})} \le C_{\delta} \qquad \forall t > 0$$

Similarly, denoting by B_R the ball of radius R, we have the following bounds:

(3.9)
$$\|\partial_t p_m\|_{L^1((0,T)\times\Omega^\delta\cap B_R)} \le C_{\delta,R,T}$$

(3.10)
$$\|\partial_{x_i} p_m\|_{L^1((0,T) \times \Omega^{\delta} \cap B_R)} \le C_{\delta,R,T}$$

Proof. Proceeding as in [PQV14], we differentiate the first equation in (2.1) with respect to time and multiply it by $sign(\partial_t \rho_m)$ and use Kato's inequality to obtain

(3.11)
$$\partial_t |\partial_t \rho_m| - \Delta(m\rho_m^{m-1}|\partial_t \rho_m|) \le \lambda |\partial_t \rho_m| + \rho_m |\partial_t \lambda| \quad \text{in } \Omega.$$

We cannot simply integrate this equation over Ω because of the boundary condition on ∂K . Instead, given a large ball B_R such that $K \subset B_R$, we introduce the function φ such that $\varphi = 0$ on ∂K , $\varphi = 1$ on ∂B_R , $\Delta \varphi = 0$ in $B_R \setminus K$ and we extend this function by 1 outside B_R . This function satisfies

 $\varphi|_{\partial K} = 0, \quad \Delta \varphi \le 0 \text{ in } \Omega, \quad \varphi > 0 \text{ in } \Omega.$

Multiplying (3.11) by φ and integrating over Ω , and using the fact that $\varphi|_{\partial K} = 0$ and $m\rho_m^{m-1}|\partial_t\rho_m| = \frac{m}{m-1}f^{\frac{1}{m-1}}\partial_t f$ on ∂K , we deduce

$$\begin{split} \frac{d}{dt} \int_{\Omega} |\partial_t \rho_m| \varphi \, dx &\leq \int_{\Omega} m\rho_m^{m-1} |\partial_t \rho_m| \Delta \varphi \, dx - \int_{\partial K} m\rho_m^{m-1} |\partial_t \rho_m| \nabla \varphi \cdot \nu \, dS \\ &+ \int_{\Omega} \lambda |\partial_t \rho_m| \varphi \, dx + \int_{\Omega} \rho_m |\partial_t \lambda| \varphi \, dx \\ &\leq C + \Lambda \int_{\Omega} |\partial_t \rho_m| \varphi \, dx + \int_{\Omega} \rho_m |\partial_t \lambda| \varphi \, dx. \end{split}$$

Since $\partial_t \rho_m(0) = \Delta \rho_{in}^m + \lambda \rho_m^0$, the bound (2.8) implies $\|\partial_t \rho_m(0)\|_{L^1(\Omega)} \leq C$ and using (3.4), we deduce

$$\|\partial_t \rho_m(t)\varphi\|_{L^1(\Omega)} \le C(T) + C \int_0^T e^{\Lambda(T-t)} \int_{B_{R_0+CT}} |\partial_t \lambda(x,s)| \, dx \, dt$$

and (3.7) follows from (2.3) and the fact that $\min_{\Omega^{\delta}} \varphi > 0$ for all $\delta > 0$ (by the strong maximum principle).

To get an estimate on $\partial_t p_m$, we want to take advantage of the term $\int_{\Omega} m \rho_m^{m-1} |\partial_t \rho_m| \Delta \varphi \, dx$ in the inequality above. We thus define, for $\eta > 0$, φ_η such that $\varphi_\eta = 0$ on ∂K , $\varphi_\eta = 1$ on ∂B_R , $\Delta \varphi_\eta = -\eta$ in $B_R \setminus K$ and we extend this function by 1 outside B_R .

Given R, we claim that $\Delta \varphi_{\eta} \leq 0$ in Ω if η is sufficiently small (depending on R). Indeed, Hopf's Lemma implies $x \cdot \nabla \varphi_0 > 0$ on ∂B_R , so the C^1 -convergence of φ_{η} to φ_0 implies $x \cdot \nabla \varphi_{\eta} \geq 0$ on ∂B_R .

Proceeding as above, we get:

$$\frac{d}{dt} \int_{\Omega} |\partial_t \rho_m| \varphi_\eta \, dx + 4\eta \int_{\Omega} m \rho_m^{m-1} |\partial_t \rho_m| \, dx \le C(T)$$

Integrating in t, we deduce that for all $\delta > 0$, R > 0 and T, there exists $C_{\delta,R,T}$ such that

$$\int_0^T \int_{\Omega^\delta \cap B_R} m\rho_m^{m-1} |\partial_t \rho_m| \, dx \, dt \le C_{\delta,R,T}.$$

Finally, we write

$$\begin{split} \int_0^T \int_{\Omega^\delta \cap B_R} |\partial_t p_m| \, dx \, dt &= \int_0^T \int_{\Omega^\delta \cap B_R} m\rho_m^{m-2} |\partial_t \rho_m| \, dx \, dt \\ &\leq \int_0^T \int_{\Omega^\delta \cap B_R \cap \{\rho_m < 1/2\}} m(1/2)^{m-2} |\partial_t \rho_m| \, dx \, dt \\ &\quad + 2 \int_0^T \int_{\Omega^\delta \cap B_R \cap \{\rho_m > 1/2\}} m\rho_m^{m-1} |\partial_t \rho_m| \, dx \, dt \\ &\leq \int_0^T \int_{\Omega^\delta} |\partial_t \rho_m| \, dx \, dt + C_{\delta,R,T} \end{split}$$

which gives (3.9).

We proceed similarly for the bound on $\partial_{x_i}\rho_m$. The only difference is that we do not have $\partial_{x_i}\rho_m|_{\partial K} = 0$, so we have an additional boundary term to worry about. More precisely, differentiating the first equation in (2.1) with respect to x_i , multiplying it by $\operatorname{sign}(\partial_{x_i}\rho_m)$ and using Kato's inequality, we obtain

(3.12)
$$\partial_t |\partial_{x_i} \rho_m| - \Delta(m\rho_m^{m-1} |\partial_{x_i} \rho_m|) \le \lambda |\partial_{x_i} \rho_m| + |\partial_{x_i} \lambda| \rho_m \quad \text{in } \Omega.$$

With the same cut-off function φ as above, we get

$$\begin{split} \frac{d}{dt} \int_{\Omega} |\partial_{x_i} \rho_m| \varphi \, dx &\leq \int_{\Omega} m\rho_m^{m-1} |\partial_{x_i} \rho_m| \Delta \varphi \, dx - \int_{\partial K} m\rho_m^{m-1} |\partial_{x_i} \rho_m| \nabla \varphi \cdot \nu \, dS \\ &+ \int_{\Omega} \lambda |\partial_{x_i} \rho_m| \varphi \, dx + \int_{\Omega} |\rho_m| |\partial_{x_i} \lambda| \, dx \\ &\leq \int_{\partial K} \rho_m |\partial_{x_i} p_m| |\nabla \varphi \cdot \nu| \, dS + \Lambda \int_{\Omega} |\partial_{x_i} \rho_m| \varphi \, dx + \int_{\Omega} |\rho_m| |\partial_{x_i} \lambda| \, dx \end{split}$$

To conclude, we thus note that the estimate (3.2) gives a bound on the normal derivative of p_m on ∂K , while the condition $p_m|_{\partial K} = \frac{m}{m-1}f$ together with the regularity assumptions (2.4) implies that

the tangential derivatives of p_m are uniformly bounded on ∂K . We deduce that $|\partial_{x_i} p_m||_{\partial K} \leq C$ and so (using (2.8), (3.4) and (2.3)):

$$\frac{d}{dt} \int_{\Omega} |\partial_{x_i} \rho_m| \varphi \, dx \le C(T) \qquad \forall t \in (0,T)$$

Hence

$$\|\partial_{x_i}\rho_m(t)\varphi\|_{L^1(\Omega)} \le \|\partial_{x_i}\rho_m(0)\|_{L^1(\Omega)} + C(T).$$

and (3.8) now follows from (2.8).

3.5. Passing to the limit. We denote

$$\Omega_k := \{ x \in \Omega \, ; \, \operatorname{dist}(x, K) > 1/k, \ |x| \le k \}.$$

Lemma 3.6 implies that ρ_m and p_m are bounded in BV($\mathbb{R}_+ \times \Omega_k$) for all k and thus converge (up to a subsequence) strongly in $L^1([0,k] \times \Omega_k)$. By a diagonal extraction process, we can thus find subsequences (still denoted ρ_m and p_m) and functions ρ_{∞} , p_{∞} such that ρ_m (resp. p_m) converges to ρ_{∞} (resp. p_{∞}) strongly in $L^1_{loc}(\mathbb{R}_+ \times \Omega)$.

Next, we note that

$$\rho_m p_m = \left(\frac{m-1}{m}\right)^{\frac{1}{m-1}} p_m^{\frac{m}{m-1}}$$

passing to the limit (using the a.e. convergence) yields $\rho_{\infty}p_{\infty} = p_{\infty}$ and thus

(3.13) $(\rho_{\infty} - 1)p_{\infty} = 0 \quad \text{a.e. } \mathbb{R}_{+} \times \Omega,$

which gives the Hele-Shaw condition $p_{\infty}(x,t) \in P_{\infty}(\rho_{\infty}(x,t))$ a.e. in $\mathbb{R}_{+} \times \Omega$.

Similarly, we have

$$\rho_m^m = \left(\frac{m-1}{m}p_m\right)^{\frac{m}{m-1}} \to p_\infty \quad \text{a.e. } \mathbb{R}_+ \times \Omega.$$

Since ρ_m^m is bounded in BV($\mathbb{R}_+ \times \Omega_k$), the convergence holds in L^1_{loc} as well. Rewriting (2.1) as

$$\partial_t \rho_m = \Delta p_m + \lambda \rho_m$$

and passing to the limit, we deduce

$$\partial_t \rho_\infty = \Delta p_\infty + \lambda \rho_\infty \qquad \text{in } \mathcal{D}'(\mathbb{R}_+ \times \Omega).$$

3.6. Bounds on the gradient of p_m and convergence of ρ_m .

Lemma 3.7. There exists a constant C independent of m such that

(3.14)
$$\int \int_{Q_T} |\nabla p_m|^2 \, dx \, dt \le CT$$

Furthermore, $\{\rho_m\}_{m\in\mathbb{N}}$ is relatively compact in $C^s(0,T; H^{-1}(\Omega))$ for all $s \in (0,1/2)$.

Proof. Integrating the equation for the pressure (1.4) yields

$$\frac{d}{dt} \int_{\Omega} p_m \, dx = -(m-2) \int_{\Omega} |\nabla p_m|^2 \, dx + (m-1) \int_{\partial K} p_m \nabla p_m \cdot \nu dS + (m-1) \int_{\Omega} \lambda p_m \, dx.$$

Using (3.2) and the fact that $p_m = \frac{m}{m-1}f$ on ∂K we deduce

$$\int_{\Omega} |\nabla p_m|^2 \, dx \le -\frac{1}{m-2} \frac{d}{dt} \int_{\Omega} p_m \, dx + \frac{m-1}{m-2} C$$

Integrating in time and using (3.6) we deduce (3.14). Using (2.1), we deduce that

 $\partial_t \rho_m$ is bounded in $L^2(0,T; H^{-1}(\Omega))$.

Since ρ_m is bounded in $L^{\infty}(0,T;L^1(\Omega))$ and in $L^{\infty}(0,T;L^{\infty}(\Omega))$, we also have

$$\rho_m$$
 is bounded in $L^{\infty}(0,T;L^2(\Omega))$.

Since $H^{-1}(\Omega)$ is compactly embedded in $L^2(\Omega)$, Lions-Aubin Lemma (see for example [Lio69, Ama00]) implies that $\{\rho_m\}$ is relatively compact in

$$C^{s}(0,T; H^{-1}(\Omega))$$
 for all $s \in (0, 1/2)$.

Estimate (3.14) implies in particular that $\nabla p_{\infty}(\cdot, t) \in L^2(\Omega)$ for a.e. t > 0: it will be useful in the proof of Theorem 2.7.

The compactness of $\{\rho_m\}_{m\in\mathbb{N}}$ in $C^s(0,T; H^{-1}(\Omega))$ implies $\rho_{\infty} \in C^s([0,\infty); H^{-1}(\Omega))$. Furthermore, (3.1) and (3.5) implies that $\rho_m(t)$ is bounded in $L^1(\Omega) \cap L^{\infty}(\Omega)$ and thus converges, up to a subsequence, weakly in $L^{\infty}(\Omega)$ to $\rho_{\infty} \in [0,1]$. We will see later in Section (5) that the limit density is unique (Proposition 5.1), which implies that the whole original subsequence converge to ρ_{∞} weakly in $L^{\infty}(\Omega)$.

4. Proof of Theorem 2.7 and Proposition 2.8, 2.9

In this section, we use the notation ρ_m and p_m even though we are only considering convergent subsequences. Let us first introduce a lemma to be used in the proof of Theorem 2.7.

Lemma 4.1. For all $t_0 \ge 0$, $p^+(\cdot, t_0) \in H^1(\Omega)$ and

$$\int_{\Omega} |\nabla p^+(x,t_0)|^2 \, dx \le \liminf_{\delta \to 0} \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 \, dx \, dt.$$

Proof. First, using (4.5) with a nonnegative test function $v \in H^1(\Omega)$ supported in U (see (3.3)) and satisfying the boundary condition on ∂K , we get:

$$\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 \, dx \, dt \le C,$$

for some constant C independent of δ and so the limit exists and is finite. Given $T(x) \in (D(\Omega))^n$, we can write

$$\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} p_{\infty} \operatorname{div} T \, dx \, dt = -\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} \nabla p_{\infty} \cdot T \, dx \, dt \le \left(\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 \, dx \, dt\right)^{1/2} \|T\|_{L^2(\Omega)},$$

and we can pass to the limit $\delta \to 0$, using (2.10), to get

$$\int_{\Omega} p^+(x,t_0) \operatorname{div} T(x) \, dx \le \left(\liminf_{\delta \to 0} \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 \, dx \, dt \right)^{1/2} \|T\|_{L^2(\Omega)},$$
It follows.

and the result follows.

Proof of Theorem 2.7. Given $t_0 \ge 0$ and a function v(x) in E_{t_0} , and using the equation for the pressure (1.4) and density (2.1) we can write, in $\mathcal{D}'(\mathbb{R}_+)$,

(4.1)

$$\int_{\Omega} \nabla p_m \cdot \nabla p_m - \rho_m \nabla p_m \cdot \nabla v - \lambda(p_m - v) \, dx$$

$$= -\frac{1}{m-1} \left[\frac{d}{dt} \int_{\Omega} p_m \, dx - \int_{\Omega} |\nabla p_m|^2 \, dx \right] + \frac{d}{dt} \int_{\Omega} v \rho_m \, dx - \int_{\Omega} \lambda(\cdot, t) v(\rho_m - 1) \, dx$$

$$+ \int_{\partial \Omega} [p_m - \rho_m v] \nabla p_m \cdot \nu \, dS.$$

Formally at least, it is not difficult to see that the variational formulation of the obstacle problem (2.12) follows by passing to the limit $m \to \infty$ and taking $t = t_0$ in (4.1). The rest of the proof is devoted to making this limit rigorous to derive (2.12) (for all $t_0 > 0$).

First, using the boundary condition, we note that the last term is equal to

$$\int_{\partial\Omega} \left[\frac{m}{m-1} f(x,t) - f(x,t)^{\frac{1}{m-1}} f(x,t_0) \right] \nabla p_m \cdot \nu \, dS$$

and thus satisfies (using (2.4)):

(4.2)
$$\lim_{m \to \infty} \sup_{m \to \infty} \left| \int_{\partial \Omega} [p_m - \rho_m v] \nabla p_m \cdot \nu \, dS \right| \le C \int_{\partial \Omega} |f(x, t) - f(x, t_0)| \, dS$$
$$\le C |t - t_0|.$$

Next, it is clear that our a priori estimates do not allow us to pass to the limit in (4.1) pointwise in time. So, given $\delta > 0$, we integrate (4.1) with respect to $t \in (t_0, t_0 + \delta)$ and pass to the limit $m \to \infty$. The left hand side of (4.1) satisfies

$$\liminf_{m \to \infty} \int_{t_0}^{t_0 + \delta} \int_{\Omega} \nabla p_m \cdot \nabla p_m - \rho_m \nabla p_m \cdot \nabla v - \lambda (p_m - v) \, dx \, dt$$
$$\geq \int_{t_0}^{t_0 + \delta} \int_{\Omega} |\nabla p_\infty|^2 - \nabla p_\infty \cdot \nabla v - \lambda (p_\infty - v) \, dx \, dt,$$

where we used in particular the fact that $\rho_m \nabla p_m$ is bounded in $L^2(Q_T)$ by (3.14) and thus converges weakly to ∇p_{∞} (since $\rho_m \nabla p_m = \nabla \rho_m^m$ converges to ∇p_{∞} in $\mathcal{D}'(Q_T)$). Using (3.14) and (3.6) to control the first term in the right hand side of (4.1) and (4.2) for the last term, we deduce

$$\int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 - \nabla p_{\infty} \cdot \nabla v - \lambda(p_{\infty} - v) \, dx \, dt$$
(4.3)

$$\leq \liminf_{m \to \infty} \int_{\Omega} v(x) [\rho_m(x, t_0 + \delta) - \rho_m(x, t_0)] \, dx + \int_{t_0}^{t_0 + \delta} \int_{\Omega} \lambda(\cdot, t) v(x) (1 - \rho_\infty(x, t)) \, dx \, dt + \mathcal{O}(\delta^2).$$

Formally, the first term in the right hand side is non-positive because $v\rho_{\infty}(\cdot, t_0 + \delta) \leq v$ while $v\rho_{\infty}(\cdot, t_0) = v$ (this is where we use the fact that $v \in E_{t_0}$). In order to make this rigorous, we first note that

$$\frac{d}{dt} \int_{\Omega} v(x)\rho_m(x,t) \, dx = -\int_{\Omega} \rho_m \nabla p_m \cdot \nabla v \, dx + \int_{\partial K} \rho_m v \nabla p_m \cdot \nu \, dS + \int_{\Omega} \lambda(\cdot,t)\rho_m v \, dx$$

and so (using the fact that $\rho_m^{m-1}|_{\partial K} = v|_{\partial K} = f$):

$$\left|\frac{d}{dt}\int_{\Omega}v(x)\rho_m(x,t)\,dx\right| \leq \int_{\Omega}|\nabla p_m(x,t)||\nabla v(x)|\,dx + \int_{\partial K}f^{\frac{1}{m-1}}(x,t)f(x,t_0)|\nabla p_m\cdot\nu|dS + \Lambda\int_{\Omega}\rho_m v(x)\,dx$$

The first term in the right hand side is bounded in $L^2(0,T)$ (using (3.14)), and the second term is bounded in $L^{\infty}(0,T)$ (using (3.2)). We deduce that the function $t \mapsto \int_{\Omega} v_m(x)\rho_m(x,t) dx$ is bounded in $H^1(0,T) \subset C^{1/2}[0,T]$ and thus converges (up to a subsequence) uniformly in [0,T]. Since $\int_{\Omega} v(x)\rho_m(x,t) dx$ converges to $\int_{\Omega} v(x)\rho_{\infty}(x,t) dx$ in $\mathcal{D}'(\mathbb{R}_+)$, we have

$$\int_{\Omega} v(\cdot)\rho_m(\cdot,t) \, dx \to \int_{\Omega} v(\cdot)\rho_\infty(\cdot,t) \, dx \text{ locally uniformly in } \mathbb{R}_+.$$

Consequently

$$\liminf_{m \to \infty} \int_{\Omega} v(x) [\rho_m(x, t_0 + \delta) - \rho_m(x, t_0)] dx = \int_{\Omega} v(x) [\rho_\infty(x, t_0 + \delta) - \rho_\infty(x, t_0)] dx \le 0,$$

where we used the fact that $v(x)\rho_{\infty}(x,t_0) = v(x)$ (since $v \in E_{t_0}$) and $\rho_{\infty} \leq 1$.

Going back to (4.3), we deduce (using the fact that $v(x)(1 - \rho_{\infty}(x, t)) \ge 0)$:

(4.4)

$$\int_{t_0}^{t_0+\delta} \int_{\Omega} |\nabla p_{\infty}|^2 - \nabla p_{\infty} \cdot \nabla v - \lambda(p_{\infty}-v) \, dx \, dt \le \Lambda \int_{t_0}^{t_0+\delta} \int_{\Omega} v(x)(1-\rho_{\infty}(x,t)) \, dx \, dt + \mathcal{O}(\delta^2),$$

To prove the result, it remains to divide by δ and pass to the limit $\delta \to 0$. We first use Young's inequality to rewrite (4.4) as:

$$\frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} \frac{1}{2} |\nabla p_{\infty}|^2 - \lambda p_{\infty} \, dx \, dt \\
\leq \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} \frac{1}{2} |\nabla v|^2 - \lambda(\cdot, t) v \, dx \, dt + \frac{\Lambda}{\delta} \int_{t_0}^{t_0+\delta} \int_{\Omega} v(x) (1 - \rho_{\infty}(x, t)) \, dx dt + \mathcal{O}(\delta)$$

$$4.5)$$

(4

$$\leq \int_{\Omega} \frac{1}{2} |\nabla v|^2 - \int_{\Omega} \left(\frac{1}{\delta} \int_{t_0}^{t_0 + \delta} \lambda(\cdot, t) dt \right) v \, dx + \frac{\Lambda}{\delta} \int_{t_0}^{t_0 + \delta} \langle 1 - \rho_{\infty}(x, t), v(x) \rangle_{H^{-1}, H^1} dt + \mathcal{O}(\delta).$$

Since $\rho_{\infty} \in C([0,\infty); H^{-1}(\Omega))$ and $v \in E_{t_0}$, we have

$$\lim_{\delta \to 0} \frac{1}{\delta} \int_{t_0}^{t_0 + \delta} \langle 1 - \rho_{\infty}(x, t), v(x) \rangle_{H^{-1}, H^1} dt = \langle 1 - \rho_{\infty}(x, t_0), v(x) \rangle_{H^{-1}, H^1} = 0 \quad \forall t_0 \ge 0$$

and we can use (2.10) (and a similar inequality for λ) to pass to the limit in the terms involving λ .

From Lemma 4.1, we deduce that

$$\int_{\Omega} \frac{1}{2} |\nabla p^{+}(x, t_{0})|^{2} - \lambda^{+}(x, t_{0})p^{+}(x, t_{0}) dx \leq \int_{\Omega} \frac{1}{2} |\nabla v(x)|^{2} - \lambda^{+}(x, t_{0})v(x) dx \qquad \forall t_{0} > 0,$$

which implies that $p^+(\cdot, t)$ is indeed a solution of (2.11) for every $t \ge 0$.

The derivation of (2.12) is classical (given $u \in E_t$ and $\varepsilon > 0$, take $v = p + \varepsilon(u - p)$ in (2.11) and pass to the limit $\varepsilon \to 0$). The uniqueness of p^* follows from (2.12): if p_1 and p_2 are two solutions, then by plugging in each other as test functions we obtain

$$\int_{\Omega} |\nabla(p_1 - p_2)|^2 dx = 0,$$

and thus $p_1 = p_2$.

Proof of Proposition 2.9. For any ball $B_r(x_0) \in \mathcal{O}(t)$, we have $\langle \varphi, (1-\rho_{\infty}(t)) \rangle = 0$ for any $\varphi \in$ $\mathcal{D}(B_r(x_0))$, and so p solves the classical obstacle problem in $B_r(x_0)$. The usual theory (see [Caf98]) implies that $p \in C^{1,1}(B_{r/2}(x_0))$ and satisfies $\Delta p = \lambda \chi_{\{p>0\}}$ in $B_{r/2}(x_0)$. The proposition follows.

5. Uniqueness of the limit solution and comparison principle

In this section we establish the uniqueness for the limit problem in a general bounded domain D of \mathbb{R}^n (with smooth boundary): Given a continuous function $g \ge 0$ defined on $\partial D \times [0,T]$ and $\bar{\rho} \geq 0$ a nonnegative function in D satisfying $0 \leq \bar{\rho} \leq 1$, we consider the problem

(5.1)
$$\begin{cases} \partial_t \rho = \Delta p + \lambda \rho, & \text{in } D \times (0, T], \quad p \in P_{\infty}(\rho) \text{ a.e. in } D \times (0, T]; \\ p = g & \text{on } \partial D \times [0, T]; \\ \rho(t = 0) = \bar{\rho} & \text{in } D. \end{cases}$$

A weak solution of (5.1) is a set of functions $(\rho, p) \in L^{\infty}(D \times (0, T]) \times L^{2}(0, T; H^{1}(D))$ satisfying (5.1) in the sense of distribution.

In particular, the condition $p \in P_{\infty}(\rho)$ implies that $0 \le \rho \le 1$ and $p(1-\rho) = 0$ a.e. in $D \times (0,T]$ and for any smooth, compactly supported test function $\psi : \Omega \to \mathbb{R}$ with $\psi(\cdot,T) = 0$ and $\psi = 0$ on $\partial D \times [0,T]$ we have

(5.2)
$$\int_{D\times[0,T]} (\rho\psi_t + p\Delta\psi + \lambda\rho\psi) dx dt = -\int_D \rho(\cdot,0)\psi(\cdot,0) dx + \int_0^T \int_{\partial D} g\partial_\nu \psi dS dt.$$

We then have the following result, which implies in particular Proposition 2.5:

Proposition 5.1. Suppose $\lambda \in L^2([0,T]; H^1(D))$, then there is at most one weak solution (ρ, p) of (5.1).

Furthermore, if (ρ_i, p_i) for i = 1, 2 are two pairs of weak solutions of (5.1) with boundary data g_i and initial data $\bar{\rho}_i$ and if $\bar{\rho}_1 \leq \bar{\rho}_2$ in D and $g_1 \leq g_2$ on $\partial D \times [0, T]$, then $\rho_1 \leq \rho_2$ a.e. in $D \times [0, T]$.

Proof. To show the uniqueness we follow the Hilbert dual argument developed in [PQV14]. Since the proof is largely parallel, we will only remark on necessary modifications due to the presence of the fixed boundary ∂K .

Suppose ψ is a nonnegative test function. Let us denote $D_T := D \times (0, T)$. Taking the differences of the weak formulation (5.2) for (ρ_i, p_i) for i = 1, 2, we have

$$\begin{split} \int \int_{D_T} [(\rho_1 - \rho_2)\partial_t \psi + (p_1 - p_2)\Delta\psi + \lambda(\rho_1 - \rho_2)] \\ &= -\int_D (\bar{\rho}_1 - \bar{\rho}_2)(x)\psi(x, 0)dx + \int_0^T \int_{\partial D} (g_1 - g_2)\partial_\nu \psi dSdt \\ &\geq \int_0^T \int_{\partial D} (g_1 - g_2)\partial_\nu \psi dSdt. \end{split}$$

Thus

(5.3)
$$\int \int_{D_T} (\rho_1 - \rho_2 + p_1 - p_2) [A\partial_t \psi + B\Delta \psi + \lambda A\psi] dx dt \ge \int_0^T \int_{\partial D} (g_1 - g_2) \partial_\nu \psi dS dt.$$

where ν denotes the outward normal at ∂D and

$$A = \frac{\rho_1 - \rho_2}{\rho_1 - \rho_2 + p_1 - p_2}, \qquad B = \frac{p_1 - p_2}{\rho_1 - \rho_2 + p_1 - p_2}$$

As in [PQV14] we define A = 0 whenever $\rho_1 = \rho_2$ (even when $p_1 = p_2$) and B = 0 when $p_1 = p_2$ (even when $\rho_1 = \rho_2$). Note that $A, B \in [0, 1]$ due to the fact that $\rho(1 - p) = 0$.

Let now G be a compactly supported and nonnegative smooth function in $D \times [0, T]$. As in [PQV14] the idea is to solve the dual problem

(5.4)
$$\begin{cases} A\partial_t \psi + B\Delta\psi + \lambda\psi = -AG & \text{in} \quad D \times [0,T);\\ \psi = 0 & \text{on} \quad \partial D \times [0,T];\\ \psi(\cdot,T) = 0. & \text{in} \quad D. \end{cases}$$

If A and B were strictly positive, by backward-in-time maximum principle, one can verify that ψ is nonnegative. Thus it follows that $\partial_{\nu}\psi \leq 0$ on $\partial D \times [0,T]$. Thus going back to (5.3) and using the fact that $g_1 \leq g_2$, it follows that

(5.5)
$$\int \int_{D_T} (\rho_1 - \rho_2) (-AG) \ge 0$$

Since G is arbitrary nonnegative smooth function, we conclude that $\rho_1 \leq \rho_2$ a.e. in $D \times [0, T]$.

However, A and B can be degenerate, so the argument requires the approximation of the dual problem (5.4), by a regularized uniformly parabolic, Dirichlet boundary value problem (see [PQV14] for detailed description of this approximation). As in [PQV14], we then pass to the limit in the regularization to deduce (5.5). The assumption $\lambda \in L^2([0,T]; H^1(D))$ is necessary to ensure that the regularized problem produces small errors.

To show uniqueness, suppose that (ρ_i, p_i) are two solutions of (5.1) with the same boundary condition f and initial condition g. Then $\rho_1 = \rho_2$ follows from the density ordering property obtained above. Once we have this, the difference of the weak equations yield

$$\int \int_{D_t} (p_1 - p_2) \Delta \psi dx dt = 0.$$

Now as in [PQV14] we can choose ψ to approximate $p_1 - p_2$ to conclude that $p_1 = p_2$ a.e. in D_T .

Remark 5.2. It is not immediately clear that the pressure satisfy the ordering property (i.e. $p_1 \leq p_2$ in Proposition 5.1). However, the characterization of the pressure given in Proposition 2.9 implies that the pressure ordering follows from the density ordering.

Now let us state two consequences of this proposition, based on the comparison principle for (2.1). First let us discuss our original problem with $\Omega := \mathbb{R}^n \setminus K$. Recall that from Lemma 3.3 that the support of ρ_m lies in $B_{\overline{R}+C(T)}$ for given time range $0 \le t \le T$. Therefore, setting $R(T) := \overline{R} + C(T)$, their limit solution $(\rho_{\infty}, p_{\infty})$ is a weak solution of (5.1) with $D := B_{R(T)} \setminus K$, g = f on ∂K and g = 0 on $\partial B_{R(T)}$. Therefore we have the following corollary:

Corollary 5.3. Given T > 0, any weak solution of (5.1) with $D := B_{R(T)} \setminus K$, g = f on ∂K , g = 0 on $\partial B_{R(T)}$ and initial data $\bar{\rho} = \rho^0$ is the $L^1(Q_T)$ - limit of the functions (ρ_m, p_m) solutions of (2.1). In particular, it follows that the pressure ordering property is true in this setting.

The next observation will be useful, when we construct radial limit solutions with explicit free boundary motion laws.

Corollary 5.4 (Comparison Principle). Let $(\rho_{\infty}, p_{\infty})$ be the limit solution of (2.1) in $\Omega \times [0, T]$. If D is a domain with smooth boundary that does not intersect K and if (ρ_1, p_1) is a weak solution of (5.1) in $D \times [t_1, t_2]$, then the following holds: If $p_{\infty} \leq p_1$ on $\partial D \times [t_1, t_2]$ and $\rho_{\infty} \leq \rho_1$ on $t = t_1$, then $p_{\infty} \leq p_1$ and $\rho_{\infty} \leq \rho_1$ in $D \times [t_1, t_2]$.

Proof. Since D does not intersect K, it is easy to check that $(\rho_{\infty}, p_{\infty})$ is a weak solution of (5.1) in $D \times [t_1, t_2]$ with initial data $\rho_{\infty}(\cdot, t_1)$ and fixed boundary data given as the trace of p_{∞} on $\partial D \times [t_1, t_2]$ (such trace exists a.e. in time since $p_{\infty}(\cdot, t)$ is in $H^1(D)$ a.e. t > 0). Now we can conclude from Proposition 5.1.

6. Proof of Proposition 2.10

In the sequel, we write p(t) instead of $p_{\infty}(t)$ for the unique solution of the obstacle problem (2.11). We also recall that $\mathcal{P}(t) = \{p(t) > 0\}$. We first show that $\sup \mu_t \subset \partial \mathcal{P}(t) \setminus \mathcal{O}(t)$: For all smooth test functions $\varphi \in \mathcal{D}(\Omega)$, by definition of μ_t we have

$$\mu_t(\varphi) = \int_{\Omega} (-\nabla p \cdot \nabla \varphi + \lambda(\cdot, t) \chi_{\mathcal{P}(t)} \varphi) \, dx$$

Clearly, if φ is supported in $\{p(\cdot, t) = 0\}$, the fact that $p \in H^1(\Omega)$ implies that $\nabla p = 0$ a.e. in $\{p = 0\}$ and thus $\mu_t(\varphi) = 0$. And if φ is supported in $\mathcal{O}(t)$, (2.14) implies

$$u_t(\varphi) = 0.$$

Since $\mathcal{O}(t)$ is an open set, we deduce that

$$\operatorname{supp}(\mu_t) \cap \operatorname{Int}(\{p(t) = 0\}) = \emptyset, \qquad \operatorname{supp}(\mu_t) \cap \mathcal{O}(t) = \emptyset.$$

On the other hand note that $\operatorname{Int}(\mathcal{P}(t)) \subset \mathcal{O}(t)$. Indeed if p(t) > 0 in $B_{\delta}(x_0)$, then $1 - \rho_{\infty}(t) = 0$ a.e. in $B_{\delta}(x_0)$ and so $\int_{B_{\delta}(x_0)} (1 - \rho_{\infty}(t)) dx = 0$. It follows that $x_0 \in \mathcal{O}(t)$. Thus we can conclude that μ_t is supported in $\partial \mathcal{P}(t) \setminus \mathcal{O}(t)$ as claimed in Proposition 2.10.

Next we show that μ_t is nonnegative. Define the function

$$Q_{\delta}(s) := \begin{cases} \frac{s}{\delta} & \text{if } s \in [0, \delta] \\ 1 & \text{if } s \ge \delta. \end{cases}$$

For any test function $\varphi \in \mathcal{D}(\Omega)$ satisfying $0 \leq \varphi(x) \leq 1$, we write

$$\mu_t(\varphi) = \int_{\Omega} -\nabla p \cdot \nabla \varphi + \lambda \chi_{\mathcal{P}(t)} \varphi \, dx$$

=
$$\int_{\Omega} -\nabla p \cdot \nabla (\varphi Q_{\delta}(p)) + \lambda \varphi Q_{\delta}(p) \, dx + \langle \Delta p, \varphi(1 - Q_{\delta}(p)) \rangle + \int_{\Omega} \lambda \chi_{\mathcal{P}(t)} \varphi(1 - Q_{\delta}(p)) \, dx.$$

Using (2.12) with $u = p - \delta \varphi Q_{\delta}(p)$ (which satisfies $p \ge u \ge p(1 - \varphi) \ge 0$ and is thus admissible) the first integral is non-negative. Next note that

$$\langle \Delta p, \varphi(1 - Q_{\delta}(p)) \rangle = \int (\nabla p \cdot \nabla \varphi(Q_{\delta}(p) - 1) + \nabla p \cdot \varphi Q_{\delta}'(p) \nabla p) dx$$

The second term in above equality is nonnegative since Q_{δ} is increasing. For the first term, we note that $\nabla \varphi(Q_{\delta}(p)-1)$ converges a.e. to $\nabla \varphi \chi_{\{p=0\}}$. Lebesgue dominated convergence theorem implies that it converges in L^2 and thus the first term converges to zero since $\nabla p = 0$ a.e. in $\{p=0\}$.

Thus

$$u_t(\varphi) \ge \int_{\Omega} \lambda \chi_{\mathcal{P}(t)} \varphi(1 - Q_{\delta}(p)) \, dx.$$

Finally, we have $\chi_{\mathcal{P}(t)}(1 - Q_{\delta}(p)) \to 0$ a.e. in Ω when $\delta \to 0$. Sending $\delta \to 0$ and using Lebesgue dominated convergence theorem, we can conclude that $\mu_t(\varphi) \ge 0$ and the result follows.

7. The velocity law

In this section we determine the velocity law of the congested zone $\{\rho_{\infty} = 1\}$ for the limit solution $(\rho_{\infty}, p_{\infty})$ by using comparison principle and barriers, as in the usual viscosity solutions approach. First we will define the relevant notion of barriers and prove that the usual comparison with barriers holds for our limit solution $(\rho_{\infty}, p_{\infty})$ (see Corollary 7.3-7.4). In Section 7.2 we show that in the radial symmetric case, the barriers we construct are indeed classical solutions.

7.1. Comparison with barriers. The difficulty in making (2.16) rigorous is the lack of regularity of the pressure or density interface ($\partial \mathcal{P}$ or $\partial \Sigma$) and the lack of monotonicity of its motion. In this section, we construct sub- and super-solutions of the limiting problem (2.1) to be used as barrier in a viscosity solution type approach.

Let B_r be a ball in Ω , and let D be either $\Omega \setminus \overline{B}_r$ or B_r . For a given time interval $[t_1, t_2] \subset [0, \infty)$ we consider a function (the pressure) $\phi \in C_c(\overline{D} \times [t_1, t_2])$ such that $\{\phi(t) > 0\}$ is monotone (increasing or decreasing) and an initial density $\rho_1(x)$ satisfying $\rho_1 = 1$ in $\{\phi(t_1) > 0\}$. We assume that $\{\phi(t) > 0\}$ and $\rho_1(x)$ are such that the external density ρ_{ϕ}^E , defined below, satisfies

(7.1)
$$\rho_{\phi}^{E}(x,t) < 1 \text{ in } \{\phi = 0\}$$

This external density $\rho_{\phi}^{E}(x,t)$ solves the equation $\partial_{t}\rho = \lambda\rho$ in the (decongestion) set $\{\phi = 0\}$ together with appropriate boundary conditions. This leads to the following definitions:

If $\{\phi(t) > 0\}$ is **increasing** ("expanding solution"), then for all $x \notin \{\phi(t_1) > 0\}$, we define t(x) = the last time that $\phi(x, t) = 0$ (with $t(x) = t_2$ is $\phi(x, t_2) = 0$) and set

$$\rho_{\phi}^{E}(x,t) = \rho_{1}(x) \exp\left(\int_{t_{1}}^{t} \lambda(x,s) \, ds\right) \qquad \text{for all } t < t(x)$$

(condition (7.1) is satisfied if $\rho_1(x)$ is small enough in $\{\phi(t_1) = 0\}$).

If $\{\phi(t) > 0\}$ is **decreasing** ("contracting solution"), then for all $x \notin \{\phi(t_2) > 0\}$, we define t(x) = the first time that $\phi(x,t) = 0$ (with $t(x) = t_1$ is $\phi(x,t_1) = 0$) and set

$$\rho_{\phi}^{E}(x,t) = \rho_{1}(x) \exp\left(\int_{t(x)}^{t} \lambda(x,s) \, ds\right) \quad \text{for all } t > t(x)$$

(condition (7.1) requires $\rho_1(x)$ to be small enough in $\{\phi(t_1) = 0\}$, but since $\rho_1 = 1$ in $\{\phi(t_1) > 0\}$, it also requires $\exp\left(\int_{t(x)}^t \lambda(x,s) \, ds\right) < 1$ for $x \in \{\phi(t_1) > 0\}$).

In both cases, we define the density in $D \times (t_1, t_2)$ by

(7.2)
$$\rho_{\phi}(x,t) := \chi_{\{\phi(t)>0\}}(x) + \rho_{\phi}^{E}(x,t)(1-\chi_{\{\phi(t)>0\}}(x)) = \begin{cases} 1 & \text{in } \{\phi>0\}\\ \rho_{\phi}^{E}(x,t) & \text{in } \{\phi=0\} \end{cases}$$

We then have:

Proposition 7.1. With the notation above, assume that (ρ_{ϕ}, ϕ) are such that (a) $\phi \in C^1(\overline{\{\phi > 0\}}) \cap C^2_{loc}(\{\phi > 0\})$ and $\Gamma := \partial \{\phi > 0\}$ is C^2 in space and C^1 in time. (b) ϕ satisfies

(7.3)
$$\begin{cases} -\Delta \phi \leq \lambda & \text{in } \{\phi > 0\};\\ (1 - \rho_{\phi}^E) V_{\phi} \leq |\nabla \phi| & \text{on } \partial \{\phi > 0\}, \end{cases}$$

where V_{ϕ} denotes the normal velocity of the interface $\partial \{\phi > 0\}$. Then (ρ_{ϕ}, ϕ) is a weak subsolution of the limiting problem (5.1) in $D \times [t_1, t_2]$, namely

$$\partial_t \rho_\phi \leq \Delta \phi + \lambda \rho_\phi \text{ in } D \times (t_1, t_2), \qquad \phi \in P_\infty(\rho_\phi) \text{ a.e. in } D \times (t_1, t_2)$$

where the first equation holds in the sense that for every smooth, compactly supported test function $\psi: D \times (t_1, t_2) \to \mathbb{R}$ with $\psi(\cdot, t_2) = 0$ and $\psi(\cdot, t) = 0$ on $\partial D \times [t_1, t_2]$ we have

(7.4)
$$\int_{D\times[t_1,t_2]} (\rho_{\phi}\psi_t + \phi\Delta\psi + \lambda\rho_{\phi}\psi)dxdt \ge -\int_D \rho_1(x)\psi(\cdot,t_1)dx + \int_{t_1}^{t_2} \int_{\partial B} \phi\partial_{\nu}\psi dSdt.$$

Similarly, we have

Proposition 7.2. With the notation above, assume that (ρ_{ϕ}, ϕ) are such that (a) $\{\phi(\cdot, t) > 0\} \in \Omega$ for all $t, \phi \in C^1(\overline{\{\phi > 0\}}) \cap C^2_{loc}(\{\phi > 0\})$ and the interface $\Gamma := \partial \{\phi > 0\}$ is C^2 in space and C^1 in time.

(b) ϕ satisfies

(7.5)
$$\begin{cases} -\Delta \phi \ge \lambda & \text{in } \{\phi > 0\};\\ (1 - \rho_{\phi}^E) V_{\phi} \ge |\nabla \phi| & \text{on } \partial \{\phi > 0\}. \end{cases}$$

Then (ρ_{ϕ}, ϕ) is a supersolution of the limiting problem (5.1) in $D \times [t_1, t_2]$, namely

$$\partial_t \rho_\phi \ge \Delta \phi + \lambda \rho_\phi \text{ in } D \times (t_1, t_2), \qquad \phi \in P_\infty(\rho_\phi) \text{ a.e. in } D \times (t_1, t_2).$$

(with the corresponding weak formulation as in (7.4))

Note that for the contracting barrier, we have $V_{\phi} \leq 0$ and $\rho_{\phi}^{E} = 1$ on $\partial \{\phi(t) > 0\}$ and so the free boundary condition reduces to $|\nabla \phi| \ge 0$ for subsolution and $|\nabla \phi| = 0$ for supersolution.

Proof of Proposition 7.1. We denote $S(t) := \{\phi(\cdot, t) > 0\} = \{\rho(\cdot, t) = 1\}$ and $\Gamma(t) = \partial S(t) \cap D$. We also denote ν as the outward normal of the boundary of either $\Gamma(t)$ or ∂D with respect to the domain S(t). With these notations, we have

$$\int_{D} \phi \Delta \psi \, dx = \int_{S(t)} \phi \Delta \psi \, dx \ge -\int_{S(t)} \lambda \psi \, dx - \int_{\partial S(t)} \psi \nabla \phi \cdot \nu \, dS + \int_{\partial B} \phi \partial_{\nu} \psi dS$$
$$\ge -\int_{S(t)} \lambda \psi \, dx + \int_{\Gamma(t)} \psi |\nabla \phi| \, dS + \int_{\partial B} \phi \partial_{\nu} \psi dS,$$

where we used the fact that $\phi = 0$ and $\nabla \phi = |\nabla \phi| \nu$ on $\Gamma(t)$.

$$\begin{split} \int_{D} \phi \psi_{t} \, dx &= \int_{S(t)} \psi_{t} \, dx + \int_{D \setminus S(t)} \rho_{\phi}^{E} \psi_{t} \, dx \\ &= \frac{d}{dt} \int_{S(t)} \psi \, dx - \int_{\Gamma(t)} V_{\phi} \psi \, dS + \int_{D \setminus S(t)} (\rho_{\phi}^{E} \psi)_{t} \, dx - \int_{D \setminus S(t)} (\rho_{\phi}^{E})_{t} \psi \, dx \\ &= \frac{d}{dt} \int_{D} \rho_{\phi} \psi \, dx - \int_{\Gamma(t)} V_{\phi} (1 - \rho^{E}) \psi \, dS - \int_{D \setminus S(t)} (\rho_{\phi}^{E})_{t} \psi \, dx \\ &\geq \frac{d}{dt} \int_{D} \rho_{\phi} \psi \, dx - \int_{\Gamma(t)} |\nabla \phi| \psi \, dS - \int_{D \setminus S(t)} (\rho^{E})_{t} \psi \, dx \end{split}$$

Using the fact that

(7.6)
$$(\rho^E)_t = \lambda \rho^E \text{ in } \{\phi = 0\}$$

and the definition of ρ_{ϕ} , we deduce

$$\int_{D} (\rho_{\phi}\psi_{t} + \phi\Delta\psi)dx \ge -\int_{D} \lambda\rho_{\phi}\psi dx + \frac{d}{dt}\int_{D} \rho\psi dx + \int_{\partial B} \partial_{\nu}\psi dS,$$

where by integrating with respect to $t \in (t_{1}, t_{2}).$

and we conclude by integrating with respect to $t \in (t_1, t_2)$.

The proof of Proposition 7.2 is parallel. Note that it is not necessary to work with barriers such that the set $\{\phi(\cdot,t) > 0\}$ is monotone: we chose to do so because the definition of ρ_{ϕ}^{E} is more manageable in that case.

Combining Proposition 7.1 with the comparison principle for weak solutions of the limiting problem (Corollary 5.4) we get:

Corollary 7.3. Let (ρ_{ϕ}, ϕ) be as in Proposition 7.1 (sub-solution). If (i) $\rho_1 \le \rho(\cdot, t_1)$ in *D*, (so in particular $\{\phi(\cdot, t_1) > 0\} \subset \{\rho_{\infty}(\cdot, t_1) = 1\}$); (ii) $\phi \leq p_{\infty} \text{ on } \partial B_r \times [t_1, t_2],$ then $\rho_{\phi} \leq \rho_{\infty}$ in $D \times [t_1, t_2]$. In particular

$$\{\phi(\cdot, t) > 0\} \subset \{\rho_{\infty}(\cdot, t) = 1\}$$
 for all $t \in [t_1, t_2]$.

Formally, this corollary says that a classical subsolution of the viscosity law (satisfying (7.3)) cannot touch ρ_{∞} from below. In other words, ρ_{∞} satisfies the motion law

 $(1 - \rho_{\infty}^{E})V_{\infty} \geq |\nabla p_{\infty}|$ in a viscosity sense.

Similarly, Proposition 7.2 implies:

Corollary 7.4. Let (ρ_{ϕ}, ϕ) be as in Proposition 7.2 (super-solution). If

(i) $\rho_1 \ge \rho(\cdot, t_1)$ in D (so in particular $\{\rho_{\infty}(\cdot, t_1) = 1\} \subset \{\phi(\cdot, t_1) > 0\}$) (ii) $\phi \ge p_{\infty}$ on $\partial B_r \times [t_1, t_2]$ Then $\rho_{\phi} \ge \rho_{\infty}$ in $D \times [t_1, t_2]$. In particular

 $\{\phi(\cdot, t) > 0\} \supset \{\rho_{\infty}(\cdot, t) = 1\}$ for all $t \in [t_1, t_2]$.

As above, this result should be interpreted as saying that ρ_{∞} satisfies

 $(1 - \rho_{\infty}^{E})V_{\infty} \leq |\nabla p_{\infty}|$ in a viscosity sense.

Typically, for free boundary problems such "barrier property" allows us to introduce a notion of viscosity solutions which describes the pointwise behavior of the interface via comparison with barriers (see e.g.[KP18]). It is thus natural to ask whether our weak solutions coincide with viscosity solutions. While we suspect that viscosity solutions theory can be established for our problem, answering this question would require a different set-up of function spaces, and we do not pursue this question here to keep our investigation focused.

7.2. The radial symmetric case. In this section we show that the free boundary velocity law holds in the classical sense in the radial setting as long as $\partial_t \lambda$ does not change signs too often. To simplify our discussion we further assume that λ is non-positive, since construction of radial barriers for positive λ has been carried out in [KP18].

We thus assume that $K = B_1$ and that the boundary data is constant (we can take f = 1 without loss of generality) and for simplicity we take $\lambda = \lambda(t) \leq 0$ independent of x monotone C^1 function of t. The analysis could be extended to radial symmetric functions $\lambda(|x|, t) \leq 0$ such that $\partial_t \lambda$ changes sign a finite number of time in the interval [0, T].

In this setting, we construct compactly supported, radial sub and super solutions of (2.9) in $Q_T := \{|x| \ge 1\} \times [0, T].$

For a given R > 1, let us define $\phi_R(\cdot, t)$ as a solution of the Dirichlet boundary problem in $1 \le |x| \le R$:

(7.7)
$$-\Delta \phi = \lambda(t) \text{ in } |x| < R, \quad \phi = 0 \text{ on } |x| = R, \quad \text{and } \phi = 1 \text{ on } |x| = 1.$$

Note that this function will take negative value if R is large (depending on λ).

For a given $R_0 > 1$, we assume that the initial density ρ_0 equals 1 on $1 \le |x| < R_0$ and is strictly less than 1 and Lipschitz in $|x| \ge R_0$. We assume that R_0 is small enough so that the initial pressure $\phi_{R_0}(\cdot, 0)$ is nonnegative. We then define the external density in the region $|x| \ge R$ by

$$\rho^{E}(|x|,t) := \rho_{0}(|x|) \exp\left(\int_{0}^{t} \lambda(s) ds\right) < 1 \text{ in } |x| \ge R_{0},$$

then $\rho^E(\cdot, t)$ is Lipschitz continuous. It is also straightforward to check that the function $\partial_r \phi_R(R)$ is Lipschitz continuous for $R_0 < R < \infty$. Thus we can solve the following ODE for $0 \le t \le T$:

(7.8)
$$R'(t) = F(R(t), t), \text{ where } F(R, t) := \frac{(\partial_r \phi_R)_-(R, t)}{1 - \rho^E(R, t)}, \qquad R(0) = R_0.$$

Note that $\partial_r \phi_R(\cdot, t) \ge 0$ if and only if the function $\phi_R(\cdot, t)$ has a negative minimum in $1 \le |x| \le R$. Indeed if $\partial_r \phi_R(R, t) < 0$ and $\phi_R(\cdot, t)$ takes negative minimum, from the radial symmetry of $\phi_R(\cdot, t)$ it follows that the function has a local positive maximum for some x such that 1 < |x| < R, which contradicts the subharmonicity of $\phi_R(\cdot, t)$.

So as long as $\phi_{R(t)}$ is a non-negative function, we have $\partial_r \phi_R(\cdot, t) < 0$ and R'(t) > 0 (which provides is an expanding solution of the limiting problem. and we can show that it happened when the function $t \mapsto \lambda(t)$ is decreasing.

Case 1: $t \to \lambda(t)$ is increasing: In this case, we can define

(7.9)
$$\phi(\cdot, t) := \phi_{R(t)}(\cdot, t) \text{ for } 0 \le t \le T$$

and we claim that ϕ stays nonnegative for all times.

To show this, suppose that $\phi(\cdot, t)$ has a negative minimum at some time $t = t_0$. Then by continuity of R(t), the same is true for $\phi(\cdot, s)$ for s sufficiently close to t_0 . Hence from above discussion we have R'(t) = 0 in a small time interval $[t_0 - \epsilon, t_0]$. Suppose we choose ϵ such that $\phi(\cdot, t_0 - \epsilon)$ no longer has negative minimum. This must be true at least with $\epsilon = t_0$ due to our assumption. But since $\lambda(t_0 - \epsilon) < \lambda(t)$ and $R(t_0 - \epsilon) = R(t)$ for $s = t_0 - \epsilon$, we have $\phi(x, t_0 - \epsilon) \leq \phi(x, t_0)$, which is a contradiction to our choice of ϵ .

Hence we have shown our claim, and it follows from (7.8) and Propositions 7.1 -7.2 that ϕ is an expanding solution of (2.9) for all $t \ge 0$.

Case 2: $t \to \lambda(t)$ is non-increasing: In this case, $\phi(\cdot, t)$ might take negative value for some positive time. We thus define

$$t^* := \sup\{t \in [0, T] : \phi(\cdot, t) \ge 0 \text{ in } 1 \le |x| < R(t)\}.$$

If $t^* = \infty$ then we can define ϕ by (7.9) as above. We thus assume that $t^* < \infty$. The same arguments as above implies that $|D\phi| = 0$ at $(R(t^*), t^*)$. Since λ is non-increasing, it follows that $\phi_{R(t^*)}(\cdot, t)$ turns negative for $t > t^*$. For $t \ge t^*$ we define $\tilde{R}(t)$ as the unique boundary point of $\{\psi(\cdot, t) > 0\}$, where $\psi(\cdot, t)$ solves the obstacle problem

$$-\Delta \psi = \lambda(\cdot, t) \chi_{\{\psi > 0\}}$$
 in $1 < |x| < R(t^*)$, with $\psi = 1$ on $|x| = 1$.

We then define

(7.10)
$$\phi(\cdot, t) := \phi_{R(t)}(\cdot, t) \text{ for } 0 \le t \le t^*, \quad \phi(\cdot, t) := \phi_{\tilde{R}(t)}(\cdot, t) \text{ for } t^* \le t \le T.$$

Since λ is non-increasing, so is \hat{R} and $|D\phi|(\hat{R}(t),t) = 0$. It follows that ϕ is a contracting solution for $t^* \leq t \leq T$.

Below is the summary of our conclusion:

Lemma 7.5.

If $t \to \lambda(t)$ is increasing, then the function ϕ defined by (7.9) is an expanding solution for $0 \le t \le T$. If $t \to \lambda(t)$ is non-increasing, then the function ϕ defined in (7.10) is an expanding solution for $0 \le t \le t^*$ and is a contracting solution for $t^* \le t \le T$.

Due to the uniqueness of the limit problem we can now completely characterize the limiting profile of radial solutions for λ that are monotone C^1 function of time.

Proposition 7.6. Assume that $K = B_1$, f = 1 and that $t \mapsto \lambda(t)$ is a monotone C^1 function. Let ρ_0^m be a radially symmetric function satisfying the conditions of Assumption 2.2. Then the limit $(\rho_{\infty}, p_{\infty})$ given by Theorem 2.4 is radially symmetric and satisfies

$$\begin{cases} \Delta p_{\infty} + \lambda = 0 & \text{in } \{p_{\infty} > 0\};\\ (1 - \rho_{\infty}^{E})V \le |\nabla p_{\infty}| & \text{on } \partial\{p_{\infty} > 0\}. \end{cases}$$

Furthermore

- (a) If $t \mapsto \lambda(t)$ is increasing, then $\{p_{\infty} > 0\}$ is always expanding $(\rho_{\infty}^{E} < 1, |\nabla p_{\infty}| > 0 \text{ and } V > 0 \text{ on } \partial\{p_{\infty} > 0\})$
- (b) If $t \mapsto \lambda(t)$ is non-increasing, then there exists a time $t^* \in [0,T]$ such that $\{p_{\infty} > 0\}$ is expanding for $0 \le t \le t^*$ and contracting for $t^* \le t \le T$ $(\rho_{\infty}^E = 1, |\nabla p_{\infty}| = 0 \text{ and } V \le 0 \text{ on } \partial\{p_{\infty} > 0\})$

7.3. Continuous expansion of the congested zone. As an application of comparison principle (Corollary 5.4) with a radial barrier, we show that the congested zone does not expand discontinuously over time. Note that it may shrink discontinuously even if λ is smooth, for instance due to topological changes. Note also that if λ is nonnegative, the expansion may not be continuous due to the nucleation of congested zones created by the growth of external densities.

Corollary 7.7. If (ρ, p) is a limit solution in $\Omega \times [0, T]$ and $\lambda \in C(Q_T) \cap L^2([0, T]; H^1(\Omega))$ is negative, then

(7.11)
$$\overline{\{p>0\} \cap Q_T} = \overline{\{p>0\}} \cap Q_T = \overline{\{\rho=1\}} \cap Q_T \text{ for any } T>0.$$

Proof. We denote

$$S_1 := \overline{\{p > 0\} \cap Q_T}, \qquad S_2 := \overline{\{p > 0\}} \cap Q_T,$$

Since $S_1 \subset S_2$ by definition, we only need to show that $S_2 \subset S_1$ in order to prove the first equality.

Given $x_0 \notin S_1$ there exists r > 0 such that

$$B_{2r}(x_0) \times [T - r, T) \subset \{p = 0\}.$$

We claim that $B_{r/2}(x_0)$ lies in $\{p(\cdot, T) = 0\}$. This proves that $(x_0, T) \notin S_2$, hence $S_2 \subset S_1$.

To show that $B_r(x_0) \subset \{p(\cdot, T) = 0\}$, we use a barrier argument in $\Sigma := B_r(x_0) \times [T - \epsilon, T)$ for a sufficiently small $\epsilon > 0$ as follows. Due to Proposition 2.10 we have $\rho_t = \lambda \rho$ in $B_{2r}(x_0) \times [T - r, T)$, and thus

$$\rho < a(\lambda, r) < 1$$
 in $B_{2r}(x_0) \times [T - r/2, T)$.

Let us construct an expanding supersolution in Σ as follows. Let ϕ_0 solve

$$\begin{split} -\Delta\phi_0 &= \Lambda \text{ in } \{r < |x| < 2r\}, \quad \phi_0 = 0 \text{ on } \{|x| = r\}, \quad \text{ and } \phi_0 = M := \|p\|_{L^{\infty}(\Omega \times [0,T])} \text{ on } \{|x| = 2r\}, \\ \text{ and let } \phi(\cdot,t) &:= \phi_{R(t)} \text{ defined by } (7.7) \text{ with } \lambda = \Lambda \text{ where } R(t) \text{ solves} \end{split}$$

$$R'(t) = \frac{|D\phi|(R(t), t)|}{1 - a \exp^{(\Lambda t)}} \quad \text{for } 0 \le t \le \epsilon, \quad \text{with } R(0) = r.$$

Then ϕ is an expanding supersolution in Σ with fixed boundary data M on $\partial B_{2r}(x_0)$ and initial data $\rho_0 = \chi_{r < |x| < 2r} + a\chi_{|x| \le r}$. Corollary 5.4 now applies to show that $p(\cdot, T) \le \phi(\cdot, T)$. Choosing $\epsilon = \epsilon(M, a)$ sufficiently small so that $R(T) \le R(0) + \frac{r}{2}$, it follows that $\phi(\cdot, T) = 0$ in $B_{r/2}(x_0)$ and we can conclude.

It remains to show the second equality of the Corollary. Note that we have $\{p > 0\} \subset \{\rho = 1\}$, and thus their closures are also ordered. On the other hand we showed above that if x_0 lies outside of $\overline{\{p > 0\}}$ then ρ is strictly less than one in a small neighborhood of x_0 , and thus it is outside of $\overline{\{\rho = 1\}}$. The result follows.

8. MONOTONE INCREASING SOLUTIONS

In this section we suppose that $\lambda \in L^2([0,T]; H^1(\Omega))$ is non-decreasing in time. We first show that in this setting, if the density starts as a characteristic function, the pressure only increases over time.

Lemma 8.1. Let Σ_0 be a bounded subset of \mathbb{R}^n which contains K. Suppose that $\rho_0 = \chi_{\Sigma_0 \setminus K}$ and that $\Sigma_0 \setminus K$ coincides with the initial pressure support $\{p_0 > 0\}$, where p_0 solves (2.12) with $\rho_{\infty}(\cdot, t)$ replaced by ρ_0 . If (ρ, p) is the limit solution given by Theorem 2.4 with initial data ρ_0 , then ρ and p are monotone increasing with respect to t.

Proof. Let B_R contain the support of Σ_0 . We claim that (ρ_0, p_0) is a stationary subsolution of (5.1) with $D = B_R \setminus K$ and with boundary data f. To verify this claim, using the monotonicity of λ over time, it is enough to check that

(8.1)
$$\int_{D} -\nabla p_0 \nabla \psi + \lambda(\cdot, 0) \rho_0 \psi dx \ge 0$$

for any nonnegative test function $\psi \in C_0^{\infty}(D)$. Since $\rho_0 = \chi_{\{p_0>0\}}$, the question boils down to the nonnegativity of the measure $\mu_0 := \Delta p_0 + \lambda(\cdot, 0)\chi_{\{p_0>0\}}$. This follows the same proof of showing $\mu_t \geq 0$ in Proposition 2.10, see section 5.

With the claim and the comparison principle for (5.1) (Proposition 5.1), it follows that

(8.2)
$$\rho(x,0) \le \rho(x,\epsilon) \text{ for all } \epsilon > 0.$$

Note that, since λ is non-decreasing in time, $\rho(\cdot, t - \epsilon)$ is a subsolution of (5.1) for any $\epsilon > 0$. Thus by comparison principle and (8.2) it follows that

$$\rho(x, t - \epsilon) \le \rho(x, t)$$
 for any $t > \epsilon > 0$,

and we conclude that ρ increases for all times. p accordingly increases by its definition.

Corollary 8.2. Let (ρ, p) be the weak solution of (5.1) in $\Omega \times [0, \infty)$ with the fixed boundary data p = f > 0 and the initial data $\rho_0 \in BV$. Then $\Sigma(t) := \{p(\cdot, t) > 0\}$ increases in time, and is a set of finite perimeter for a.e. t > 0. Moreover for all $t \ge 0$

(8.3)
$$\rho(\cdot,t) = \chi_{\Sigma(t)} + \rho^E \chi_{\mathbb{R}^n \setminus \Sigma(t)}, \text{ where } \rho^E(x,t) := \rho_0 \exp^{\int_0^t \lambda(x,s) ds}.$$

Proof. We claim that the pressure support $\Sigma(t) := \{p(\cdot, t) > 0\}$ increases over time. For any $t_0 > 0$, Let us call ρ^* be the weak solution of (5.1) with the initial data $\chi_{\Sigma(t_0)}$, and with the same fixed boundary data f for the pressure. Then ρ^* increases in time due to Lemma 8.1. From the monotonicity of ρ^* and Proposition 5.1, we have

(8.4)
$$\chi_{\Sigma(t_0)} \le \rho^*(\cdot, t) \le \rho(\cdot, t_0 + t) \text{ for all } t > 0.$$

It follows that $\Sigma(t)$ increases over time. It follows from Proposition 1.5 that $\rho_t = \lambda \rho$ in $\overline{\Sigma(t)}^C \times [0, t]$ for any T > 0, and thus we can conclude (8.3). Lastly $\Sigma(t)$, is a set of finite perimeter for a.e. t > 0 since $\rho \in BV(\Omega)$ for a.e. t > 0 and ρ has jump discontinuity on the boundary of $\Sigma(t)$ due to (8.3).

APPENDIX A. TUMOR GROWTH MODEL WITH NUTRIENT

In [PQV14] (see also [DP21]), the following model for tumor growth is studied:

(A.1)
$$\begin{cases} \partial_t \rho_m - \operatorname{div}\left(\rho_m \nabla p_m\right) = \rho_m G(p_m, c_m) & x \in \mathbb{R}^n, \ t \ge 0\\ \partial_t c_m - \Delta c_m + \rho_m H(c_m) = (c_B - c_m) K(p_m)\\ c_m(x, t) \to c_B \text{ for } x \to \infty \end{cases}$$

where

$$p_m = \frac{m}{m-1}\rho_m^{m-1}.$$

In this system, the evolution of the cell population density $\rho_m \ge 0$ is coupled to the concentration of nutrients $c_m \ge 0$ by the cell division rate G(p, c). Importantly, this function satisfies

$$\partial_p G < -\beta < 0$$

(see [PQV14] for a complete list of the assumptions necessary to get a good existence and uniqueness framework as well as the appropriate estimates to pass to the limit).

It is proved in [PQV14] that $\rho_m(x,t)$, $p_m(x,t)$ and $c_m(x,t)$ converge strongly in $L^1(Q_T)$ (for all T > 0) to $\rho_{\infty}, p_{\infty}, c_{\infty}$ in $BV(Q_T)$ which solves the system

(A.2)
$$\begin{cases} \partial_t \rho_{\infty} - \operatorname{div} \left(\rho_{\infty} \nabla p_{\infty} \right) = \rho_{\infty} G(p_{\infty}, c_{\infty}) & x \in \mathbb{R}^n, \ t \ge 0\\ \partial_t c_{\infty} - \Delta c_{\infty} + \rho_{\infty} H(c_{\infty}) = (c_B - c_{\infty}) K(p_{\infty})\\ c_{\infty}(x, t) \to c_B \text{ for } x \to \infty \end{cases}$$

with the Hele-Shaw relation $p_{\infty} \in P_{\infty}(\rho_{\infty})$.

Remarkably, the solution of this system is unique, and one would like to interpret the system as a weak form of some geometric Hele-Shaw type free boundary problem. For this one needs to identify the pressure p_{∞} as solution of an elliptic equation in $\{\rho_{\infty} = 1\}$.

In [DP21], it is proved that p_{∞} solves the complementarity condition

$$p_{\infty}(\Delta p_{\infty} + G(p_{\infty}, c_{\infty})) = 0$$
 in $\mathcal{D}'(Q)$.

This condition says that p_{∞} solves an elliptic equation in $\{p_{\infty}\}$ and is proved by deriving additional estimates on p_m .

We will show below that the approach used in this paper can be used to characterize $p_{\infty}(\cdot,t)$ as the unique solution of an obstacle problem. First, we summarize the estimates proved in [PQV14]:

Lemma A.1. Under the assumptions listed in [PQV14], the following holds for all T > 0:

- $\rho_m(t)$ is uniformly compactly supported for $t \in [0,T]$;
- $|\nabla p_m|$ is bounded in $L^2(Q_T)$
- $0 \le p_m \le p_M, 0 \le \rho_m \le \left(\frac{m-1}{m}p_M\right)^{\frac{1}{m-1}}, 0 < c_m < c_B$ $\rho_m, p_m \text{ and } c_B c_m \text{ are bounded in } BV(Q_T)$
- ρ_m , p_m and $c_B c_m$ converge strongly in L^1 and almost everywhere to ρ_{∞} , p_{∞} and $c_B c_{\infty}$.

Furthermore, proceeding as in Lemma 3.7, it is not difficult to show that $\{\rho_m\}_{m\in\mathbb{N}}$ is relatively compact in $C^s(0,T; H^{-1}(\mathbb{R}^n))$ for all $s \in (0,1/2)$ and thus that $\rho_{\infty} \in C(0,T; H^{-1}(\mathbb{R}^n))$.

Finally, since p_{∞} and $c_B - c_{\infty}$ are in $BV(Q_T)$, we can define the trace $p^+(\cdot, t)$ and $c^+(\cdot, t)$ for all t > 0 as in (2.10). We can then prove the following result:

Proposition A.2. For all t > 0, let E_t denote the space

$$E_t = \{ v \in H^1(\mathbb{R}^n) \cap L^1(\mathbb{R}^n) ; \ v(x) \ge 0, \ \langle v, 1 - \rho_\infty(t) \rangle_{H^1, H^{-1}} = 0 \}.$$

Then for all t > 0, the function $x \mapsto p^+(x,t)$ is the unique solution of the minimization problem:

(A.3)
$$\begin{cases} p \in E_t \\ \int_{\mathbb{R}^n} \frac{1}{2} |\nabla p|^2 - \mathcal{G}(p, c^+) \, dx \le \int_{\mathbb{R}^n} \frac{1}{2} |\nabla v|^2 - \mathcal{G}(v, c^+) \, dx \qquad \forall v \in E_t \end{cases}$$

where \mathcal{G} is the (concave) function such that $\partial_p \mathcal{G}(p,c) = G(p,c)$ and $\mathcal{G}(0,c) = 0$. Furthermore p_{∞} satisfies the complementarity condition

(A.4)
$$p_{\infty}(\Delta p_{\infty} + G(p_{\infty}, c_{\infty})) = 0 \quad in \ \mathcal{D}'(\mathbb{R}^n \times (0, \infty)).$$

As mentioned in the introduction (see Proposition 2.8), if the complementarity condition (A.4) is known to hold, then one can derive the variational formulation (A.3) from the weak equation (A.2). In particular, this complementarity condition was derived for this particular model in [DP21] by using a generalized Aronson-Bénilan estimate and the $L^2(W^{1,4})$ estimate on the pressure (but our proof here does not require either of these estimates).

Proof. First we recall the equation for the pressure p_m :

(A.5)
$$\partial_t p_m = (m-1)p_m(\Delta p_m + G(p_m, c_m)) + |\nabla p_m|^2.$$

We then proceed as in the proof of Theorem 2.7: Given $t_0 > 0$ and a function v(x) in E_{t_0} , we use the equation for the pressure (A.5) and density (A.2) to write:

$$\begin{split} \int_{\mathbb{R}^n} \nabla p_m \cdot \nabla p_m - \rho_m \nabla p_m \cdot \nabla v - \mathcal{G}(p_m, c_m) + \mathcal{G}(v, c_m) \, dx \\ &= -\frac{1}{m-1} \left[\frac{d}{dt} \int_{\mathbb{R}^n} p_m \, dx - \int_{\Omega} |\nabla p_m|^2 \, dx \right] \\ &+ \frac{d}{dt} \int_{\mathbb{R}^n} v \rho_m \, dx \\ &+ \int_{\mathbb{R}^n} p_m G(p_m, c_m) - \rho_m v G(p_m, c_m) - \mathcal{G}(p_m, c_m) + \mathcal{G}(v, c_m) \, dx \end{split}$$

in $\mathcal{D}'(\mathbb{R}_+)$. Using the concavity of \mathcal{G} to write

$$\mathcal{G}(v, c_m) - \mathcal{G}(p_m, c_m) \le G(p_m, c_m)(v - p_m)$$

we deduce

$$\int_{\mathbb{R}^n} \nabla p_m \cdot \nabla p_m - \rho_m \nabla p_m \cdot \nabla v - \mathcal{G}(p_m, c_m) + \mathcal{G}(v, c_m) \, dx = -\frac{1}{m-1} \left[\frac{d}{dt} \int_{\mathbb{R}^n} p_m \, dx - \int_{\mathbb{R}^n} |\nabla p_m|^2 \, dx \right] \\ + \frac{d}{dt} \int_{\mathbb{R}^n} v \, \rho_m \, dx + \int_{\mathbb{R}^n} (1 - \rho_m) \, v \, G(p_m, c_m) \, dx$$

We can now proceed as in the proof of Theorem 2.7: Integrating this equality with respect to $t \in [t_0, t_0 + \delta)$ and using the weak L^2 convergence of ∇p_m and $\rho_m \nabla p_m$ to ∇p , we get

$$\begin{split} &\int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} |\nabla p_{\infty}|^2 - \nabla p_{\infty} \cdot \nabla v - \mathcal{G}(p_{\infty}, c_{\infty}) + \mathcal{G}(v, c_{\infty}) \, dx \, dt \\ &\leq \int_{\mathbb{R}^n} v(x) [\rho_{\infty}(x, t_0 + \delta) - \rho_{\infty}(x, t_0)] \, dx + \int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} (1 - \rho_{\infty}) v G(p_{\infty}, c_{\infty}) \, dx \, dt \\ &\leq \|G(p_{\infty}, c_{\infty})\|_{L^{\infty}} \int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} v(1 - \rho_{\infty}) \, dx \, dt \end{split}$$

(where we used the fact that $v(x)\rho_{\infty}(x,t_0) = v(x)$ and $v(x)\rho_{\infty}(x,t) \leq v(x)$ for all t)

Finally, dividing by δ and using Young's inequality, we rewrite the inequality as

$$\begin{split} \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} \frac{1}{2} |\nabla p_{\infty}|^2 - \mathcal{G}(p_{\infty}, c_{\infty}) \, dx \\ &\leq \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} \frac{1}{2} |\nabla v|^2 - \mathcal{G}(v, c_{\infty}) \, dx \, dt + \frac{C}{\delta} \int_{t_0}^{t_0+\delta} \int_{\mathbb{R}^n} v(1-\rho_{\infty}) \, dx \, dt \\ &\leq \int_{\mathbb{R}^n} \frac{1}{2} |\nabla v|^2 - \frac{1}{\delta} \int_{t_0}^{t_0+\delta} \mathcal{G}(v, c_{\infty}) \, dt \, dx + \frac{C}{\delta} \int_{t_0}^{t_0+\delta} \langle v, 1-\rho_{\infty} \rangle_{H^1, H^{-1}} \, dx \, dt. \end{split}$$

The continuity of $t \mapsto \langle v, 1 - \rho_{\infty} \rangle_{H^1, H^{-1}}$ and the fact that $v \in E_t$ implies that the last term converges to zero as $\delta \to 0$. We can now conclude as in the proof of Theorem 2.7.

Finally, given a test function $\varphi \in \mathcal{D}(\mathbb{R}^n \times (0,\infty))$, we take $v = p_\infty + \varepsilon(p_\infty \varphi) = p_\infty(1 + \varepsilon \varphi)$ in (A.3), with $|\varepsilon|$ small enough so that $1 + \varepsilon \varphi \ge 0$. Passing to the limit $\varepsilon \to 0^-$ and $\varepsilon \to 0^+$ yields

$$\int_{\mathbb{R}^n} \nabla p_\infty \cdot \nabla (p_\infty \varphi) - G(p_\infty, c_\infty) p_\infty \varphi \, dx = 0$$
²⁹

and (A.4) follows.

APPENDIX B. THE COMPLEMENTARITY CONDITION

Proof of Proposition 2.8. We note that $\partial_t \rho = \Delta p + \lambda \rho \in L^2(0,T; H^{-1}(\Omega))$. Given $u \in E_t$, we have $p - u \in L^2(0,T; H^1_0(\Omega))$ and so we can write (in $\mathcal{D}'(\mathbb{R}_+)$):

(B.1)
$$\langle \partial_t \rho, (p-u) \rangle_{H^{-1}, H^1_0} = \langle \Delta p + \lambda \rho, p-u \rangle_{H^{-1}, H^1_0} = -\int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda \rho (p-u) \, dx.$$

Next, proceeding as in the beginning of the proof of Lemma 8.1 (using the comparison principle for the limiting problem, Proposition 5.1), we can show that $\rho = 1$ in $U \times \mathbb{R}_+$ for some neighborhood U of K and that supp p is bounded in $\Omega \times [0, T]$. In particular, $\partial_t \rho$ vanishes in $U \times \mathbb{R}_+$. Taking a smooth function $\phi(x)$ which is equal to 1 in supp $p \setminus (U \times [0, T])$ and vanishes on ∂K , we can write

$$\begin{split} \langle \partial_t \rho, (p-u) \rangle_{H^{-1}, H^1_0} &= \langle \partial_t \rho, (p-u) \phi \rangle_{H^{-1}, H^1_0} \\ &= \langle \Delta p + \lambda \rho, p \phi \rangle_{H^{-1}, H^1_0} - \langle \partial_t \rho, u \phi \rangle_{H^{-1}, H^1_0} \\ &= \langle p(\Delta p + \lambda \rho), \phi \rangle_{\mathcal{D}', \mathcal{D}} - \langle \partial_t \rho, u \phi \rangle_{H^{-1}, H^1_0} \\ &= \langle p(\Delta p + \lambda \rho), \phi \rangle_{\mathcal{D}', \mathcal{D}} - \frac{d}{dt} \int_{\Omega} \rho u \phi \, dx \\ &= -\frac{d}{dt} \int_{\Omega} \rho u \phi \, dx \qquad \text{in } \mathcal{D}'(\mathbb{R}_+) \end{split}$$

where we used the fact that $\langle p(\Delta p + \lambda \rho), \phi \rangle_{\mathcal{D}', \mathcal{D}} = 0$ (this is the complementarity condition). Using (B.1), we deduce

$$\int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda \rho(p-u) \, dx = \frac{d}{dt} \int_{\Omega} \rho u \phi \, dx \qquad \text{in } \mathcal{D}'(\mathbb{R}_+).$$

Using the fact that $\rho(x,t)p(x,t) = p(x,t)$, we deduce:

$$\begin{split} \int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda (p-u) \, dx \, dt &= \int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda \rho (p-u) \, dx \, dt + \int_{\Omega} \lambda (1-\rho) u \, dx \, dt \\ &\leq \frac{d}{dt} \int_{\Omega} \rho u \phi \, dx + \Lambda \int_{\Omega} (1-\rho) u \, dx \, dt \end{split}$$

Integrating with respect to $t \in [t_0, t_o + \delta]$, we get

$$\begin{split} \int_{t_0}^{t_0+\delta} \int_{\Omega} \nabla p \cdot \nabla (p-u) - \lambda (p-u) \, dx \, dt &\leq \int_{\Omega} (\rho(t_0+\delta) - \rho(t_0)) u \phi \, dx + \Lambda \int_{t_0}^{t_0+\delta} \int_{\Omega} (1-\rho) u \, dx \, dt \\ &\leq \int_{\Omega} (\rho(t_0+\delta) - 1) u \phi \, dx + \Lambda \int_{t_0}^{t_0+\delta} \int_{\Omega} (1-\rho) u \, dx \, dt \\ &\leq \Lambda \int_{t_0}^{t_0+\delta} \int_{\Omega} (1-\rho) u \, dx \, dt \end{split}$$

and the result now follows by proceeding as in the proof of Theorem 2.7.

References

- [AKY14] Damon Alexander, Inwon Kim, and Yao Yao. Quasi-static evolution and congested crowd transport. Nonlinearity, 27(4):823–858, 2014.
- [Ama00] Herbert Amann. Compact embeddings of vector-valued Sobolev and Besov spaces. volume 35(55), pages 161–177. 2000. Dedicated to the memory of Branko Najman.
- [BC81] Philippe Bénilan and Michael G Crandall. The continuous dependence on φ of solutions of $u_t \Delta \varphi(u) = 0$. Indiana University Mathematics Journal, 30(2):161–177, 1981.
- [BGHP84] M Bertsch, ME Gurtin, D Hilhorst, and LA Peletier. On interacting populations that disperse to avoid crowding: The effect of a sedentary colony. *Journal of Mathematical Biology*, 19(1):1–12, 1984.
- [BH86] M Bertsch and D Hilhorst. A density dependent diffusion equation in population dynamics: stabilization to equilibrium. *SIAM Journal on Mathematical Analysis*, 17(4):863–883, 1986.
- [BKM09] Ivan Blank, Marianne Korten, and Charles Moore. The hele-shaw problem as a "mesa" limit of stefan problems: Existence, uniqueness, and regularity of the free boundary. Transactions of the American Mathematical Society, 361(3):1241–1268, 2009.
- [Caf98] L. A. Caffarelli. The obstacle problem revisited. J. Fourier Anal. Appl., 4(4-5):383-402, 1998.
- [CF87] Luis A Caffarelli and Avner Friedman. Asymptotic behavior of solutions of $u_t = \Delta u^m$ as $m \to \infty$. Indiana University mathematics journal, 36(4):711–728, 1987.
- [DP21] Noemi David and Benoît Perthame. Free boundary limit of a tumor growth model with nutrient. Journal de Mathématiques Pures et Appliquées, 155:62–82, 2021.
- [EHK086] CM Elliott, MA Herrero, JR King, and JR Ockendon. The mesa problem: Diffusion patterns for $u_t = \Delta u^m$ as $m \to \infty$. IMA journal of applied mathematics, 37(2):147–154, 1986.
- [Giu84] Enrico Giusti. Minimal surfaces and functions of bounded variation, volume 80 of Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984.
- [GQ03] Omar Gil and F Quirós. Boundary layer formation in the transition from the porous media equation to a hele-shaw flow. Annales de l'IHP Analyse non linéaire, 20(1):13–36, 2003.
- [Kim03] Inwon Kim. Uniqueness and existence results on the hele-shaw and the stefan problems. Archive for Rational Mechanics and Analysis, 168(4):299–328, 2003.
- [KM14] Inwon Kim and Antoine Mellet. Liquid drops sliding down an inclined plane. Trans. Amer. Math. Soc., 366(11):6119–6150, 2014.
- [KP18] Inwon Kim and Norbert Požár. Porous medium equation to hele-shaw flow with general initial density. Transactions of the American Mathematical Society, 370(2):873–909, 2018.
- [KPW19] Inwon Kim, Norbert Požár, and Brent Woodhouse. Singular limit of the porous medium equation with a drift. Advances in Mathematics, 349:682–732, 2019.
- [Lio69] J.-L. Lions. Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod; Gauthier-Villars, Paris, 1969.
- [MPQ17] Antoine Mellet, Benoît Perthame, and Fernando Quiros. A hele–shaw problem for tumor growth. *Journal of Functional Analysis*, 273(10):3061–3093, 2017.
- [MRCS10] Bertrand Maury, Aude Roudneff-Chupin, and Filippo Santambrogio. A macroscopic crowd motion model of gradient flow type. *Mathematical Models and Methods in Applied Sciences*, 20(10):1787–1821, 2010.
- [Mur07] James D Murray. Mathematical biology: I. An introduction, volume 17. Springer Science & Business Media, 2007.
- [PQV14] Benoît Perthame, Fernando Quirós, and Juan Luis Vázquez. The Hele-Shaw asymptotics for mechanical models of tumor growth. Arch. Ration. Mech. Anal., 212(1):93–127, 2014.
- [QV99] Fernando Quirós and Juan Luis Vazquez. Asymptotic behaviour of the porous media equation in an exterior domain. Annali della Scuola Normale Superiore di Pisa-Classe di Scienze, 28(2):183–227, 1999.
- [San18] Filippo Santambrogio. Crowd motion and evolution pdes under density constraints. ESAIM: Proceedings and Surveys, 64:137–157, 2018.
- [V07] Juan Luis Vázquez. *The porous medium equation*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2007. Mathematical theory.
- [Wit97] Thomas P Witelski. Segregation and mixing in degenerate diffusion in population dynamics. *Journal of Mathematical Biology*, 35(6):695–712, 1997.

DEPARTMENT OF MATHEMATICS, TEXAS STATE UNIVERSITY, SAN MARCOS, TX *Email address*: nestor@txstate.edu

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA $\mathit{Email}\ address:\ \texttt{ikim@math.ucla.edu}$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD *Email address*: mellet@umd.edu