

A HEURISTIC FOR DISCRETE MEAN VALUES OF THE DERIVATIVE OF THE RIEMANN ZETA FUNCTION

CHRISTOPHER HUGHES, GREG MARTIN, AND ANDREW PEARCE-CRUMP

ABSTRACT. Shanks conjectured that $\zeta'(\rho)$, where ρ ranges over non-trivial zeros of the Riemann zeta function, is real and positive in the mean. We present a history of this problem, including a generalisation to all higher-order derivatives $\zeta^{(n)}(s)$, for which the sign of the mean alternatives between positive for odd n and negative for even n . Furthermore, we give a simple heuristic that provides the leading term (including its sign) of the asymptotic formula for the average value of $\zeta^{(n)}(\rho)$.

1. INTRODUCTION

Shanks' conjecture, which dates from 1961, states that

$\zeta'(\rho)$ is real and positive in the mean

as $\rho = \beta + i\gamma$ ranges over non-trivial zeros of the Riemann zeta function $\zeta(s)$. More recently this assertion has been generalised to higher derivatives:

On average, $\zeta^{(n)}(\rho)$ is positive if n is odd, but negative if n is even.

The aim of this paper is to show how one can heuristically derive this Generalised Shanks' Conjecture, indeed in quantitative form, almost immediately from an explicit formula known as the Landau–Gonek Theorem.

The Riemann zeta function $\zeta(s)$ has infinitely many non-trivial zeros, which are numbers $\rho = \beta + i\gamma$ with $0 < \rho < 1$ that satisfy $\zeta(\rho) = 0$. The locations of these non-trivial zeros, as we have known for a century and a half (since Riemann), are intimately connected with the distribution of prime numbers. While impressive calculations were done by hand over the years (culminating in Titchmarsh and Comrie's verification [18] that the first thousand zeros of $\zeta(s)$ lie on the critical line $\beta = \frac{1}{2}$), the advent of electronic computers in the middle of the 20th century provided a significant boost to computations of the non-trivial zeros.

Shanks first made his conjecture [17] when he was reviewing Haselgrove's tables [10] of numerical values of the Riemann zeta function. He plotted the graph of $t \mapsto \zeta(\frac{1}{2} + it)$ and noticed that the way this curve approaches the origin, mainly through the third and fourth quadrants, suggests the phase of $\zeta'(\frac{1}{2} + i\gamma)$ is close to zero in the mean and thus $\zeta'(\frac{1}{2} + i\gamma)$ is positive and real in the mean. This observation stands in contrast to the general behaviour of the function $\zeta'(\frac{1}{2} + it)$, whose mean value tends quickly to 0. We present a different graph to the one Shanks created: Figure 1 shows the graph of $\zeta'(\rho)$ for the first 100,000 zeros of $\zeta(s)$, which clearly displays symmetry and a bias towards the positive side of the complex plane.

In 1985, Conrey, Ghosh and Gonek [3] were looking for a straightforward proof of the fact that there are infinitely many simple zeros of $\zeta(s)$. To this end, they used the




FIGURE 1. Scatterplot of $\zeta'(\rho)$ for the first 100,000 zeros of the zeta function

Cauchy–Schwarz inequality to write

$$\left| \sum_{0 < \gamma \leq T} \zeta'(\rho) \right|^2 \leq \sum_{0 < \gamma \leq T}^* 1 \sum_{0 < \gamma \leq T} |\zeta'(\rho)|^2$$

where the star on the middle sum denotes counting only the simple zeros. To complete the proof they used the asymptotic for the discrete second moment on the right-hand side, which was proved by Gonek [7] in 1984, and so only needed to find the leading order behaviour of the sum on the left-hand side. They finished their proof that there are infinitely many simple zeros of $\zeta(s)$ by showing that

$$\sum_{0 < \gamma \leq T} \zeta'(\rho) = \frac{T}{4\pi} (\log T)^2 + O(T \log T), \quad (1.1)$$

which proved Shanks' conjecture as an additional benefit.

As a side note, this method was not strong enough to show that a positive proportion of the non-trivial zeros of $\zeta(s)$ are simple (a fact that goes back to work of Levinson [15]). However, later in 1998 they were able [4] to prove, by modifying the method to include mollifiers, that at least $\frac{19}{27}$ of these non-trivial zeros are simple assuming the Riemann Hypothesis and the Generalised Lindelöf Hypothesis. In 2013, Bui and Heath-Brown [2] removed the assumption of the Generalised Lindelöf Hypothesis through careful use of the generalised Vaughan identity.

Before we describe the generalisation of Shanks' conjecture to higher derivatives, it is worth describing the ongoing research progress on the distribution of $\zeta'(\rho)$. In 1994 Fujii [5] found explicit lower order terms for the asymptotic formula (1.1) (he later corrected [6] a slight error in the lowest-order coefficient when writing a paper where he combined $\zeta'(\rho)$ with the Landau–Gonek Theorem). The full correct asymptotic is given by

$$\sum_{0 < \gamma \leq T} \zeta'(\rho) = \frac{T}{4\pi} \left(\log \frac{T}{2\pi} \right)^2 + (-1 + C_0) \frac{T}{2\pi} \log \frac{T}{2\pi} + (1 - C_0 - C_0^2 + 3C_1) \frac{T}{2\pi} + E(T)$$

where C_0 and C_1 are coefficients in the Laurent expansion of $\zeta(s)$ about $s = 1$. (Unconditional effective bounds for $E(T)$ are known, and a savings of a power of T can be shown under the Riemann Hypothesis. The best known unconditional error term can be found in [11]). In 2021 Kobayashi [13] gave a similar result for Dirichlet L -functions.

In 2010 Trudgian [19] gave an alternative proof of Shanks' conjecture, based on Shanks' observation of the connection between $\arg \zeta'(\rho)$ and the so-called Gram's Law. Although care needs to be taken in the definition of the argument (a priori it is only defined up to a multiple of 2π), Trudgian was able to show that

$$\sum_{0 < \gamma \leq T} \arg \zeta'(\rho) \ll_{\varepsilon} T^{\varepsilon}$$

for every $\varepsilon > 0$.

We are now ready to explore the generalisation of Shanks' Conjecture to $\zeta^{(n)}(s)$, the n th derivative of $\zeta(s)$, for every positive integer n . This extension, which we call the Generalised Shanks' Conjecture, states that $\zeta^{(n)}(\rho)$ is real and positive in the mean if n is even, while $\zeta^{(n)}(\rho)$ is real and negative in the mean if n is odd.




FIGURE 2. Scatterplot of $\zeta^{(n)}(\rho)$ for $n = 2$ and $n = 3$ for the first 100,000 zeros of the zeta function

In 2011, using ideas similar to those in Conrey, Ghosh and Gonek's proof of Shanks' original conjecture, Kaptan, Karabulut and Yıldırım [12] found the main term for the summatory function of $\zeta^{(n)}(\rho)$. Specifically, they showed that

$$\sum_{0 < \gamma \leq T} \zeta^{(n)}(\rho) = \frac{(-1)^{n+1}}{n+1} \frac{T}{2\pi} (\log T)^{n+1} + O(T(\log T)^n) \quad (1.2)$$

which clearly implies the Generalised Shanks' Conjecture as stated above. They have also proven a similar result for derivatives of Dirichlet L -functions, with the same leading-order asymptotic term. (Note that Shanks' Conjecture was first conjectured and later proved, in the usual order of events, but that we still call the established assertion Shanks' "Conjecture". This nomenclature leads to an unusual situation where the extension is reasonably labeled "Shanks' Generalised Conjecture" even though its first appearance in the literature was as a proven result!)

In 2022 the first and third author [11] investigated the Generalised Shanks' Conjecture and found lower order terms for the asymptotic formula (1.2). The result can be stated

in the form

$$\sum_{0 < \gamma \leq T} \zeta^{(n)}(\rho) = \frac{(-1)^{n+1}}{n+1} \frac{T}{2\pi} \left(\log \frac{T}{2\pi} \right)^{n+1} + \frac{T}{2\pi} \mathcal{P}_n \left(\log \frac{T}{2\pi} \right) + E_n(T)$$

where $\mathcal{P}_n(x)$ is a polynomial of degree n described explicitly in [11]; as with Fujii's result, unconditional effective bounds for the error term $E_n(T)$ are known, and a power savings in T can be shown under the Riemann Hypothesis. Furthermore, the third author [16] has also extended Fujii's combination (mentioned earlier) of $\zeta'(s)$ and the Landau–Gonek Theorem to all derivatives of $\zeta(s)$.

All of the results we have described have complicated proofs, and it is unclear that any of them provide an intuitive explanation for the alternating signs in the Generalised Shanks' Conjecture, or indeed of the positive sign in Shanks' original conjecture. We now present a simple heuristic that explains these signs, and indeed recovers (nonrigorously) the leading-order term in the asymptotic formula (1.2).

2. HEURISTIC FOR THE RIEMANN ZETA FUNCTION

We begin with some preliminaries concerning the Riemann zeta function, which for $\Re(s) > 1$ can be written as the convergent Dirichlet series

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}.$$

Since Dirichlet series converge locally uniformly we may differentiate term by term; and since the derivative of $\frac{1}{m^s} = e^{-s \log m}$ is $(-\log m)m^{-s}$, we see that

$$\zeta^{(n)}(s) = (-1)^n \sum_{m=1}^{\infty} \frac{(\log m)^n}{m^s}$$

for $\Re(s) > 1$. Writing $s = \sigma + it$ with $\sigma > 1$, a simple application of Euler–Maclaurin yields

$$\zeta^{(n)}(s) = (-1)^n \sum_{m \leq t} \frac{(\log m)^n}{m^s} + O(t^{-\sigma} (\log t)^n). \quad (2.1)$$

We state a corollary of the Landau–Gonek Theorem, found in [9], concerning sums of the form X^ρ over the non-trivial zeros of the Riemann zeta function. This corollary is the main result that the heuristic relies upon. We call this result the Landau–Gonek Theorem for the sake of brevity in this paper, but note it is given in a more general form in the cited papers. It was proved by Landau [14] for fixed X and made uniform by Gonek [8, 9]. We note that while our quoted result requires the assumption of the Riemann hypothesis, the general result does not.

Lemma 1. *Under the Riemann Hypothesis, for $T > 1, m \in \mathbb{N}$ with $m \geq 2$ and ρ a non-trivial zero of the Riemann zeta function $\zeta(s)$,*

$$\sum_{0 < \gamma \leq T} m^{-\rho} = -\frac{T}{2\pi} \frac{\Lambda(m)}{m} + O(\log(2mT) \log \log(3m)), \quad (2.2)$$

where $\Lambda(m)$ is the von Mangoldt function, which therefore only contributes when m is a prime power.

2.1. The heuristic. We now state the short heuristic argument which yields the Generalised Shanks' Conjecture as stated in Section 1. We ignore error terms for the large part so the equal signs are not 'true' equalities but show the general argument. (But see Section 2.2.)

Remark. As we were writing this paper, we became aware that the heuristic presented in this paper can be found in [9] in a different context, where it can be applied rigorously.

Throughout this section we write $\rho = \beta + i\gamma$ for a non-trivial zero of $\zeta(s)$.

Using the approximate formula (2.1) for $\zeta(s)$ and the Landau–Gonek Theorem (2.2), we may write for $n \geq 1$,

$$\begin{aligned} \sum_{0 < \gamma \leq T} \zeta^{(n)}(\rho) &\approx (-1)^n \sum_{0 < \gamma \leq T} \sum_{m \leq \gamma} (\log m)^n m^{-\rho} \\ &= (-1)^n \sum_{m \leq T} (\log m)^n \sum_{m < \gamma \leq T} m^{-\rho} \\ &\approx (-1)^{n+1} \frac{T}{2\pi} \sum_{m \leq T} \frac{(\log m)^n \Lambda(m)}{m} - (-1)^{n+1} \frac{1}{2\pi} \sum_{m \leq T} (\log m)^n \Lambda(m) \end{aligned} \quad (2.3)$$

To finish the heuristic, we need to sum the series in the last line. By Chebyshev's Theorem [1],

$$C(x) = \sum_{m \leq x} \frac{\Lambda(m)}{m} = \log x + O(1)$$

so by partial summation, we have

$$\begin{aligned} \sum_{m \leq T} \frac{(\log m)^n \Lambda(m)}{m} &= C(T)(\log T)^n - n \int_1^T \frac{C(x)(\log x)^{n-1}}{x} dx \\ &= \frac{1}{n+1} (\log T)^{n+1} + O((\log T)^n). \end{aligned}$$

and similarly

$$\sum_{m \leq T} (\log m)^n \Lambda(m) = O(T(\log T)^n)$$

Combining this with our argument above, we have

$$\sum_{0 < \gamma \leq T} \zeta^{(n)}(\rho) = (-1)^{n+1} \frac{T}{2\pi} \frac{1}{n+1} (\log T)^{n+1} + O(T(\log T)^n),$$

as $T \rightarrow \infty$, which is the leading order the asymptotic result for the Generalised Shanks' Conjecture.

Remark. Note that this result assumes $n \geq 1$. When $n = 0$ the LHS is trivially zero, whilst the RHS, as written, is not. This is explained by noticing that when $n \geq 1$ the term when $m = 1$ in the sum (2.3) is not present, since $\log(1)^n = 0$. However, if $n = 0$ this term is present and is not accounted for in the Landau–Gonek Theorem, which holds for $m \geq 2$ only. The $m = 1$ term in that case clearly contributes $N(T)$, which perfectly cancels the $-\frac{T}{2\pi} \log T + O(T)$ term coming from the calculation given above.

Remark. This argument can be applied *mutatis mutandis* for Dirichlet L -functions, yielding exactly the same leading-order behaviour as for the Riemann zeta function. This result was already known [12, 13].

2.2. The error terms. In the previous section in the last part of equation (2.3) we ignored the error terms coming from the Landau–Gonek formula (2.2). Including them one can see that they contribute

$$\ll \sum_{m \leq T} (\log m)^n \log \log(3m) \log T \ll T(\log T)^{n+1} \log \log T$$

which dominates the main term! However, these are worst-case point-wise estimates and take no account of any potential cancellation when averaged, so it is likely that when summed over m the true error is smaller and the main term is correct. (Indeed that is what is proved, by different methods, in [3, 12] for $n = 1$ and $n \geq 2$ respectively).

The point is that whilst this method must remain a heuristic, it is a much quicker approach to find the main term in the overage of $\zeta^{(n)}(\rho)$ and gives some sense of why the Generalised Shanks’ Conjecture holds true, that is why the mean of $\zeta^{(n)}(\rho)$ is real and positive / negative.

Remark. In [11] the first and third authors prove that the true value of the mean of $\zeta^{(n)}(\rho)$ (that is, the main term and all the lower-order terms up to a power saving) is given by

$$\sum_{\substack{\ell, m \\ \ell m \leq \frac{T}{2\pi}}} \Lambda(m) (\log m)^n$$

which shows that the heuristic in the previous section can only yield the main term since the last line of equation (2.3) differs from the true value by $O(T(\log T)^n)$.

ACKNOWLEDGEMENTS

The second author suggested the heuristic in this paper at the PIMS CRG ‘Moments of L -functions’ workshop in Prince George, British Columbia to the third author. The third author would like to thank the group, the LMS and the University of York for their generous funding.

REFERENCES

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, 1st ed., Springer, 1976.
- [2] H. M. Bui and D. R. Heath-Brown, *On simple zeros of the Riemann zeta-function*, Bull. Lond. Math. Soc. **45** (2013), 953–961.
- [3] J. B. Conrey, A. Ghosh, and S. M. Gonek, *Simple zeros of zeta functions*, Colloque de Théorie Analytique des Nombres “Jean Coquet” (Marseille, 1985), Publ. Math. Orsay, vol. 88, Univ. Paris XI, Orsay, 1988, pp. 77–83.
- [4] ———, *Simple zeros of the Riemann zeta-function*, Proc. Lond. Math. Soc. (3) **76** (1998), 497–522.
- [5] A. Fujii, *On a conjecture of Shanks*, Proc. Japan Acad. A **70** (1994), 109–114.
- [6] ———, *On the Distribution of Values of the Derivative of the Riemann Zeta Function at Its Zeros. I*, Proc. Steklov Inst. Math. **276** (2012), 51–76.
- [7] S. M. Gonek, *Mean values of the Riemann zeta-function and its derivatives*, Invent. math. **75** (1984), 123–141.
- [8] ———, *A formula of Landau and mean values of $\zeta(s)$* , Topics in analytic number theory (Austin, Tex., 1982), Univ. Texas Press, Austin, TX, 1985, pp. 92–97.

- [9] ———, *An explicit formula of Landau and its applications to the theory of the zeta-function*, A tribute to Emil Grosswald: number theory and related analysis, Contemp. Math., vol. 143, Amer. Math. Soc., Providence, RI, 1993, pp. 395–413.
- [10] C.B. Haselgrove in collaboration with J.C.P. Miller, *Mathematical Tables: Tables of the Riemann zeta function* Royal Society Mathematical Tables No. 6, Cambridge University Press, (1960).
- [11] C. Hughes and A. Pearce-Crump, *A discrete mean-value theorem for the higher derivatives of the Riemann zeta function*, J. Number Theory **241** (2022), 142–164.
- [12] D. Kaptan, Y. Karabut, and C. Yildirim, *Some mean value theorems for the Riemann zeta-function and Dirichlet L-functions.*, Comment. Math. Univ. St. Pauli **60** (2011), 83–87.
- [13] H. Kobayashi, *On a certain sum of the derivatives of Dirichlet L-functions*, arXiv:2003.05629, 2021.
- [14] E. Landau, *Über die Nullstellen der Zetafunktion*, Math Annalen **71** (1911), 548–564.
- [15] N. Levinson, *More than one-third of zeros of Riemann’s zeta-function are on $\sigma = \frac{1}{2}$* , Adv. in Math. **13** (1974) 383–436.
- [16] A. Pearce-Crump, *A further generalisation of sums of higher derivatives of the Riemann zeta function*, arXiv:2111.02756, 2021.
- [17] D. Shanks, *Review of ‘Tables of the Riemann zeta function’ by C.B. Haselgrove in collaboration with J.C.P. Miller*, Math. Comp. (1961), 84–86.
- [18] E. C. Titchmarsh, *The Zeros of the Riemann Zeta-Function*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, The Royal Society, **157** (1936): 261—263.
- [19] T. Trudgian, *On a conjecture of Shanks*, J. Number Theory **130** (2010), 2635–2638.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK, YO10 5DD, UNITED KINGDOM
Email address: christopher.hughes@york.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BRITISH COLUMBIA,
V6T 1Z2, CANADA
Email address: gerg@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK, YO10 5DD, UNITED KINGDOM
Email address: aepc501@york.ac.uk