

A HISTOCHEMICAL STUDY OF PHAGOCYTTIC AND ENZYMATIC FUNCTIONS OF RABBIT MONONUCLEAR AND POLYMORPHONUCLEAR EXUDATE CELLS AND ALVEOLAR MACROPHAGES

I. Survey and Quantitation of Enzymes, and States of Cellular Activation

ARTHUR M. DANNENBERG, JR., M.D., MARVIN S. BURSTONE, D.D.S., PAUL C. WALTER, and JUNE W. KINSLEY

From The Henry Phipps Institute, Department of Public Health and Preventive Medicine, University of Pennsylvania School of Medicine, Philadelphia, and the National Cancer Institute, Bethesda

ABSTRACT

The cytochrome oxidase (CO), aminopeptidase (AMP), succinic dehydrogenase (SD), acid phosphatase, esterase, and alkaline phosphatase of rabbit mononuclear (MN) and polymorphonuclear (PMN) peritoneal exudate cells and pulmonary alveolar macrophages (AM) — air dried on Mylar strips — were characterized by histochemical techniques with respect to stability, activators, inhibitors, and pH optima. A granule count method was established for the quantitation of these enzymes. For the acid phosphatase of MN, in which the most precise results were obtained, time, pH, substrate, and inhibitor curves resembled those commonly obtained biochemically. Five of these enzymes were usually more active in AM than MN, whereas the sixth, alkaline phosphatase, was not present in either cell type. AM also tended to consume more oxygen than MN and to divide more frequently. Since the most active cells in the population would be first involved in the host's defense against microbial agents, a comparison was made of the 10 per cent of the AM and MN with the highest enzymatic activities. No differences were found in the granule counts that were not reflected by the means. However, within a given AM population, cells containing ingested dust particles seemed to have higher enzymatic activities than those without particles. MN had greater acid phosphatase and SD activities than PMN and consumed more oxygen, but the CO, AMP, and esterase activities of both types of cells were of similar magnitude. PMN showed high alkaline phosphatase activity; MN showed none. A survey of the histochemical literature indicates that a positive correlation between the enzymatic and phagocytic activities of both MN and PMN exists *in vivo*.

Mononuclear phagocytes (MN)¹ are essential components of the reticuloendothelial system—the chief defense of the host against many microorganisms and their products. Similar functions

are served by polymorphonuclears (PMN) in the blood and in areas of inflammation, and by alveolar macrophages (AM) in the lung. The present report (a) characterizes six enzymes of rabbit MN

¹The abbreviations used in this report are as follows: MN, mononuclear(s), or mononuclear peritoneal exudate cells; PMN, polymorphonuclear(s) or polymorphonuclear peritoneal exudate cells; AM,

alveolar macrophage(s) of the lung; ip, intraperitoneal; CO, cytochrome oxidase; AMP, aminopeptidase; SD, succinic dehydrogenase; Tris, tris(hydroxymethyl)aminomethane.

and PMN peritoneal exudate cells and alveolar macrophages by utilizing *unfixed* Mylar strip preparations in aqueous histochemical reagents, (b) establishes the granule count method for the quantitation of such enzymes, and (c) utilizes this technique to compare the enzyme contents of MN and AM, and MN and PMN. The following paper (3) evaluates the role of these enzymes in the phagocytic process.

MATERIALS AND METHODS

Collection and Quantitation of MN, PMN, and AM

MN exudates were elicited in rabbits by the intraperitoneal (ip) injection of 35 ml of heavy mineral oil (u.s.p.). Five to 6 days later, the animals were killed by exsanguination, and their peritoneal cavities washed with citrate-saline (0.4 per cent sodium citrate in 0.85 per cent sodium chloride) or Hanks' solution (4) containing 0.01 per cent heparin. The exudate cells were filtered through gauze, collected by centrifugation, and either used as such or washed once with physiological saline (PSS) or Hanks' solution. Total counts ranged from 30 to 150 million cells per exudate, with a mean of about 80 million. Differential cell counts employing Wright's stain ranged from 80 to 99 per cent MN, with a mean of about 95 per cent. A high percentage of the mononuclear cells were of medium size with a fairly basophilic cytoplasm that sometimes contained mineral oil vacuoles. This type of cell proved most active in the phagocytosis of particles, and for this reason was selected for all the studies on MN exudates reported below. The other cells in these exudates were ignored; e.g., larger, paler MN, PMN, and occasional lymphocytes and mesothelial cells.

PMN exudates were elicited by a method adapted from that of Hirsch (5), namely, the ip injection of 250 to 500 ml of 0.1 per cent glycogen (Nutritional Biochemicals Corporation, Cleveland) in PSS, and the collection of cells 4 (or 18) hours later in a manner similar to that used for MN. Total PMN counts averaged above 100 million cells. Differential cell counts averaged 85 to 95 per cent PMN.

Alveolar macrophages were obtained by a method adapted from that of Myrvik *et al.* (6). Specifically, about 35 ml of PSS, citrate-saline, or heparinized Hanks' solution were injected intratracheally after removal of the lungs from a recently sacrificed rabbit. The fluid was then allowed to drain out. This procedure was repeated and the two washings were pooled, centrifuged and counted in a manner similar to that used for the peritoneal exudates. Total counts averaged about 20 million cells; differential counts averaged 98 per cent AM and 2 per cent PMN.

The *viabilities* of the exudate cells were determined by the trypan blue and eosin methods. These methods, adapted from others (7-10), consisted of mixing, on a microscope slide, one drop of exudate, one drop of autologous serum, and a third of a drop of 1 per cent trypan blue or 0.5 per cent eosin Y in PSS, covering the mixture with a coverslip, and incubating it at 37°C for 10 minutes. Examination under the microscope revealed stained cells (dead) and unstained cells (alive). AM and MN averaged 93 to 99 per cent viability with trypan blue. PMN averaged about 6 per cent lower. With eosin the viabilities of all three types of cells appeared 5 to 10 per cent less than with trypan blue. High degrees of viability were also found in phagocytosis studies on AM, MN, and PMN (3), which have been shown by Tullis (11) to be a more rigorous criterion than dye-permeability tests.

Preparation of Mylar Strips

The cell suspensions were smeared on Mylar² strips (12, 13), employing the end of a microscope slide. The strips were air-dried at room temperature for about 20 minutes; no fixation was employed. They then were placed in the refrigerator (5°C) until samples were removed for the determination of succinic dehydrogenase (usually within 3 hours, though sometimes on the next day). After this, the strips were stored in the deep freezer (-25°C). Aminopeptidase and cytochrome oxidase were usually assayed within 3 days, but the activities of the other 3 enzymes described in this report (esterase, acid phosphatase, and alkaline phosphatase) seemed more stable. Although they were usually assayed within a week, they were readily demonstrable in the Mylar preparations after several weeks of storage.

In contrast with preparations on glass, unfixed exudate cells adhered rather well to the Mylar strips during their incubation face down on the surface of the reagent solutions, during their subsequent dips in distilled water, and during their counterstaining with half-strength Ehrlich's acid alum hematoxylin (frequently omitted). Occasionally, however, there was some loss of cells in certain areas. This loss can be decreased, if necessary, by the addition of serum to the exudate cells before they are smeared. After histochemical staining, the strips were air-dried, mounted face up in glycerol gelatin on glass slides, and covered by a coverslip.

Measurement of Oxygen Consumption

The oxygen consumption of cells was determined manometrically in standard Warburg flasks by the

² DuPont Mylar, type 50C, 0.25 mil thick (13), is available from Brownell Distributors, Inc., New York. It resembles cellophane or Saran wrap.

method described by Umbreit *et al.* (14). Although the experiments varied, the following amounts of materials are representative of those employed: In the main compartment were phagocytes in citrate-saline (0.5 ml), 1 per cent glucose in PSS (0.6 ml), autologous serum (0.3 ml), and serum from a rabbit immunized against yeast (3) and diluted 1:6 with PSS (0.8 ml). The pH was around 7.0. In each of the two side arms were 0.4 ml of boiled yeast suspension (3). In the center well was 0.20 ml of 20 per cent KOH. The yeast was employed in the phagocytosis studies presented in another report (3); the endogenous respiration of the various phagocytes is presented in this report. Controls made up similarly, but with no phagocytes in the flask, showed no oxygen uptake.

EXPERIMENTAL DATA AND RESULTS

Histochemical Studies on Six MN, PMN, and AM Enzymes Whose Activities May Reflect the Metabolic Potentialities of These Cells

CYTOCHROME OXIDASE (CO)

This enzyme was demonstrated in MN, PMN, and AM by the histochemical method of Burstone (15, 16). Naphthol AS-L3G³ or 8-amino-1,2,3,4-tetrahydroquinoline⁴ (10 mg) and *p*-aminodiphenylamine⁵ (10 mg) were dissolved in 0.5 ml of ethanol and made up to 50 ml with 0.1 M Tris buffer (tris(hydroxymethyl)aminomethane) at pH 7.4. The solution was then filtered into small beakers, and Mylar strips of exudate cells were floated on the surface for 1 to 2 hours of incubation at room temperature. A brown-black or blue-black color resulted, depending on the reagents used. The strips were then fixed for 1 hour in a solution containing 10 per cent of each of the following materials: cobalt acetate, formalin, and 0.2 M acetate buffer, (pH 5.2). They were then washed in water and mounted in glycerol gelatin. Figs. 1, 13, and 14 show representative results.

³ Pfister Chemical Works, Inc., Ridgefield, New Jersey. (See reference 16 for chemical name).

⁴ Eastman Organic Chemicals, Distillation Products Industries, Rochester, New York, (practical grade).

⁵ British Drug Houses, Ltd., Poole, England, available from the Earling Corporation, Cambridge, Massachusetts. (See reference 16 for chemical name and formula, synonyms, and alternate sources of supply.)

The color in MN and AM was considerably augmented by cytochrome C in a final concentration of 0.2 mg per ml (see Table I). This substance was used in all routine CO determinations; in fact, it enabled the demonstration of this oxidase after the Mylar preparations had been stored in the deep freezer for several days. Catalase had no apparent effect on the enzyme's activity, whereas 10 per cent neutral formalin destroyed it within 5 minutes (see Hannibal *et al.*, 17). Other inhibitors are listed in Table I. These observations suggest that cytochrome oxidase, rather than peroxidase (18), is demonstrated by this reaction in rabbit MN and AM. Our studies with PMN, however, were not extensive enough to favor one enzyme above the other.

No precise quantitation of this enzyme was possible, because of the variability in the amount of color developed in adjacent cells of the same exudate and in adjacent areas of the same Mylar strip.

AMINOPEPTIDASE (AMP)

This enzyme was demonstrated in MN, PMN, and AM by the histochemical method of Burstone and Folk (19) which had been modified from Gomori (20). DL-alanyl- β -naphthylamide⁶ and fast garnet GBC^{7, 8} (each 1 mg per ml) were used as substrate and coupler, respectively, in 0.1 M Tris buffer (pH 7.1). The solution was filtered and Mylar strips incubated on its surface for 1 to 2 hours at room temperature. Figs. 2 and 3 show representative, though rather heavily stained, results for MN and AM. The effects of certain inhibitors on this enzyme are presented in Table I.

Because of a certain amount of non-specific yellow staining, (the specific color is orange-brown), this method was not too satisfactory for quantitative studies. In addition, with storage the color occasionally fades and the dye reorganizes into a coarse precipitate.

⁶ Synthesized by Dr. M. S. Burstone (19); commercially available from Mann Research Lab., New York.

⁷ Dyestuff Chemical and Pigments Division, General Aniline and Film Corporation, New York.

⁸ The Azoic Diazo Number, Color Index Number, structural formula, chemical name, molecular weight and synonyms of this stable diazonium salt are listed by Lillie (21).

FIGURE 1

Cytochrome oxidase activity of rabbit polymorphonuclear (PMN) and mononuclear (MN) exudate cells. The densely stained cell is an MN; the rest are PMN. In other microscopic fields PMN stained as densely as MN. Glycogen exudate. $\times 1000$.

FIGURE 2

Aminoamidase activity of MN. One of the more densely stained groups of cells is depicted. Mineral oil exudate. $\times 1000$.

FIGURE 3

Aminoamidase activity of alveolar macrophages (AM) from the same rabbit that provided the MN of Fig. 2. Both the AM and MN were incubated simultaneously in the same histochemical reagents. One of the more densely stained groups of cells is depicted. $\times 1000$.

FIGURE 4

Succinic dehydrogenase activity of MN and PMN. The two cells with the most dye granules are MN. The rest are PMN. All but a few of the PMN depicted are unstained. An MN can usually be differentiated from a PMN by observing its nucleus at different levels of focus. Glycogen exudate, counterstained with half-strength Ehrlich's acid alum hematoxylin. $\times 1000$.

FIGURE 5

Succinic dehydrogenase activity of MN. One of the more densely stained groups of cells is depicted. Mineral oil exudate. $\times 1000$.

FIGURE 6

Succinic dehydrogenase activity of AM from the same rabbit that provided the MN in Fig. 5. The cells represented in both of these figures were incubated simultaneously in the same histochemical reagents. One of the more densely stained groups of cells is depicted. $\times 1000$.

FIGURE 7

Acid phosphatase activity of PMN and MN. In general, the MN are more darkly stained. Four are present (at 1, 2, 3, and 7 o'clock respectively). Glycogen exudate, counterstained with half-strength Ehrlich's acid alum hematoxylin. $\times 1000$.

FIGURE 8

Esterase activity of MN. One of the more densely stained groups of cells is depicted. Mineral oil exudate. $\times 1000$.

FIGURE 9

Esterase activity of AM from the same rabbit that provided the MN in Fig. 8. The cells represented in both of these figures were incubated simultaneously in the same histochemical reagents. One of the more densely stained groups of cells is depicted. $\times 1000$.

FIGURE 10

Alkaline phosphatase activity of PMN and MN. The two very pale cells to the right of the center are MN. The rest are PMN. Glycogen exudate, counterstained with half-strength Ehrlich's acid alum hematoxylin. $\times 1000$.

FIGURE 11

Esterase activity of PMN and MN. The four largest cells are MN; the rest are PMN. In this example, the MN stained slightly darker than the PMN, but in other examples the two groups stained with equal intensities. Glycogen exudate. $\times 1000$.

FIGURE 12

AM that have ingested autologous red blood corpuscles (rbc) *in vivo*. The center cell contains three rbc, the lower cell contains one. A few dust particles are also visible in the center cell. Wright's stain. $\times 1000$.




TABLE I
*The Effect of Inhibitors and Activators on Certain Histochemically Demonstrable
 Enzymes of MN, PMN, and AM*

| Enzyme | Inhibitor (I) or activator (A) | Amount of inhibition or activation* | | |
|------------------------|--|-------------------------------------|------|------|
| | | MN | PMN | AM |
| Cytochrome oxidase | A Cytochrome C‡ (0.2 mg per ml) | ++++ | ++ | ++++ |
| | I NaCN (0.001 M) | ++ | ++ | |
| | I Na ₂ S (0.01 M) | ++ | +++ | |
| | I Sodium azide (0.01 M) | ++ | 0 | |
| | I Catalase (0.02 per cent) | 0 | 0 | 0 |
| Aminopeptidase | I KCN (0.01 M) | + | + | + |
| | I CuSO ₄ (0.01 M) | ++ | ++++ | ++ |
| | I CuSO ₄ (0.0015 M) | +++ | +++ | +++ |
| Succinic dehydrogenase | I Sodium malonate (0.008 M) | ++ | + | + |
| | I <i>p</i> -chloromercuribenzoate (0.0002 M) | +++ | ++++ | +++ |
| Acid phosphatase | I Sodium molybdate (0.005 M) | ++++ | +++ | ++++ |
| | I Sodium arsenilate (0.001 M) | 0 | | 0 |
| | I NaF (0.01 M) | +++ | +++ | ++ |
| | A MnCl ₂ (0.002 M) | ± | | |
| Alkaline phosphatase | I Sodium molybdate (0.005 M) | | + | |
| | I Sodium arsenilate (0.001 M) | | + | |
| | I NaF (0.01 M) | | + | |
| | I Sodium iodoacetate (0.002 M) | | + | |
| | A MnCl ₂ (0.002 M) | | 0 | |
| Esterase | I Eserine salicylate (0.01 M) | 0 | ± | 0 |
| | I Sodium azide (0.01 M) | ++++ | ++++ | +++ |
| | I Diisopropyl fluorophosphate (0.001 M)§ | +++ | ++++ | ++ |
| | I NaF (0.01 M) | ++ | ++++ | 0 |

* 0, no inhibition (or activation); +++++, complete or almost complete inhibition (or activation); others (+, ++, +++) were in between these extremes. Each category represents the results of 2 to 5 experiments.

‡ Cytochrome C, Type II, from horse heart, Sigma Chemical Company, St. Louis.

§ Merck and Co., Inc., Chemical Division, Rahway, New Jersey.

SUCCINIC DEHYDROGENASE (SD)

Dehydrogenase activity was demonstrated in MN, PMN, and AM, essentially by the method of Nachlas *et al.* (22). To 10 mg of Nitro-BT⁹ (2,2'-di-*p*-nitrophenyl-5,5'-diphenyl-3,3'-(3,3'-dimethoxy-4,4'-biphenylene) ditetrazolium chloride) dissolved in 0.5 ml of ethanol was added 500 mg of sodium succinate in 40 ml of 0.1 M Tris

⁹ Dajac Laboratories, Borden Company Chemical Division, Philadelphia.

buffer at pH 7.4. The solution was then filtered into small beakers, and the Mylar strips were incubated on the surface for 0.5 to 2.5 hours at 37°C.

The dehydrogenase of these phagocytes seemed to be a rather unstable enzyme, and so it was generally assayed on the same day that the exudates were withdrawn from the animal. Figs. 4, 5, and 6 show representative results in PMN, MN, and AM. For both MN and PMN the pH optimum was around 7 to 8. At pH 5.7 and 8.9, the activity

was somewhat reduced. The pH optimum for AM was not determined. The effects of certain inhibitors on this reaction are listed in Table I.

When sodium succinate was omitted from the Nitro-BT reagent solution, only a rare formazan granule appeared in MN and AM. With PMN, however, the number of formazan granules formed in the absence of substrate seemed to equal the number formed in its presence. Thus our studies on the SD activity of PMN probably do not concern this specific substrate, but rather an as yet undefined substrate within these cells.

The formazan granules formed by the reduction of Nitro-BT showed more variation in size than the dye granules produced by the other enzymes. An additional complication arose from the *instability* of this dehydrogenase. In phagocytes injured during their collection or during the preparation of the Mylar smear, SD activity was poorly preserved, whereas in morphologically intact MN and AM the dehydrogenase frequently showed high activity.¹⁰ In fact, Wachstein and Meisel (23, 24) consider the activity of this enzyme a most sensitive indicator of cell damage.

ACID PHOSPHATASE

This enzyme was demonstrated in MN, PMN, and AM by the method of Burstone (25, 26). Five mg of naphthol AS-BI phosphate¹¹ was dissolved in 0.5 ml of dimethylformamide. Fifty ml of 0.1 M acetate buffer (pH 5.2) and 10 mg of fast red violet LB^{8, 12} were added and also two drops of 10 per cent MnCl₂ solution as a possible activator. The somewhat opalescent solution was then filtered, and the Mylar strips incubated on its surface at room temperature for 10 to 60 minutes. Figs. 7, 16, and 17 show representative results in PMN, MN, and AM.

The azo dye granules formed in MN usually were discrete, fairly uniform in size, and almost never small enough to create a diffuse appearance. This enzyme was rather stable, for Mylar strips of

¹⁰ Injured and morphologically intact phagocytes can be readily differentiated in Wright's stained-preparations. The injured cell appears larger (presumably because it has absorbed fluid), its nucleus is less basophilic, and its edge is less distinctly demarcated.

¹¹ Synthesized by Dr. M. S. Burstone (25, 26); commercially available from Sigma Chemical Company, St. Louis.

¹² Verona Dyestuffs, Verona Chemical Co., Union, New Jersey.

MN still showed activity after several months of storage in the deep freezer (-25°C). Because of these properties, MN acid phosphatase was used to establish the granule count method described in the next section, where time, pH, substrate, and inhibitor (NaF) relationships are presented.

The dye deposits in AM and PMN were both diffuse and granular and could not be quantitated as precisely as those in MN. Table I presents the effects of inhibitors on all three types of phagocyte.

The method for alkaline phosphatase (see below), with the addition of MnCl₂, was used to construct a pH curve for PMN phosphatase. It had two peaks: one at pH 4.7 to 5.2, and one, 1.5 times as large, at pH 8.3 to 8.9. Some activity was present, however, from pH 4.1 to 8.9 which was the entire pH range tested.

ALKALINE PHOSPHATASE

This enzyme does not seem to be present in rabbit MN or AM, but only in PMN. It was demonstrated by the method of Burstone (25) employing 5 mg of naphthol AS-MX phosphate,¹¹ 0.5 ml of dimethylformamide, 10 mg of fast red violet LB,^{8, 12} and 50 ml of Tris buffer (0.1 M) at pH 8.3. The procedure was similar to that for acid phosphatase, described above. Incubation times varied from 2 to 15 minutes at room temperature. The type of granule produced was rather fine and somewhat diffuse (Fig. 10), so that *quantitative* studies on this enzyme were less precise.

At incubation times of 2 and 3 minutes, pH curves showed an optimal alkaline phosphatase activity above 8.0. Follette *et al.* (27) found biochemically an optimum between 8.3 and 8.9. The effects of certain inhibitors and activators are presented in Table I.

ESTERASE

The activity of this enzyme was demonstrated in MN, PMN, and AM by the method of Burstone (28). Naphthol AS-D acetate¹³ (5 mg) was dissolved in 0.5 ml of dimethylformamide, and 50 ml of 0.1 M Tris buffer at pH 7.1 was added, followed by fast garnet GBC⁷ (10 mg). The mixture was then filtered, and the cell preparations incubated

¹³ Naphthol AS-D acetate was synthesized by Dr. M. S. Burstone (29) and used instead of naphthol AS-LC acetate. It is commercially available from Sigma Chemical Company, St. Louis.

on the surface of the filtrate for 3 to 40 minutes at room temperature. Figs. 8, 9, and 11 show representative results for MN, AM, and PMN.

The azo dye granules produced ranged in size. Some of them were so small that a diffuse appearance resulted, whereas some were as large as *Rickettsia* or even larger. Quantitative studies on this enzyme, therefore, were not very precise. The esterase of both intact and injured MN, PMN, and AM seemed rather stable in the deep freezer. Its pH optimum was around 7.0 to 7.6 for both MN and PMN; the optimum for AM was not evaluated.

Table I presents the effect of certain inhibitors on the esterase activity of these three cell types.

Histochemical Quantitation by the Granule Count Method

STUDIES WITH MN ACID PHOSPHATASE

The acid phosphatase reaction of MN produced discrete dye granules of fairly uniform size and therefore seemed suitable for the development of a new quantitative procedure which we have termed the granule count method. Granule counts were made in the following manner: A unit granule approximately 0.3μ in diameter (Fig. 21) was first selected, and granules of other sizes were designated as fractions or multiples of this unit. The total number of granule units per

FIGURE 13

Cytochrome oxidase activity of MN. One of the more densely stained groups of cells is depicted. Mineral oil exudate. $\times 1000$.

FIGURE 14

Cytochrome oxidase activity of AM from the same rabbit that provided the MN in Fig. 13. The cells represented in both of these figures were incubated simultaneously in the same histochemical reagents. One of the more densely stained groups of cells is depicted. $\times 1000$.

FIGURE 15

Acid phosphatase activity of AM. The center cell contains the most "dust" as well as enzyme. However, some dust and other foreign particles are present in all of the cells shown. $\times 1000$.

FIGURE 16

Acid phosphatase activity of MN. One of the more densely stained groups of cells is depicted. Mineral oil exudate. $\times 1000$.

FIGURE 17


Acid phosphatase activity of AM from the same rabbit that provided the MN in Fig. 16. The cells represented in both of these figures were incubated simultaneously in the same histochemical reagents. One of the more densely stained groups of cells is depicted. $\times 1000$.

FIGURE 18

An alveolar macrophage with 2 nuclei, indicative of cell division. Wright's stain. $\times 1000$.

FIGURES 19 TO 23

The effect of incubation time on MN acid phosphatase activity. The MN represented in these figures were incubated 4, 8, 16, 20, and 45 minutes, respectively, at room temperature. They are the same cells that are represented by black dots in Fig. 24. The smaller granules in Fig. 21 were considered unity for purposes of quantitation. Two of these are marked with arrows. Not all the granules in a given cell are visible at a single level of focus. Mineral oil exudate. $\times 1500$.



cell was then considered a measure of its enzyme content.

As a test of the reliability of this method, the acid phosphatase reaction of MN was run at varying time, pH, and substrate and inhibitor concentrations. The results for varying time are depicted in Figs. 19 to 23 and plotted in Fig. 24. The others are plotted in Figs. 25 to 27. These results resemble curves obtained for enzymes by biochemical techniques. The standard deviation of the mean for a given point on a given curve was usually less than

10 per cent of its numerical value, provided that a single investigator did all the counting.

It seems, therefore, that the granule count method is a valid quantitative procedure and can be used to compare *different* Mylar preparations. However, it is most suitable for the study of individual cells or of different types of cells in the *same* Mylar preparation, *e.g.* cells that have ingested particles and those that have not (see reference 3), or MN and PMN in the same exudate (see below).

STUDIES WITH OTHER ENZYMES

Although these studies could not be so precise as those just plotted for MN acid phosphatase, they




FIGURE 24

Effect of incubation time on acid phosphatase activity of MN. ● and ○ represent the results of two independent investigators and two separate experiments, adjusted to a common base line.




FIGURE 26

Effect of substrate concentration on acid phosphatase activity of MN. ● and ○ represent the results of two independent investigators, adjusted to a common base line.




FIGURE 25

Effect of pH on acid phosphatase activity of MN. ● and ○ represent the results of two independent investigators, adjusted to a common base line.




FIGURE 27

Effect of concentration of NaF inhibitor on acid phosphatase activity of MN. ● and ○ represent the results of two independent investigators, adjusted to a common base line.

did follow a similar pattern. There was a straight line relationship between time and the number of granule units resulting from the succinic dehydrogenase and esterase activities of MN and PMN and the acid and alkaline phosphatase activities of PMN. The effect of pH on these enzymes was reported in the previous section. No time and pH studies were made on AM.

A Comparison of MN to AM and PMN with Respect to Certain Histochemically Demonstrable Enzymes and Oxygen Consumption

Peritoneal and alveolar macrophages represent two varieties of mononuclear phagocytes. The former (MN) are mobilized into the peritoneal cavity by mineral oil, a stimulus that is present 5 to 6 days. The latter (AM) are continuously mobilized into the pulmonary air spaces in response to the inhalation of dust and bacteria, and possibly to the extravasation of occasional red corpuscles (Fig. 12). Although AM resemble MN in general morphology, they are slightly larger, occasionally have a somewhat eosinophilic cytoplasm,¹⁴ and may contain dust particles. MN, on the other hand, may contain vacuoles filled with mineral oil.

MN and PMN represent the two major types of phagocyte in the body. The former clear the blood as a part of the reticuloendothelial system and are active in areas of chronic inflammation. The latter circulate in the blood, and emigrate to areas of acute inflammation.

Because of the differences in function of all three kinds of phagocyte, it is of interest to determine whether some of their enzymes are qualitatively or quantitatively different and to compare their oxygen consumption as a measure of their general activity.

HISTOCHEMICAL STUDIES ON MN AND AM

COMPARISON OF AM AND MN ENZYMES: AM and MN were removed from the same rabbit at the time of death, smeared on Mylar strips, incubated for identical periods in histochemical reagents, and mounted side by side on the same slide. The results were read by two methods. In the first, the paired strips were studied alternately until the amount of color of the cells in the AM

¹⁴ Possibly from the previous ingestion of red blood corpuscles (30); they contain ferritin (31).

strip relative to that of the cells in the MN strip could be determined (Table II). In the second, the amount of granular and diffuse dye material in each cell was assigned numerical values (Table III). A different series of rabbits was used with each method in order to extend the number of observations.

From these tables which present the results, it is apparent that AM showed more enzymatic activity than MN in over half of the cases. In the rest they were approximately equal, except for rare instances in which the reverse occurred. Figs. 2 and 3; 5 and 6; 8 and 9; 13 and 14; and

TABLE II
*A Histochemical Comparison of Various Enzyme Activities of AM and MN from the Same Rabbit—Survey Method**

| Rabbit No. (Series I) | Enzyme | AM | MN | Ratio: AM/MN |
|-----------------------|------------------------|------|-----|--------------|
| 1 | Cytochrome oxidase | ++++ | 1/2 | 8 |
| 2 | | ++ | ++ | 1 |
| 3 | | ++++ | 1/2 | 8 |
| 4 | | ++ | ++ | 1 |
| 8 | | ++ | + | 2 |
| 12 | | ++++ | ++ | 2 |
| 13 | ++ | ++ | 1 | |
| 1 | Aminopeptidase | ++ | + | 2 |
| 4 | | ++ | + | 2 |
| 5 | | + | + | 1 |
| 8 | | ++++ | + | 4 |
| 11 | | ++++ | + | 4 |
| 12 | | ++ | ++ | 1 |
| 2 | Acid phosphatase | +++ | + | 3 |
| 8 | | ++++ | 1/2 | 8 |
| 9 | | ++ | + | 2 |
| 10 | | ++ | + | 2 |
| 5 | Esterase | ++ | + | 2 |
| 6 | | ++++ | + | 4 |
| 7 | | ++++ | + | 4 |
| 10 | | ++++ | + | 4 |
| 12 | | ++ | + | 2 |
| 13 | | ++ | + | 2 |
| 1 | Succinic dehydrogenase | ++++ | + | 4 |
| 6 | | ++++ | + | 4 |
| 7 | | ++ | ++ | 1 |
| 8 | | ++ | + | 2 |

* See text

TABLE III

A Histochemical Comparison of Various Enzyme Activities of AM and MN from the Same Rabbit—Granule Count Method

| Rabbit No. (Series II) | Enzyme | No. of cells evaluated | | Average No. of histochemical dye granules per cell* | | | | | | Ratio: AM/MN for | | |
|------------------------|------------------------|------------------------|-----|---|-----|--|-----|------------------------------|-----|------------------|-----------------|--------------|
| | | AM | MN | In the darkest staining 10 per cent of the cells | | In the darkest staining 20 per cent of the cells | | In 100 per cent of the cells | | Top 10 per cent | Top 20 per cent | 100 per cent |
| | | | | AM | MN | AM | MN | AM | MN | | | |
| II | Cytochrome oxidase | 50 | 55 | 711 | 640 | 693 | 606 | 543 | 386 | 1.1 | 1.1 | 1.4 |
| III | | 53 | 52 | 650 | 671 | 616 | 616 | 384 | 376 | 1.0 | 1.0 | 1.0 |
| V | | 50 | 50 | 692 | 474 | 670 | 456 | 527 | 301 | 1.5 | 1.5 | 1.8 |
| I | Aminopeptidase | 100 | 100 | 200 | 178 | 190 | 173 | 157 | 159 | 1.1 | 1.1 | 1.0 |
| II | | 54 | 59 | 602 | 520 | 633 | 487 | 423 | 294 | 1.2 | 1.3 | 1.4 |
| V | | 50 | 65 | 621 | 402 | 588 | 372 | 439 | 171 | 1.5 | 1.6 | 2.6 |
| II | Succinic dehydrogenase | 41 | 62 | 206 | 313 | 191 | 318 | 106 | 179 | 0.7 | 0.6 | 0.6 |
| III | | 100 | 100 | 105 | 103 | 99 | 92 | 58 | 59 | 1.0 | 1.1 | 1.0 |
| V | | 118 | 134 | 108 | 72 | 89 | 55 | 27 | 14 | 1.5 | 1.6 | 1.9 |
| VI | | 50 | 50 | 356 | 133 | 318 | 119 | 184 | 67 | 2.7 | 2.7 | 2.8 |
| I | Acid phosphatase | 100 | 100 | 54 | 37 | 48 | 35 | 78 | 22 | 1.5 | 1.4 | 3.6 |
| II | | 55 | 43 | 206 | 50 | 185 | 30 | 70 | 18 | 4.1 | 6.2 | 3.9 |
| IV | | 100 | 100 | 113 | 90 | 114 | 80 | 184 | 48 | 1.3 | 1.4 | 3.8 |
| V | | 61 | 61 | 321 | 373 | 229 | 349 | 115 | 171 | 0.9 | 0.7 | 0.7 |
| I | Esterase | 100 | 100 | 132 | 54 | 149 | 48 | 123 | 27 | 2.4 | 3.1 | 4.6 |
| II | | 81 | 145 | 588 | 526 | 531 | 499 | 372 | 271 | 1.1 | 1.1 | 1.4 |
| IV | | 100 | 80 | 532 | 493 | 495 | 448 | 351 | 304 | 1.1 | 1.1 | 1.2 |
| V | | 66 | 55 | 495 | 683 | 468 | 640 | 232 | 269 | 0.7 | 0.7 | 0.9 |

* When diffuse dye material was present in a cell, its equivalent in dye-granules was estimated and added to the number of granules present. This was accomplished by assigning the diffuse staining to one of 5 intensities that ranged from 75 to 375 granules. Succinic dehydrogenase of AM and MN and acid phosphatase of MN rarely were manifest by diffuse staining, whereas the other enzymes commonly were. This explains why many of the granule counts of the latter group appear high.

16 and 17 show representative results from rabbits whose AM had distinctly higher enzymatic activity. With the exception of succinic dehydrogenase, all the enzymes studied in AM were represented by dye material that was partly diffuse and therefore were only amenable to semiquantitative estimation.

COMPARISON OF THE ENZYMATICALLY MOST ACTIVE CELLS IN AM AND MN PREPARATIONS: A group of AM or MN may be characterized by its most active members, for these

are the cells that ingest the foreign particles in the alveolar air or the blood stream, as the case may be, not the sluggish cells. For a comparison of the enzymatically most active AM and MN, studies were made of the 5, 10, 15, and 20 per cent of the cell population with the highest enzyme content. The results at the top 10 and 20 per cent levels typify the others and are included in Table III. From this, it is apparent that most of the AM:MN ratios were not appreciably altered at higher enzyme levels.

TABLE IV
Correlations between the Number of Dust* Particles in AM and Their Enzymatic Activities

| Rabbit No. | Enzyme | No. of cells evaluated | Type of cell evaluated | Average No. of histochemical dye granules per cell | Average No. of dust* particles per cell |
|------------|-------------------|------------------------|------------------------|--|---|
| 14 | Esterase | 66 | Top 10 per cent‡ | 240 | 33 |
| | | | Top 20 per cent | 220 | 34 |
| | | | Bottom 80 per cent | 75 | 13 |
| 15 | Acid phosphatase | 55 | Top 10 per cent | 78 | 30 |
| | | | Top 20 per cent | 65 | 12 |
| | | | Bottom 80 per cent | 15 | 6 |
| 16 | Acid phosphatase§ | 100 | Top 20 per cent | 48 | 0.0 |
| | | | Bottom 80 per cent | 21 | 0.5 |

* Some of these "dust" particles may be hemosiderin granules¹⁴ and other material.

‡ "Top 10 per cent" refers to the 10 per cent of the cells that contained the most dye granules.

§ In this experiment there seemed to be too few dust particles to make a reliable study.

TABLE V
Oxygen Consumption of PMN, MN, and AM

| Rabbit No. and differential cell count (in parentheses)* | | | No. of cells per Warburg flask (in millions) | | | μ l O ₂ consumed per hour per million cells | | | Ratio: MN:AM | |
|--|---------|---------|--|------|------|--|-----|-----|--------------|------|
| PMN | MN | AM | PMN | MN | AM | PMN | MN | AM | | |
| 1 (91) | 6 (90) | 6 (100) | 34.0 | 6.0 | 6.0 | 0.47 | 2.2 | 3.2 | 1:1.5 | |
| 2 (85) | 7 (96) | 7 (99) | 21.3 | 7.2 | 7.2 | 0.56 | 2.2 | 2.2 | 1:1.0 | |
| 3 (88) | 8 (99) | 8 (100) | 23.6 | 18.0 | 18.0 | 0.25 | 1.8 | 2.4 | 1:1.3 | |
| 2 (85) | 9 (85) | 9 (100) | 213. | 8.7 | 9.8 | 0.32 | 1.7 | 4.3 | 1:2.5 | |
| 3 (88) | 10 (93) | 10 (99) | 236. | 3.8 | 3.8 | 0.21 | 7.4 | 6.8 | 1:0.92 | |
| 4 (81) | | | 180. | | | 0.29 | | | | |
| 5 (91) | | | 200. | | | 0.29 | | | | |
| Average | | | | | | 0.34 | 3.1 | 3.8 | 1:1.4 | |
| Median | | | | | | 0.29 | 2.2 | 3.2 | 1:1.3 | |
| Ratio: Average PMN:MN:AM | | | | | | 1 | : | 9.1 | : | 11.2 |
| Median PMN:MN:AM | | | | | | 1 | : | 7.6 | : | 11.0 |

* For example, the cell preparations listed on the top line contained, respectively, 91 per cent PMN, 90 per cent MN, and 100 per cent AM. The values listed for oxygen consumption represent the cell preparation as a whole with no adjustment for its differential cell count.

Throughout the course of these studies, it was observed that AM with the highest enzyme activities frequently contained the most "dust" particles (Fig. 15). In the two experiments where these factors were adequately quantitated, this correlation was indeed found to be the case (Table IV). It is of interest that AM containing dust particles

are among the first cells to ingest inhaled tubercle bacilli (32).

OXYGEN CONSUMPTION OF AM AND MN

Not only do AM tend to have higher levels of certain enzymes than MN, but they also tend to

TABLE VI
*A Histochemical Comparison of Various Enzyme Activities of PMN with Those of MN
in the Same Glycogen Exudate*

| Enzyme | Rabbit No. | No. of MN studied | No. of PMN studied | Average No. of histochemical granules per cell | | Ratio: MN/PMN |
|------------------------|------------|-------------------|--------------------|--|----------|---------------|
| | | | | MN | PMN | |
| Cytochrome oxidase | 1 | 6 | 33 | 54 | 33 | 1.6 |
| | 2 | 25 | 150 | 43 | 43 | 1.0 |
| | 3 | Numerous | Numerous | ++ | ++ | 1.0 |
| | 4 | Numerous | Numerous | +++ | ++ | 1.5 |
| Aminopeptidase | 1 | 26 | 105 | 73 | 89 | 0.8 |
| | 2 | 30 | 75 | 78 | 50 | 1.6 |
| | 5 | 35 | 138 | 60 | 104 | 0.6 |
| | 3 | Numerous | Numerous | ++ | ++ | 1.0 |
| | 4 | Numerous | Numerous | ++ | ++ | 1.0 |
| Succinic dehydrogenase | 1 | 20 | 84 | 95 + 100* | 32 | 6.1 |
| | 2 | 51 | 255 | 94 + 100* | 93 | 2.1 |
| | 4 | 50 | 50 | 47 | 13 | 3.6 |
| | 6 | 20 | 50 | 95 + 100* | 28 | 7.0 |
| | 7 | Numerous | Numerous | ++++ | + | 4.0 |
| | 8 | Numerous | Numerous | +++++ | + | 5.0 |
| Acid phosphatase | 8 | 19 | 50 | 47 | 6 | 7.8 |
| | 9 | 20 | 50 | 56 | 15 | 3.7 |
| | 10 | 20 | 50 | 61 | 20 | 3.1 |
| | 11 | 30 | 70 | 80 | 21 | 3.8 |
| | 12 | 35 | 72 | 125 | 31 | 4.0 |
| Alkaline phosphatase | 1 | 19 | 84 | 0 | 50 | 0.0 |
| | 2 | 30 | 187 | 4‡ | 46 | 0.1 |
| | 6 | 52 | 53 | 9‡ | 45 | 0.2 |
| | 8 | 25 | 60 | 8‡ | 32 | 0.3 |
| | 10 | 25 | 50 | 7‡ | 36 | 0.2 |
| | 11 | 25 | 50 | 6‡ | 36 | 0.2 |
| Esterase | 1 | 15 | 24 | 37 | 36 | 1.0 |
| | 8 | 25 | 45 | 11 | 12 | 0.9 |
| | 11 | 25 | 50 | 19 + 32* | 15 + 38* | 1.0 |
| | 12 | 30 | 40 | 12 + 18* | 13 + 26* | 0.8 |
| | 13 | 17 | 110 | 52 | 32 | 1.6 |

* Estimate in granules of the amount of diffuse dye material present in the cells. Such diffuse color was rare with succinic dehydrogenase. Its presence in three of the samples listed may have been due to the use of glycogen to produce these exudates.

‡ These values probably represent MN *acid* phosphatase acting suboptimally at this alkaline pH.

have higher rates of oxygen consumption (Table V). The respiration of MN, expressed in microliters of oxygen per hour per million cells, averaged 2 to 3, and that of AM, 3 to 4. Comparable figures of Myrvik (33) for rabbits were approximately 1.5 and 4 μ l, respectively; and comparable figures

from Karnovsky's laboratory (34, 35) for guinea pigs were 1.4 and 8.2 μ l, respectively.

An additional indication of greater AM activity was their more frequent cell division (*cf.* Lurie, 36), which we observed in Wright's stained preparations, but did not quantitate (Fig. 18).

HISTOCHEMICAL STUDIES ON MN AND PMN

Since MN and PMN from different rabbits in response to different irritants are not strictly comparable (*cf.* Elberg, 37), a histochemical comparison was made of these cells when they were simultaneously present in the same 18-hour glycogen exudate. In this type of exudate, MN and PMN show high viabilities both by the dye tests (see Materials and Methods) and by their ability to phagocytize particles (3).

The results are presented in Table VI and Figs. 1, 4, 7, 10, and 11. From these it is apparent that MN had greater acid phosphatase and succinic dehydrogenase activities, and PMN greater alkaline phosphatase activity (MN showed none). Their cytochrome oxidase, aminopeptidase, and esterase activities were of similar magnitude.

OXYGEN CONSUMPTION OF MN AND PMN

Table V indicates that rabbit MN exudates have a higher oxygen consumption than PMN. PMN consumed about 0.3 μ l of oxygen per hour per million cells and MN consumed from 2 to 3 μ l. Comparable figures of Stähelin, Suter, and Karnovsky (8, 34) for PMN and MN exudates of guinea pigs were 0.3 and 1.0 μ l, respectively; and comparable figures of Frei *et al.* (38) for PMN and MN of human blood were 0.15 and 0.5 μ l, respectively.

Exudate MN are slightly larger than PMN, but not enough to account for their greater enzymatic activities and oxygen consumption. Their motility as observed by time-lapse cinematography (39) was also greater. These three parameters indicate that MN are metabolically and functionally more active cells.

DISCUSSION

This paper reports our attempts to quantitate by the granule count technique several histochemically demonstrable enzymes in rabbit mononuclear and polymorphonuclear exudate cells (MN and PMN) and pulmonary alveolar macrophages (AM). We were successful with acid phosphatase in MN and partially successful with succinic dehydrogenase in all three types of phagocyte. With the others (cytochrome oxidase, aminopeptidase, and esterase) the method became semi-quantitative in nature, because part of the color

produced was diffuse and not amenable to granule counting.

The drying of the cells on the Mylar strips evidently caused rupture of such structures as lysosomes and mitochondria whose enzymes became more freely distributed throughout the cytoplasm. This redistribution enabled the quantitative procedures herein described, for the sites of enzymatic activities as evidenced by the dye granules became dependent on the enzyme concentration rather than on the number of organelles (*cf.* Doyle, 40). The granule count method therefore may not be applicable to cells fixed in formol calcium or other agents that seem to maintain organelle structure.

Other methods have been introduced for the histochemical quantitation of enzymes in individual cells. Benditt and Arase (41) studied the time, substrate, and inhibitor relationships of a chymotrypsin-like enzyme in mast cells by matching color intensities. Similar methods were employed by Hopsu and Glenner (42) to evaluate the trypsin-like enzymes of mast cells, and by Kaplow (43) and Quigley *et al.* (44) to evaluate the alkaline phosphatase activity of PMN in blood smears. The use of particle sizing and particle counting was described by Eränkö in his text on quantitative histochemical methods (45), but their application to enzymatic reactions was not discussed.

Survey of Literature

There have been few if any previous histochemical studies on the enzymes of rather pure populations of phagocytes, but there have been many such studies on these phagocytes in blood and bone marrow films and in tissue sections.

Baker and Klapper (46) demonstrated cytochrome oxidase in foreign body giant cells (which are closely related to MN). Hannibal *et al.* (17) found strongly positive reactions in human MN and PMN, which they believed were due to *both* the G-Nadi reaction for cytochrome oxidase and the M-Nadi reaction for peroxidase. Positive reactions in blood and bone marrow cells have also been described by others (47-49).

The aminopeptidase of human blood leukocytes has been evaluated by Ackerman (50), and identified in tissue MN (19, 51) and PMN (19).

Dehydrogenase activity specific for succinate has been described in PMN, MN, and lymphocytes of human blood and bone marrow (52-56). Ackerman (54) was able to show mitochondrial

localization of succinic dehydrogenase in viable blood cells by exposing them to a non-ionic, surface-active agent. Balogh and Cohen (56) studied this enzyme in *unfixed* human blood leukocytes by means of phase microscopy.

Acid phosphatase has been demonstrated by histochemical methods in the MN of man (57), rabbits, mice, and rats (58), and chickens (59), but has not been found in the MN of guinea pigs (58). This enzyme is found in the lysosomes, phagosomes, and pinocytotic vacuoles of cells (60-66).

Among the first histochemical reports on the alkaline phosphatase of PMN are those of Gomori (67), Wachstein (68), and Wislocki and Dempsey (69). Many other investigators have since confirmed the presence of alkaline phosphatase in the PMN of blood (see references 70 and 71). It is of interest that Grogg and Pearse (58) demonstrated alkaline phosphatase in PMN of rats, guinea pigs, and rabbits, but could not detect it in PMN of mice. It was absent in the MN of most species. A rather complete biochemical characterization of purified alkaline phosphatase from human leukocytes was made by Trubowitz *et al.* (72). This enzyme is low in chronic myeloid leukemia, and rather high in leukemoid states (73-76), a finding that may be of considerable diagnostic importance.

There are many reports that describe esterases in tissue phagocytes. Among these are the following: for PMN, (references 58, 77-81), for MN (58, 77, 80-91), and for AM (82, 87, 89, 90, 92). With few exceptions, the inhibitor studies reported in the literature (77, 82-88, 92) agree with our results. AM and MN appear to contain both the inhibitor-sensitive and inhibitor-resistant varieties of esterase described by Shnitka and Seligman (93). Increased MN esterase activity was found in the lesions of cutaneous tuberculosis (83, 85), BCG infection (91), leprosy (85, 91), and leishmaniasis (94). In infectious leukocytosis circulating PMN likewise showed increased esterase activity (80).

Our findings that AM contained higher concentrations of enzymes than MN are in agreement with other investigators. Myrvik *et al.* (95, 96) found that AM had 2 to 5 times more lysozyme, β -glucuronidase, and acid phosphatase. Cohn and Wiener (65) have confirmed these results and added an acid-acting proteinase and ribonuclease to the list. Using histochemical methods Nachlas and Seligman (89) and Chessick (90) observed

higher esterase activity in AM than in Kupffer cells.

With respect to our observations on the enzymes of PMN and MN, there are both conflicting and confirmatory reports. The biochemical studies of Frei *et al.* (38) showed that human blood MN had over twice the esterase activity of PMN, whereas the histochemical studies of Ackerman (81) showed the reverse. In glycogen exudates from rabbits we found that MN and PMN contained equal amounts of esterase. Our studies on their phosphatases are in agreement with others (58, 67, 68). The peak we observed in the acid range of the pH curve for PMN phosphatase seems to be of special significance, because such a peak determined biochemically (97) might have arisen from contaminating MN.

In Vivo Activation of PMN

Ludwik Fleck has termed this phenomenon "leukergy" and found that circulating PMN were more active following certain stress situations, *i.e.* infections, trauma, and endotoxin injection (98, 99). With such PMN, the cells' "stickiness," motility, phagocytic ability, and activity in destroying staphylococci were increased, as were their contents of alkaline phosphatase and glycogen (100-106). Other investigators have undoubtedly studied similar phenomena by evaluating under various conditions PMN alkaline phosphatase (43, 68, 73, 107-110), "lipase" (esterase) (111), peroxidase (112), and glycogen contents (112). A stress-induced adrenal response may be at least partially responsible for PMN activation, for ACTH and 17-hydroxy-corticosteroids increased PMN alkaline phosphatase levels in man (109, 110).

It seems from the references just listed that PMN circulate in two states of activation: (a) a relatively inactive PMN with low alkaline phosphatase content, and (b) an activated, probably more mature (113) PMN with high alkaline phosphatase content. Evidently conditions of stress increase the proportion of activated PMN in the blood. Circulating PMN are only a small fraction of those stored in the bone marrow, spleen, and lung. A study of these reserves may be necessary to elucidate the mechanisms responsible for PMN activation (*cf.* reference 114).

In Vivo Activation of MN

Since the original work of Lurie (36), there have been many investigations of this nature

(115-120), for MN are the functional element of the reticuloendothelial system. Among the stimuli employed were live, attenuated, or dead bacteria, endotoxin, drum trauma, and thyroid hormone. Among the indices of MN activation were phagocytic ability and resistance to shock-producing and infectious agents.

In spite of the extensive work on activation of the reticuloendothelial system, there have been relatively few reports on accompanying changes in the enzyme content of MN. Lurie (36) has demonstrated that MN from tuberculous animals are more active phagocytically than controls. Such MN also have higher dehydrogenase (121) and acid phosphatase activities (58, 121-123). Connective tissue histiocytes of mice that were most phagocytic for several substances *in vivo* were highest in acid phosphatase, aminopeptidase, adenosine triphosphatase, and β -glucuronidase (124). Connective tissue MN of man that contained hemosiderin were highest in esterase activity (88), and those of guinea pigs that probably ingested particles of adjuvant were highest in lipase activity (125).

Most of the other studies concerned acid phosphatase whose activity was increased (*a*) in Kupffer cells most active in the phagocytosis of particles from the blood stream (63, 123, 126, 127), (*b*) in MN in contact with various foreign bodies placed subcutaneously in mice and rats (128), (*c*) in the reticular cells of the rabbit appendix that had ingested the cell debris caused by radiation damage (129), (*d*) in reticuloendothelial cells of rat thymus

and lymph nodes following the intravenous injection of killed typhoid bacteria (63) and in those of the liver following the ip injection of *Salmonella typhimurium* (130), (*e*) in mouse macrophages that had ingested red corpuscles in the peritoneal cavity (62), and (*f*) in alveolar phagocytes containing the most dust particles (this report). In addition, both the acid phosphatase and phagocytic activities of Kupffer cells can be increased by endotoxins (123, 126, 127).

The following paper (3) will present studies on *in vitro* activation of phagocytes and suggest a theory that seems to unify current knowledge concerning their enzymatic and phagocytic functions.

We are indebted to Dr. Max B. Lurie and Dr. Stuart Mudd for reviewing this manuscript, and to Lawrie Winning, Peter Zappasodi, and John McGuire for the photomicrographs.

This investigation began during the month spent by the senior author with Dr. M. S. Burstone at the National Institute of Dental Research, Bethesda. It was supported by contract Nonr-551(24) with the Office of Naval Research and grant AI-02048 from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, United States Public Health Service.

The results were presented in part at the annual meeting of the Histochemical Society on April 9, 1960 in New York City (1), and at the annual meeting of the American Association of Immunologists on April 12, 1961 in Atlantic City, New Jersey (2).

Received for publication, July 20, 1962.

BIBLIOGRAPHY

1. DANNENBERG, A. M., JR., BURSTONE, M. S., and WILDHACK, J. A., Phagocytic and enzymatic activities of rabbit mononuclear and polymorphonuclear exudate cells: a histochemical study, *J. Histochem. and Cytochem.*, 1960, **8**, 318, abstract.
2. DANNENBERG, A. M., JR., and WALTER, P. C., A histochemical study of phagocytic and enzymatic functions of rabbit mononuclear and polymorphonuclear exudate cells and alveolar macrophages, *Fed. Proc.*, 1961, **20**, 264.
3. DANNENBERG, A. M., JR., WALTER, P. C., and KAPRAL, F. A., A histochemical study of phagocytic and enzymatic functions of rabbit mononuclear and polymorphonuclear exudate cells and alveolar macrophages. II. The effect of particle ingestion on enzyme activity; two phases of *in vitro* activation, *J. Immunol.*, 1963, **90**, 448.
4. HANKS, J. H., and WALLACE, R. E., Relation of oxygen and temperature in the preservation of tissues by refrigeration, *Proc. Soc. Exp. Biol. and Med.*, 1949, **71**, 196.
5. HIRSCH, J. G., Phagocytin: a bactericidal substance from polymorphonuclear leucocytes, *J. Exp. Med.*, 1956, **103**, 589.
6. MYRVIK, Q. N., SOTO LEAKE, E., and FARISS, B., Studies on pulmonary alveolar macrophages from the normal rabbit: A technique to procure them in a high state of purity, *J. Immunol.*, 1961, **86**, 128.
7. ROUS, P., and JONES, F. S., The protection of pathogenic microorganisms by living tissue cells, *J. Exp. Med.*, 1916, **23**, 601.
8. STÄHELIN, H., SUTER, E., and KARNOVSKY,

- M. L., Studies on the interaction between phagocytes and tubercle bacilli. I. Observations on the metabolism of guinea pig leucocytes and the influence of phagocytosis, *J. Exp. Med.*, 1956, **104**, 121.
9. SCHREK, R., A method for counting the viable cells in normal and malignant cell suspensions, *Am. J. Cancer*, 1936, **28**, 389.
 10. HANKS, J. H., and WALLACE, J. H., Determination of cell viability, *Proc. Soc. Exp. Biol. and Med.*, 1958, **98**, 188.
 11. TULLIS, J. L., Preservation of leucocytes, *Blood*, 1953, **8**, 563.
 12. BURSTONE, M. S., and FLEMING, T. J., A new technique for the histochemical study of smears, *J. Histochem. and Cytochem.*, 1959, **7**, 203.
 13. JOHNSTON, M. E., Mylar film as a histologic tissue support, *J. Histochem. and Cytochem.*, 1960, **8**, 139.
 14. UMBREIT, W. W., BURRIS, R. H., and STAUFFER, J. F., *Manometric Techniques*, Minneapolis, Burgess Publishing Co., 3rd edition, 1957.
 15. BURSTONE, M. S., New histochemical techniques for the demonstration of tissue oxidase (cytochrome oxidase), *J. Histochem. and Cytochem.*, 1959, **7**, 112.
 16. BURSTONE, M. S., Histochemical demonstration of cytochrome oxidase with new amine reagents, *J. Histochem. and Cytochem.*, 1960, **8**, 63.
 17. HANNIBAL, M. J., NACHLAS, M. M., and SELIGMAN, A. M., Histochemical localization of cytochrome oxidase in human tissues and neoplasms, *Cancer*, 1960, **13**, 1008.
 18. AGNER, K., Verdoperoxidase. A ferment isolated from leucocytes, *Acta Physiol. Scand.*, 1941, **2**, suppl. 8, 1.
 19. BURSTONE, M. S., and FOLK, J. E., Histochemical demonstration of aminopeptidase, *J. Histochem. and Cytochem.*, 1956, **4**, 217.
 20. GOMORI, G., Chromogenic substrates for aminopeptidase, *Proc. Soc. Exp. Biol. and Med.*, 1954, **87**, 559.
 21. LILLIE, R. D., Preferred common names, formulae, Colour Index references and synonyms of stable diazonium salts used in histochemistry, *J. Histochem. and Cytochem.*, 1959, **7**, 281.
 22. NACHLAS, M. M., TSOU, K. C., DE SOUZA, E., CHENG, C. S., and SELIGMAN, A. M., Cytochemical demonstration of succinic dehydrogenase by the use of a new p-nitrophenyl substituted ditetrazole, *J. Histochem. and Cytochem.*, 1957, **5**, 420.
 23. WACHSTEIN, M., and MEISEL, E., Influence of experimental renal damage on histochemically demonstrable succinic dehydrogenase activity in the rat, *Am. J. Path.*, 1954, **30**, 147.
 24. WACHSTEIN, M., and MEISEL, E., A comparative study of enzymatic staining reactions in the rat kidney with necrobiosis induced by ischemia and nephrotoxic agents (mercurhydrin and DL-serine), *J. Histochem. and Cytochem.*, 1957, **5**, 204.
 25. BURSTONE, M. S., Histochemical comparison of naphthol AS-phosphates for the demonstration of phosphatases, *J. Nat. Cancer Inst.*, 1958, **20**, 601.
 26. BURSTONE, M. S., Histochemical demonstration of acid phosphatases with naphthol AS-phosphates, *J. Nat. Cancer Inst.*, 1958, **21**, 523.
 27. FOLLETTE, J. H., VALENTINE, W. N., and REYNOLDS, J., A comparison of human leukocyte phosphatase activity toward sodium β -glycerophosphate, adenosine 5'-phosphate and glucose 1-phosphate, *Blood*, 1959, **14**, 415.
 28. BURSTONE, M. S., The cytochemical localization of esterase, *J. Nat. Cancer Inst.*, 1957, **18**, 167.
 29. BURSTONE, M. S., Esterase activity of developing bones and teeth, *Arch. Path.*, 1957, **63**, 164.
 30. FLOREY, H., *Lectures on General Pathology*, Philadelphia, W. B. Saunders Company, 1954, 102.
 31. KARRER, H. E., The ultrastructure of mouse lung: the alveolar macrophage, *J. Biophysic. and Biochem. Cytol.*, 1958, **4**, 693.
 32. LURIE, M. B., unpublished observations.
 33. MYRVIK, Q., unpublished observations.
 34. KARNOVSKY, M. L., Metabolic basis of phagocytic activity, *Physiol. Rev.*, 1962, **42**, 143.
 35. OREN, R., FARNHAM, A. E., SAITO, K., MILOFSKY, E., and KARNOVSKY, M. L., Metabolic patterns in three types of phagocytizing cells, *J. Cell Biol.*, **17**, 487.
 36. LURIE, M. B., Studies on the mechanism of immunity in tuberculosis. The mobilization of mononuclear phagocytes in normal and immunized animals and their relative capacities for division and phagocytosis, *J. Exp. Med.*, 1939, **69**, 579.
 37. ELBERG, S. S., MASCARENHAS, P., and FONG, J., Response of peritoneal exudate cells to *Brucella melitensis*. Influence of nature of inflammatory irritant, *Proc. Soc. Exp. Biol. and Med.*, 1960, **104**, 295.
 38. FREI, J., BOREL, C., HORVATH, G., CULLITY, B., and VANNOTTI, A., Enzymatic studies in

- the different types of normal and leukemic human white cells, *Blood*, 1961, 18, 317.
39. DOUGHERTY, T. F., and SCHNEEBELI, G. L., Inflammation, A motion picture, produced at the University of Utah, that will eventually be available through the Library of Congress.
 40. DOYLE, W. L., Discussion in Eleventh Annual Meeting of the Histochemical Society, *J. Histochem. and Cytochem.*, 1960, 8, 319.
 41. BENDITT, E. P., and ARASE, M., Enzyme kinetics in a histochemical system, *J. Histochem. and Cytochem.*, 1958, 6, 431.
 42. HOPUSU, V. K. and GLENNER, G. G., A histochemical study of the enzyme kinetics of trypsin-like amidase and esterase activities in human mast cells, *J. Histochem. and Cytochem.*, 1962, 10, 674, abstract.
 43. KAPLOW, L. S., A histochemical procedure for localizing and evaluating leucocyte alkaline phosphatase activity in smears of blood and marrow, *Blood*, 1955, 10, 1023.
 44. QUIGLEY, H. J., DAWSON, E. A., HYUN, B. H., and CUSTER, R. P., The activity of alkaline phosphatase in granular leukocytes during pregnancy and the puerperium: a preliminary report, *Am. J. Clin. Path.*, 1960, 33, 109.
 45. ERÄNKÖ, O., Quantitative Methods in Histology and Microscopic Histochemistry, Boston, Little, Brown and Company, 1955, 74-86.
 46. BAKER, B. L., and KLAPPER, Z. F., Oxidative enzymes in the foreign body giant cell, *J. Histochem. and Cytochem.*, 1961, 9, 713.
 47. HOFFMANN, G. T., ROTTINO, A., and STERN, K. G., Demonstration by the Nadi reaction of cytochrome oxidase activity in cells of the lymphoid and myeloid series obtained from normal individuals and patients suffering from Hodgkin's and other diseases, *Blood*, 1951, 6, 1051.
 48. RYERSON, D. L., Does the M-nadi test locate cytochrome oxidase in leukocytes?, *Anat. Rec.*, 1954, 118, 348.
 49. REBUCK, J. W., Enhanced Nadi reaction in lymphocytes in inflammation, *Fed. Proc.*, 1952, 11, 424.
 50. ACKERMAN, G. A., Histochemical demonstration of aminopeptidase activity in the leukocytes of blood and bone marrow, *J. Histochem. and Cytochem.*, 1960, 8, 386.
 51. BRAUNSTEIN, H., FREIMAN, D. G., THOMAS, W., JR., and GALL, E. A., A histochemical study of the enzymatic activity of lymph nodes. II. Further investigation of normal and hyperplastic lymph nodes, *Cancer* 1962, 15, 130.
 52. DE SOUZA, E. J., and KOTHARE, S. N., A method for the cytochemical demonstration of succinic dehydrogenase in human leucocytes, *J. Histochem. and Cytochem.*, 1959, 7, 77.
 53. MORRISON, J. H., and KRONHEIM, S., The cytochemical demonstration of succinic dehydrogenase in mouse leukocytes, *J. Histochem. and Cytochem.*, 1962, 10, 402.
 54. ACKERMAN, G. A., Histochemical demonstration of dehydrogenase activity in the cells of normal human blood and bone marrow, *J. Biophysic. and Biochem. Cytol.*, 1960, 8, 61.
 55. QUAGLINO, D., and HAYHOE, F. G. J., Acetone fixation for the cytochemical demonstration of dehydrogenases in blood and bone marrow cells, *Nature*, 1960, 187, 85.
 56. BALOGH, K., JR., and COHEN, R. B., Histochemical demonstration of diaphorases and dehydrogenases in normal human leukocytes and platelets, *Blood*, 1961, 17, 491.
 57. GOMORI, G., Distribution of acid phosphatase in the tissues under normal and under pathologic conditions, *Arch. Path.*, 1941, 32, 189.
 58. GROGG, E., and PEARSE, A. G. E., The enzymic and lipid histochemistry of experimental tuberculosis, *Brit. J. Exp. Path.*, 1952, 33, 567.
 59. WEISS, L. P., and FAWCETT, D. W., Cytochemical observations on chicken monocytes, macrophages and giant cells in tissue culture, *J. Histochem. and Cytochem.*, 1953, 1, 47.
 60. DE DUVE, C., Lysosomes, a new group of cytoplasmic particles, in *Subcellular Particles* (T. Hayashi, editor), New York, The Ronald Press Co., 1959, 128-159.
 61. NOVIKOFF, A. B., and ESSNER, E., The liver cell. Some new approaches to its study, *Am. J. Med.*, 1960, 29, 102.
 62. ESSNER, E., An electron microscopic study of erythrophagocytosis, *J. Biophysic. and Biochem. Cytol.*, 1960, 7, 329.
 63. BARKA, T., SCHAFFNER, F., and POPPER, H., Acid phosphatase and the reticuloendothelial system, *Lab. Inv.*, 1961, 10, 590.
 64. COHN, Z. A., and HIRSCH, J. G., The isolation and properties of the specific cytoplasmic granules of rabbit polymorphonuclear leukocytes, *J. Exp. Med.*, 1960, 112, 983.
 65. COHN, Z. A., and WIENER, E., personal communication on the lysosomes and hydrolases of alveolar and peritoneal macrophages.
 66. BIRNS, M., The localization of acid phosphatase activity in the ameba, *Chaos chaos, Exp. Cell Research*, 1960, 20, 202.
 67. GOMORI, G., The distribution of phosphatase in normal organs and tissues, *J. Cell. and Comp. Physiol.*, 1941, 17, 71.
 68. WACHSTEIN, M., Alkaline phosphatase activity in normal and abnormal human blood and

- bone marrow cells, *J. Lab. and Clin. Med.*, 1946, **31**, 1.
69. WISLOCKI, G. B., and DEMPSEY, E. W., Observations on the chemical cytology of normal blood and hemopoietic tissues, *Anat. Rec.*, 1946, **96**, 249.
 70. PEARSE, A. G. E., *Histochemistry*, Boston, Little, Brown and Company, 2nd edition, 1960, 401.
 71. ACKERMAN, G. A., *Cytochemistry of the lymphocytes*, in *The Lymphocyte and Lymphocytic Tissue*, (J. W. Rebeck, editor), New York, Paul B. Hoeber, Inc., 1960, 28.
 72. TRUBOWITZ, S., FELDMAN, D., MORGENSTERN, S. W., and HUNT, V. M., The isolation, purification and some properties of the alkaline phosphatase of human leucocytes, *Biochem. J.*, 1961, **80**, 369.
 73. VALENTINE, W. N., and BECK, W. S., Biochemical studies on leukocytes. I. Phosphatase activity in health, leucocytosis and myelocytic leukemia, *J. Lab. and Clin. Med.*, 1951, **38**, 39.
 74. MOLONEY, W. C., and LANGE, R. D., Cytologic and biochemical studies on the granulocytes in early leukemia among atom bomb survivors, *Texas Rept. Biol. and Med.*, 1954, **12**, 887.
 75. LEONARD, B. J., Alkaline phosphatase in the white cells in leukaemia and leukaemoid reactions, *Lancet*, 1958, **1**, 289.
 76. KOLER, R. D., SEAMAN, A. J., OSGOOD, E. E., VAN BELLINGHEN, P., Myeloproliferative diseases. Diagnostic value of the leukocyte alkaline phosphatase test, *Am. J. Clin. Path.*, 1958, **30**, 295.
 77. MOLONEY, W. C., MCPHERSON, K., and FLIEGELMAN, L., Esterase activity in leukocytes demonstrated by the use of naphthol AS-D chloroacetate substrate, *J. Histochem. and Cytochem.*, 1960, **8**, 200.
 78. GOMORI, G., Chloracyl esters as histochemical substrates, *J. Histochem. and Cytochem.*, 1953, **1**, 469.
 79. BARNETT, R. J., and SELIGMAN, A. M., Histochemical demonstration of esterase by production of indigo, *Science*, 1951, **114**, 579.
 80. WACHSTEIN, M., and WOLF, G., The histochemical demonstration of esterase activity in human blood and bone marrow smears, *J. Histochem. and Cytochem.*, 1958, **6**, 457.
 81. ACKERMAN, G. A., Histochemical demonstration of indoxyl acetate esterase activity in normal human blood and bone marrow, *Lab. Inv.*, 1960, **9**, 298.
 82. GOMORI, G., and CHESICK, R. D., Histochemical studies of inhibition of esterases, *J. Cell. and Comp. Physiol.*, 1953, **41**, suppl. 1, 51.
 83. FINDLAY, G. H., The simple esterases of human skin, *Brit. J. Dermat.*, 1955, **67**, 83.
 84. PEARSON, B., and DEFENDI, V., A comparison between the histochemical demonstration of non-specific esterase activity by 5-bromo-indoxyl acetate, α -naphthyl acetate and naphthol AS acetate, *J. Histochem. and Cytochem.*, 1957, **5**, 72.
 85. WELLS, G. C., Esterases in cutaneous granulomata, *Brit. J. Dermat.*, 1957, **69**, 415.
 86. BRAUNSTEIN, H., FREEMAN, D. G., and GALL, E. A., A histochemical study of the enzymatic activity of lymph nodes, *Cancer*, 1958, **11**, 829.
 87. WACHSTEIN, M., MEISEL, E., and FALCON, C., Histochemistry of thiolacetic acid esterase: A comparison with nonspecific esterase with special regard to the effect of fixatives and inhibitors on intracellular localization, *J. Histochem. and Cytochem.*, 1961, **9**, 325.
 88. MONIS, B., and WEINBERG, T., Cytochemical study of esterase activity of human neoplasms and stromal macrophages, *Cancer*, 1961, **14**, 369.
 89. NACHLAS, M. N., and SELIGMAN, A. M., The comparative distribution of esterase in the tissues of five mammals by a histochemical technique, *Anat. Rec.*, 1949, **105**, 677.
 90. CHESICK, R. D., Histochemical study of the distribution of esterases, *J. Histochem. and Cytochem.*, 1953, **1**, 471.
 91. STEIGLEDER, G. K., and SCHULTIS, K., Die Bedeutung des Nachweises unspezifischer Esterasen in Bindegewebszellen der Haut, *Arch. Klin. u. exp. Dermat.*, 1957, **204**, 448.
 92. MALATY, H. A., and BOURNE, G. H., The histochemistry of simple esterase, *Acta Anat.*, 1955, **24**, 337.
 93. SHNITKA, T. K., and SELIGMAN, A. M., Role of esterase inhibition on localization of esterase and the simultaneous cytochemical demonstration of inhibitor sensitive and resistant enzyme species, *J. Histochem. and Cytochem.*, 1961, **9**, 504.
 94. WELLS, G. C., Hydrolysing enzymes in macrophages, *J. Inv. Dermat.*, 1958, **31**, 83.
 95. MYRVIK, Q. N., SOTO LEAKE, E., and FARISS, B., Lysozyme content of alveolar and peritoneal macrophages from the rabbit, *J. Immunol.*, 1961, **86**, 133.
 96. MYRVIK, Q. N., SOTO LEAKE, E., and GONZALEZ-OJEDA, D., Enzymatic differences between normal rabbit alveolar macrophages and oil-induced peritoneal macrophages, *Bact. Proc.*, 1962, 79.

97. CRAM, D. M., and ROSSITER, R. J., Phosphatase of rabbit polymorphonuclear leucocytes, *Canad. J. Research*, 1949, **27**, 290.
98. FLECK, L., Recent investigations on leukergy, *Texas Rept. Biol. and Med.*, 1956, **14**, 424.
99. FLECK, L., La leukergie et son rôle dans l'immunité, *Ann. Inst. Pasteur*, 1957, **92**, 380.
100. FLECK, L., and MURCZYNSKA, Z., Leukergy, *Texas Rept. Biol. and Med.*, 1947, **5**, 156.
101. FLECK, L., Über Leukergie, *Acta haemat.*, 1952, **8**, 282.
102. FLECK, L., and SZCZYGIELSKA, J., The immunological significance of leukergy, I, *Ann. Univ. Mariae Curie-Skłodowska, Lublin-Polonia*, 1950, **5**, sect. D, 507.
103. FLECK, L., and SZCZYGIELSKA, J., The immunological significance of leukergy, II, *Ann. Univ. Mariae Curie-Skłodowska, Lublin-Polonia*, 1950, **5**, sect. D, 523.
104. FLECK, L., and MURCZYNSKA, Z., The phenomenon of leukergy, *Arch. Path.*, 1949, **47**, 261.
105. FLECK, L., and LILLE-SZYSZKOWICZ, I., Leukergy and the metabolism of leukocytes, International Society of Haematology, Proceedings of the Sixth International Congress (1956), New York, Grune and Stratton, Inc., 1958, 157.
106. FLECK, L., and LILLE-SZYSZKOWICZ, I., Leukergy and the glycogen content in leukocytes, *Bull. Acad. Polon. Sc., Cl. II.*, 1955, **3**, 137.
107. VACCARI, F., SABOTTO, B., and MANZINI, E., Alkaline phosphatase activity of leukocytes in shock, *Blood*, 1955, **10**, 730.
108. WILTSHAW, E., and MALONEY, W. C., Histochemical and biochemical studies on leukocyte alkaline phosphatase activity, *Blood*, 1955, **10**, 1120.
109. VALENTINE, W. N., FOLLETTE, J. H., HARDIN, E. B., BECK, W. S., and LAWRENCE, J. S., Studies on leukocyte alkaline phosphatase activity: relation to "stress" and pituitary-adrenal activity, *J. Lab. and Clin. Med.*, 1954, **44**, 219.
110. VALENTINE, W. N., FOLLETTE, J. H., SOLOMON, D. H., and REYNOLDS, J., The relationship of leukocyte alkaline phosphatase to "stress", to ACTH, and to adrenal 17-OH-corticosteroids, *J. Lab. and Clin. Med.*, 1957, **49**, 723.
111. ANTONIOLI, J. A., Metabolism of the white blood cell inside and outside the blood stream, in *Ciba Found. Study Group 10*, 1961, 92.
112. FREIDLIN, I. S., The change in phagocytic activity and in certain cytochemical reactions of the leukocytes of guinea-pig peritoneal exudate during immunization, *Bull. Exp. Biol. and Med. (USSR)*, 1961, **52**, 80, (translation: 1962, **52**, 1056).
113. TRUBOWITZ, S., FELDMAN, D., BENANTE, C., and HUNT, V. M., The alkaline phosphatase content of human leukocytes following marrow suppression by nitrogen mustard, *J. Lab. Clin. and Med.*, 1958, **52**, 259.
114. TRUBOWITZ, S., MOSCHIDES, E., and FELDMAN, D., Alkaline phosphatase activity of the polymorphonuclear leukocyte in rapidly induced leukopenia and leukocytosis, *J. Lab. and Clin. Med.*, 1961, **57**, 747.
115. HELLER, J. H., Consulting Editor, The reticulo-endothelial system (RES), *Ann. New York Acad. Sc.*, 1960, **88**, 1.
116. HALPERN, B. N., editor, Physiopathology of the Reticulo-Endothelial System: A Symposium, Springfield, Illinois, Charles C. Thomas, 1957.
117. HELLER, J. H., editor, Reticuloendothelial Structure and Function, New York, The Ronald Press Co., 1960.
118. Fort Detrick Symposium, Nonspecific resistance to infection, *Bact. Rev.*, 1960, **24**, 1.
119. SHILO, M., Nonspecific resistance to infections, *Ann. Rev. Microbiol.*, 1959, **13**, 255.
120. GÖZSY, B., and KÁTÓ, L., Studies on Phagocytic Stimulation, Montreal, Thérien Frères Limitée, 1957.
121. ALLISON, M. J., ZAPPASODI, P., and LURIE, M. B., Metabolic studies on mononuclear cells from rabbits of varying genetic resistance to tuberculosis. II. Studies on cells from BCG-vaccinated animals, *Am. Rev. Respirat. Dis.*, 1962, **85**, 364.
122. SUTER, E., and HULLIGER, L., Nonspecific and specific cellular reactions to infections, *Ann. New York Acad. Sc.*, 1960, **88**, 1237.
123. THORBECKE, G. J., OLD, L. J., BENACERRAF, B., and CLARKE, D. A., A histochemical study of acid and alkaline phosphatase in mouse livers during various conditions modifying activity of the reticuloendothelial system, *J. Histochem. and Cytochem.*, 1961, **9**, 392.
124. HOSODA, S., and NAKAMURA, K., A new method for histochemical observation of the subcutaneous tissue, *J. Histochem. and Cytochem.*, 1960, **8**, 72.
125. VOGEL, F. S., A lipolytic enzyme in reactive histiocytes of guinea pigs with experimental encephalomyelitis, *J. Exp. Med.*, 1951, **93**, 305.
126. HOWARD, J. G., ROWLEY, D., and WARDLAW, A. C., Investigations on the mechanism of stimulation of non-specific immunity by bacterial lipopolysaccharides, *Immunology*, 1958, **1**, 181.

127. HOWARD, J. G., Activation of the reticulo-endothelial cells of the mouse liver by bacterial lipopolysaccharide, *J. Path. and Bact.*, 1959, 78, 465.
128. GEDIGK, P., and BONTKE, E., Über die Enzymaktivität in Fremdkörpergranulationsgewebe, *Arch. Path. Anat. u. Physiol.*, 1957, 330, 538.
129. DOYLE, W. L., The distribution of phosphatase in the rabbit appendix after x-irradiation, *Am. J. Anat.*, 1950, 87, 79.
130. BÖHME, D., WACHSTEIN, M., and SCHNEIDER, H. A., Behavior of acid phosphatase in the reticulo-endothelial system of genetically susceptible and resistant mice infected with *S. typhimurium*, *Am. J. Path.*, 1961, 39, 103.