

# A late Miocene dust shower from the break-up of an asteroid in the main belt

Kenneth A. Farley<sup>1</sup>, David Vokrouhlický<sup>2</sup>, William F. Bottke<sup>3</sup> & David Nesvorný<sup>3</sup>

Throughout the history of the Solar System, Earth has been bombarded by interplanetary dust particles (IDPs), which are asteroid and comet fragments of diameter  $\sim 1\text{--}1,000\ \mu\text{m}$ . The IDP flux is believed to be in quasi-steady state: particles created by episodic main belt collisions or cometary fragmentation replace those removed by comminution, dynamical ejection, and planetary or solar impact. Because IDPs are rich in  $^3\text{He}$ , seafloor sediment  $^3\text{He}$  concentrations provide a unique means of probing the major events that have affected the IDP flux and its source bodies over geological timescales<sup>1–4</sup>. Here we report that collisional disruption of the  $>150\text{-km}$ -diameter asteroid that created the Veritas family  $8.3 \pm 0.5$  Myr ago<sup>5</sup> also produced a transient increase in the flux of interplanetary dust-derived  $^3\text{He}$ . The increase began at  $8.2 \pm 0.1$  Myr ago, reached a maximum of  $\sim 4$  times pre-event levels, and dissipated over  $\sim 1.5$  Myr. The terrestrial IDP accretion rate was overwhelmingly dominated by Veritas family fragments during the late Miocene. No other event of this magnitude over the past  $\sim 10^8$  yr has been deduced from main belt asteroid orbits. One remarkably similar event is present in the  $^3\text{He}$  record 35 Myr ago, but its origin by comet shower<sup>1</sup> or asteroid collision<sup>6</sup> remains uncertain.

After release from their comet or asteroid parent bodies, IDPs spiral towards the Sun under the effects of non-gravitational forces including Poynting–Robertson (P-R) and solar wind drags<sup>7</sup>. P-R drag occurs when dust grains revolving around the Sun absorb solar photons and then reradiate the energy in all directions. At the same time, implantation of solar wind ions enriches IDPs in  $^3\text{He}$ . If these particles avoid intense frictional heating during atmospheric entry, they can reach the Earth's surface with their  $^3\text{He}$  intact<sup>8</sup>. We have obtained  $^3\text{He}$  data on sediments spanning the past 70 Myr (refs 1, 9, 10), with new data in the interval 3–38 Myr ago reported here. We analysed two pelagic carbonate cores: ODP Site 757 in the Indian Ocean over this entire interval, and Site 926 in the Atlantic Ocean in the late Miocene.

$^3\text{He}$  measurements indicate the IDP flux is characterized by a somewhat bumpy continuum punctuated by sharp peaks at 8.2 and 35 Myr ago (Fig. 1). The older of these peaks, in the late Eocene, has been described from a different locality<sup>1</sup>. This peak is well above the average of the past 70 Myr and is coincident with the formation of the two largest terrestrial impact craters of the Cenozoic era: Popagai and Chesapeake Bay. The simultaneous increase in the dust and large body flux, and the match between the duration of the dust spike and that predicted for the ejection timescale of long period comets, were taken as evidence for a comet shower, perhaps produced by a close stellar encounter<sup>1</sup>. However the composition of impact melt at Popagai crater suggests an L-chondrite impactor<sup>6</sup>, implying that asteroids rather than comets may produce the spikes in IDP flux. Although several other episodes of elevated flux have been hinted at,


none have yet been confirmed. Possible connections between the late Miocene and Eocene events are further discussed in Supplementary Information.

To assess the distribution and temporal evolution of the late Miocene (8.2 Myr ago) peak, we studied the event at higher temporal resolution (Fig. 2) at Site 757 and also at Site 926. A  $^3\text{He}$  flux peak beginning 8.2 Myr ago and with nearly identical relative magnitude (factor of  $\sim 4$  above pre-event values) and duration ( $\sim 1.5$  Myr) is apparent at both sites. The only major distinction between the records is that the flux at Site 926 is about three times higher than at Site 757. This probably reflects the effects of sediment focusing, which is known to occur at Site 926 (ref 11). Given the similarity of the  $^3\text{He}$  peak at these two sites, and the fact that the peak does not correspond to dramatic changes in sediment composition or sedimentation rate (Supplementary Information), it seems unlikely that it is a sedimentation artefact. Furthermore, at both sites the flux peak corresponds to peaks in  $^3\text{He}$  concentration,  $^3\text{He}/^4\text{He}$  ratio and  $^3\text{He}/\text{non-carbonate fraction}$  (Supplementary Fig. S1). These observations indicate an increase in the IDP flux<sup>9</sup>. Thus we conclude that this  $^3\text{He}$  peak, like that in the late Eocene, is a global signature of an IDP-producing astronomical event.

Although the late Miocene and late Eocene  $^3\text{He}$  peaks are similar in duration and magnitude (Fig. 1), there is one important difference. Unlike the late Eocene with its two large impact craters that demand an increase in the large body flux coincident with the IDP spike, no late Miocene craters have yet been found. Apparently the late Miocene event was not accompanied by an asteroid or comet shower. This suggests the need for a mechanism capable of increasing the flux of IDPs striking Earth without affecting the flux of larger bodies. A likely candidate is the disruption of the diameter  $D > 150$  km asteroid that produced the Veritas family, a cluster of fragments on similar orbits at 3.17 AU. The Veritas event was the largest asteroid disruption in the past  $10^8$  yr (refs 5, 12); resulting collisions still produce as much as 10% of all Solar System near-ecliptic dust<sup>13–15</sup>. The age of the family,  $8.3 \pm 0.5$  Myr (ref. 5) was determined by tracking the orbits of Veritas family members backwards in time to their formation (for details, see Supplementary Fig. S4), and coincides with the onset of the late Miocene  $^3\text{He}$  spike.

When the parent body of Veritas disrupted, it ejected almost half of its mass in the form of fragments ranging from micrometre-sized dust grains to multi-kilometre asteroids<sup>5</sup>. These bodies then experienced dynamical evolution according to size. The evolution of small fragments ( $D = 1\text{--}1,000\ \mu\text{m}$ ) was dominated by planetary perturbations and non-gravitational forces, which caused them to spiral inwards towards the Sun. In contrast, larger fragments ( $D > 1,000\ \mu\text{m}$ ) were trapped in the main belt unless they could reach a chaotic resonance capable of placing them onto a planet-crossing orbit. The nearest resonances capable of producing an

<sup>1</sup>Division of Geological and Planetary Sciences, California Institute of Technology, MS 170-25, Pasadena, California 91125, USA. <sup>2</sup>Institute of Astronomy, Charles University, V Holešovickách 2, 180 00 Prague 8, Czech Republic. <sup>3</sup>Department of Space Studies, Southwest Research Institute, 1050 Walnut Street, Suite 400, Boulder, Colorado 80302, USA.



**Figure 1** | Two periods of elevated  $^3\text{He}$  flux, at  $\sim 35$  Myr ago (late Eocene) and  $\sim 8$  Myr ago (late Miocene), indicate intervals of enhanced accumulation rate of IDPs. Open and filled symbols are individual new  $^3\text{He}$  measurements from ODP Site 757 (central Indian Ocean,  $17^\circ 01.458' \text{ S}$ ,  $88^\circ 10.899' \text{ E}$ ). Lines are 3-point running means through the data points, taken to minimize the effects of occasional sampling of large individual

IDPs<sup>10</sup>. The grey segments of the running mean line indicate the late Miocene event (highlighted by open symbols), and the previously reported late Eocene peak from the Italian Apennines<sup>1</sup>. Cretaceous to mid-Tertiary running mean data are from ref. 10. Details of the new samples, analytical methods, data and age models are provided in Supplementary Information.

asteroid shower are  $\sim 0.1$  AU from the Veritas family<sup>16</sup> (for example, the 9:4, 11:5 or 2:1 mean motion resonances with Jupiter). This distance would either require huge ejection velocities from Veritas (that are not observed) or extremely long drift times via Yarkovsky thermal forces (which would fail to produce a spike of impactors). (Yarkovsky drift occurs when small asteroids absorb solar photons, heat up and then reradiate the energy away in a non-isotropic manner after a short delay<sup>7</sup>.) Moreover, these resonances are very unlikely to produce Earth impactors<sup>17,18</sup>. Thus, the event that formed the Veritas family almost certainly did not produce an asteroid shower on Earth, so the absence of craters of this age is not surprising.

To investigate under what conditions the Veritas collision might produce a dust spike similar to that defined by the  $^3\text{He}$  record, we developed a statistical Monte Carlo model to track the collisional and dynamical evolution of particles formed by the disruption of Veritas. (Model details can be found in Supplementary Information). The results of our code were calibrated by modelling the evolution of dust in several latitudinal bands observed by the Infrared Astronomical Satellite (IRAS; see ref. 15 for details). To compare our model results with the  $^3\text{He}$  data, we calculated the flux of  $D = 10 \mu\text{m}$  particles reaching 1 AU. Particles of about this size can escape intense atmospheric entry heating and He loss, and currently dominate the  $^3\text{He}$  flux to Earth<sup>8</sup>. Since we do not yet have a model of  $^3\text{He}$  implantation in IDPs, or one for heating and helium retention during atmospheric entry of IDPs produced during the dust spike, here we simply compare our model 1 AU flux with the shape and duration of the  $^3\text{He}$  peak.


Small particles from the Veritas family were assumed to reach Earth through P-R and solar wind drags. The production rate of the first-generation particles that were started at 3.17 AU was defined using a 'broken' cumulative power-law size frequency distribution (SFD) with two slopes, index  $\alpha_1$  at smaller sizes and  $\alpha_2$  at larger sizes. The SFD extends from  $D = 10 \mu\text{m}$  to 1 cm. We assume the SFD decays exponentially with time from collisional evolution and radiation drag forces. This causes the diameter of the knee between

$\alpha_1$  and  $\alpha_2$  to increase with time. The rate of IDP disruptions was defined as a function of diameter  $D$  and heliocentric distance  $R$  (ref. 19). When a particle disrupts, we replace it with a swarm of fragments that follow a power-law SFD. We assumed that the mass of the largest fragment was half that of the parent particle. The power law index of the fragment size distribution was determined by mass conservation<sup>20</sup>. Thus, in our simulations, we follow several generations of particles produced by a collisional cascade; typical runs track the histories of  $10^8$ – $10^9$  particles.

We find that  $D = 10 \mu\text{m}$  IDPs from Veritas reach 1 AU in  $\sim 40$  kyr, typically shorter than their collisional lifetime (20–200 kyr, depending on model assumptions<sup>19</sup>). This means the  $^3\text{He}$  signal at Earth would be extremely brief unless these particles are continuously replenished in some fashion. The Veritas break-up, however, produced a SFD of fragments. We note that  $D = 1$ – $5$  mm particles have short collisional lifetimes<sup>19</sup> ( $< 100$  kyr), such that their fragments not only replenish the  $D = 10 \mu\text{m}$  population but also create intermediate-size particles that also dynamically evolve and disrupt over time. This means that the  $^3\text{He}$  signal was produced by fragments from a collisional cascade that was fed new material by disruption events occurring both near the Veritas source region and *en route* to Earth.

The late Miocene event allows us to glean insights into the SFD of particles produced by the break-up and how it changes with time. We found that the initial break-up of the Veritas family probably produced a swarm of IDPs that dominated the main belt population by at least an order of magnitude for  $\sim 1$  Myr. Our best fit to the shape and the decay time of the  $^3\text{He}$  peak comes from using  $\alpha_1 = -2.5$  and  $\alpha_2 = -3.3$  (Fig. 2). We found that  $\alpha_2$  values significantly shallower or steeper than our best fit value produce  $^3\text{He}$  peaks that are longer or shorter, respectively, than those observed.

Today, collisions in the Veritas family produce one of the prominent dust bands observed by infrared telescopes, and also contribute at least  $5 \times 10^6$  kg per year to the terrestrial IDP flux<sup>15</sup>. Our modelling suggests that the IDP flux from the Veritas family will continue to decay for several tens of Myr until it reaches a collisional steady state



**Figure 2 | Concordant  $^3\text{He}$  data from two widely separated seafloor sites support a global increase in IDP flux at 8.2 Myr ago that can be attributed to the asteroid collision that produced the Veritas family.** **a, b.** The extraterrestrial  $^3\text{He}$  flux peak (grey line, with small symbols) through the Late Miocene event is similar at Site 757 (**a**) and Site 926 (**b**); western equatorial Atlantic;  $3^\circ 43.148' \text{ N}$ ,  $42^\circ 54.507' \text{ W}$ ). The modelled  $10\text{-}\mu\text{m}$  IDP flux following the Veritas collision is shown by the black curves. The model dust spike was positioned at 8.25 Myr ago and scaled to align with the  $^3\text{He}$  peak. The fast rise time and  $\sim 1.5$  Myr decay time observed in the  $^3\text{He}$  record at both sites are well matched by the model. The inferred time of Veritas break-up is indicated (see Supplementary Fig. S4 for details). The symbol at  $\sim 8$  Myr ago indicates transition from illite (I) to kaolinite (K) clays at Site 926, a proxy for climate change.

and takes on the same approximate shape of the overall main belt size distribution for  $D < 5$  km bodies<sup>21</sup>.

The late Miocene event roughly coincides with cosmic ray exposure ages of 7–8 Myr on many H-chondrite meteorites<sup>22</sup>. The connection between these observations, however, is unclear. As described previously, the nearest powerful resonances are not only  $\sim 0.1$  AU from Veritas family members, but they are also highly inefficient at producing Earth impactors. Moreover, mineralogical and spectroscopic differences between Veritas family members and the H-chondrites indicate that the latter almost certainly did not originate on the former<sup>23</sup>. We also find it highly unlikely that the projectile that produced the Veritas family was the source of the H-chondrites, partly for the reasons above, but also because the H-chondrites do not show evidence for significant shocks at 7–8 Myr ago<sup>24</sup>. If the two events are indeed linked, we postulate that Veritas family members disrupted a well-positioned fragment from the H-chondrite parent body shortly after the family-forming event took place.

Previous work has suggested a possible link between the IDP accretion rate and Earth's climate<sup>25</sup>. Correlations between extraterrestrial  $^3\text{He}$  in sediments and global climate in the Quaternary period may support this suggestion<sup>9</sup> but also may be an artefact of climate-induced changes in sedimentation<sup>26</sup>. Modest global cooling and strengthening of the Asian monsoon occurred in the late Miocene<sup>27</sup>. At Site 926 there is a sharp transition from kaolinite-rich to illite-rich sediment<sup>28</sup>, occurring within the  $^3\text{He}$  peak but 200 kyr after its onset (Fig. 2). This transition may document a change from warm humid to cold dry continental weathering. Although the relative timing of these events is suggestive, we caution that a compelling link between the events cannot be established until a plausible mechanism is found by which IDPs can change climate.

Received 21 June; accepted 31 October 2005.

1. Farley, K. A., Montanari, A., Shoemaker, E. M. & Shoemaker, C. S. Geochemical

evidence for a comet shower in the Late Eocene. *Science* **280**, 1250–1253 (1998).

- Takayanagi, M. & Ozima, M. Temporal variation of  $^3\text{He}/^4\text{He}$  in deep-sea sediment cores. *J. Geophys. Res.* **92**, 12531–12538 (1987).
- Farley, K. A. Cenozoic variations in the flux of interplanetary dust recorded by  $^3\text{He}$  in a deep-sea sediment. *Nature* **376**, 153–156 (1995).
- Kortenkamp, S. & Dermott, S. A 100,000-year periodicity in the accretion rate of interplanetary dust. *Science* **280**, 874–876 (1998).
- Nesvorný, D., Bottke, W. F., Levison, H. F. & Dones, L. Recent origin of the solar system dust bands. *Astrophys. J.* **591**, 486–497 (2003).
- Tagle, R. & Claeys, P. Comet or asteroid shower in the late Eocene? *Science* **305**, 492 (2004).
- Burns, J. A., Lamy, P. L. & Soter, S. Radiation forces on small particles in the Solar-System. *Icarus* **40**, 1–48 (1979).
- Farley, K. A., Love, S. G. & Patterson, D. B. Atmospheric entry heating and helium retentivity of interplanetary dust particles. *Geochim. Cosmochim. Acta* **61**, 2309–2316 (1997).
- Patterson, D. B. & Farley, K. A. Extraterrestrial He-3 in seafloor sediments: Evidence for correlated 100 kyr periodicity in the accretion rate of interplanetary dust, orbital parameters, and Quaternary climate. *Geochim. Cosmochim. Acta* **62**, 3669–3682 (1998).
- Mukhopadhyay, S., Farley, K. & Montanari, A. A 35 Myr record of helium in pelagic limestones: implications for interplanetary dust accretion from the early Maastrichtian to the Middle Eocene. *Geochim. Cosmochim. Acta* **65**, 653–669 (2001).
- Shackleton, N. J., Curry, W. B., Richter, C. & Bralower, T. J. Ceara Rise. *Proc. ODP Sci. Res.* **154**, 1–552 (1997).
- Bottke, W. F. *et al.* The fossilized size distribution of the main asteroid belt. *Icarus* **175**, 111–140 (2005).
- Low, F. J. *et al.* Infrared cirrus—new components of the extended infrared-emission. *Astrophys. J.* **278**, L19–L22 (1984).
- Dermott, S. F., *et al.* in *Interplanetary Dust* (eds Grün, E., Gustafson, B. A. S., Dermott, S. F. & Fechtig, H.) 569–639 (Springer, Berlin, 2001).
- Nesvorný, D., Vokrouhlický, D., Bottke, W. F. & Sykes, M. Physical properties of asteroid dust bands and their sources. *Icarus* (submitted).
- Morbidelli, A. & Nesvorný, D. Numerous weak resonances drive asteroids toward terrestrial planet orbits. *Icarus* **139**, 295–308 (1999).
- Gladman, B. J. *et al.* Dynamical lifetimes of objects injected into asteroid belt resonances. *Science* **277**, 197–201 (1997).
- Bottke, W. F. *et al.* Debaised orbital and absolute magnitude distribution of the near-earth objects. *Icarus* **156**, 399–433 (2002).
- Grün, E., Zook, H. A., Fechtig, H. & Giese, R. H. Collisional balance of the meteoritic complex. *Icarus* **62**, 244–272 (1985).
- Nolan, M. C. & Greenberg, R. Stochastic-evolution of asteroids to produce the ordinary chondrites. *Meteoritics* **24**, 310 (1989).
- Bottke, W. F. *et al.* in *Dynamics of Populations of Planetary Systems* (eds Knezevic, Z. & Milani, A.) 357–376 (IAU Colloquium 197, Cambridge Univ. Press, Cambridge, UK, 2005).
- Graf, T. & Marti, K. Collisional history of H chondrites. *J. Geophys. Res.* **100**, 21247–21263 (1995).
- Burbine, T. H., McCoy, T. J., Meibom, A., Gladman, B. & Keil, K. in *Asteroids III* (eds Bottke, W. F., Cellino, A., Paolicchi, P. & Binzel, R. P.) 653–667 (Univ. Arizona Press, Tucson, 2002).
- Bogard, D. D. Impact ages of meteorites—A synthesis. *Meteoritics* **30**, 244–268 (1995).
- Muller, R. A. & MacDonald, G. J. Glacial cycles and astronomical forcing. *Science* **277**, 215–218 (1997).
- Marcantonio, F. *et al.* Extraterrestrial  $^3\text{He}$  as a tracer of marine sediment transport and accumulation. *Nature* **383**, 705–707 (1996).
- Gupta, A. K., Singh, R. K., Joseph, S. & Thomas, E. Indian Ocean high-productivity event (10–8 Ma): Linked to global cooling or to the initiation of the Indian monsoons? *Geology* **32**, 753–756 (2004).
- Curry, W. B., *et al.* *ODP Init. Rep. Leg 154* 1–1111, (1995).

**Supplementary Information** is linked to the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

**Acknowledgements** Financial support for this project was provided by NASA's Planetary Geology & Geophysics program (W.F.B., D.N. and K.A.F.). Financial and travel support for D.V. was provided by the Czech Republic grant agency and NSF's COBASE program. We also thank D. Durda, A. Morbidelli, M. Sykes and S. Mukhopadhyay for several discussions, and S. Goldstein and J. Burns for comments and suggestions.

**Author Contributions** K.A.F. measured  $^3\text{He}$  in the seafloor sediments. D.N. determined the age of the Veritas family using numerical integration methods. D.V., W.F.B. and D.N. constructed the Monte Carlo dust evolution code and analysed the results.

**Author Information** Reprints and permissions information is available at [npg.nature.com/reprintsandpermissions](http://npg.nature.com/reprintsandpermissions). The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to K.A.F. ([farley@gps.caltech.edu](mailto:farley@gps.caltech.edu)).