



3-25-2017

## A Local Leaky-Box Model for the Local Stellar Surface Density-Gas Surface Density-Gas Phase Metallicity Relation

Guangtun Ben Zhu  
*Johns Hopkins University*

Jorge K. Barrera-Ballesteros  
*Johns Hopkins University*

Timothy M. Heckman  
*Johns Hopkins University*

Nadia L. Zakamska  
*Johns Hopkins University*

Sebastian F. Sánchez  
*Universidad Nacional Autónoma de México, Mexico*

*See next page for additional authors*

Follow this and additional works at: [https://uknowledge.uky.edu/physastron\\_facpub](https://uknowledge.uky.edu/physastron_facpub)

 Part of the [Stars, Interstellar Medium and the Galaxy Commons](#)

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

---

### Repository Citation

Zhu, Guangtun Ben; Barrera-Ballesteros, Jorge K.; Heckman, Timothy M.; Zakamska, Nadia L.; Sánchez, Sebastian F.; Yan, Renbin; and Brinkman, Jonathan, "A Local Leaky-Box Model for the Local Stellar Surface Density-Gas Surface Density-Gas Phase Metallicity Relation" (2017). *Physics and Astronomy Faculty Publications*. 550.

[https://uknowledge.uky.edu/physastron\\_facpub/550](https://uknowledge.uky.edu/physastron_facpub/550)

This Article is brought to you for free and open access by the Physics and Astronomy at UKnowledge. It has been accepted for inclusion in Physics and Astronomy Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@sv.uky.edu](mailto:UKnowledge@sv.uky.edu).

---

# A Local Leaky-Box Model for the Local Stellar Surface Density-Gas Surface Density-Gas Phase Metallicity Relation

Digital Object Identifier (DOI)

<https://doi.org/10.1093/mnras/stx740>

## Notes/Citation Information

Published in *Monthly Notices of the Royal Astronomical Society*, v. 468, issue 4, p. 4494-4501.

This article has been accepted for publication in *Monthly Notices of the Royal Astronomical Society* ©: 2017 The Authors. Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved.

The copyright holders have granted the permission for posting the article here.

## Authors

Guangtun Ben Zhu, Jorge K. Barrera-Ballesteros, Timothy M. Heckman, Nadia L. Zakamska, Sebastian F. Sánchez, Renbin Yan, and Jonathan Brinkman

# A local leaky-box model for the local stellar surface density–gas surface density–gas phase metallicity relation

Guangtun Ben Zhu,<sup>1</sup>★† Jorge K. Barrera-Ballesteros,<sup>1</sup> Timothy M. Heckman,<sup>1</sup>  
Nadia L. Zakamska,<sup>1,2</sup> Sebastian F. Sánchez,<sup>3</sup> Renbin Yan<sup>4</sup> and Jonathan Brinkmann<sup>5</sup>

<sup>1</sup>Department of Physics and Astronomy, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

<sup>2</sup>Deborah Lunder and Alan Ezekowitz, Founders' Circle Member, Institute for Advanced Study, Einstein Dr., Princeton, NJ 08540, USA

<sup>3</sup>Instituto de Astronomía, Universidad Nacional Autónoma de México, A.P. 70-264, 04510 México, D.F., Mexico

<sup>4</sup>Department of Physics and Astronomy, University of Kentucky, 505 Rose St., Lexington, KY 40506-0057, USA

<sup>5</sup>Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349, USA

Accepted 2017 March 23. Received 2017 March 22; in original form 2016 September 12

## ABSTRACT

We revisit the relation between the stellar surface density, the gas surface density and the gas-phase metallicity of typical disc galaxies in the local Universe with the SDSS-IV/MaNGA survey, using the star formation rate surface density as an indicator for the gas surface density. We show that these three local parameters form a tight relationship, confirming previous works (e.g. by the PINGS and CALIFA surveys), but with a larger sample. We present a new local leaky-box model, assuming star-formation history and chemical evolution is localized except for outflowing materials. We derive closed-form solutions for the evolution of stellar surface density, gas surface density and gas-phase metallicity, and show that these parameters form a tight relation independent of initial gas density and time. We show that, with canonical values of model parameters, this predicted relation match the observed one well. In addition, we briefly describe a pathway to improving the current semi-analytic models of galaxy formation by incorporating the local leaky-box model in the cosmological context, which can potentially explain simultaneously multiple properties of Milky Way-type disc galaxies, such as the size growth and the global stellar mass–gas metallicity relation.

**Key words:** galaxies: abundances – galaxies: evolution – galaxies: spiral – galaxies: star formation.

## 1 INTRODUCTION

Over the past few decades, a standard cosmological model of structure formation emerged in a series of major observational and theoretical advances (e.g. White & Rees 1978). However, most of these studies have largely focused on the global properties of galaxies (e.g. Kauffmann, White & Guiderdoni 1993; Springel et al. 2005; Somerville & Davé 2015).

Recent integral-field-unit (IFU) spectroscopic surveys from the ground (e.g. Bacon et al. 2001; Rosales-Ortega et al. 2010; Sánchez et al. 2012), high-spatial resolution deep imaging surveys with the *Hubble Space Telescope* (e.g. Scoville et al. 2007; Koekemoer et al. 2011) and high-resolution hydrodynamical simulations (e.g. Vogelsberger et al. 2014; Hopkins et al. 2014) have shifted the focus of the investigations of galaxy formation to small-scale astrophysics and to the relationships between local and global properties of galaxies. In particular, the MaNGA survey (Bundy et al. 2015) in SDSS-IV (Blanton et al. 2017) is obtaining IFU spectroscopy for

about 10 000 nearby galaxies and will provide the largest sample of galaxies with kpc-scale resolved optical spectroscopy, enabling systematic investigations of local properties and also their correlations with global parameters. In this paper, using the MaNGA data obtained in the first two years, we investigate the relation between the stellar surface density ( $\Sigma_*$ ), gas surface density ( $\Sigma_{\text{gas}}$ ) and gas-phase metallicity ( $Z$ ) in typical disc galaxies, using the star-formation rate (SFR) surface density ( $\Sigma_{\text{SFR}}$ ) as a proxy for  $\Sigma_{\text{gas}}$ . In particular, we show that a simple leaky-box model can explain well the observed relation between these parameters and propose a new way of thinking about disc galaxy formation.


The rest of the paper is organized as follows. In Section 2 and 3, we describe the data we use and the observed relation. We present the local leaky-box model in Section 4. In Section 5, we outline a global semi-analytic model for disc galaxy formation. We summarize our results in Section 6. When necessary, we assume the  $\Lambda$ CDM cosmogony, with  $\Omega_\Lambda = 0.7$ ,  $\Omega_m = 0.3$  and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 2 DATA

The SDSS-IV/MaNGA IFU survey uses the BOSS spectrographs (Smee et al. 2013) on the 2.5-m SDSS telescope (Gunn et al. 2006) at

\* E-mail: gz323@pha.jhu.edu

† Hubble Fellow.



**Figure 1.** Left-hand panel: The observed  $\Sigma_*$ – $Z$  relation of star-forming regions in typical disc galaxies. The contours enclose 90 per cent of the sub-samples with highest (blue), intermediate (green) and lowest (red) SFR surface density. Right-hand panel: The observed  $\Sigma_*$ – $\Sigma_{\text{SFR}}$ – $Z$  relation (grey-scale, equation 10), assuming  $\mathcal{R} = 0.3$ ,  $\epsilon = 0.0004$ ,  $k = 2.2$  and  $\eta = 1$ . The dashed line shows the relation with the best-fitting yield  $y = 0.003$ . The error bars show the typical measurement uncertainties, 0.06 dex for metallicity and 0.15 dex for stellar and SFR surface density.

the Apache Point Observatory. Detailed description of the MaNGA surveys are available in Bundy et al. (2015, overview), Drory et al. (2015, instrumentation), Law et al. (2015, 2016, observation, data reduction) and Yan et al. (2016a,b, calibration, survey design). We use the fourth internal data release of the MaNGA survey (MPL-4), which includes 1390 galaxies observed as of June 2015.

For our purposes, we are interested in typical disc galaxies and we select our sample and use the same data as we did in Barrera-Ballesteros et al. (2016). We select 653 disc galaxies spanning stellar masses between  $10^{8.5}$  and  $10^{11} M_\odot$ . The data cubes include about 507 000 star-forming spaxels with spatial resolution ranging from  $\sim 1.5$  to  $\sim 2.5$  kpc. For the parameter measurements, we use the estimates from the PIPE3D pipeline (Sánchez et al. 2016). PIPE3D estimated the stellar mass at a given spaxel by fitting the underlying stellar continuum with spectral templates taken from MIUSCAT SSP library (Vazdekis et al. 2012), assuming a Salpeter (1955) initial mass function (IMF). The pipeline also took into account of dust attenuation (Calzetti 2001). We estimated SFR using the dust attenuation-corrected flux of  $H\alpha$ . We have also corrected the surface densities for the inclination effect (see Barrera-Ballesteros et al. 2016). For gas-phase metallicity, we use the O3N2 indicator based on the  $[O III] \lambda 5008$  and  $[N II] \lambda 6584$  ratio (e.g. Marino et al. 2013). For more details regarding the data and the survey, we refer the reader to references above.

### 3 THE LOCAL $\Sigma_*$ – $\Sigma_{\text{SFR}}$ – $Z$ RELATION

Early works (e.g. Edmunds & Pagel 1984; Vila-Costas & Edmunds 1992) have already suggested that there exists a relationship between the local stellar surface density and the gas-phase metallicity. More recently, the PINGS and CALIFA surveys have presented conclusive evidence for such a relationship (Rosales-Ortega et al. 2012; Sánchez et al. 2013). In Barrera-Ballesteros et al. (2016), we presented further evidence with the MaNGA survey. Rosales-Ortega et al. (2012) and Sánchez et al. (2013) further showed that including the local SFR surface density indicates that the three

parameters together form a tight relationship. Our objective is to revisit this relation with a larger sample and then devise a local chemical evolution model for its interpretation.

In the left-hand panel of Fig. 1, we show the  $\Sigma_*$ – $Z$  relation (the same as in fig. 2 of Barrera-Ballesteros et al. 2016). In addition, we divide the star-forming regions into three sub-samples with the highest, intermediate and lowest SFR surface density and show their distributions in blue, green and red contours, respectively. We find that these three parameters,  $\Sigma_*$ ,  $\Sigma_{\text{SFR}}$  and  $Z$ , form a tight correlation with each other. We therefore confirm the findings by Rosales-Ortega et al. (2012) with the PINGS survey (Rosales-Ortega et al. 2010), who used luminosity surface density as a proxy for stellar surface density and  $H\alpha$  equivalent width for specific SFR, and also the recent results with the derived physical parameters from the larger CALIFA survey (Sánchez et al. 2013).

The gas-phase metallicity is the ratio of the amount of heavy elements (in our case, oxygen) to the total amount of gas in the galaxy, i.e.  $Z = \Sigma_{\text{metal}}/\Sigma_{\text{gas}}$ . Both metals and stars are integrated products of the star-formation history, while the SFR is closely correlated to the amount of gas available, through the Kennicutt–Schmidt (K–S) law (Schmidt 1959; Kennicutt 1998). The relations between the three parameters must therefore be closely related to the local star-formation history. In the next section, we present a leaky-box model of the local star-formation history and chemical evolution and show that it can naturally explain our observation.

### 4 THE LOCAL LEAKY-BOX MODEL

We assume a disc galaxy grows inside out (e.g. Larson 1976; Matteucci & Francois 1989; Governato et al. 2007; Pilkington et al. 2012; Gibson et al. 2013, among others), and gas falls in on to the outskirts, collapses and triggers star formation.<sup>1</sup> In this scenario all processes – star formation and metal production – are

<sup>1</sup> We note, if we start with a disc of gas right from the beginning, our analysis still applies.

localized within the same region except for the outflowing gas. These assumptions enable us to construct a model of the localized star-formation history and chemical evolution, which we describe in detail below.

If gas is accreted on to the galaxy with initial gas surface density  $\Sigma_0 \equiv \Sigma_{\text{gas}}(t_0)$  at accretion time  $t_0$ , we can define a total surface density as

$$\begin{aligned}\Sigma_{\text{tot}}(t) &= \Sigma_*(t) + \Sigma_{\text{gas}}(t) + \Sigma_{\text{out}}(t) \\ &= \Sigma_{\text{tot}}(t_0) \\ &= \Sigma_0,\end{aligned}\quad (1)$$

where  $\Sigma_{\text{gas}}(t)$  and  $\Sigma_*(t)$  are the surface densities of gas and long-lived stars at a given time  $t$ , respectively. For convenience, we have defined  $\Sigma_{\text{out}}(t)$  to represent the would-be density of the expelled gas should it stay within the same area, even though it can be anywhere in the circumgalactic/intergalactic media. If there is no outflow (i.e.  $\Sigma_{\text{out}} = 0$ ), we have a closed-box model. There has been ample evidence showing that star-forming galaxies exhibit ubiquitous outflows (e.g. Lynds & Sandage 1963; Bland & Tully 1988; Heckman, Armus & Miley 1990; Shapley et al. 2003; Rupke, Veilleux & Sanders 2005; Martin & Bouché 2009; Weiner et al. 2009; Rubin et al. 2014; Zhu et al. 2015, among others). Outflows also help explain the large amount of metals found outside galaxies in the circum-/inter-galactic media (e.g. Bergeron 1986; Steidel et al. 2010; Tumlinson et al. 2011; Borthakur et al. 2013; Stocke et al. 2013; Bordoloi et al. 2014; Werk et al. 2014; Zhu et al. 2014, among others). We here therefore assume a leaky-box model.

Another assumption of our model is that the expelled gas does not fall back on to the galaxy. Theoretical studies have suggested at least a fraction of the expelled gas would be re-accreted (e.g. Oppenheimer et al. 2010; Bower, Benson & Crain 2012; Brook et al. 2012; Marasco, Fraternali & Binney 2012; Henriques et al. 2013; Christensen et al. 2016). If some of the expelled gas falls right back on to the same region, its effect is equivalent to a smaller outflow rate and our model still applies. If some of the expelled gas gets mixed with gas outside and falls back in on to the outskirts, the formalism applies as well since the recycled gas does not invalidate the locality. If a significant fraction of the expelled gas is spread out and falls back over the whole galaxy (e.g. as in the galaxy fountain model, Marasco et al. 2012), it may have a non-negligible effect on the chemical evolution. This last scenario is more complicated than our simple model can yet address and we leave it for future work.

With the assumptions above, the total surface density defined above stays constant over the cosmic time ( $=\Sigma_0$ ). This synthetic density,  $\Sigma_{\text{tot}}(t)$ , includes the outflowing gas, while the total density within the disc would only include the gas and stars in the disc [ $\Sigma_*(t) + \Sigma_{\text{gas}}(t)$ ]. The constancy of this density and the direct connection between the amount of outflowing gas and the instantaneous SFR make it possible to derive a closed-form solution of the full chemical evolution history, as described below.

The SFR surface density is related to the gas surface density through the K–S law as

$$\Sigma_{\text{SFR}} \equiv \frac{1}{1-\mathcal{R}} \frac{d\Sigma_*(t)}{dt} = \epsilon \Sigma_{\text{gas}}^k(t), \quad (2)$$

where  $\mathcal{R}$  is the ‘return fraction’, i.e. the fraction of the stellar mass formed that is assumed to be instantaneously returned to the gas from short-lived massive stars, and  $\epsilon$  is the effective star formation efficiency and  $k$  is the K–S index. Note  $\epsilon$  is not unitless and its dimension depends on  $k$ . Following convention, we express  $\Sigma_*$  and

$\Sigma_{\text{gas}}$  in unit of  $\text{M}_{\odot} \text{pc}^{-2}$ , while  $\Sigma_{\text{SFR}}$  in unit of  $\text{M}_{\odot} \text{kpc}^{-2}$ . We also expect there is a threshold below which star formation cannot continue, and we assume this threshold to be  $10 \text{M}_{\odot} \text{pc}^{-2}$  (e.g. Skillman 1987; Schaye 2004; Leroy et al. 2008).

In global models, the outflow rate is usually assumed to be proportional to the total SFR (e.g. Springel & Hernquist 2003; Dalla Vecchia & Schaye 2008), and we extend this assumption to our local model. The outflow rate is related to the SFR through

$$\frac{d\Sigma_{\text{out}}(t)}{dt} = \eta \Sigma_{\text{SFR}} = \frac{\eta}{1-\mathcal{R}} \frac{d\Sigma_*(t)}{dt}, \quad (3)$$

where  $\eta$  is the mass loading factor and we assume it is constant (e.g. Springel & Hernquist 2003; Heckman et al. 2015).

Combining the above equations gives the relation between gas consumption rate, SFR surface density and gas surface density as

$$\frac{d\Sigma_{\text{gas}}(t)}{dt} = -(1 + \frac{\eta}{1-\mathcal{R}}) \frac{d\Sigma_*(t)}{dt} \quad (4)$$

$$= -(1 - \mathcal{R} + \eta) \epsilon \Sigma_{\text{gas}}^k(t), \quad (5)$$

from which we can solve for the full star-formation history, including  $\Sigma_{\text{gas}}(t)$ ,  $\Sigma_*(t)$ ,  $\Sigma_{\text{SFR}}(t)$ ,  $\Sigma_{\text{out}}(t)$ , mass-weighted age of the stars, etc. In particular, assuming  $k > 1$ ,  $\Sigma_{\text{gas}}(t)$  is given by

$$\Sigma_{\text{gas}}^{1-k}(t) = \Sigma_0^{1-k} - (1 - \mathcal{R} + \eta) \epsilon (1 - k)(t - t_0). \quad (6)$$

We can now derive the chemical evolution of this leaky-box model. The metallicity ( $Z \equiv \Sigma_{\text{metal}}/\Sigma_{\text{gas}}$ ) growth rate is given by

$$\begin{aligned}\frac{dZ(t)}{dt} &= \frac{1}{\Sigma_{\text{gas}}(t)} \frac{d\Sigma_{\text{metal}}(t)}{dt} - \frac{\Sigma_{\text{metal}}(t)}{\Sigma_{\text{gas}}^2(t)} \frac{d\Sigma_{\text{gas}}(t)}{dt} \\ &= \frac{1}{\Sigma_{\text{gas}}(t)} \left( \frac{d\Sigma_{\text{metal}}(t)}{dt} - Z(t) \frac{d\Sigma_{\text{gas}}(t)}{dt} \right),\end{aligned} \quad (7)$$

where  $\Sigma_{\text{metal}}$  is the surface density of metals in the gas. If  $y$  is the total metal mass yield that a stellar population releases into the ISM normalized by the mass locked up in long-lived stars, the amount of new metals that stay in the gas in the galaxy is given by the total yield minus that locked in stars and expelled along with outflows:

$$\begin{aligned}\frac{d\Sigma_{\text{metal}}(t)}{dt} &= y \frac{d\Sigma_*(t)}{dt} - Z(t) \left( \frac{d\Sigma_*(t)}{dt} + \frac{d\Sigma_{\text{out}}(t)}{dt} \right) \\ &= \left( y - Z(t) - Z(t) \frac{\eta}{1-\mathcal{R}} \right) \frac{d\Sigma_*(t)}{dt} \\ &= \left( y - Z(t) - Z(t) \frac{\eta}{1-\mathcal{R}} \right) \\ &\quad \times \frac{-1}{1 + \eta/(1-\mathcal{R})} \frac{d\Sigma_{\text{gas}}(t)}{dt},\end{aligned} \quad (8)$$

where we have assumed the metallicity in the outflowing gas is the same as in the ISM at the time.


The metallicity growth rate is then given by

$$\frac{dZ(t)}{dt} = \frac{d\Sigma_{\text{gas}}(t)}{\Sigma_{\text{gas}}(t) dt} \left( \frac{y}{1 + \eta/(1-\mathcal{R})} \right). \quad (9)$$

Eliminating  $dt$  gives the dependence of the metallicity on  $\Sigma_0$  and  $\Sigma_{\text{gas}}(t)$ :

$$\begin{aligned}Z(t) - Z_0 &= \frac{y}{1 + \eta/(1-\mathcal{R})} \log \frac{\Sigma_0}{\Sigma_{\text{gas}}(t)} \\ &= \frac{\log(10)y}{1 + \eta/(1-\mathcal{R})} [\log_{10} \Sigma_0 - \log_{10} \Sigma_{\text{gas}}(t)].\end{aligned} \quad (10)$$





**Figure 2.** Radial dependence of the local  $\Sigma_*$ – $\Sigma_{\text{SFR}}$ – $Z$  relation. Left-hand panel: Regions within  $r_{\text{eff}}$ . Right-hand panel: Regions outside  $r_{\text{eff}}$ . The dashed lines are the same as in Fig. 1.

We have thus derived the local version of the well-known global leaky-box model of chemical evolution (e.g. Tinsley 1980), which has been used to study the global mass–metallicity relation (e.g. Zahid et al. 2014; Belfiore, Maiolino & Bothwell 2016). We assume  $Z_0$  is 0.1 per cent of the solar value, though as long as it is lower than 1 per cent solar, it has no effect on any of our conclusions.

Based on the assumptions of the model (equations 1 and 3), we can also calculate  $\Sigma_0$  as

$$\Sigma_0 = \Sigma_{\text{gas}}(t) + \left(1 + \frac{\eta}{1 - \mathcal{R}}\right) \Sigma_*(t), \quad (11)$$

and the metallicity can now be fully determined if we can observe  $\Sigma_*$  and  $\Sigma_{\text{gas}}$  and if we know  $\eta$  and  $y$ . This  $\Sigma_*$ – $\Sigma_{\text{gas}}$ – $Z$  relation is a fundamental relation predicted by the local leaky-box model.

Now if we assume the K–S law (equation 2) holds and we can measure  $\Sigma_{\text{SFR}}$ , we can estimate the gas density  $\Sigma_{\text{gas}}(t)$  with


$$\Sigma_{\text{gas}}(t) = \left(\frac{\Sigma_{\text{SFR}}(t)}{\epsilon}\right)^{1/k}. \quad (12)$$

In principle, we can constrain the parameters ( $\mathcal{R}$ ,  $y$ ,  $\eta$ ,  $\epsilon$ ,  $k$ ) directly using the observation. The model, however, is non-linear and the parameters are degenerate with each other. For example, the yield  $y$  and the loading factor  $\eta$  are degenerate in the amplitude, thus a closed-box model (with  $\eta = 0$ ) with high yield can also fit the data well. A robust modelling therefore requires careful treatments of the completeness (as a function of the observables). In this first work, we choose to investigate the relation using a fiducial model with values calibrated from the literature. In particular, we first fix the return fraction  $\mathcal{R}$  to be 0.3 for a Salpeter IMF (e.g. Tinsley 1980; Madau & Dickinson 2014). We use  $\epsilon = 0.0004$  and  $k = 2.2$  for the K–S law in normal spiral galaxies (e.g. Misiriotis et al. 2006; Bigiel et al. 2008). The K–S law is observed to be non-linear. For normal galaxies, the slope is  $k \sim 2.2$  when total gas surface density is considered, and is smaller ( $k \sim 1.2$ ) if only molecular gas density is included (e.g. Wong & Blitz 2002; Boissier et al. 2003; Luna et al. 2006). For starburst galaxies, the K–S law is shallower (e.g. Bigiel et al. 2008). As we are interested in the total gas density for

typical star-forming galaxies, we here adopt a linear K–S relation with  $k = 2.2$  and take the amplitude from Bigiel et al. (2008). For the mass loading factor  $\eta$ , we set it to be 1, a choice consistent with suggestions by past studies (e.g. Martin 1999; Veilleux, Cecil & Bland-Hawthorn 2005; Schaye et al. 2010; Heckman et al. 2015). The right-hand panel of Fig. 1 shows the observed relation with these choices.

Fixing these three values ( $\epsilon = 0.0004$ ,  $k = 2.2$  and  $\eta = 1$ ), we fit the normalization for the metal yield and obtain  $y \sim 0.003$ . This yield is for oxygen ( $^{16}\text{O}$ ), and the total metal yield is larger by about a factor of 2,  $y_{\text{total}} \sim 0.006$ . The values above are for a Salpeter IMF. For a Chabrier or Kroupa IMF (Kroupa 2001; Chabrier 2003), the oxygen and total metal yield would be about 0.0045 and 0.009, respectively. We plot this best-fitting relation with the dashed line. We find it remarkable that, with these canonical values, we obtain a tight  $\Sigma_*$ – $\Sigma_{\text{gas}}/\Sigma_{\text{SFR}}$ – $Z$  relation, and the fiducial model matches the observation very well. Our best-fitting metal yield is at the lower end of the theoretical estimates (e.g. Henry, Edmunds & Köppen 2000; Kobayashi et al. 2006; Zahid et al. 2012; Vincenzo et al. 2016). As it is degenerate with the mass loading factor ( $\eta$ ), if we choose a larger  $\eta$ , we will get a larger yield.

To take a further look at this local relation, we separate the parent spaxel samples by their galactocentric distance and the stellar mass of their host galaxy. In Fig. 2, we plot the local relation for star-forming regions outside (left-hand panel) and within (right-hand panel) the effective radius. In Fig. 3, we show the relation for low-mass (left-hand panel) and high-mass (right-hand panel) galaxies. The dash lines in all panels are the same as in Fig. 1. We show the best-fitting local relation fits well the data of all the sub-samples. We observe a weak dependence of the relation on the galactocentric distance and stellar mass: Regions at larger radius and in more massive galaxies tend to be distributed above the best-fitting relation with higher metallicity. We suspect that this weak dependence may be caused by some of the simple assumptions we made in the model: constant yield and mass loading factor, no recycled gas and metals, and no radial mixing. We leave detailed investigation for future work.



**Figure 3.** Mass dependence of the local  $\Sigma_*$ – $\Sigma_{\text{SFR}}$ – $Z$  relation. Left-hand panel: Regions in host galaxies with  $M_* < 10^{10} M_\odot$ . Right-hand panel: Regions in host galaxies with  $M_* > 10^{10} M_\odot$ . The dashed lines are the same as in Fig. 1.

As similar in the global leaky-box model, given an initial gas surface density  $\Sigma_0$ , the leaky-box model fully describes the local star-formation history and chemical evolution. In Fig. 4, we show for the fiducial model the predicted evolutionary tracks of metallicity for different  $\Sigma_0$  as a function of  $\Sigma_{\text{gas}}$ ,  $\Sigma_*$  and  $\log_{10} \Sigma_0 / \Sigma_{\text{gas}}$ . Each line shows that as time increases, the metallicity and stellar surface density increase, while the gas surface density decreases. We show that the evolution of metallicity, stellar and gas surface density, as well as their relations, are strong functions of the initial gas surface density, while the  $\Sigma_*$ – $\Sigma_{\text{gas}}$ – $Z$  relation (bottom) does not depend on either time or  $\Sigma_0$  and is a fundamental relation predicted by the local leaky-box model.

Since the local leaky-box model is fully determined by the initial gas surface density  $\Sigma_0$ , for any typical disc galaxy, if we can determine the initial surface density at the accretion time at any given radius, we can connect the small-scale astrophysics with the large-scale cosmological context. We briefly discuss how to expand the local model to a cosmological inside-out growth model in the next section.

Some of the earlier works have presented similar ideas of localized star-formation history and chemical evolution (e.g. Rosales-Ortega et al. 2012; Fu et al. 2013; Sánchez et al. 2013; Carton et al. 2015; Ho et al. 2015; Kudritzki et al. 2015). In particular, Ho et al. (2015) and Carton et al. (2015) extended a global gas regulatory model (Lilly et al. 2013) by ignoring radial mass transfer, which is also an assumption of our model, and showed that it could reproduce the radial metallicity profile for a large fraction of disc galaxies in their samples. They used global parameters (total stellar mass, total SFR) except for the metallicity in their models to reconstruct the observed density/metallicity gradient from resolved IFU observations. Although they did not provide a formalism for the localized star-formation history as we did, they presented new ideas to connect the global properties of the galaxy with the local ones. The model we suggest below outlines a way to integrate these ideas presented in their pioneering works and our local leaky-box model to build a typical disc galaxy analytically in the cosmological context.

## 5 THE COSMOLOGICAL INSIDE-OUT GROWTH MODEL

Suppose the dark matter accretion rate of a given dark matter halo (with mass  $M_{\text{DM}}$ ) at a given time ( $t$ ) is

$$\dot{M}_{\text{DM}} \equiv \frac{dM_{\text{DM}}(t)}{dt} = \dot{M}_{\text{DM}}(M_{\text{DM}}, t), \quad (13)$$

which is a function of  $M_{\text{DM}}$  and  $t$  and can be calibrated from simulations (e.g. Wechsler et al. 2002; Correa et al. 2015), the gas accretion rate (on to the galaxy) is then given by

$$\dot{M}_{\text{gas}}(t) \equiv \frac{dM_{\text{gas}}(t)}{dt} = \lambda f_b \dot{M}_{\text{DM}}(M_{\text{DM}}, t), \quad (14)$$

where  $f_b$  is the cosmic ratio of baryon mass to dark matter and  $\lambda$  is the fraction of baryons that fall all the way in on to the galaxy.


We assume the newly accreted gas only stays on the outskirts and the galaxy grows from inside out. In this case, the gas accretion rate is naturally connected to the size growth of the galaxy  $\dot{R}(t)$  and the initial surface density at the galaxy-size radius at the accretion time  $\Sigma_0(R)$ :

$$\dot{M}_{\text{gas}}(t) = n h(R) 2\pi R(t) \frac{dR}{dt} \quad (15)$$

$$= \Sigma_0(R) 2\pi R(t) \dot{R}, \quad (16)$$

where  $n$  is the volume density when gas starts to form stars and must be closely connected to the star formation density threshold for giant molecular clouds,  $R(t)$  is the galaxy size at  $t$ ,  $h(R)$  is the initial scale height at  $R$  and  $\Sigma_0(R)$  is the initial total surface density at  $R$ .

If we can calibrate  $\dot{M}_{\text{gas}}(t)$  with simulations, we can infer the radial profile of the initial density  $\Sigma_0(R)$  from the size growth of the galaxy  $\dot{R}$ , and vice versa. In particular, if we know the size  $R(t)$  and its growth rate  $\dot{R}(t)$  of a typical disc galaxy (e.g. van Dokkum et al. 2013; van der Wel et al. 2014), by applying the local leaky-box model, we can fully derive the radial profiles of  $\Sigma_{\text{gas}}(r, t)$ ,  $\Sigma_*(r, t)$ ,  $\Sigma_{\text{SFR}}(r, t)$ ,  $Z(r, t)$  and mass-weighted stellar age  $\langle t_* \rangle(r, t)$ , where



**Figure 4.** The predicted evolutionary tracks of the local star-formation history as a function of  $\Sigma_0$  (in  $\text{M}_{\odot} \text{pc}^{-2}$ ). For each track, time increases from left- to right-hand sides and from bottom to top,  $\Sigma_{\text{gas}}$  decreases with time, while  $\Sigma_*$  and  $Z$  increase with time. Top: the  $\Sigma_{\text{gas}}-Z$  relation. We have reversed the order of  $\Sigma_{\text{gas}}$  for display purposes. Middle: the  $\Sigma_*-Z$  relation. Bottom: the  $\Sigma_*-\Sigma_{\text{gas}}-Z$  relation, as given by Equation 10. All tracks with different  $\Sigma_0$  overlap for this relation. The black dashed line is the same as the magenta dashed line in the right panel of Fig. 1, showing the range probed by the MaNGA survey, slightly shifted downwards for clarity.

$r < R(t)$ . IFU surveys such as CALIFA and MaNGA have started to obtain these radial profiles for a large sample of disc galaxies (e.g. Pérez et al. 2013; Sánchez et al. 2013). Galactic surveys, such as RAVE (Steinmetz et al. 2006) and APOGEE (Majewski et al. 2015), have also started to provide chemical gradient measurements of Galactic stars (e.g. Boeche et al. 2013; Hayden et al. 2014; Ness et al. 2016), lending support to an inside-out growth scenario for our own Milky Way. We can also compare the relations among the above parameters and their dependence on global properties should we observe a large sample of systems, such as the stellar mass/SFR (in)dependence of the  $\Sigma_*$ - $Z$  relation observed in our previous paper (e.g. Barrera-Ballesteros et al. 2016) and the relation between global stellar mass, SFR and central-region metallicity (e.g. Lara-López et al. 2010; Mannucci et al. 2010; Sánchez et al. 2013; Salim et al. 2014, 2015; Bothwell et al. 2016). We therefore expect that a full semi-analytical model can be compared with observations directly, not only for global properties as previous-generation models, but also for local and structural properties revealed by IFU spectroscopic and deep high-spatial resolution imaging surveys. We leave the full modelling for future work.

## 6 CONCLUSIONS

With the most recent data from the MaNGA survey, we have confirmed a tight relation between the stellar surface density, gas surface density and gas-phase metallicity. We introduced a new local leaky-box model, in which star formation and metal production are localized within the same region except for the outflowing gas. With this model, we derived closed-form solutions for the evolution of stellar surface density, gas surface density and gas-phase metallicity, and showed that they follow a tight relation regardless of initial gas density and time. We further demonstrated that, with canonical values for the model parameters, the closed-form relation predicted by the model matches the observed one well. Our local leaky-box model therefore provided a natural explanation for the relationship between local parameters by the recent IFU observations and suggested a new look at the evolution of typical disc galaxies like our own Milky Way. We briefly introduced how to build a cosmological semi-analytical inside-out growth model that can take into account of the small-scale astrophysics by including the localized star-formation history.

We can further refine and improve the local leaky-box model. For example, if we can observe the gas density (e.g. as in the DiskMass Survey, Martinsson et al. 2013), then we can investigate the local relation directly without the assumption of the Kennicutt–Schmidt law. The current local leaky-box model also neglects several possible effects. We have assumed the parameters ( $\epsilon$ ,  $k$ ,  $\eta$ ,  $y$ ) are all constant. In reality, the K–S index depends on  $\Sigma_{\text{gas}}$  (e.g. Bigiel et al. 2008), and the mass loading factor must also depend on  $\Sigma_{\text{SFR}}$  (Heckman et al. 2015) and also the local and/or global gravitational potential. It is believed that radial migration of stars and gas happens on some level (e.g. Haywood 2008), though it is yet unclear how important it is in the general evolution of disc galaxies. The expelled gas can also be recycled back to the galaxy (e.g. Oppenheimer et al. 2010; Christensen et al. 2016). Mergers can also affect the distribution of metals (e.g. Rupke, Kewley & Barnes 2010). In addition, the model we described does not address the formation and evolution of bulges and bars at the centre. It is also a statistical model and neglects structures such as spiral arms. We expect these open issues to be the focuses of future investigations.



On a larger scale, the outflow component can be connected to quenching due to stellar/supernova feedback. The cosmological inside-out growth model with the localized star-formation history is a natural next step of the gas regulatory model used for global evolution of galaxies (e.g. Bouché et al. 2010; Lilly et al. 2013). Instead of adding more gas to the total gas reservoir, the inside-out growth model simplifies the physical treatments as it adds new gas to the outskirts without interfering with the (local) reservoir on the inside.

The MaNGA survey is continuing its operation and will provide us with six times more data by the end of the survey. With such a large data set, we will be able to investigate not only the local properties with IFU data themselves, but also the correlations between them and global properties and large-scale structures. Together with the rapid development of high-resolution hydrodynamical simulations and new analytical models as the one described in this paper, we are entering a new era of galaxy formation and evolution where we can now connect directly small-scale astrophysics with the cosmological context in both observation and theory.

## ACKNOWLEDGEMENTS

GBZ acknowledges support provided by NASA through Hubble Fellowship grant #HST-HF2-51351 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract NAS 5-26555. We thank an anonymous referee for many constructive comments that have helped improve this paper.

Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is [www.sdss.org](http://www.sdss.org).

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

## REFERENCES

Bacon R. et al., 2001, *MNRAS*, 326, 23  
 Barrera-Ballesteros J. K. et al., 2016, *MNRAS*, 463, 2513  
 Belfiore F., Maiolino R., Bothwell M., 2016, *MNRAS*, 455, 1218  
 Bergeron J., 1986, *A&A*, 155, L8

Bigiel F., Leroy A., Walter F., Brinks E., de Blok W. J. G., Madore B., Thornley M. D., 2008, *AJ*, 136, 2846  
 Bland J., Tully B., 1988, *Nature*, 334, 43  
 Blanton M. R., 2017, preprint ([arXiv:1703.00052](https://arxiv.org/abs/1703.00052))  
 Boeche C. et al., 2013, *A&A*, 559, A59  
 Boissier S., Prantzos N., Boselli A., Gavazzi G., 2003, *MNRAS*, 346, 1215  
 Bordoloi R. et al., 2014, *ApJ*, 796, 136  
 Borthakur S., Heckman T., Strickland D., Wild V., Schiminovich D., 2013, *ApJ*, 768, 18  
 Bothwell M. S., Maiolino R., Peng Y., Ciccone C., Griffith H., Wagg J., 2016, *MNRAS*, 455, 1156  
 Bouché N. et al., 2010, *ApJ*, 718, 1001  
 Bower R. G., Benson A. J., Crain R. A., 2012, *MNRAS*, 422, 2816  
 Brammer G. B. et al., 2012, *ApJS*, 200, 13  
 Brook C. B., Stinson G., Gibson B. K., Roškar R., Wadsley J., Quinn T., 2012, *MNRAS*, 419, 771  
 Bundy K. et al., 2015, *ApJ*, 798, 7  
 Calzetti D., 2001, *PASP*, 113, 1449  
 Carton D. et al., 2015, *MNRAS*, 451, 210  
 Chabrier G., 2003, *PASP*, 115, 763  
 Christensen C. R., Davé R., Governato F., Pontzen A., Brooks A., Munshi F., Quinn T., Wadsley J., 2016, *ApJ*, 824, 57  
 Correa C. A., Wyithe J. S. B., Schaye J., Duffy A. R., 2015, *MNRAS*, 450, 1514  
 Dalla Vecchia C., Schaye J., 2008, *MNRAS*, 387, 1431  
 Drory N. et al., 2015, *AJ*, 149, 77  
 Edmunds M. G., Pagel B. E. J., 1984, *MNRAS*, 211, 507  
 Fu J. et al., 2013, *MNRAS*, 434, 1531  
 Gibson B. K., Pilkington K., Brook C. B., Stinson G. S., Bailin J., 2013, *A&A*, 554, A47  
 Governato F., Willman B., Mayer L., Brooks A., Stinson G., Valenzuela O., Wadsley J., Quinn T., 2007, *MNRAS*, 374, 1479  
 Gunn J. E. et al., 2006, *AJ*, 131, 232  
 Hayden M. R. et al., 2014, *AJ*, 147, 116  
 Haywood M., 2008, *MNRAS*, 388, 1175  
 Heckman T. M., Armus L., Miley G. K., 1990, *ApJS*, 74, 833  
 Heckman T. M., Alexandroff R. M., Borthakur S., Overzier R., Leitherer C., 2015, *ApJ*, 809, 147  
 Henriques B. M. B., White S. D. M., Thomas P. A., Angulo Raul E., Guo Q., Lemson G., Springel V., 2013, *MNRAS*, 431, 3373  
 Henry R. B. C., Edmunds M. G., Köppen J., 2000, *ApJ*, 541, 660  
 Ho I.-T., Kudritzki R.-P., Kewley L. J., Zahid H. J., Dopita M. A., Bresolin F., Rupke D. S. N., 2015, *MNRAS*, 448, 2030  
 Hopkins P. F., Kereš D., Oñorbe J., Faucher-Giguère C.-A., Quataert E., Murray N., Bullock James S., 2014, *MNRAS*, 445, 581  
 Kauffmann G., White S. D. M., Guiderdoni B., 1993, *MNRAS*, 264, 201  
 Kennicutt R. C. Jr., 1998, *ApJ*, 498, 541  
 Kobayashi C., Umeda H., Nomoto K., Tominaga N., Ohkubo T., 2006, *ApJ*, 653, 1145  
 Koekemoer A. M. et al., 2011, *ApJS*, 197, 36  
 Kroupa P., 2001, *MNRAS*, 322, 231  
 Kudritzki R.-P., Ho I.-T., Schruha A., Burkert A., Zahid H. J., Bresolin F., Dima G. I., 2015, *MNRAS*, 450, 342  
 Lara-López M. A. et al., 2010, *A&A*, 521, L53  
 Larson R. B., 1976, *MNRAS*, 176, 31  
 Law D. R. et al., 2015, *AJ*, 150, 19  
 Law D. R. et al., 2016, *AJ*, 152, 83  
 Leroy A. K., Walter F., Brinks E., Bigiel F., de Blok W. J. G., Madore B., Thornley M. D., 2008, *AJ*, 136, 2782  
 Lilly S. J., Carollo C. M., Pipino A., Renzini A., Peng Y., 2013, *ApJ*, 772, 119  
 Luna A., Bronfman L., Carrasco L., May J., 2006, *ApJ*, 641, 938  
 Lynds C. R., Sandage A. R., 1963, *ApJ*, 137, 1005  
 Madau P., Dickinson M., 2014, *ARA&A*, 52, 415  
 Majewski S. R. et al., 2015, preprint ([arXiv:1509.05420](https://arxiv.org/abs/1509.05420))  
 Mannucci F., Cresci G., Maiolino R., Marconi A., Gnerucci A., 2010, *MNRAS*, 408, 2115

- Marasco A., Fraternali F., Binney J. J., 2012, *MNRAS*, 419, 1107
- Marino R. A. et al., 2013, *A&A*, 559, A114
- Martin C. L., 1999, *ApJ*, 513, 156
- Martin C. L., Bouché N., 2009, *ApJ*, 703, 1394
- Martinsson T. P. K., Verheijen M. A. W., Westfall K. B., Bershadsky M. A., Andersen D. R., Swaters R. A., 2013, *A&A*, 557, A131
- Matteucci F., Francois P., 1989, *MNRAS*, 239, 885
- Misiriotis A., Xilouris E. M., Papamastorakis J., Boumis P., Goudis C. D., 2006, *A&A*, 459, 113
- Ness M., Hogg D. W., Rix H.-W., Martig M., Pinsonneault M. H., Ho A. Y. Q., 2016, *ApJ*, 823, 114
- Oppenheimer B. D., Davé R., Kereš D., Fardal M., Katz N., Kollmeier J. A., Weinberg D. H., 2010, *MNRAS*, 406, 2325
- Pérez E., Cid Fernandes R. et al., 2013, *ApJ*, 764, L1
- Pilkington K. et al., 2012, *A&A*, 540, A56
- Rosales-Ortega F. F., Kennicutt R. C., Sánchez S. F., Díaz A. I., Pasquali A., Johnson B. D., Hao C. N., 2010, *MNRAS*, 405, 735
- Rosales-Ortega F. F., Sánchez S. F., Iglesias-Páramo J., Díaz A. I., Vílchez J. M., Bland-Hawthorn J., Husemann B., Mast D., 2012, *ApJ*, 756, L31
- Rubin K. H. R., Prochaska J. X., Koo D. C., Phillips A. C., Martin C. L., Winstrom L. O., 2014, *ApJ*, 794, 156
- Rupke D. S., Veilleux S., Sanders D. B., 2005, *ApJS*, 160, 115
- Rupke D. S. N., Kewley L. J., Barnes J. E., 2010, *ApJ*, 710, L156
- Salim S., Lee J. C., Ly C., Brinchmann J., Davé R., Dickinson M., Salzer J. J., Charlot S., 2014, *ApJ*, 797, 126
- Salim S., Lee J. C., Davé R., Dickinson M., 2015, *ApJ*, 808, 25
- Salpeter E. E., 1955, *ApJ*, 121, 161
- Sánchez S. F. et al., 2012, *A&A*, 538, A8
- Sánchez S. F. et al., 2013, *A&A*, 554, A58
- Sánchez S. F. et al., 2016, *Rev. Mex. Astron. Astrofis.*, 52, 21
- Schaye J., 2004, *ApJ*, 609, 667
- Schaye J. et al., 2010, *MNRAS*, 402, 1536
- Schmidt M., 1959, *ApJ*, 129, 243
- Scoville N. et al., 2007, *ApJS*, 172, 1
- Shapley A. E., Steidel C. C., Pettini M., Adelberger K. L., 2003, *ApJ*, 588, 65
- Skillman E. D., 1987, *NASA Conference Publication*, 2466
- Smee S. A. et al., 2013, *AJ*, 146, 32
- Somerville R. S., Davé R., 2015, *ARA&A*, 53, 51
- Springel V., Hernquist L., 2003, *MNRAS*, 339, 289
- Springel V. et al., 2005, *Nature*, 435, 629
- Steidel C. C., Erb D. K., Shapley A. E., Pettini M., Reddy N., Bogosavljević M., Rudie G. C., Rakic O., 2010, *ApJ*, 717, 289
- Steinmetz M. et al., 2006, *AJ*, 132, 1645
- Stoeck J. T., Keeney B. A., Danforth C. W., Shull J. M., Froning C. S., Green J. C., Penton S. V., Savage B. D., 2013, *ApJ*, 763, 148
- Tinsley B. M., 1980, *Fundam. Cosm. Phys.*, 5, 287
- Tremonti C. A. et al., 2004, *ApJ*, 613, 898
- Tumlinson J. et al., 2011, *Science*, 334, 948
- van der Wel A. et al., 2014, *ApJ*, 788, 28
- van Dokkum P. G. et al., 2013, *ApJ*, 771, L35
- Vazdekis A., Ricciardelli E., Cenarro A. J., Rivero-González J. G., Díaz-García L. A., Falcón-Barroso J., 2012, *MNRAS*, 424, 157
- Veilleux S., Cecil G., Bland-Hawthorn J., 2005, *ARA&A*, 43, 769
- Vila-Costas M. B., Edmunds M. G., 1992, *MNRAS*, 259, 121P
- Vincenzo F., Matteucci F., Belfiore F., Maiolino R., 2016, *MNRAS*, 455, 4183
- Vogelsberger M. et al., 2014, *Nature*, 509, 177
- Wechsler R. H., Bullock J. S., Primack J. R., Kravtsov A. V., Dekel A., 2002, *ApJ*, 568, 52
- Weiner B. J. et al., 2009, *ApJ*, 692, 187
- Werk J. K. et al., 2014, *ApJ*, 792, 8
- White S. D. M., Rees M. J., 1978, *MNRAS*, 183, 341
- Wong T., Blitz L., 2002, *ApJ*, 569, 157
- Yan R. et al., 2016, *AJ*, 151, 8
- Yan R., Bundy K., Law D. R. et al., 2016, *AJ*, 152, 197
- Zahid H. J., Dima G. I., Kewley L. J., Erb D. K., Davé R., 2012, *ApJ*, 757, 54
- Zahid H. J., Dima G. I., Kudritzki R.-P., Kewley L. J., Geller M. J., Hwang H. S., Silverman J. D., Kashino D., 2014, *ApJ*, 791, 130
- Zhu G. et al., 2014, *MNRAS*, 439, 3139
- Zhu G. B. et al., 2015, *ApJ*, 815, 48

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.