

HNO 2019 · 67 (Suppl 1):S10–S42
<https://doi.org/10.1007/s00106-019-0633-7>
 Published online: 7 March 2019
 © Springer Medizin Verlag GmbH, ein Teil von Springer Nature 2019



R. F. F. Cima^{1,2} · B. Mazurek³ · H. Haider⁴ · D. Kikidis⁵ · A. Lapira⁶ · A. Noreña⁷ · D. J. Hoare^{8,9}

¹ Faculty of Psychology and Neuroscience, Department of clinical Psychological Science, Experimental Health Psychology, Maastricht University, Maastricht, The Netherlands

² Adelante Centre of Expertise in Rehabilitation and Audiology, Hoensbroek, The Netherlands

³ Tinnituscenter, Charité—Universitätsmedizin Berlin, Berlin, Germany

⁴ ENT department, Hospital Cuf Infante Santo, Lisbon, Portugal

⁵ 1st Department of Otolaryngology, National and Kapodistrian University of Athens, Athens, Greece

⁶ Malta University, Valetta, Malta

⁷ Sensory systems and neuroplasticity, Aix-Marseille Université, Marseille, France

⁸ National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, Ropewalk House, University of Nottingham, Nottingham, United Kingdom

⁹ Otology and Hearing Group, Division of Clinical Neuroscience, School of Medicine, University of Nottingham, Berlin, Germany

A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment


Contributors

Adrian Agius, Alain Londero, Allison Clarke, Angela Brandão, Annick Gilles, Antonio Romero-Garcia, Aristides Sismanis, Berthold Langguth, Chris van den Dries, Christopher R. Cederroth, Daniela Ivansic, David Stockdale, Deborah Hall, Diogo Ribeiro, Domenico Cuda, Don McFerran, Dyon Scheijen, Elizabeth Marks, Emile de Kleine, Fioretti Alessandra, Gerhard Hesse, Giovanna Baracca, Gordon Sun, Helen Pryce, Isabel Diges, Jan Bulla, Joaquim Ferreira, Joost van Tongeren, Julia Hodson, Kneginja Richter, Loes Schenk-Sandbergen, Luca Del Bo, Lucy Handscomb, Magdalena Sereda, Mary Mitchell, Matthijs Killian, Michael Golenhofen, Myriam Westcott, Niklas Karl Edvall, Nuno T Cunha, Paris Conostas, Paul Van de Heyning, Peter Byrom, Pim van Dijk, R. Arnold, R. Hofman, Remo Arts, Saba Battelino, Sandra Cummings, Sarah Michiels, Susanne Nemholt Rosing, Thomas Nikolopoulos, Tijana Bojić, Tobias Kleinjung, Tony Kay, Vasco de Oliveira, William Sedley, and Yahav Oron.

- Date: 18 February 2018
- Version: 1.3
- Date of revision: 18 February 2023

Contents

		3.1	Introduction
		3.2	Minimum patient assessment
Chapter 1	General introduction		
1.1	Motivation for the guideline	3.3	Further assessment
1.2	Aim of the guideline	3.4	Assessment by questionnaires
1.3	Delineation of the guideline	Chapter 4	Treatment options and referral pathways
1.4	Intended users of the guideline	4.1	Available treatments and evidence
1.5	Main classifications for subjective tinnitus	4.2	Referral options and criteria, triage and a stepwise proposal
1.6	Epidemiology of tinnitus	Chapter 5	Patient information and support
1.7	Pathophysiology of tinnitus		
1.8	Mechanisms of tinnitus awareness and distress	5.1	Confirming knowledge and dispelling myths. Key messages to prepare the patient for treatment and beyond
1.9	Theoretical models of tinnitus in more detail	5.2	Information that should be given to patients
Chapter 2	Methods	5.3	Further information and support
2.1	Introduction	Appendix A	Pulsatile tinnitus management
2.2	Preparatory work		
2.3	Development of the European clinical guideline		
2.4	Consultation rounds and consensus		
Chapter 3	Diagnostics, assessments, and outcomes		



Members of the Guideline Steering Group			
Member (role)	Position and institute(s)/affiliations	Clinical field/research area of interest	Memberships and conflict of interest
Rilana F.F. Cima, PhD (Chair of Steering Group)	Assistant Professor, Maastricht University, Clinical Psychological Science, Section of Behavioural Medicine Clinical Psychologist/Research Coordinator, Adelante Centre for Rehabilitation and Audiology <i>The Netherlands</i>	Tinnitus, CBT, clinical trials, experimental studies, fear-avoidance, associative learning, patient-profiling, standards in health care	Member of the Dutch National Guideline Committee for Tinnitus Expert on advisory board of the Superior Health Council of Belgium: Prevention, Diagnosis and Therapy of Tinnitus Rilana Cima is funded by the Innovational Research Incentives Scheme “Veni-grant” from The Netherlands Organization for Scientific Research for her research in tinnitus Chair of the TINNET Project Working Group I: Clinical
Haúla Haider, MD	ENT/Lecturer at Nova Medical School ENT Consultant at Hospital Cuf Infante Santo <i>Portugal</i>	Tinnitus ENT surgery PhD student with thesis on: Presbycusis, tinnitus and quality of life	Member of ENT Portuguese Association, Member of Otoneurologic Portuguese Association Has received funding from JmelloSaude for PhD Co-Chair of the TINNET Project Working Group I: Clinical
Dimitris Kikidis, MD, PhD	Otolaryngologist, Clinical Fellow and Research Associate, University of Athens <i>Greece</i>	Neurotology, tinnitus, balance disorders, ear surgery	Has received funding as Co-Principal Investigator in two European-funded projects, one for public health policy in regard to hearing aids and one for regeneration of outer hair cells of the cochlea
Alec Lapira, MD	ENT Specialist/Audiological Physician Lecturer Malta University Adjunct Professor, Nova SE University <i>Malta</i>	Tinnitus, balance disorders, voice rehabilitation	Member of the ENT Association of Malta Fellow of the American Academy of Audiology Member of the International Society of Audiology Member of the International Association of Physicians in Audiology
Birgit Mazurek, MD, PhD	Head and Director of the Tinnitus Centre Charité—Universitätsmedizin Berlin <i>Germany</i>	Tinnitus, ENT, CBT, tinnitus research	Member of the German ENT Society Member of the Berlin ENT Society Member of the German Society of Audiology Member of the ADANO Member of the German Tinnitus-Liga e. V. and scientific board member Head of the German Tinnitus Foundation Charité Member of the EUTI
Arnaud Noreña, PhD	Professor, Head of Team: Sensory Systems and Neuroplasticity <i>France</i>	Rehabilitation medicine, otolaryngology, speech and language pathology, psychophysics Psychophysics	Member of French National Centre for Scientific Research
Derek J. Hoare, MRes, PhD	Associate Professor in Hearing Sciences, NIHR Nottingham Biomedical Research Centre, Hearing Sciences, Division of Clinical Neuroscience, University of Nottingham <i>UK</i>	Tinnitus, hyperacusis, audiology, paediatrics, psychological therapy, evidence synthesis	Chair of the British Society of Audiology Tinnitus and Hyperacusis Special Interest Group. Was site PI on a clinical trial of Acoustic CR® Neuromodulation Derek Hoare is funded by the National Institute for Health Research (NIHR) Biomedical Research Centre programme. However, the views expressed in this document are those of the authors and not necessarily those of the NIHR, the NHS or the Department of Health and Social Care

- Appendix B Method for the development of an implementation plan
- Appendix C Somatosensory tinnitus
- Appendix D Characteristics and psychometrics of existing tinnitus health-related quality-of-life instruments
- Appendix E Cognitive therapy and cognitive behavioural therapy

Chapter 1: General introduction

1.1 Motivation for the guideline

Tinnitus involves the percept of a sound or sounds in the ear or head without an external source. Most individuals experiencing tinnitus have a neutral reaction to the percept. However, for some it becomes a problem. Bothersome (distressing) tinnitus might be better described as a negative emotional and auditory experience, associated with, or described in terms of, actual or potential physi-

cal or psychological harm [10]. Objective tinnitus is defined as the perception of a sound which has a physical source generated in or near the ear. An external observer can perceive objective tinnitus. Subjective tinnitus does not involve an identifiable sound source so cannot be heard by examination. It is caused by anomalous activity in the auditory system. Subjective tinnitus is a highly complex condition with a multifactorial origin and, therefore, heterogeneous patient profiles. In most people, tinnitus is not traceable to medical causes. In

Table 1 Frequent tinnitus comorbidities

Hearing and vestibular disorders	Hearing loss (H90.5) Disturbance of auditory perception (H93.2) Hyperacusis (H93.2) Vestibular disorders (H81)
Mood disorders	Adjustment disorder (ICD-10: F43.2) Dysthymia (ICD-10: F34.1) Depressive episode (ICD-10: F32.0, F32.1, F32.2, F32.3): Recurrent depressive episodes (ICD-10: F33.0, F33.1, F33.2, F33.3)
Anxiety disorders	Phobic disorders (ICD-10: F40.) e. g., specific phobia (ICD-10: F40.2) Anxiety disorder (ICD-10: 41.) Generalised anxiety disorder (ICD-10: F41.1) Anxiety and depressive disorder, mixed (ICD-10: F41.2)
Reaction to severe stress and adjustment disorders	Acute reaction to burdening (F43.0) Posttraumatic stress disorder (ICD-10: F43.1) Somatoform disorders Somatisation disorder (ICD-10: F45.0) Hypochondriasis (ICD-10: F45.2) Psychological factors and behavioural factors in another classified disease (ICD 19: F54) Insomnia (G47)

most cases there is no available curative treatment [7, 45]. Standard treatment, assessment, and referral-trajectories are poorly defined, not well established, and often insufficient. The lack of standard guidelines likely leads to untreated, under-, as well as over-treated patients [15, 21, 26, 34, 56]. This leads to increasing complaints, prolonged suffering, and loss of societal participation, health-care overuse, and endless referral trajectories, resulting in an enormous psychological, societal, and economic burden in Europe. There is therefore a need for a European harmonised guideline for the assessment and treatment of tinnitus. Through development and implementation of this guideline, we anticipate that assessment and treatment of tinnitus will be significantly more effective, leading to reduced suffering and frustration for patients, their families, and clinicians alike.

1.2 Aim of the guideline

The main goal of this guideline is to establish uniformity in the assessment and treatment of adult patients with subjective tinnitus. In addition, this guideline aims to establish consistency in policy to optimise referral trajectories and reduce over- and under-assessment and treatment. Guidance for detailed clinical definition and characterisation of cases is

also included. Experts from different disciplines from across Europe have joined forces to develop standardisation procedures for easy, practicable, and meaningful patient profiling. The guideline should be used as a tool to support shared decision-making with patients to facilitate individualised care.

1.3 Delineation of the guideline

Tinnitus is a common auditory symptom, which may result in serious burden particularly when there are comorbidities. Tinnitus can present in many forms. It is necessary that clinicians identify all relevant tinnitus-related factors during tinnitus assessment. Treatment should be proposed based on an assessment that accounts for tinnitus as part of a complex system with intricate interactions between its constituent factors. A classification protocol should identify tinnitus clinically relevant patient profiles and offer a rational path to individualised treatment.

1.4 Intended users of the guideline

This guideline was established for every health professional involved in tinnitus assessment and treatment, including but not limited to: general practitioners; ear, nose, and throat doctors (ENT); neurologists;

audiologists; psychiatrists; psychologists; maxillofacial doctors; nurses; therapists; basic and clinical researchers. It was designed to be accessible to researchers, tinnitus patients and their significant others, patient organisations, policy-makers, and any other relevant stakeholders.

1.5 Main classifications for subjective tinnitus

Acute, sub-acute, and chronic tinnitus

Tinnitus is acute if the patient has experienced it for less than 3 months and is considered sub-acute after 3 months. It is termed “chronic” when the patient has experienced it for 6 months or more. Assessing whether tinnitus is acute or chronic is relevant to the choice of certain treatments.

Possible comorbidities

Comorbidities can be pre-existent or induced by tinnitus. Frequently psychological, psychosomatic, and/or psychiatric comorbidities are associated with tinnitus [42, 68, 77]. Anxiety, depression, and insomnia are commonly found in patients with tinnitus. The higher the level of distress, the more likely comorbid disorders are present [24, 44]. On suspicion of psychological comorbidity, further assessment and treatment should be undertaken by appropriate specialists (psychologist, psychosomatic specialist, psychiatrist, or neurologist). A list of comorbidities that may be observed in tinnitus patients is given in [Table 1](#).

Profile of severity

Tinnitus severity can be graded using multi-item tinnitus questionnaire scores where there is a valid grading system provided. It can also be (more subjectively) graded according to structured medical history. For example, the structured tinnitus interview by Biesinger et al. [8] allows tinnitus severity to be classified into one of four grades based on level and frequency of impairment (see [Table 2](#) for an adapted version of this grading system).

Table 2 Tinnitus grading system

Grade	Description
1	No distress, no impairment
2	Tinnitus impairs (e. g. emotion, cognition, attention, task performance) occasionally/ occurs under stressful situations and mainly in silence
3	Tinnitus regularly impairs (e. g. emotion, cognition, attention, task performance) occurs in several situations
4	Tinnitus constantly leads to impairment (e. g. in emotion, cognition, attention, task and daily life interference) occurs in all situations

Characteristics of tinnitus

Tinnitus can be any sound but it is typically ringing, buzzing, hissing, or tonal. Some patients experience multiple sounds. For some the sound is persistent in quality and for others it changes. It can be constant or intermittent, and heard in one or both ears or inside the head. The sound might be pulsatile (either synchronous with the heartbeat or not) or non-pulsatile, and both may be objective or subjective (see Appendix A for further information on pulsatile tinnitus).

1.6 Epidemiology of tinnitus

Studies of tinnitus prevalence are not free of methodological problems; they use different definitions of tinnitus, or address selected age or professional groups. Most studies report conservative tinnitus prevalence rates to be between 10 and 19% of adults [6, 23]. The phantom auditory sensation of tinnitus is chronically perceived by roughly one in three of elderly adults [1, 59]. Tinnitus is commonly associated with hearing loss, noise exposure, ageing and stress [27, 31, 55, 58, 67] and less often with other otologic, neurologic, infectious, and drug-related effects and other comorbidities [76]. In the tinnitus population the prevalence of hyperacusis (reduced tolerance to everyday sounds) is 40–86% according to different studies [2, 35].

1.7 Pathophysiology of tinnitus

Most of our knowledge on tinnitus pathophysiology originates from animal research. Firstly, tinnitus is related to “aberrant” neural activity (that is not produced by physically measurable sounds from the environment) that is generated at some

level of the auditory system. Secondly, the tinnitus-related signal is interpreted as an auditory percept and can be associated with distress. Tinnitus has been suggested to result from an increased firing rate, increased synchrony between neural discharges, or an aberrant pattern of oscillatory activity (possibly resulting in an increase of neural firing synchrony; [62]). In most cases, tinnitus is believed to be associated with some degree of cochlear damage. There can be cochlear damage not detected by a standard audiogram. Indeed, it has been shown that high-threshold cochlear fibres (that are activated well above absolute threshold) can be uncoupled from their corresponding inner hair cells and/or degenerated after moderate noise trauma [50, 51, 73].

One can distinguish between the peripheral and central mechanisms that can result in the generation of the tinnitus-related activity. Many mechanisms may account for cochlear tinnitus (related to cochlear aberrant spontaneous activity): activation of N-methyl-D-aspartate receptors [25, 66], shift in the operating point of outer hair cells [19, 48, 61], decoupling of outer hair cell stereocilia [49], and increase in the endo-cochlear potential [63]. Central forms of tinnitus are then thought to result from aberrant neural discharges produced by central changes which may be triggered by hearing loss. Many central mechanisms that can account for the generation of the tinnitus-related activity have been proposed [20]. The vast repertoire of mechanisms involved in homeostatic plasticity may account for central hyperactivity after hearing loss. The reorganisation of the cortical tonotopic map may also play a role in tinnitus generation, by increasing synchrony of discharges between cortical neurons. This mechanism is likely

not involved in tinnitus in people without hearing loss [43] or those with a pantonal hearing loss.

Somatosensory inputs can modulate neural activity in the cochlear nucleus [69–71]. The excitatory modulation of these inputs on the auditory pathway is enhanced after hearing loss and can be responsible for the noise-induced central hyperactivity (potentially associated with tinnitus) after noise trauma [18, 75].

Finally, the thalamus may play a direct role in generating tinnitus-related activity. The reduction of sensory inputs is thought to hyperpolarise thalamic neurons, which leads to the generation of bursts of action potentials in the thalamus and cortex. This activity can be maintained by the cortico-thalamic pathway and the activation of the reticular nucleus sending inhibitory inputs back to specific nuclei of the thalamus [52, 53].

1.8 Mechanisms of tinnitus awareness and distress

Tinnitus is not necessarily associated with distress. Indeed, 6–20% of people with tinnitus consider it bothersome [59]. This indicates a partial dichotomy between mechanisms generating the tinnitus-related signal and those responsible for tinnitus distress. It has been proposed that non-auditory brain regions evaluate the tinnitus-related signal. If the tinnitus-related activity is not negatively reinforced, this neural activity is blocked from reaching conscious perception (i. e. habituation occurs). If, on the other hand, the tinnitus-related activity is negatively reinforced (e. g. because tinnitus is concomitant to a life event or associated with negative thoughts), the limbic and autonomic nervous systems are activated, attention is directed to tinnitus, and tinnitus-related distress develops. This model postulates that tinnitus awareness and tinnitus-related distress result from the mechanisms of conditioned reflex [37, 40]. Another model suggests that tinnitus-related activity (enhanced activity in the gamma frequency range) becomes a conscious percept only if it is associated with co-activation of self-awareness and salience brain networks, i. e. anterior cingulate




Fig. 1 ▲ The neurophysiological model of tinnitus




Fig. 2 ▲ The cognitive model of tinnitus

cortex, ventromedial prefrontal cortex, insula, amygdala and parahippocampus [17]. A somewhat different model suggests that irrelevant information is tuned out by an inhibitory gating mechanism. Whereas the thalamic reticular nucleus, activated by subcallosal regions (nucleus accumbens and ventromedial prefrontal cortex), is supposed to block the activity of sensory thalamus through

its inhibitory influence, in this model it is supposed that the inhibitory gating mechanism may not block irrelevant information properly (including the tinnitus-related activity) if part of the circuit such as the subcallosal regions is lesioned [46, 47, 64].

1.9 Theoretical models of tinnitus in more detail

Habituation theory

Habituation theory [29] proposes that the negative interpretation of the tinnitus signal, and related heightened autonomic arousal levels, leads to dysfunctional cognitive processing and thereby distress. Hallam et al. [29] proposed that for most people repeated perception of the tinnitus sound teaches them that it is not worthy of attentional resources. In other words, to function effectively, the brain selects which stimuli to pay attention to and which stimuli to ignore. Hallam purported that most people learn that the tinnitus sound is of low informational value and thus not requiring a reaction. Consequently, tinnitus does not pose a problem for most people living with it. However, tinnitus-related *distress* does occur when these attentional processes are malfunctioning, which is more likely at times of increased stress and arousal, which in turn strains cognitive resources.

Habituation theory has remained largely theoretical, although tinnitus treatment approaches such as relaxation therapy, attention diversion techniques (directing attention away from tinnitus), and stress reduction by means of cognitive restructuring methods (aimed at altering beliefs about the tinnitus) have been based on its main premises. To treat tinnitus distress (or facilitate habituation to tinnitus), it was recommended that stress and arousal levels be reduced and to try and change the meaning of the tinnitus signal for the patient [28]. Research to date has yielded mixed evidence for the validity of habituation theory [7].

Neurophysiological model

The neurophysiological model of chronic tinnitus (■ Fig. 1) is based on the premise that conditioned fear responses elicited by the tinnitus sound are the cause of the tinnitus becoming bothersome [36, 37]. This reasoning stems from animal research in which conditioning paradigms were used to induce tinnitus-like fearful behaviour in rats [38, 39]. The neurophysiological model distinguishes be-




Fig. 3 ◀ The fear-avoidance model of tinnitus (based on the fear-avoidance model of chronic pain [74])

tween three stages. Stage 1 involves generation of the auditory stimulus in the auditory periphery. Stage 2 involves detection of the tinnitus-related signal; and Stage 3 involves the perception evaluation of tinnitus. The neurophysiological model is mainly a model of tinnitus generation/detection, based on neurophysiological mechanisms.

Cognitive model

McKenna et al. [57] propose a conceptual cognitive (behavioural) model of tinnitus distress whereby negative cognitive misinterpretations of the tinnitus signal, distress, and bodily arousal are provoked leading to inaccurate evaluations of sensory activity and distorted perceptions (Fig. 2). It is proposed that the resulting hypervigilance and distorted perception of the tinnitus sound contribute to a feedback cycle that exacerbates negative thinking and thus distress. The model attributes a fundamental role to the negative evaluation of tinnitus. The negative evaluation of the tinnitus can be viewed as comprising primary and secondary appraisals. For example, a person might initially appraise the tinnitus as being threatening to their health, and then make a secondary appraisal of their ability or inability to cope with it.

Evidence exists that cognitive processes, such as interpretation, attention, and memory, are indeed involved in chronic tinnitus suffering [5, 16, 60, 65, 72], and the validity of the cognitive model is currently being tested psychometrically [30].

Fear-avoidance model

The fear-avoidance model of chronic tinnitus (Fig. 3) offers explanatory predictions about both the cognitive processes as well as the behavioural mechanisms of tinnitus [11, 12]. It predicts that individuals perceiving the tinnitus signal are subject to automatic emotional and sympathetic responses. These symptoms are misinterpreted as harmful or threatening. If the signal persists, the coinciding threatening (alarm) states, which indicate malignance of the signal, elicit conditioned (both classic and operant) fear responses, i.e. fear, increased attention, and safety-seeking (avoidance and escape behaviours). Safety behaviours become negatively reinforced through instant decreased fear, which is adaptive in the acute phase. In other words, by avoiding or not exposing themselves to tinnitus-related perceptions, patients learn that their fear instantly diminishes. However, through persistent avoidance of the tin-

nitus, tinnitus-eliciting, or tinnitus-increasing stimuli, fear and fear responses such as hypervigilance and safety-seeking are maintained. Avoidance behaviours subsequently lead to task interference and functional disability [9, 33, 41, 54]. The maintained high threat value of the tinnitus leads to increased tinnitus severity and distress, feeding into an endless cycle of increased disability [14].

A feature of the fear-avoidance model is that it predicts both a maladaptive pathway and an alternative more adaptive pathway. In the more adaptive pathway, a positive or neutral evaluation of the tinnitus (the system accepts it as being benign) results in no fear or low fear of tinnitus, distress, or avoidance behaviours.

These models differ in explaining how classic and operant learning principles contribute to tinnitus suffering. Nonetheless, there exists a large conceptual overlap between them. In both the neurophysiological model and McKenna's conceptual cognitive (behavioural) model it is hypothesised that effortful conscious alteration of negative interpretations will decrease arousal and distress because of tinnitus. Both models place less importance on the behavioural processes. The fear avoidance model, which is based on




Fig. 4 ▲ Individual stages in the development of current guideline: a roadmap. WG working group, MT management team

associative learning principles, offers both explanatory predictions about cognitive processes of change and predictions about behavioural mechanisms. This fear-avoidance approach integrates previous cognitive conjectures within a behavioural framework. This may prove helpful, both for discovering new venues for investigations and as a means of discovering why the cognitive as well as the behavioural treatment approaches are consistently found to be beneficial. It also offers a means of discerning which components work best and for whom.

Empirical data support the fear-avoidance model. Accumulating evidence indicates that a cognitive behavioural treatment, based on this fear-avoidance notion—which targets re-appraisal of, and exposure to, the tinnitus-sound—significantly reduces tinnitus distress, decreases tinnitus suffering, and improves quality and daily functioning

of tinnitus patients [3, 4, 13, 120]. However, the cause–effect relationships of specific learning mechanisms are still unknown [11, 22, 32, 41].

Chapter 2 Methods

2.1 Introduction

Tinnitus is a complex condition with a multifactorial origin. Consensus on clinically relevant patient profiles, standard treatment, assessment, and referral trajectories has not been reached thus far. Additionally, inconsistent results in tinnitus studies, in experimental research, clinical trials, observational and cross-sectional research, represent a barrier to efficient standards in health care for tinnitus. Even though chronic tinnitus complaints represent enormous socio-economic relevance [13, 80–82], research funding is still limited. Hence, there is

increasing need for a European guideline for the assessment and treatment of tinnitus patients. The purpose of this chapter is to describe the methods used to develop this guideline.

Duration and validity

This guideline consists of five chapters containing several modules. Some or all these modules may need revision or extension in the future. The current guideline steering group aims to pursue re-assessment and maintenance of the guideline within 5 years of the initial publication.

Implementation

During the development of this guideline, the practicality and feasibility of the implementation of recommendations across Europe were considered at every stage (■ Fig. 4). Consensus was reached at the beginning of the project that a Eu-

Table 3 Summary of barriers and facilitators identified from a pan-European survey

Barriers	
1	Lack of knowledge about or non-existence of specialised tinnitus clinics or teams makes it difficult for tinnitus patients to find their way to the most appropriate professionals in a country
2	Lack of time or other resources for adequate counselling
3	Lack of time or other resources for professionals responsible for tinnitus patients to be able to adequately assess the distress level of tinnitus patients
4	Lack of multidisciplinary teams and/or availability of psychologists in southern and eastern European countries
5	High variation in available treatment options; more medical–pharmacological treatment in southern and eastern countries. Counselling–rehabilitative approaches more available in northern countries. When many treatment avenues are seen as viable, it may be difficult to reach consensus on what works for whom
6	The use of self-report instruments is much less common in southern and eastern countries
7	There are differences in how patients pay for treatment. If regulatory bodies in health care in a country are unwilling or unable to hold to the restrictions or recommendations stated in a guideline, the chances of implementation of this guideline drastically decrease
Facilitators	
1	Common ground in expert opinion that tinnitus is a central auditory symptom. This offers options for discussions on the definition of tinnitus in a European guideline
2	Consensus across regions on what conditions are relevant to or associated with tinnitus. Harmonisations such as these are to be highlighted where possible to facilitate implementation of a standard guideline
3	Although some minor differences in procedures were reported, most experts agree that otoscopy and pure tone audiometry are used. This finding facilitates discussions on diagnostics to include in the guidelines
4	The most commonly used questionnaire irrespective of region is the Tinnitus Handicap Inventory. This might facilitate discussions on assessment methods to recommend within a guideline
5	The percentage of respondents satisfied with current tinnitus health care in their country in southern and eastern Europe was low; less than half of respondents reported they were satisfied. Health-care professionals are likely to be positive towards progressive guidelines and towards changes in health care for tinnitus

European guideline was to be as adaptive as possible, with the possibility that it can be tailored to resources and needs dictated by the participating countries¹. The practicability of these recommendations and guidelines will need testing beyond the initial publication of the guideline; a method and criteria for the development of an implementation plan are provided in Appendix B.

¹ Belgium, Cyprus, Czech Republic, Denmark, Finland, France, Germany, Greece, Israel, Italy, Lithuania, Malta, The Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovenia, Spain, Sweden, Switzerland, Turkey, United Kingdom.

Responsibility

The responsibility to maintain the guideline, for reassessment purposes and possible future revisions or extensions, lies with Drs. Rilana Cima and Derek Hoare.

2.2 Preparatory work

Activities, meetings, and studies performed for the development of this European guideline were supported by the TINNET—COST Action BM1306 (2014–2018), which aimed to establish a pan-European tinnitus network (researchresearch.net). Consensus was reached in the preparatory phase that the main aim of the TINNET Working Group 1 (WG1) was to develop a consensus-based meaningful, adaptive, and actionable European guideline for the

assessment and treatment of tinnitus patients.

European guideline steering group formation

In 2015 a selection of WG1 members of TINNET were nominated to become members of a guideline steering group for the development of this guideline. The group consists of representatives of all specialties and fields thought to be stakeholders in the clinical practice of tinnitus health care across Europe. All steering group members are responsible for the integral text of this guideline.

Clinical demands and bottleneck analysis (Stage 1)

During the first preparatory (consensus) meeting with the WG1 members of TINNET, the chair (R.C.) inventoried clinical demands and possible bottlenecks. After this first meeting a roadmap was created for the agreed different stages of development of this guideline (■ Fig. 4).

Understanding the state of the art (Stages 2 & 3): a pan-European survey

Tinnitus remains a scientific and clinical problem. In spite of increasing knowledge about its management and treatment, little impact on clinical practice has been observed. There is evidence that prolonged, obscure and indirect referral trajectories persist in usual tinnitus care.

It is widely acknowledged that efforts to change professional practice are more successful if barriers are identified and implementation activities are systematically tailored to the specific determinants of practice. The first step towards the development of meaningful and actionable European guidelines for the assessment and treatment of tinnitus patients involved scoping the existence and current knowledge of standards in tinnitus care. Here, we addressed this by performing a pan-European survey of experts, clinicians, and policy-makers to gain insight into the status quo of expert views on treatment and assessment, standards, and guidelines in tinnitus health care throughout Europe.

Table 4 Summary of recommendations from a systematic review of clinical practice guidelines for tinnitus

<i>Summary of recommendations regarding the assessment of subjective tinnitus</i>	
–	Conduct a physical examination to exclude possible (physical) causes of tinnitus
–	Conduct an audiological assessment
–	Establish the degree to which a patient experiences subjective tinnitus as bothersome or distressing using a validated and reliable questionnaire such as the Tinnitus Questionnaire, Tinnitus Handicap Inventory, or Tinnitus Functional Index
–	In situations where patients appear to be experiencing a degree of distress or difficulties related to living with tinnitus, consider making a referral for an assessment by a psychologist or psychiatrist
–	Variations exist in recommendations regarding the use of imaging studies (e. g. magnetic resonance imaging)
<i>Summary of therapeutic recommendations regarding the treatment of subjective tinnitus</i>	
–	Provide information about tinnitus and treatment options
–	Use hearing aids only when patients have a diagnosed hearing loss
–	Specialised cognitive behavioural therapy for tinnitus should be offered to patients
–	There is a lack of consensus on the use (or otherwise) of sound therapy for tinnitus
–	Prescribed and herbal medicines and dietary supplements should not be used for the sole treatment of tinnitus
–	Transcranial magnetic stimulation is not recommended as a treatment for tinnitus

Table 5 Levels of evidence (Oxford)

Level	Therapy/prevention, aetiology/harm
1a	Systematic review (SR) of randomised controlled trials (RCTs)
1b	Individual RCT (with narrow confidence intervals)
1c	All or none effects
2a	SR (with homogeneity) of cohort studies
2b	Individual cohort study (including low-quality RCT; e. g. <80% follow-up)
2c	“Outcomes” research; ecological studies
3a	SR (with homogeneity) of case–control studies
3b	Individual case–control study
4	Case series (and poor-quality cohort and case–control studies)
5	Expert opinion without explicit critical appraisal, or based on physiology, bench research or “first principles”

Survey results showed that there are significant differences in reports on national health-care structure, tinnitus definitions, characteristics of the tinnitus patient, and management, treatment, and diagnostic options, particularly notable when comparing northern, southern, and eastern European countries. Results indicate seven important barriers to be taken into consideration if a European clinical guideline is to be implementable (Table 3). Additionally, five facilitators were defined. Overall, the findings of this study confirm the need for a European guideline to provide consistency and promote equity of access to services for all tinnitus patients.

Systematic review of existing guidelines

A systematic review was performed to collect all available guidelines and identify fields of agreement and remaining open questions about tinnitus assessment and treatment. The tinnitus assessments (diagnostics and measures), processes and treatment options recommended by the respective guidelines were compared and summarised. Methods are described in brief here and are reported in full by Fuller et al. [79].

Selecting guidelines for the review.

Guidelines were considered eligible for inclusion if they fit the definition, and no

publication date or language restrictions were imposed. Guidelines on objective tinnitus, referral pathways, or a specific type of assessment or treatment procedure were excluded. Guidelines were identified through a systematic search for “tinnitus” and “guideline” using Medline, PubMed, and the Cumulative Index to Nursing and Allied Health Literature, as well as the EMBASE databases. In addition to these, the National Guideline Clearinghouse (www.guideline.gov), National Institute for Health and Clinical Excellence (NICE; <https://www.nice.org.uk/>), Guideline International Network (GIN; <http://www.g-i-n.net/>), and Google were searched, and a hand-search of reference lists of any included guidelines was undertaken. International experts were also contacted to ask if they were aware of any guidelines that had not already been identified from the search results. Final searches were conducted on 2 May 2016. Two reviewers independently screened each search result. Five documents fitting the definition of guidelines were included, from the USA, Germany, Sweden, The Netherlands, and Denmark. Data extraction was conducted in a structured way by two independent reviewers and the quality of each guideline was evaluated using the AGREE II tool [78].

Findings. The absence of guidelines for most countries contributes to the explanation for the variations that exist in assessment and treatment of tinnitus internationally. Across guidelines, differences in recommended assessment procedures tend to relate to specific techniques, questionnaires, diagnostic tests, or types of scanning techniques rather than to the assessment of tinnitus severity, hearing loss, psycho-social problem(s), and the presence of severe physical pathology causing the tinnitus. Consensus exists on the need to exclude a physical cause of tinnitus, conducting an audiometric assessment of the patient, using standardised questionnaires to measure degrees of tinnitus-related distress and, when relevant, making referrals for further psychological assessment. Summary recommendations emerging from the systematic review are provided in Table 3.

Table 6 Levels of recommendation

Strength of recommendation	Valence of recommendation	Description
Strong recommendation Most or all individuals will be best served by the recommended course of action	For	Level 1a, 1b, or 2a evidence that the desirable effects of an intervention outweigh its undesirable effects
	Against	Level 1a, 1b, or 2a evidence that the undesirable effects of an intervention outweigh its desirable effects
Weak recommendation Not all individuals will be best served by the recommended course of action	For	Level 2b, 2c, or 3a evidence that the desirable effects of an intervention probably outweigh its undesirable effects
	Against	Level 2b, 2c, or 3a evidence that the undesirable effects of an intervention probably outweigh its desirable effects
No recommendation	Not applicable	Only level 3b, 4, or 5 evidence available, or highest level of evidence shows no clear trade-off between desirable effects of an intervention and its undesirable effects, or individual patient's reactions to undesirable effects are likely to be so different that it makes little sense to think about typical values and preferences

2.3 Development of the European clinical guideline

Results from the survey and the systematic review of guidelines were used to select and evaluate the modules included in Chap. 3, “Diagnostics, assessments, and outcomes,” and Chap. 4, “Treatment options and referral pathways”.

Inclusion

We included assessment methods and diagnostic tools where there was evidence of their use and indication that they are clinically useful (see Chap. 3 for further details). Treatment methods (treatment elements or protocols, devices, or procedures) were included if high-level research evidence (randomised clinical trials, meta-analyses) was available and/or a need existed that the guideline informed clinical practice (i.e. practices identified in the survey results; see [Table 4](#)).

Exclusion

Treatment methods (treatment elements or protocols, devices, or procedures) were not included when not supported by high-level research evidence and when only weak evidence was available (case reports, case control, and retrospective studies), and/or the need existed that the guideline informed clinical practice

(i.e. practices identified in the survey results). Further assessment methods, diagnostic tools, treatment methods (treatment elements or protocols, devices, or procedures) may be included in future revisions or extensions of this guideline if new research evidence becomes available or a need to inform new practices is identified.

Levels of evidence

Quality of evidence for the treatment methods (treatment elements or protocols, devices, or procedures) included in Chap. 4 was guided by the Oxford Centre of Evidence method² (see [Table 5](#)).

Levels of recommendation

The recommendation level for each treatment method (treatment elements or protocols, devices, or procedures) included in Chap. 4 was guided by the GRADE system³ ([Table 5](#)), whereby only high-level evidence (Levels 1a–2b; randomised controlled trials [RCTs] and systematic reviews) was considered in making a recommendation for or against treatment. Where only low-

level evidence existed (2c–5), we automatically make a judgment of “no recommendation” (see [Table 6](#)).

2.4 Consultation rounds and consensus

Consensus method on the drafts of the clinical European guideline

Initial drafts of this guideline were subjected to review in two consultation rounds. The comments from each round were aggregated and addressed across three consensus meetings of the steering group (held in July and November 2017, and in January 2018).

Consultation round 1 involved only members of TINNET, being:

- Members of WG1
- Members of the TINNET management committee
- Representatives of all other TINNET working groups

Consultation round 2 involved stakeholders outside TINNET, being:

- Patient representatives/associations
- Professional associations/national professional representatives
- Committee members of existing national guidelines
- Policy-makers
- EU commission members

Contributors in both consultation rounds were invited to comment on consecutive drafts of the guideline. For each comment submitted, contributors were asked to rate whether they felt their comment reflected what they considered to be an essential change to the guideline (rated as 1), or was a recommend change (rated as 2), or was simply a note for the author to consider (rated as 3).

Consensus meetings

In the consensus meetings, steering members voted whether to implement a change to the guideline based on the comment. Contributors received feedback on all comments. The feedback included whether the change was made or not. Where changes were not made in response to a comment (i.e. there was author consensus not to imple-

² <http://www.cebm.net/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/>.

³ <http://www.gradeworkinggroup.org/>.

Table 7 Some known conditions associated with tinnitus

Site of affection	Associated conditions
Outer ear	Wax blockage of outer ear channel Otitis of outer ear channel Obliterative exostoses
Middle ear	Otitis media with effusion Otosclerosis
Inner ear	Presbycusis Noise-induced hearing loss Ménière's disease Sudden hearing loss Acoustic neuromas
Muscular	Palatal myoclonus Tensor tympani myoclonus Patency of eustachian tube
Cardiovascular	Glomus jugular or glomus tympanicum Mitral or aortic stenosis
Pharmacological	Benzodiazepines withdrawal Induction from ototoxic drugs
Metabolic	Hyperthyroidism Diabetes Hypertension
Haematologic	Anaemia
Arthrogenous	Dysfunction of temporomandibular joint Cervical dysfunction
Impairment of cognitive–emotional reaction system	Concentration disturbance – Attentional problems – Memory deficit – Executive function deficit Loss of control/helplessness/resignation Dysfunctional thoughts: catastrophising – Fear reactions – Safety behaviours (avoidance)
Psychological/psychiatric	– Psychological trauma – Distress – Major affective event – Depression – Anxiety/panic disorder
Trauma	– Traumatic brain injury – Neurosurgical

ment a change based on the comment), a rationale for the decision was provided.

In total, 395 comments on the first draft were submitted in consultation round 1 by 33 within-TINNET contributors. In the second consultation round, an additional 25 outside-TINNET contributors submitted 265 comments on the second draft.

Authorisation

All contributors were asked to indicate if they did not want their details to be included in the contributor list of the final version of the guideline.

Chapter 3 Diagnostics, assessments, and outcomes

3.1 Introduction

Many factors can contribute to the onset of tinnitus. It is a symptom associated with multiple medical disorders (Table 7). In addition to hearing-related causes, other potential causes must be individually identified or excluded. In most cases, the aetiology of tinnitus is unknown, and many clinical approaches are dedicated to helping patients cope with their tinnitus rather than to treating the cause [37, 62, 84].

When referring patients for a specific diagnostic algorithm, for example in the case of acute tinnitus, the medical necessity as well as financial cost must be considered in each individual case. Instead of using rigid diagnostic steps, it is recommended to choose a diagnostic pathway based on the patient's history and on basic diagnostics (for an example of a diagnostic pathway, see the Tinnitus Research Initiative flowchart at www.tinnitusresearch.org).

The following sections outline recommended progressive levels of diagnostics and assessment of tinnitus. The content and structure of these sections were informed by multiple sources in two steps. First, the authors reviewed diagnostics and assessments that are currently used across Europe [15], and the recommended diagnostics and assessments included in existing national clinical practice guidelines (USA, Denmark, Sweden, Germany, and The Netherlands; Fuller et al. [79]). An initial proposal was thereafter agreed by consensus of the authors based on knowledge of current use in clinical practice specifically for tinnitus, and therefore of the need to provide endorsement of those procedures considered safe and clinically useful (and exclude those that were considered not). This section was heavily revised according to comments received in both rounds of expert consultation.

3.2 Minimum patient assessment

A comprehensive patient history is the foundation of diagnosis, accurately grading tinnitus severity and identifying relevant comorbidities. To exclude treatable medical conditions, e.g. otitis media, otosclerosis, wax blockage of outer ear channel, or acoustic neuroma among other putative causes, a detailed tinnitus characterisation should include whether tinnitus is objective (can be heard by an external observer) or subjective (can only be heard by the patient), perceptual characteristics of the tinnitus sound (tonality, pitch, loudness), temporal properties (pulsatile or not, constant, intermittent, or fluctuating), location (in one or both ears, or in the head), and severity (assessed by score on suggested

measures, see Sect. 3.4 “Assessment by questionnaires”).

It is crucial to perform a detailed clinical examination/history of the tinnitus patient. There are causal diagnostics and severity-oriented diagnostics.

Tinnitus history and clinical examination

1. Tinnitus history

- Onset: Since when is tinnitus perceived; what are important associated clinical factors (noise trauma, stress, recent events, acute illness, other); was it a sudden onset or did symptoms start gradually with a continuous increase?
 - Modulation: Can the tinnitus percept be modulated by: orofacial, cervical or eye movements, head positions, movements of the jaw, tension of jaw muscles, physical exertions?
 - Severity/impact of the tinnitus: Is the tinnitus bothersome/interfering with daily life (sleep difficulties, task interruptions, fearful reactions, cognitive-attentional problems, negative affect). A questionnaire should be used to establish the degree to which a patient experiences subjective tinnitus as bothersome or distressing (see Sect. 3.4 for more details). Furthermore, the level of tinnitus awareness is of importance: Can tinnitus be perceived only in silence or also in noise; is the tinnitus easily masked or amplified by ordinary background noise; are there changes in tinnitus loudness?
2. Thorough audiological history and prioritisation: Assessment of hearing loss, perceived “ear fullness” (pressure), sensitivity to normal sound (or hyperacusis), problems in balance/dizziness/vertigo
 3. Medical history: ear, nose and throat, orthopaedic, cervical, dental, jaw, internal medicine (thyroid, hypertension, anaemia), mental disorders (psychological, psychiatric)
 4. Presence of comorbidities/drug history/medications; ototoxic drugs (e.g. chemotherapy, antimalarial drugs); long-term pharmacological

consumption (e.g. antidepressants, anxiolytics)

5. Relevant personal history, occupational history, hobbies/leisure activities, noise exposure, head/neck trauma, social support status, education, recent life events

Essential primary diagnostic steps

Conduct a thorough physical–medical assessment to exclude possible (physical) causes of tinnitus:

Complete ear, nose, and throat examination, especially otoscopy (preferably micro-otoscopy) to exclude presence of wax, tympanic membrane rupture, otitis media with effusion, chronic otitis media, retro-tympanic mass or any other pathology.

Special consideration should be given in rare tinnitus causes (e.g. palatal myoclonus, temporomandibular joint [TMJ] disorders).

A comprehensive diagnostic investigation should include pure tone audiometry, speech audiometry, and evaluation of the perceptual quality of tinnitus (e.g. loudness, pitch, and minimum masking estimations), sound tolerance, tympanometry, and acoustic reflex including auscultation of the ear and carotid artery in pulsatile tinnitus, as clinically indicated. Care must be taken in performing loudness-based tests, particularly where there is evidence of recent fluctuations in loudness or intensity of the patient’s tinnitus.

3.3 Further assessment

Further investigations or referrals in special cases

Only to be considered if clinically indicated:

- Auditory brainstem responses and/or magnetic resonance imaging (MRI) in cases of unilateral tinnitus and/or asymmetric hearing loss consider
- High-frequency audiometry in cases of tinnitus with normal hearing at standard (conversational) frequencies
- Further sound tolerance assessment (e.g. loudness discomfort level) for sound sensitivity grading or hearing aid settings

- Residual inhibition to evaluate short-term effects of sound on the tinnitus
- Transient-evoked otoacoustic emissions and/or distortion product otoacoustic emissions in cases of normal standard audiogram and suspicion of cochlear dysfunction
- Caloric testing, and vestibular evoked myogenic potential, as indicated in cases of dizziness, vertigo, or balance problems
- Functional cervical diagnostics in a quiet environment for detecting tinnitus modulations in somatosensory tinnitus. Consider imaging of cervical spine in cervical pathology associated with somatosensory tinnitus (see Appendix C for further information on somatosensory tinnitus)
- Dental examination (including TMJ) in a quiet environment for detecting tinnitus modulations in TMJ dysfunction or bruxism
- MRI of the brain in abnormal auditory brainstem response or abnormal vestibular evoked myogenic potential

3.4 Assessment by questionnaires

Tinnitus severity in terms of distress/impact

Tinnitus severity can be defined as a function of the level of distress or impact that tinnitus has on the individual. For a small proportion of patients (5–8%) tinnitus is severely distressing and therefore disabling [83]. Since distress refers to the general aversive state, instruments to measure this construct usually include sub-domains which are hypothesised to contribute to tinnitus severity. These instruments are therefore hybrid in that they measure several concepts as a means of capturing tinnitus distress.

There are several instruments in use for assessing level of severity of tinnitus complaints. In a review on disease-specific health-related quality-of-life (HR-QoL) instruments used to measure outcomes in tinnitus trials, six commonly used HR-QoL tinnitus instruments were identified [89].

The Tinnitus Handicap Inventory (THI; [94]) was developed to measure the impact of tinnitus on daily life. It has three subscales; functional, emotional,

and catastrophic responses to the tinnitus. Both the overall questionnaire and the functional and emotional subscales show good internal consistency. However, a unifactorial structure was found in subsequent validation studies [90].

The Tinnitus Questionnaire (TQ; [88]) has six domains; emotional distress, cognitive distress, intrusiveness, auditory and perceptual difficulties, sleep disturbances, and somatic complaints because of the tinnitus. The TQ items are internally consistent; the subscales lack internal consistency, however.

The Tinnitus Reaction Questionnaire (TRQ; [95]) was developed to measure distress related to tinnitus. It has four sub-scales: general distress, interference, severity, and avoidance of the tinnitus.

The Tinnitus Severity Index (TSI; [92]) was introduced as a measure of how much tinnitus negatively impacts a patient's life and how bothersome patients perceive their tinnitus to be. Two items specifically measure how much tinnitus interferes with daily life activities.

The Tinnitus Handicap Questionnaire (THQ; [91]) was intended to measure patients' perceived degree of handicap due to tinnitus. The THQ has three domains: physical health/emotional status/social consequences, hearing and communication, and personal viewpoint on tinnitus. Seven items specifically address the interference of the tinnitus on daily activities; four of which address hearing difficulties, two items address social interactions and one item addresses sleep difficulties because of the tinnitus. The THQ subscales fail on internal consistency however.

The Tinnitus Severity Questionnaire (TSQ; [85]) is a short, unified measure, with two items specifically addressing interference of the tinnitus, one item on sleeping habits and one on impairment of concentration.

More recently, the Tinnitus Functional Index (TFI) was developed as a new measure of the severity and negative impact of tinnitus, both for use as a diagnostic tool and for measuring treatment-related changes in tinnitus [93]. The TFI is a multi-domain questionnaire, measuring tinnitus-related distress/severity as a function of pre-

dominantly psychological constructs, such as attention, worry, anxiety, depression as well as the more functional constructs such as hearing, social life, and activity level [86].

The TQ and the THI are widely used in clinical practice and clinical trials [87]. Additionally, almost all existing clinical practice guidelines [79] recommend using the Hospital Anxiety and Depression Scale [96] to assess negative affect coinciding with or reactionary to tinnitus.

Appendix D summarises the characteristics and psychometric properties of the seven tinnitus-specific instruments described here.

Recommendation

Tinnitus patients who report complaints/show decompensation (grade 2 and higher; see Table 2) should be evaluated with at least one measure of tinnitus-related disability, such as the TQ or THI.

Chapter 4 Treatment options and referral pathways

4.1 Available treatments and evidence

Informed knowledge

Clinicians should educate patients with tinnitus about treatment strategies. For an extended presentation of the information that should be conveyed, see Chap. 5.

Drug/pharmacological

Weak recommendation against

There is no evidence for the effectiveness of drug treatments specifically for tinnitus but evidence for potentially significant side effects. Recommendation is based on systematic reviews and randomised trials.

It is common that treatment of acute tinnitus is given according to treatment of acute sudden hearing loss. However, in both cases the evidence base for treatment is scarce [99]. Therefore, if tinnitus occurs acutely without hearing loss, the standard cortisone therapy is not recom-

mended. Present psychosomatic factors can play a decisive role [118]. Therapeutic approaches such as intratympanic steroid treatment have no effect on tinnitus [168]. Any increase in tinnitus severity or distress in chronic tinnitus should not be treated as new-onset tinnitus. This should be regarded and treated as a fluctuation of chronic tinnitus [119].

For chronic tinnitus, many classes of drugs have been used or trialled, including various anti-arrhythmics, anticonvulsants, anxiolytics, glutamate receptor antagonists, antidepressants, muscle relaxants, and others [45], with little evidence of benefit over harm [169]. The Cochrane review of antidepressants for tinnitus [100] identified six RCTs (610 patients) on the topic. Only one study was judged to be of high quality. This study compared the effect of Paroxetine (a serotonin re-uptake inhibitor) with placebo, finding no significant difference in effect between groups. No effect was seen for trazadone (serotonin antagonist and re-uptake inhibitor) and a small effect was seen for tricyclic antidepressants, but the reviewers concluded this could have been due to methodological issues in the studies. Side effects were commonly reported including sedation, sexual dysfunction, and dry mouth. Nonetheless, antidepressants are often successfully applied in the treatment accompanying depression and anxiety, not for improvement of the tinnitus.

Jufas and Wood [131] provided a systematic review of benzodiazepines for tinnitus also finding six clinical trials which examined the use of diazepam, oxazepam, and clonazepam. There were mixed results across studies and methodological issues which reduced confidence in the estimate of effect they reported. Thus, they concluded that benzodiazepine use for subjective tinnitus does not have a robust evidence base and that these drugs must be used with caution because of serious side effects.

No drug can generally be recommended for the treatment of chronic tinnitus. However, psychiatric comorbidities associated with tinnitus (anxiety, depression) may need drug treatment. Antidepressants should not be

prescribed to tinnitus patients without the diagnosis of depression.

Hearing loss interventions

Cochlear implants. Despite the relatively limited number of cochlear implant users, there are many studies of their effects on tinnitus. In a systematic review of the effects of cochlear implantation on tinnitus in patients with bilateral hearing loss, five studies were found which reported on changes in tinnitus after implantation [157]. Based on tinnitus questionnaire scores, the review found total suppression of tinnitus in 30–37% of patients, a decrease in 29–72% of patients, no change in 0–30% of patients, and a worsening in 0–25% of patients across studies. Of course, RCTs are not applicable in this context. Small case-control studies (3b) have shown the efficacy of cochlear implantation in patients with unilateral deafness and persistent, bothersome tinnitus. Hence, larger studies are necessary to confirm these findings.

No recommendation for (tinnitus); recommendation for (deafness)

Cochlear implantation is recommended only for patients meeting the hearing loss criteria for candidacy. Recommendation for tinnitus based on evidence for safety but low-level evidence of effectiveness

Hearing aids. Hearing loss is one of the most prevalent chronic diseases and causes of disability [176]. The consequences of hearing loss in the overall health condition of the people suffering from it are significant. It has been suggested that the reduced physical and mental activity and secondary social isolation caused by hearing loss [98] increase the risk of cognitive decline/dementia [138], mental illness [140] and depression [105, 140]. Although tinnitus has been strongly associated with hearing loss, the degree of hearing loss cannot linearly predict tinnitus severity. Only 50% of patients with hearing loss experience tinnitus including many patients who are profoundly deaf [133, 156]. In

addition, 10% of tinnitus sufferers have normal pure tone audiometry [161, 162]. Furthermore, tinnitus occurs in different percentages in groups of patients with various causes of hearing loss ranging from 30% (in ototoxicity) to 90% (in acoustic trauma). The significant benefit of hearing aids for hearing difficulties have been demonstrated in RCTs [111, 128, 144].

It has also been suggested that hearing aids reduce tinnitus awareness, and thereby stress [109], and reduce central auditory gain [152] and homeostatic hyperactivity [177], implicating them in tinnitus. It has been hypothesised [109] that increasing bandwidth (the frequency range of sounds amplified) may improve effectiveness. Combination hearing aids (including amplification and sound generator in the same device) are another option for patients who may benefit from both amplification and passive sound stimulation. Yet there is minimal high-level evidence for the efficacy of hearing aids for tinnitus in systematic reviews; Hoare et al. [124] included just one RCT [151] which found hearing aids to be beneficial but equally effective to sound generators for tinnitus. Two subsequent RCTs compared hearing aids with combination hearing aids [116] and conventional hearing aids with combination hearing aids or deep-fit hearing aids [117] in patients with hearing loss and tinnitus. Both trials concluded that all devices offered some equivalent benefit for tinnitus. Hesse [119] included lower-level evidence studies in their systematic review but found study results to be contradictory and concluded that convincing prospective studies are required.

Weak recommendation for

Hearing aids are recommended for the management of hearing loss and should be considered as an option for patients with tinnitus and hearing loss. Recommendation is based on evidence of effectiveness and safety in RCTs of hearing aids for hearing loss and tinnitus, and systematic reviews considering hearing aids for tinnitus.

Hearing aids should not be offered to tinnitus patients without hearing loss. Tinnitus might be a parameter to be considered in hearing aid fitting and consequent relevant decision-making.

Neurostimulation

Neurostimulation treatments are hypothesised to alter tinnitus-generating neural firing. They can be invasive or non-invasive, and use electromagnetic, electrical, or sound stimuli. However, the precise neural mechanism by which changes occur at both local and network levels is not fully understood [123, 158]. Moreover, with non-invasive treatments, the precise area of the brain to be stimulated is unknown. Non-invasive treatments include transcranial electrical stimulation, vagus nerve stimulation (transcutaneous), repetitive transcranial magnetic stimulation (rTMS), and acoustic coordinated reset (CR) neuromodulation. Invasive treatments include vagus nerve stimulation (implantable device), cortical surface stimulation, and deep brain stimulation.

Transcranial electrical stimulation. Transcranial direct current stimulation (tDCS) delivers low-level direct current (about 0.5–2 mA) via scalp electrodes to the cortex. Thereby, some current is conducted through the scalp and some flows into the cerebral cortex where it is hypothesised to increase or decrease cortical excitability (depending on the polarity) in the brain regions where it is applied. It was first proposed as a treatment for tinnitus by Fregni et al. [112]. The most recent systematic review of tDCS included 17 studies but only two RCTs [166]. It concluded that there was insufficient evidence to determine whether tDCS was effective for tinnitus. The review called for further RCTs of tDCS and studies involving variations to the stimulation protocol. Many RCTs of tDCS have subsequently been conducted [101, 102, 149, 163, 172], which report it to be safe but with little or no effect on tinnitus.

Transcranial alternating current stimulation (tACS) involves the delivery of alternating current (constant polarity changes) between electrodes placed on

the skin over cortical regions of interest. It is hypothesised to affect up- and down-regulation of synapses, possibly affecting change in oscillatory cortical activity. There are few studies investigating tACS. One randomised study concluded there are no effects on tinnitus [172].

No recommendation

There is evidence for safety but no evidence for the effectiveness of transcranial electrical stimulation for tinnitus. Recommendation is based on systematic review and RCTs.

Vagus nerve stimulation. Stimulation of the vagus nerve is a means of stimulating the cholinergic nucleus basalis, which in turn has been shown to induce sustained changes in cortical organisation. By this mechanism, vagus nerve stimulation, paired with sound stimuli (to promote reorganisation in the auditory cortex), is a hypothesised treatment for tinnitus. Experimental studies have examined the safety and efficacy of vagus nerve stimulation, both direct (i.e. implanted electrode) and transcutaneous, paired with acoustic stimulation for tinnitus [108, 129, 134, 136, 178]. Tyler et al. [171] conducted a prospective randomised double-blind controlled pilot study of the effects of direct vagus nerve stimulation paired with tones on tinnitus. They reported high compliance, mild, well-tolerated adverse effects, but no significant between-group difference in tinnitus at the end of their 6-week randomisation period.

No recommendation

There is evidence for safety but insufficient evidence that vagus nerve stimulation treatments have effects on tinnitus. Recommendation is based on the lack of RCTs or systematic review.

Repetitive transcranial magnetic stimulation. Repetitive transcranial magnetic stimulation (rTMS) uses strong electric current generated within a coil to create fast-oscillating magnetic fields. When used in treatment, the coil is placed next to the head over the target brain area.

It is hypothesised that the energy from the magnetic fields penetrates the skull to cause depolarisation of the superficial cortical neurons; rTMS for tinnitus has been studied extensively. The most recent systematic review [165] included 15 studies and concluded on a significant effect of treatment. However, high variability in study design and reported outcomes was noted and thus the review concluded the need for large-scale trials and replication studies. Safety was not reported in this review. A Cochrane review [143] included five RCTs and concluded that (1) there was limited support across studies that rTMS was beneficial, and (2) there was insufficient information to conclude it was safe in the long term.

Recommendation against

No consistent evidence that repetitive transcranial magnetic stimulation is effective for tinnitus and no evidence that it is safe in the long term. Recommendation is based on systematic reviews.

Acoustic coordinated reset (CR[®]) neuromodulation. Acoustic CR[®] neuromodulation is a sound therapy involving a randomised sequence of four “phase resetting” tones adjusted to the patient’s dominant tinnitus pitch that are hypothesised to generate a lasting desynchronisation of the pathological brain rhythms causing tinnitus. A systematic review of acoustic CR[®] neuromodulation included eight studies [175]. It concluded that the available evidence indicates the treatment to be safe but that there is insufficient evidence of its effectiveness for clinical implementation of this treatment. The review also concluded that the hypothesised mechanism of effect is unproven.

No recommendation

Acoustic CR[®] neuromodulation is safe but there is no high-level evidence of effectiveness. Recommendation is based on systematic review.

Invasive neurostimulation treatments. Invasive forms of tinnitus treatment are highly experimental and span vagus

nerve stimulation with an implanted device, chronic electrical vestibulocochlear nerve stimulation, brain surface (extradural) implanted electrodes, and deep brain neural stimulator implantation. That they are invasive means they are not a viable option for widespread use. Research to date is limited to a small number of cases and in each the precise neural mechanism by which changes occur at both local and network levels is not fully understood (for a comprehensive review, see Hoare et al., [123]). There are no RCTs or systematic reviews to date.

No recommendation

There is no high-level evidence for the effectiveness or safety of invasive treatments for tinnitus. Recommendation is based on lack of RCTs or systematic review.

Cognitive behavioural therapy

Strong recommendation for

There is high-level evidence for the effectiveness and safety of CBT for tinnitus. Recommendation is based on systematic review and one further RCT.

Cognitive behavioural therapy approaches share the premise that human suffering (psychological distress) and resulting problems are based in malfunctioning information processing, emotional reactivity, and behavioural mechanisms (see Appendix E for further general information on CBT). The CBT approaches have led to a plethora of evidence-based cognitive behavioural treatments for mental and somatic health disorders [127]. Cognitive behavioural therapy is an integrative and pragmatic therapy where the aim is to modify dysfunctional behaviours and beliefs to reduce symptoms, increase daily life functioning, and ultimately promote recovery from the disorder [110]. Confusion often exists about the differences between cognitive therapy and CBT. Since CBT stems from the convergence of two distinct theoretical schools, the radical behavioural school (first wave)

and the cognitive school (second wave), CBT entails a diversity of both cognitive and behavioural principles and methods, and usually a combination of these are used in therapeutic sessions. Therefore, both cognitive and behavioural treatment elements can be found when reviewing CBT procedures in general and thus in tinnitus intervention/treatment research as well.

Cognitive behavioural theory and treatment have been applied in tinnitus research for decades and the results of the effectiveness of CBT approaches for tinnitus have been shown to vary in decreasing tinnitus severity/distress, tinnitus-related fear, tinnitus disability, and tinnitus-related cognitive problems and in improving daily life functioning [13, 120, 126, 139, 141]. Establishing the effectiveness of CBT in tinnitus health care and research is difficult because patients report to suffer in various life domains. In addition to general problems with daily functioning because of concentration difficulties and sleep deprivation, despair, depression, fear, and worry are amongst the most incapacitating. Disagreement still exists on what tinnitus-related domains and outcomes to measure, why, and how [122], and in the research literature there is as of yet no standardisation of outcome selection. Additionally, often the investigated tinnitus CBT approaches vary in number of treatment sessions, hours spent in therapy, group versus individual formats, face-to-face versus Internet- or book-based self-help therapies, combinations of different treatment elements, and tinnitus diagnostics and outcome assessments.

Since the most recent Cochrane review of CBT for tinnitus was published [139], a new Cochrane CBT review protocol has been published [22]. This review will include more recent RCTs and comply with the latest Cochrane standards.

The most recently published review of CBT interventions for tinnitus was a historical and narrative overview in which a range of study designs in addition to RCTs were included, but one in which neither a risk of bias assessment was undertaken, nor a meta-analysis conducted [84]. These methodological issues make

it harder to draw conclusions about the strength of any treatment effects and risks of bias in the evidence included in the narrative synthesis. Cognitive behavioural therapy for tinnitus (CBT4T) often includes a combination of several elements (such as education, counselling, exposure, mindfulness, relaxation, hearing rehabilitation). In a large RCT it was found that specialised CBT for tinnitus showed significant better group differences in improvement in quality of life ($d=0.24$), decreasing severity of tinnitus ($d=0.43$) and tinnitus disability ($d=0.45$), as well as decreasing depressive and anxious symptoms, when compared with general audiological counselling and diagnostics only [13].

Despite the afore-mentioned limitations and the need to be cautious about the exact effectiveness of CBT for tinnitus in general, at present, a specialised stepped-care CBT4T [13] is the only available therapeutic health-care intervention for tinnitus supported by a high-quality clinical trial. Stepped-care CBT4T has been implemented across several Dutch clinical centres as the cost-effective treatment option. Additionally, stepped-care CBT4T is generally well received by patients and is potentially a cost-effective means for reducing the reactivity [80]. The treatment can be defined as a stepped-care⁴ multimodal CBT4T approach in which audiological diagnostics, treatment and consultation as well as CBT-treatment elements are combined.

Self-help CBT interventions (Internet-based or otherwise) appear efficacious in decreasing tinnitus distress when compared with passive control conditions, and less so when compared with active face-to-face CBT treatment [147]. Additionally, treatment attrition in trials of self-help (Internet-based or otherwise) CBT interventions is high. Nonetheless, CBT in a self-help format might be a useful alternative to support tinnitus patients who are unable or unwilling to take part in a face-to-face CBT treatment.

⁴ A stepped-care approach is a framework for organising health services based on individual patients' needs, with a gradual increase in the intensity of the care at each level [174].

Where there are pragmatic barriers and/or lack of resources, an initial step of CBT treatment might be performed by a competent non-psychological professional provided there is appropriate support.

Tinnitus retraining therapy

No recommendation

There is evidence for safety but little high-level evidence for the effectiveness of TRT. Recommendation is based on availability of one RCT and two systematic reviews.

One widely used treatment is tinnitus retraining therapy (TRT), which is based on the neurophysiological model of tinnitus [37]. Tinnitus retraining therapy is a specific implementation of general tinnitus habituation therapy, which utilises directive counselling to decrease the negative tinnitus-evoked reactions and sound to decrease the strength of tinnitus signal [130]. The principal goal of TRT is to achieve habituation of tinnitus through the retraining of the brain [130, 164]. It means that owing to the high level of plasticity of the central nervous system, it is possible to reduce the responsiveness to repeated stimulation with neutral sound stimuli and through the counselling [164]. In this process, the limbic system and autonomic nervous system are the main systems responsible for negative tinnitus-evoked reactions, because those areas are activated when one stimulus is associated in the category of unpleasant or dangerous stimuli, which results in reactions of stress, anxiety, panic attack, or loss of well-being (*fight, flight, or freeze*). But, tinnitus without negative association leads to the extinction of a response to tinnitus. Thus, the goal of TRT is to prevent tinnitus from activating the limbic system and automatic nervous system—habituation of reaction—and when the habituation of reaction is fully achieved, the patient does not experience negative tinnitus-evoked reaction. After this, the cerebral cortex—habituation of perception—is automatically activated, because the brain habituates to all unim-

Table 8 Categories of tinnitus retraining therapy for patients with tinnitus and hyperacusis

Category	Impact on life	Tinnitus	Subjective hearing loss	Hyperacusis	Prolonged sound induced exacerbation	Treatment
0	Low	Present	Not present	Not present	Not present	Abbreviated version of counselling
1	High	Present	Not relevant	Not present	Not present	Full counselling and sound therapy with sound generators
2	High	Present	Significant presence	Not present	Not present	Full counselling and sound therapy with combination instruments
3	High	Not relevant	Not relevant	Present	Not present	Full counselling with stress issues related to hyperacusis and sound therapy using sounds generators and hearing aids
4	High	Not relevant	Not relevant	Present	Present	Full counselling, sound therapy with sound generators set at the threshold

portant stimuli. If the patient achieves this habituation of perception, tinnitus is blocked before it reaches the consciousness level and the patient does not hear tinnitus [164].

Based on a medical evaluation of tinnitus, patients are placed into one of five general categories that guide the treatment recommended (Table 8). Each one of five categories is associated with a specific variant of TRT treatment, and all patients receive counselling and sound therapy, with substantial differences. Sound therapy has an important role in TRT. Specifically, sound therapy acts by providing the auditory systems with constant neutral signs with sound generators, hearing aids, or background noise. This decreases the contrast between tinnitus-related neural activity and background activity. Furthermore, the sound therapy interferes with the detection of tinnitus signal and decreases the gain within the auditory pathways [119, 135, 164].

The Cochrane review of TRT [153] found only one trial that met their inclusion criteria, concluding that the trial was of low quality and no final conclusions concerning the efficacy of TRT can be drawn. The same single study was also included in a more recent systematic review of CBT and TRT, although in this review the study was rated as high quality [113].

Sound therapy

No recommendation

There is evidence for safety but little high-level evidence for the effectiveness of sound therapy. Recommendation is based on RCTs and a systematic review.

Acoustic stimulation may be the oldest approach aimed at improving tinnitus. It is at least the most “natural” one, as tinnitus patients can experience every day that an external acoustic source can mask their tinnitus. This simple and intuitive approach has been (and is still) widely used. Importantly, this approach is not aimed at treating the causes of tinnitus but simply at helping to manage the consequences of tinnitus. It is used in different ways. Other acoustic approaches have been developed to interfere with the tinnitus causes. For these methods, the assumptions relative to the tinnitus mechanisms are critical. All these methods assume that tinnitus results from central changes after hearing loss that can be reversed by appropriate acoustic stimulation.

In general, acoustic stimulation has been shown to modestly improve tinnitus condition in several independent low-quality studies. It is unclear whether acoustic stimulation might improve tinnitus through some interaction with tinnitus mechanisms, through the partial or complete masking of tinnitus, and/or through certain cognitive influences (diversion, stress management etc.). Tinnitus is a highly heterogeneous

entity and acoustic stimulation may be very beneficial for some patients while completely ineffective for others. The “central” model of tinnitus assumes that the central changes due to sensory deprivation involved in tinnitus generation are reversible. However, some changes resulting from sensory deprivation might be difficult to reverse, especially when sensory deprivation has been present for many years. It may not be possible to compensate fully for deprived inputs by means of acoustic stimulation. Indeed, the cochlea (and/or cochlear nerve) can have nonfunctional areas, which are called “dead regions”. The presence of “dead regions” prevents any acoustic stimulation from activating the auditory centres within the corresponding projecting areas.

Tinnitus masking therapy. Many studies have shown that tinnitus masking therapy (TMT) can provide some relief for certain tinnitus subjects. However, only a few of the studies included placebo controls [126] and the different studies are not always comparable, as they used different questionnaires and protocols, with some studies even using their own custom questionnaires. Henry et al. [115] compared the efficacy of TMT and TRT, finding that both methods led to self-reported improvements in tinnitus, but that TRT was superior to TMT in reducing tinnitus-related distress, especially in the group of patients for whom “tinnitus is a very big problem.” Most of the improvement induced by TMT was achieved during the first 3–6 months of treatment, while TRT induced a steady

improvement over the course of the treatment (18 months). A more recent study, however, showed that TMT and TRT had similar effects on tinnitus when both were associated with counselling [170]. Finally, a randomised controlled study showed that TRT (masker + counselling) significantly improved tinnitus handicap [151].

Neuromonics approach. Neuromonics treatment consists of an acoustic stimulation combining music and broadband noise [106, 107, 114]. The spectrum of this combination is customised to provide an equalised stimulation over the audible frequency range. In addition to providing stimulation within the deprived sensory region, the acoustic stimulation is also designed to promote relaxation and relief. These effects are reinforced and complemented by counselling. Patients undergoing Neuromonics treatment are permitted to completely mask their tinnitus in the early stages of the treatment to maximise relief and relaxation (2 months). This initial stage is also intended to maximise the amount of stimulation of the deprived sensory region. In a second stage (4 months), the patients are discouraged from masking their tinnitus to facilitate desensitisation [107]. In an RCT by the manufacturers, this method was reported to significantly improve tinnitus. The study design included two groups with different modules of Neuromonics intervention, but participants self-adjusted the prescribed treatment for what they felt worked best, such that the intervention was no longer different between groups and their data were pooled. Overall, however, they reported clinically significant changes in tinnitus severity at 6 months for 86% of Neuromonics patients. Few independent studies of Neuromonics have been conducted. Of note, Newman and Sandridge [145] compared the cost-effectiveness and cost utility of Neuromonics versus ear-level sound generators at about one third of the cost. Both interventions resulted in reduced tinnitus handicap score with no difference in improvement between groups.

Notched music stimulation. A recent approach investigated the effects of notched music on tinnitus, the notch (1 octave width) being chosen to correspond to the tinnitus pitch [148, 167]. The notched music was intended to reduce tinnitus-related cortical activity within the notch, possibly through increasing lateral inhibition [150]. After 12 months of regular listening, this approach was reported to reduce self-reported tinnitus loudness, by around two points on a ten-point scale (eight subjects were treated with the notched music). The authors interpreted this approach as reversing the “maladaptive cortical reorganisation by the notched music training”.

Customised music stimulation. It has been suggested that tinnitus may result from the central changes accompanying hearing loss [146]. An implication of this model is that an appropriate acoustic stimulation may reverse the central changes due to hearing loss, including those involved in tinnitus generation. In this context, hearing aids may improve the tinnitus condition by restoring sensory inputs thereby reversing the tinnitus-related central changes due to hearing deprivation. Recently, an RCT investigated the effects of their own product aimed at reversing the tinnitus-related central changes using a customised music stimulation [137]. Tinnitus severity was significantly reduced according to the THQ questionnaire (by 34%). Tinnitus severity estimated from the TFI, however, was not changed by the method.

Sound therapy (including masking, music, environmental sound) may be useful for acute relief purposes but is not considered an effective intervention with long-term results.

Dietary and alternative therapies

Recommendation against

There is evidence that dietary and alternative therapies (e.g. Ginkgo biloba, melatonin, zinc, or other dietary supplements) have no proven efficacy and pose potential harm in the management of tinnitus. Recommendation is

based on RCTs and systematic reviews with methodological concerns.

Ginkgo biloba. *Ginkgo biloba* is the most commonly used herbal supplement for tinnitus. The two latest systematic reviews included three RCTs on *Ginkgo biloba* for tinnitus as a primary complaint [121, 173]. A Cochrane review, first published in 2004 and most recently updated in 2013, concluded that *Ginkgo biloba* was not effective [121]. A second systematic review included five RCTs, with most trials having low methodological rigour [173]. The results were favourable toward Ginkgo, but the authors stated that a firm conclusion about efficacy was not possible. A meta-analysis pooled data from six RCTs and concluded that there was no benefit of Ginkgo over placebo [159]. *Ginkgo biloba* can interact with other blood thinners to cause serious bleeding and can worsen bleeding risk in patients with underlying clotting disorders [155].

Melatonin. Melatonin is a hormone secreted by the pineal gland that is involved with regulation of the sleep-wake cycle. Three RCTs, with a total of 193 participants, studied melatonin to treat tinnitus, and each demonstrated benefit with the greatest improvement in those patients with severe tinnitus and insomnia [154]. However, given the small number of overall patients studied and the methodological limitations, including lack of a placebo group in the largest trial, these results should be interpreted with caution. Although another study demonstrated potential benefit for patients with concomitant sleep disturbance due to tinnitus, this study lacked randomisation, blinding, or placebo control [142]. Only one study reported possible adverse effects of melatonin, which included bad dreams and fatigue [160].

Dietary supplements. Three RCTs of zinc as a treatment for tinnitus, with a total of 205 participants, showed inconsistent results [97, 103, 104]. It was suggested that benefit could be associated with underlying zinc deficiency.

Several other dietary supplements have been used for tinnitus, including lipoflavonoids, garlic, homeopathy, traditional Chinese/Korean herbal medicine, honeybee larvae, and other various vitamins and minerals. Evidence for the efficacy of these therapies for tinnitus does not exist [155].

No recommendation

There is evidence for safety but little high-level evidence for the effectiveness of acupuncture. Recommendation is based on systematic review.

Acupuncture. No recommendation can be made regarding the effect of acupuncture in patients with persistent bothersome tinnitus, based on poor-quality trials, no benefit, and minimal harm. A systematic review in 2012 on acupuncture for the treatment of tinnitus included nine RCTs, with a total of 431 participants [132]. However, this systematic review highlighted the heterogeneity among study designs as well as their methodological limitations using the Cochrane tool for assessing risk of bias. Variations in study design included types of acupuncture intervention, frequency, intensity and duration of treatment sessions, selection of other control groups, variability with blinding, and selection of outcome measures, many of which were not validated [132]. The authors concluded that the small number of RCTs of acupuncture for the treatment of tinnitus, with small sample size and methodological issues, were insufficient to make conclusions about effectiveness.

4.2 Referral options and criteria, triage, and a stepwise proposal

A stepped-care approach that provides a standard pathway based on patient need, including the disciplines involved, assessments, and treatments at each stage, is presented in [Fig. 5](#). The steps proposed in the flowchart are based on the studies executed within the framework of the current guideline and the consensus meetings held within the steering committee. The barriers and facilitators of each member-country of

the TINNET project have been taken into consideration, with a gradual increase in the intensity of the care at each level to be implemented according to health-care policy, available resources and health-care coordination within the specific country, region, or state. Suggested cut-off scores on the THI and TQ have been included as an example. For other instruments on tinnitus severity, we refer the reader to Chap. 3.

A stepwise multi-disciplinary approach

On the basis of the evidence described at present, we suggest that a CBT-based approach, whether in groups or individually, is the most evidence-based choice for effectively relieving tinnitus complaints. Tinnitus treatment aimed at the sound-perception level, such as sound therapy, including the use of hearing aids prescribed for tinnitus relief only, masking devices, ear-level sound generators, sound perceptual training, or other sound generating technology, however, has not been proven to have an additional effect on counselling or CBT, or as a standalone treatment. Nevertheless, evidence indicates the merits of audiological diagnostics, counselling, and education to decrease tinnitus suffering as well. On the basis of the current evidence, we suggest that the best tinnitus treatment strategy might be CBT based. Research suggests that next to otolaryngological/medical diagnostics, an overall CBT-based framework in tinnitus management is advisable, from audiological diagnostics (assessment of hearing and prescription of hearing aids if indicated to increase hearing function) and tinnitus counselling to psychological diagnostics and tinnitus treatment, since all studies showed benefits from some form of education, information, and/or counselling initially for all patients, and for the more severely impaired a more intensive CBT treatment. Moreover, tinnitus standard care might be best organised in a multi-disciplinary manner, using a stepped-care approach [13, 174], gradually increasing intensity of treatment in steps, so that most patients can be treated effectively with a fairly short process (diagnostics and information/

education), and additional treatment steps can be indicated for those suffering on a more severe level.

Chapter 5 Patient information and support

5.1 Confirming knowledge and dispelling myths. Key messages to prepare the patient for treatment and beyond

It is essential to successful tinnitus treatment that patients are empowered to self-care, that they are provided with reliable information and learning resources, and that they are signposted to appropriate sources of support [34]. Patients need to understand tinnitus, how distressing tinnitus is managed, and what sources of information and support are available to them beyond their treatment sessions.

The provision of information should be timely and fill gaps in knowledge, dispel myths, offer hope, and provide key messages that are a framework for treatment. Information should never be negative, e.g. “there is nothing that can be done” or “you will just have to learn to live with tinnitus”.

Having information about tinnitus can be very powerful. Many patients start their tinnitus journey by looking for information on the Internet and while there is some accurate and useful information available on the Internet, there is also a lot of very unhelpful information [202]. Patients may read “nightmare” stories on the Internet, or hear inaccurate “facts” from friends, family, or even a clinician. Dispelling such myths is crucial.

In the assessment described in Chap. 3, information is gathered about how tinnitus is affecting the patient in their daily life, about their understanding of tinnitus, and their concerns or fears surrounding it. This can be used to explore with them how their beliefs about tinnitus and the meaning they attach to it influences how they think, feel, and react to it. Health-care professionals should be compassionate to the concerns and fears expressed by patients.

There is no cure for subjective tinnitus, but patients can be treated to help them




Fig. 5 ◀ Criteria for assessment and treatment of tinnitus. CBT cognitive behavioural therapy, THI Tinnitus Handicap Inventory, TQ Tinnitus Questionnaire

habituate and to reduce the functional impact of their tinnitus. The functional impact might include sleep disturbance, difficulty concentrating, problems with hearing, and difficulty relaxing. Patients can learn to manage their reactions to tinnitus, thereby improving their quality of life.

5.2 Information that should be given to patients

Patient information and resources (e.g. decision aids) need to be sufficient to allow them to reach a shared decision with the clinician about their treatment and to fully engage themselves in the process of care [182, 208, 212]. Initial information and advice need to be given that will enable patients to have immediate strate-

gies to alleviate tinnitus. Treatments may include hearing aids, sound-generating devices, medicines, and ways to learn how to cope with tinnitus. Depending on what is available locally, tinnitus care might involve the family doctor, an ear, nose and throat (ENT) specialist or an audiological physician, hearing therapists or specially trained audiologists, clinical psychologists, nurses, or other clinical professionals [15, 21, 34, 187].

Table 9 Common myths/misconceptions and information for patients

Example myth/misconception	Example information
Having tinnitus means you will go deaf	It should be explained to the patient that this is not the case. Everyone's hearing deteriorates as they get older, but tinnitus does not <i>cause</i> hearing loss. In many cases, tinnitus is caused by pre-existing hearing loss, but not the other way around [204]
Tinnitus is temporary and will go away	It should be explained that tinnitus can be temporary, but often it is not. The attitude of "hoping" or "wishing" it will go away when it does not go away causes additional distress and monitoring, thus driving tinnitus distress [57]
Tinnitus will get worse over time	It should be explained that tinnitus can be affected by life situations that make it more bothersome in time. In general, the severity of tinnitus decreases over time [207]
Hearing aids will make the tinnitus louder	Where there is a hearing loss as well as tinnitus, and the hearing loss can be compensated by hearing aids, the hearing aids may also help to address tinnitus. Hearing aids amplify external sounds, i.e. those in our environment; hearing aids cannot and do not amplify an internal noise, which is what the tinnitus noise is [211]
Caffeine makes tinnitus worse	This may be the case for a small number of individuals; however, a clinical trial found that caffeine had no effect on tinnitus severity [213]. In fact, the same study reported a significant negative effect of caffeine withdrawal, concluding there is no evidence to justify caffeine abstinence as a therapy to alleviate tinnitus. However, the consumption of coffee does influence sleeping patterns. If there are sleeping problems associated with tinnitus, then coffee consumption should be limited
Tinnitus means there is a very serious underlying illness	Although there are many conditions that may cause or exacerbate tinnitus, it is extremely rare that it is a symptom of a serious illness [179]

Patient information and support topics should be tailored to the patient's need and what is available to them. Topics include:

- What is tinnitus? What causes and maintains it?
- Pulsatile tinnitus (follows heartbeat)
- Common misunderstandings and myths
- Hearing loss and hearing aids
- Ear wax removal
- Hyperacusis and tinnitus
- Protecting your hearing
- Habituating to tinnitus
- Relaxation
- Monitoring tinnitus
- Use of sound
- Dealing with sleep problems
- Dealing with emotional consequences of tinnitus
- Self-help and support groups

What is tinnitus? What causes and maintains it?

Even though tinnitus is a symptom and not a disease, it can lead to illness in

some people. Tinnitus should be explained to the patient as a sensation of hearing a sound in the absence of any external sound and their reaction to it. It can be reassuring for patients to describe the various ways in which people experience tinnitus. Different patients will hear different types of sound, for example ringing, buzzing, whooshing, or humming, and occasionally tinnitus has a musical quality. Tinnitus might be continuous or intermittent. It might seem like it is heard in one or both ears, inside the head, or it might be difficult for the patient to describe where it is coming from. Tinnitus is very common (about 10–19% of the population) and is reported in all age groups including children. It is more common in people who have hearing loss or other ear problems, but it can also be found in people with normal hearing. Most people find that it does not affect them in any way. Some people find it moderately annoying, while others finding it very troublesome [172, 189].

It is important to explain that for subjective tinnitus, exactly how and why it occurs is not yet fully understood but that research is ongoing. It is thought that tinnitus is the sound of activity within the auditory system, which can generally be heard in a silent environment, but in the presence of normal environmental sounds it is "filtered" out by the brain because it does not have meaning.

Although an underlying cause of tinnitus is rarely known, it is commonly linked to *hearing loss*. Age-related "wear and tear" in the inner ear is a common cause of sensorineural hearing loss [204]. It can also be linked to exposure to loud sounds such as occupational or leisure noise [180]. If relevant, it should be explained to the patient that tinnitus is also associated with disorders causing *conductive hearing loss* such as ear wax blockage and otosclerosis [204]. With any ear condition, the amount of information being sent to the brain can change. Research suggests that tinnitus results from the compensatory adaptation of the central auditory system to hearing loss.

Note that tinnitus is sometimes associated with emotional stress, use of certain medications, ear infections, ear, head or neck injuries, neurological disorders including acoustic neuroma, metabolic disorders including hypothyroidism, hyperthyroidism, and diabetes. Patients should be reassured that tinnitus is rarely an indication of a serious disorder.

Some clinicians invite patients to a *group tinnitus information session*, following which they can request individual therapy if they wish [203]. Some patients find the group session alone adequate to reassure them and no further treatment is needed. Group sessions have the advantage of considerably reducing patient waiting times until their first appointment.

Common myths and misconceptions about tinnitus

Where myths or misconceptions about tinnitus are identified during assessment, it is important to provide information to dispel those myths and misconceptions. Some of these are commonly observed in clinical practice (■ Table 9).

Hearing loss and hearing aids

Patients with tinnitus often attribute hearing problems to tinnitus and thus part of assessment is to determine how much of a patient's complaint is due to a hearing problem and how much is due specifically to the tinnitus [192]. Hearing loss is often an unnoticeable and gradual process, and many people are surprised when they are told that they have a hearing loss. For patients who have even a mild hearing loss and tinnitus, there is consensus that using hearing aids may be helpful because they amplify normal environmental sounds which otherwise may not be heard and may thus reduce the perception of tinnitus [124, 209]. Where they are available, clinicians should discuss hearing aids or combination hearing aid options.

Hyperacusis and tinnitus

Many people who experience tinnitus also report hyperacusis, an increased sensitivity to everyday sounds that makes them uncomfortable to hear [186]. A natural response can be to block out as much sound as possible, for example using ear protection, but avoiding sound can make hyperacusis, and tinnitus, worse. Often using sound in a very controlled way can improve hyperacusis [198].

Looking after your hearing

Exposure to loud noise can cause a temporary shift in hearing thresholds as well temporary tinnitus or make existing tinnitus worse. Patients should be informed that these temporary changes will likely resolve within a few days following the noise insult but that repeated episodes of noise exposure increase the likelihood that the tinnitus will become worse in the longer term [195].

The clinician should teach the patient what is “too loud”. Examples of what is too loud are:

- If you have to shout to be heard by somebody around a meter away
- If you find your hearing is dulled after exposing yourself to noise
- If you find a ringing or buzzing in your ears (tinnitus) after exposing yourself to noise
- If a sound is painfully or uncomfortably loud, stop exposure immediately

People should take care of their hearing by wearing proper ear protectors for activities that are “too loud”, e.g. using power tools or being near noisy motors for any amount of time. Ear protection is also important if the patient attends live music events or plays in a band or orchestra. Ear protection should not be used if ordinary, everyday sounds are uncomfortable [198].

Patients should also be educated to limit the amount of time and the volume when they are listening to music through earbuds or headphones. A simple 60/60 rule can be useful; never turn the volume above 60% and restrict listening time with earbud headphones to 60 min per day. At higher volumes, the amount of listening time needs to be reduced significantly. Earmuff-style or noise-cancelling headphones are preferable. For leaflets on safe listening volumes available in many languages, visit www.who.int/pbd/deafness/activities/MLS/en/.

Adaptation

A thorough clinical assessment will determine where and when tinnitus is more and less intrusive, and a follow-on discussion should include what can be done to make the patient's environment more “tinnitus friendly”, e.g. introducing some low-level sound. It is important to explain that, whilst many people have tinnitus, only some are aware of it all the time or bothered by it. This is because people often get used to the noises just as we get used to other noises around us (e.g. air conditioning, a clock ticking) in the short term, i.e. they adapt to the presence of those unimportant sounds. If our attention is focused on something else, it may be possible to “forget” the tinnitus at times and thus reduce its impact. It is a new sensation and patients need to give themselves time to adapt to it. Most people find that their tinnitus seems to settle down over time as they notice it less. There is good evidence from clinical trials that in general tinnitus becomes less bothersome over time, even without doing anything [193, 207].

Patients should be encouraged to keep doing the things they enjoy. If they start living life differently to accommodate the tinnitus, it is going to seem more of

a problem. They may need to do some things differently, however. For example, if they enjoy reading but it becomes a quiet activity where tinnitus is more noticeable they may need to start using some background music or environmental noise. The important thing is that they find ways to keep doing what they enjoy.

Relaxation

Patients should be reassured that it is perfectly normal to feel anxious or afraid because of tinnitus, especially when they first experience it [57, 214]. Tinnitus works a bit like an “emotional barometer”—it is often more intrusive when there is anxiety or fear around. Tinnitus might become more bothersome when under duress.

If patients can learn to relax more, they will be less anxious and afraid and so notice their tinnitus less. There are many forms of relaxation such as progressive muscle relaxation or deep-breathing exercises, and these can be guided by a person, audio, or written instructions (e.g. www.tinnitus.org.uk/relaxation). A relatively easy way your patient might relax is to find somewhere peaceful and just slow their breathing down (they may have some sound on in the background). A simple relaxation exercise is to take a few slow deep breaths and pay full attention to the feeling of the breath entering the body, filling the lungs, and leaving the body.

Both adaptation and relaxation *take time*. It may be useful for patients to create some catchphrases to use when tinnitus is being more bothersome, such as “Calming tinnitus takes time”.

Monitoring tinnitus

If life is planned around tinnitus it is given much importance, and this prevents the patient adapting to it. Simply put, patients should not avoid activities they think may make their tinnitus worse. Patients should not be putting their life on hold. Tinnitus does not have to control their life. Each time the patient tries to “monitor” their tinnitus they are guiding their attention to it; there is, however, no evidence that this would make tinnitus worse. They should be advised to engage in normal activities when they no-

tice themselves trying to monitor their tinnitus.

Use of sound

Patients should be advised on the use of sound to reduce the intrusiveness of tinnitus in quiet situations such as when trying to sleep, work, or to read in a quiet environment. Patients should be advised to experiment with the type and volume of different sounds to find what suits them best in different situations. Examples of what can be useful include opening a window to let in sounds, leaving a television on in the background, static noise from a radio, noise from a fan, or recorded environmental sounds produced by a bedside generator [199].

Some research suggests that sound volumes that allow for constant exposure to a very low level of tinnitus help the patient adapt to their tinnitus [126, 153]. Higher sound volumes may make tinnitus inaudible, but for some patients this may make the tinnitus more noticeable when the sounds are turned off. A wide variety of non-wearable and wearable sound therapy devices are available that can be helpful for some patients.

Addressing sleep problems

Many patients report difficulty in getting to sleep or staying asleep because of tinnitus. It can be a vicious cycle where worrying about tinnitus and/or worrying about the sleep difficulty makes it even more difficult to sleep. Whilst in a patient population most will complain of sleep problems, in the general population most people with tinnitus sleep well and their tinnitus is no different in quality from those who do not sleep well. If sleep is a problem then it may be helpful for the patient to learn about the biology of sleep, the internal biological clock, and sleeping patterns.

People who have tinnitus and sleep poorly tend to worry more at night than people with tinnitus who sleep well [183]. Patients should be advised that working through problems during waking hours is better than doing so in the middle of the night when they have nothing else to occupy them. One strategy patients might try is to insert a “worry moment” of 10–15 min into their daily schedule,

ideally in the early evening. At that time, they can write down (or think about) all the worries, plans, comments, questions that come to their mind. The good thing about our brain is that “a thought once thought, can be filed away”. In other words, the patient should know they can control their worry. Another strategy to counter worry involves “thought stopping”; patients may use methods such as simple breathing exercises (e.g. exhale three times consciously, or simply verbalising the word “STOP” out loud). The practice of relaxation and breathing exercises puts the body in a state that allows for rest, reduces worry, and increases the effectiveness of your biological clock. General advice:

- Plan participation in any sports or other high-energy activities for the afternoon or early evening (rather than later). Ritually “powering-off” in the last 2 h before bed, e.g. dimming the lights, only engaging in low-intensity energy activities, avoiding aggravating television shows or late-night discussions.
- Be aware that caffeinated drinks and alcohol can negatively affect sleep, as does smoking, and the consumption of meals within 2 h of going to bed. Warm milk has a sedating effect.
- Ensure an ambient temperature in the bedroom. Reserve the bedroom for sleeping (and intimacies). Try and limit any other activities (even reading and watching television) to as little as possible in the bedroom. Try to get into a routine of going to bed at a fixed time each night, but, at the same time, go to bed only when tired.
- When the preferred sleep pattern is disturbed (e.g. when travelling or working in shifts), take as little sleep (naps) as possible outside your normal routine.

Addressing psychological problems

The clinician should emphasise to patients that although there is no cure there are many things that they can do to make tinnitus less of a problem. There are a wide variety of psychological treatment options available, from providing

the basic information described earlier, to treatment elements that focus on certain problems with thoughts, emotions, and behaviours leading to functional difficulties (concentration, sleep, daily activities). The patient should be provided with information about the treatment options available locally.

One structured approach to tinnitus treatment is *tinnitus retraining therapy* (TRT; [130]). The basis of TRT is that tinnitus is prioritised by the brain as an important signal. This treatment uses bilateral sound generators or hearing aids at a certain sound level to try to reduce the percept of tinnitus so as to allow patients to adapt to the tinnitus sound. Sound is used in combination with directive (educational) counselling. Most clinicians who treat tinnitus use some elements of TRT (e.g. Cima et al. [13], who used the neurophysiological model to explain tinnitus) but the strict method is not frequently used because it is resource intensive and there is limited evidence for its effectiveness [153].

Cognitive behavioural therapy (CBT) is a psychological approach to treating tinnitus that has substantial high-level evidence for its effectiveness [84, 120, 125, 139]. The basic premise should be explained to the patient. It works on the principle that when the patient becomes aware of tinnitus, they respond negatively to it. For example, the patient develops a belief that something is seriously wrong with them (belief) and this leads to anxiety and fear (emotion), and unhelpful behaviours such as avoiding silence (behaviour). Cognitive behavioural therapy helps patients to recognise which beliefs or behaviours are unhelpful. They then work with their clinician (usually a psychologist, specialist audiologist, or hearing therapist) to develop different ways of responding to the tinnitus that ultimately make it less bothersome (see Appendix E).

A clinician-guided Internet-delivered version of CBT is available in some countries and there is good evidence for its effectiveness when delivered by this medium [215].

Mindfulness-based interventions, or simply *mindfulness*, have been classified as a psychological treatment aimed at

psychological distress associated with a range of complaints [196] and typically consists of up to ten group sessions. Mindfulness trains the skill of being mindful, fostering moment-to-moment awareness, and observing emotions, sensations, and cognitions non-judgmentally. The rationale for using mindfulness is that if the patient stops trying to avoid unpleasant sensations (such as tinnitus) they will be able to perceive the tinnitus without the struggle. The approach offers individuals a new way to relate to thoughts that allow them to reduce their tendency to engage in negative and catastrophic thoughts. Mindfulness for tinnitus has been tested in clinical trials with evidence that it is feasible as a treatment for tinnitus [141, 197] and may be beneficial in reducing negative emotions, rumination, and psychological difficulties [206].

Acceptance and commitment therapy (ACT) has its roots in the behavioural tradition (see Appendix E). The focus in ACT is on functional utility of thoughts and actions, and not on their “right- or wrongfulness” [190]. One of the key elements of ACT is to help experience psychological events (thoughts, perceptions, and emotions) in a non-judgmental way, not trying to change or modify those events. This leads to a more functional awareness of how thoughts, emotions, and behaviours create and maintain distress. Since *mindfulness* promotes present-moment awareness and observation in a non-judgmental way, it has become an integrated part of the ACT protocol.

Self-help

Self-help resources are shown to be useful in reducing tinnitus-related distress with mixed evidence for effects on comorbid symptoms such as depression [188]. Self-help books are widely available, and some provide complete programmes of self-help for tinnitus. For example, the book *Tinnitus: A Self-Management Guide for the Ringing in Your Ears* by Henry & Wilson [191] provides a series of exercises focused on a cognitive behaviour modification approach, and there is some from a controlled study [200] that it is beneficial. *Living with Tinnitus and Hypera-*

cusis [201] looks at strategies for living with tinnitus and hyperacusis. It includes a complete programme of self-help covering causes and mechanisms of tinnitus and hyperacusis; their impact; effective treatments; relaxation and sound therapy; relieving stress; and avoiding relapse.

Hundreds of mobile applications have been developed specifically for use in hearing health care, including for tinnitus [205]. For tinnitus, most were developed to assess the characteristics of a person's tinnitus over time, or to provide sound-enrichment options [210]. Unfortunately, we are at present unable to validate these apps or identify which might be useful.

It should be acknowledged that the patient's family and others may not understand what tinnitus is and how it might affect the patient. If that is the case, it can be helpful for them to talk to someone who has experience of tinnitus. One option would be for them to attend a tinnitus self-help group or support group. These groups facilitate an exchange of information between its members; patients can keep informed on the latest information and share treatment experiences by talking to others with similar problems [181, 184]. Groups may be lay- or clinician-led. Where possible these groups should be facilitated or attended by a clinician to prevent misinformation from being conveyed. Group activities may include lectures from different related disciplines. Clinicians should be aware of local groups or consider forming one if there is no support group locally.

Some patients may be unwilling or unable to attend a face-to-face self-help or support group. An alternative for these patients may be to take part in an online support group (OSG; [194]). These groups share many of the supposed benefits of face-to-face groups such as being a means to share experiences and seek or provide emotional support. Equally, they risk patients being exposed to inaccurate or misleading information [185]. As such, OSGs need to be appropriately moderated, ideally by a clinician.

5.3 Further information and support

National registered charities for tinnitus are a reliable source of information and support for people with tinnitus. Clinical professional organisations also provide patient information and leaflets.

- The *British Tinnitus Association* (www.tinnitus.org.uk/) provides a helpline and information leaflets (in English) that are free to all, and a quarterly magazine for its members.
- The *Ida Institute* (<http://idainstitute.com/>) has worked in collaboration with the *British Tinnitus Association* to produce a web resource (www.tinnituskit.com) providing basic information about tinnitus and how to deal with it.
- The *Charité German Tinnitus Foundation* (www.deutsche-tinnitus-stiftung-charite.de/die_stiftung/) provides online information in German and English (www.deutsche-tinnitus-stiftung-charite.de/en/tinnitus/).
- The *German Tinnitus League* (www.tinnitus-liga.de) provides information only in German.
- The *American Tinnitus Association* (www.ata.org/) provides online information and podcasts in English on tinnitus causes, treatment, and the research they support.
- The *Dutch Tinnitus Association* (www.stichtinghoormij.nl/) provides online newsletters, information and publishes about patient meetings and recent results from the research they support.
- The *Dutch ENT-Guideline* can be found here: https://richtlijndatabase.nl/richtlijn/tinnitus/behandeling_van_patienten_met_tinnitus.html (only available in Dutch).
- The *National Institute on Deafness and other Communication Disorders* provides information and PDF leaflets on hearing health and research, available in English and Spanish: www.nidcd.nih.gov/health/tinnitus
- The *French Patients' Tinnitus Association* (www.france-acouphenes.org/).

Guidelines

- The *Belgian Patients' Tinnitus Association* (www.belgiqueacouphenes.be/).
- The *Danish Association of the Hard of Hearing* (<https://hoeforeningen.dk>) provides a helpline (open 8:00–21:00) run by volunteers with tinnitus, who have been trained at The Danish Association of the Hard of Hearing on guidance, support, and the structure of the Danish Health-Care System.
- The *Italian Au-Tu (Acufene Uniti-Tinnitus United)*; www.au-tu.org) provides information in Italian both online and through meetings reporting the state of basic and clinical research on tinnitus and offers a moderated forum (for subscribers only) for tinnitus patients.
- The *Foundation for the Research of Tinnitus and Hyperacusis* (FINVAC): No online information found.

Corresponding address

Dr. R. F. F. Cima

Faculty of Psychology and Neuroscience,
Department of clinical Psychological Science,
Experimental Health Psychology, Maastricht
University
Maastricht, The Netherlands
r.cima@maastrichtuniversity.nl

Acknowledgements. A COST Action grant (BM1306) supported collaboration between the authors and the formation of the COST Action BM1306 (2014–2018) TINNET Working Group I.

Special thanks are due to Thanos Bibas, Christopher Cederroth, Thomas Fuller, and Sarah Rabau for their contribution to the initial developmental stages of this guideline.

Compliance with ethical guidelines

Conflict of interest. R.F.F. Cima, B. Mazurek, H. Haider, D. Kikidis, A. Lapira, A. Noreña, and D.J. Hoare declare that they have no competing interests.

This article does not contain any studies with human participants or animals performed by any of the authors.

Table A.1 Clinical examination in patients with pulsatile tinnitus

Examination	Rationale
Ask the patient to tick his/her finger in each pulse, while taking radial pulse	Confirm that tinnitus is pulsatile and follows heartbeat
Otoscopy: Special attention to retrotympanic mass	Possibility of glomus jugulare or tympanicum, high jugular bulb, ectopic carotid artery
Press jugular vein and ask patient whether tinnitus is alternated	Discriminate between arterial and venous pulsatile tinnitus
Use a stethoscope to listen to mastoid, external auditory meatus, neck, and chest	Look for murmur, indicative of an arteriovenous abnormality

Appendix A

Pulsatile tinnitus management

Pulsatile tinnitus follows a person's heart-beat and can be either subjective or objective [218]. It occurs in less than 5% of tinnitus patients [216]. It is commonly associated with venous and arterial abnormalities, either because of increased blood flow or stenosis. Pulsatile tinnitus can also be discriminated as venous or arterial, based on whether it disappears with pressure in the jugular vein or not.

The overall approach and assessment of patients with pulsatile tinnitus differs from that for patients with subjective tinnitus, and special clinical investigations should be implemented because serious and potentially reversible causes might be found. In this brief section, special examination aspects which should be implemented in addition to a typical and radiological evaluation algorithm are presented.

Clinical examination

A summary of the clinical examination of patients with pulsatile tinnitus is presented in Table A.1 [217, 218].

Radiological evaluation

The radiological evaluation of pulsatile tinnitus [216, 218–220] comprises:

Arterial pulsatile tinnitus

- Carotid triplex (stenosis)
- Computed tomography angiography (glomus, aneurysms, atherosclerosis, arteriovenous malformations)

Venous pulsatile tinnitus

- Magnetic resonance angiography (arteriovenous malformations, empty

sella syndrome, Arnold–Chiari malformation, Sylvius aqueduct stenosis, sigmoid sinus diverticulosis, etc.).

- When imaging is normal, consider benign intracranial hypertension, especially in patients with a high body mass index.

Appendix B

Method for the development of an implementation plan

Per recommendation, and possibly per country/region, the following points need to be considered:

- In which time frame is it expected/recommended that the recommendation is implemented across Europe?
- What are the conditions that need to be met for the recommendation to be successfully implemented?
- What are the possible barriers per country/region for the recommendation to be successfully implemented?
- What are the possible facilitators per country/region for the recommendation to be successfully implemented?
- What is the impact on health-care costs for the implementation of the recommendation?
- Who is the responsible party to undertake the actions needed for successful implementation of the recommendation?

In considering the aforementioned points, we propose the following:

- Recommendations might be categorised according to level of recommendation (strong—weak—against). All recommendations in the European guideline should be weighed,

according to the “strength” of the recommendation.

- For the strong recommendations, each point will need extensive investigation and reporting.
- For the weak recommendations, alternatives might be mentioned or considered.
- The recommendations against will be absolutely discouraged in favour of treatment/procedures with strong recommendations.

Appendix C

Somatosensory tinnitus

Tinnitus is considered somatosensory when it can be modulated by somatic stimulation or movement. It might also be considered a subtype of tinnitus that is associated with activation of the somatosensory, somato-motor, and visual-motor systems. Somatic modulation of tinnitus has been reported to be observable, when actively looked for, in up to 83% of tinnitus patients [229, 236, 239, 241]. Somatosensory tinnitus can be modulated by jaw clench [239] or eye movements [232, 235]. Temporomandibular joint dysfunction may cause somatosensory tinnitus [240]. In some cases, pulsatile tinnitus can be modulated by movement of the head or upper lateral neck [231].

Pathophysiology

Somatosensory tinnitus is hypothesised to be related to abnormal cross-modal plasticity of somatic-auditory interactions [222, 226, 227, 230, 234]. Somatic modulation of tinnitus is assumed to result from abnormal auditory neural interactions within the central nervous system [237]. Cross-modal interaction is further supported by the observation of auditory cortical area activation in deafness as a result of somatosensory stimulation [222–224].

Diagnosis

Somatosensory tinnitus should be considered when the patient can modulate the loudness or intensity of their tinnitus

by movement or tactile stimulation [221, 228, 233], and when they report [238]:

- Head or neck trauma.
- That they can modulate tinnitus quality by manipulation of teeth, jaw, or cervical spine.
- Frequent pain in head, neck, or shoulder girdle.
- Simultaneous onset of pain and tinnitus.
- Tinnitus is aggravated by incorrect body postures.
- Severe bruxism.

Treatment

Treatment for somatosensory tinnitus tends to include: oro-myofacial therapy, splinting, physical therapy, relaxation therapy, somatic modulation therapy, electrical stimulation, and local cortisone injection [225].

Appendix D

Table D.1 Characteristics and psychometrics of existing tinnitus health-related quality-of-life instruments						
Instrument ^a	Developed to	Items	Scoring	Construct validity	Reliability (test re-test)	Subscales
Tinnitus Handicap Inventory (THI)	Measure level of perceived tinnitus severity	25	(0) never (2) sometimes (4) yes	+	+	Functional, emotional, catastrophic responses
Tinnitus Questionnaire (TQ)	Measure psychological aspects of tinnitus complaint and distress	52	True Partly true Not true	+	+	Emotional distress, cognitive distress, intrusiveness, auditory perceptual difficulties, sleep disturbance, somatic complaints
Tinnitus Reaction Questionnaire (TRQ)	Measure psychological distress due to tinnitus	26	(0) not at all (4) almost all the time	+	+	General distress, interference, severity, avoidance
Tinnitus Severity Index (TSI)	Measure how much tinnitus negatively impacts a patient's life and how bothersome patients perceive their tinnitus to be	12	(0) never (4) always	–	+	None
Tinnitus Handicap Questionnaire (THQ)	Measure patients' perceived degree of handicap due to tinnitus	27	(0) strongly disagree (100) strongly agree	+	+	Physical health/emotional status/social consequences, hearing and communication, personal viewpoint
Tinnitus Severity Questionnaire (TSQ)	Measure tinnitus severity	10	0 (not affected) 4 (always affected)	–	–	None
Tinnitus Functional Index (TFI)	Measure tinnitus severity and treatment-related change	25	0 (not affected) 10 (always affected)	+	+	Intrusiveness, sense of control, concentration, sleep, hearing, relaxation, quality of life, emotion

^aSome of these questionnaires are already available in multiple languages. Hall et al. [258] provide a guide to translation and adaptation of hearing-related questionnaires

Table E.1 The “third-wave” CBT treatments**MBSR**

Mindfulness is a type of psychological treatment aimed at psychological distress, depressive symptoms, and anxiety, initially devised for individuals suffering from chronic disease. Mindfulness-based stress reduction (MBSR) was developed by Kabat-Zinn [196, 251]. MBSR protocols typically consist of up to ten group sessions. The focus lies on training of the skill of being mindful, which is a moment-to-moment awareness, and observing emotions, sensations, and cognitions non-judgmentally. Sessions are built around meditational skills, bodily exercises, and psycho-education. MBSR was developed for chronic pain sufferers and was later adapted for chronic diseases, such as heart disease, and recently adapted for tinnitus [206, 243, 250]. As a stand-alone treatment approach, mindfulness has been applied to many psychological disorders [245, 253]. Mindfulness is also an important component of other psychological treatments such as acceptance and commitment therapy (ACT), some forms of behavioural treatment, and cognitive therapy [190, 247, 255]

ACT

According to its founder [190], ACT has its roots in the behavioural tradition. ACT does not emphasise the accuracy or the content validity of cognitions and behaviours, as is the case in more cognitive approaches. ACT focuses on functional usefulness of thoughts and actions, and not on the “right- or wrong-fullness” [190, 249]. A key element of ACT is to decrease “experiential avoidance” [248]. ACT advocates experiencing psychological events (thoughts, perceptions, emotions) in a non-judgmental way, rather than trying to change or modify those events, leading to a more functional awareness of how thoughts, emotions, and behaviours create and maintain distress. Since MBSR approaches advocate present moment awareness and observation in a non-judgmental way, which results in decreased rumination and worry, it has been integrated within ACT protocols

Appendix E

Cognitive therapy and cognitive behavioural therapy

Confusion often exists about the differences between cognitive therapy and cognitive behavioural therapy (CBT), as both terms are used interchangeably. Since CBT stems from the convergence of two distinct theoretical schools, the radical behavioural school (first wave) and the cognitive school (second wave), CBT entails both cognitive and behavioural principles and methods, and usually a combination of these are used in therapeutic sessions. Cognitive behavioural theory and treatment have been applied in tinnitus research for decades [246, 252, 254] and CBT approaches for tinnitus have been repeatedly shown to be effective in decreasing tinnitus distress, anxiety, and tinnitus annoyance, and improving daily life functioning. Although there are common elements across CBT-based treatments for tinnitus, the CBT approaches investigated vary in number of treatment sessions, hours spent in therapy, group versus individual formats, face-to-face versus Internet-based self-help therapies, combinations of different treatment ele-

ments, and tinnitus diagnostics and outcome assessments.

The cognitive approach

The main premise of the cognitive approach is that cognitive factors maintain psychological distress. Typical cognitive therapeutic interventions (or so-called talking therapy) are aimed at: (1) correcting/changing “erroneous” beliefs or thought processes (cognitions), (2) current problems and thought processes, and not so much the past, and (3) advising patients to perform behavioural experiments in order to test the validity of maladaptive thoughts and beliefs [242, 244]. The cognitive model [57], in line with the cognitive tradition, posits that therapeutic strategies to change these maladaptive cognitions automatically lead to changes in emotional distress and problematic behaviours. These cognitive techniques seem to be helpful in the short term. In addition to educational counselling, techniques include but are not limited to “Socratic dialog”, thought control, rational thought formulation, exploring automatic thoughts, and testing of thoughts and beliefs through behavioural experiments.

A fear-avoidance approach

Recently, *exposure* therapy, a cognitive-behavioural treatment strategy, entered CBT treatment protocols for tinnitus [13]. Exposure therapy, also applied in CBT treatments for chronic pain [256], is a clinical application of what is called “extinction” of the association between two stimuli in classic learning theory terms. In the case of tinnitus patients, it is assumed they learn to be fearful of the tinnitus signal. That is, in the distressed tinnitus patient the initially neutral tinnitus signal became associated with sympathetic arousal (alarm detection; [37, 257]). According to fear-avoidance reasoning, the neutral tinnitus signal became a predictor of unwanted aversive states, hereby receiving a very negative value (danger). Patients interpret the signal as a sign of harm or injury, which is why they are so fearful, and selectively single out the tinnitus signal. They are eventually constantly interrupted and engaging in safety-seeking behaviour, because of these threat expectancies. These mechanisms are likewise at work in arachnophobes, for example. The spider, which is harmless, has become a sign of great danger, leading to extreme fear.

For arachnophobia, exposure procedures consist of repeated confrontations with spider-related images, objects, and eventually real-life spiders, which evoke the greatest fear of the patient. As a result of repeated exposure to the most feared stimuli, the patient learns that confrontation with spiders is not life-threatening, and therefore they are not in danger, and the fear of spiders dissipates (extinction). By analogy, the tinnitus patient is extremely fearful of perceiving the tinnitus. Even though the tinnitus is continuously present, the involuntary response is trying not to hear it, and trying to be minimally confronted with the tinnitus sound (avoidance). Patients do this by trying to control their sound environment, not thinking about it, directing their attention elsewhere, and consequently increasing their monitoring and awareness of their tinnitus. Consequently, cognitive resources are depleted, leading to task interruptions, more avoidance (safety-seeking), and eventually disruptions in

functional activities. In the long term, severe disability ensues, disrupting all life domains and leading to severe dysphoria.

This “new” forms of CBT for tinnitus typically includes the third-wave forms of therapy (see Table E.1) to enhance internal observations, increase moment-to-moment consciousness of the tinnitus, and to provide the ability of observing tinnitus-related emotions, sensations and cognitions non-judgmentally. Exposure therapy for tinnitus patients entails exposing them to their tinnitus sound, the interceptive sensations associated with the tinnitus, as well as their moment-to-moment narrative. To provide an appropriate context, exposure is performed in quiet circumstances. This way the patient experiences that the tinnitus sound is harmless, not dangerous, and listening to it in silent environments will not lead to catastrophe. They also learn that the aversive consequences are not always triggered. These experiences lead to a “neutralisation” of tinnitus by adaptation of fear expectancies; consequently, the tinnitus becomes less intrusive and bothersome, the more they engage in exposure.

Comparing the treatments

The theoretical frameworks have strong conceptual overlap, and are based on the premise that the initially neutral tinnitus signal receives an “alarm” value, through classic conditioning. In turn, this negative tinnitus valence exacerbates negative responses in cognitions, emotions and behaviours, hindering the “normal” process of habituation. Tinnitus distress ensues; which is the very negative and aversive state when processes of adaptation and the efforts thereto have failed to return the organism to equilibrium or homeostasis.

Treatment avenues sometimes seem to be contradictory. The neurophysiological model-based tinnitus retraining therapy includes extensive education and (partial) masking of the signal (avoidance of the signal, by avoiding silence at all costs), hypothesising that both are required for habituation. The habituation theory and cognitive approaches purport that thought control and attention-diver-

sion techniques (alter thoughts/beliefs about the tinnitus and actively directing attention away from the tinnitus) will be beneficial for habituation. For short-term habituation, these strategies may work. On the other hand, the fear-avoidance approach leads to an opposite treatment strategy, promoting exposure to tinnitus and eliminating avoidance tendencies (such as avoiding silence or directing attention away from tinnitus), to adjust threat expectancies and decrease fear.

References

1. Abutan BB, Hoes AW, Van Dalsen CL, Verschuure J, Prins A (1993) Prevalence of hearing impairment and hearing complaints in older adults: a study in general practice. *Fam Pract* 10(4):391–395
2. Anari M, Axelsson A, Eliasson A, Magnusson L (1999) Hypersensitivity to sound: questionnaire data, audiometry and classification. *Scand Audiol* 28(4):219–230
3. Andersson G (2002) Psychological aspects of tinnitus and the application of cognitive-behavioral therapy. *Clin Psychol Rev* 22(7):977–990
4. Andersson G, Stromgren T, Strom L, Lyttkens L (2002) Randomized controlled trial of internet-based cognitive behavior therapy for distress associated with tinnitus. *Psychosom Med* 64(5):810–816
5. Andersson G, Hesser H, Cima RF, Weise C (2013) Autobiographical memory specificity in patients with tinnitus versus patients with depression and normal controls. *Cogn Behav Ther* 42(2):116–126
6. Axelsson A, Ringdahl A (1989) Tinnitus—a study of its prevalence and characteristics. *Br J Audiol* 23(1):53–62
7. Baguley D, McFerran D, Hall D (2013) Tinnitus. *Lancet* 382(9904):1600–1607. [https://doi.org/10.1016/S0140-6736\(13\)60142-7](https://doi.org/10.1016/S0140-6736(13)60142-7)
8. Biesinger E, Heiden C, Greimel V, Lendle T, Höing R, Albigger K (1998) Strategien in der ambulanten Behandlung des Tinnitus. *HNO* 46(2):157–169
9. Blaesing L, Kroener-Herwig B (2012) Self-reported and behavioral sound avoidance in tinnitus and hyperacusis subjects, and association with anxiety ratings. *Int J Audiol* 51(8):611–617
10. Cima RFF (2017) Stress-related Tinnitus treatment protocols. In: Szczeppek A, Mazurek B (eds) *Tinnitus and stress: an interdisciplinary companion for healthcare professionals*. Springer, Cham, pp 139–172
11. Cima RFF, Crombez G, Vlaeyen JW (2011) Catastrophizing and fear of tinnitus predict quality of life in patients with chronic tinnitus. *Ear Hear* 32(5):634–641
12. Cima RFF (2013) *Tinnitus: A CBT-based approach*. Maastricht University, Maastricht The Netherlands & Catholic University of Leuven, Leuven (Joint Doctorate: Dissertation)
13. Cima RFF, Maes IH, Joore MA, Scheyen DJ, El Refaie A, Baguley DM, Vlaeyen JW et al (2012) Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *Lancet* 379(9830):1951–1959
14. Cima RFF, Vlaeyen JWS, Maes IHL, Joore MA, Anteonis LJC (2011) Tinnitus interferes with daily life activities: a psychometric examination of the Tinnitus Disability Index. *Ear Hear* 32(5):623–633
15. Cima RFF, Haider H, Mazurek B, Cederroth CR, Lapira A, Kikidis D, Noreña A (2016) Establishment of a standard for Tinnitus; patient assessment, characterization, and treatment options. Proceedings of the 10th International Tinnitus Research Initiative Conference and 1st EU Cost Action (TINNET) Conference, Nottingham, 16th–18th March
16. Conrad I, Kleinstäuber M, Jasper K, Hiller W, Andersson G, Weise C (2015) The role of dysfunctional cognitions in patients with chronic tinnitus. *Ear Hear* 36(5):e279–e289
17. De Ridder D, Elgoyhen AB, Romo R, Langguth B (2011) Phantom percepts: tinnitus and pain as persisting aversive memory networks. *Proc Natl Acad Sci USA* 108(20):8075–8080
18. Dehmel S, Cui YL, Shore SE (2008) Cross-modal interactions of auditory and somatic inputs in the brainstem and midbrain and their imbalance in tinnitus and deafness. *Am J Audiol* 17(2):S193–S209
19. Drexler M, Überfuhr M, Weddell TD, Lukashkin AN, Wiegrebe L, Krause E, Gürkov R (2014) Multiple indices of the “bounce” phenomenon obtained from the same human ears. *J Assoc Res Otolaryngol* 15(1):57–72
20. Eggermont JJ, Roberts LE (2004) The neuroscience of tinnitus. *Trends Neurosci* 27(11):676–682
21. El-Shunnar SK, Hoare DJ, Smith S, Gander PE, Kang S, Fackrell K, Hall DA (2011) Primary care for tinnitus: practice and opinion among GPs in England. *J Eval Clin Pract* 17(4):684–692
22. Fuller T, Cima R, Langguth B, Mazurek B, Waddell A, Hoare DJ, Vlaeyen JWS (2017) Cognitive behavioural therapy for tinnitus (Protocol). *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD012614>
23. Gilles A, De Ridder D, Van Hal G, Wouters K, Punte AK, Van de Heyning P (2012) Prevalence of leisure noise-induced tinnitus and the attitude toward noise in university students. *Otol Neurotol* 33(6):899–906
24. Goebel G, Fichter M (2005) Psychiatrische Komorbidität bei Tinnitus. In: *Tinnitus*. Springer, Berlin Heidelberg, pp 137–150
25. Guillon MJ, Caston J, Ruel J, Johnson RM, Pujol R, Puel JL (2003) Salicylate induces tinnitus through activation of cochlear NMDA receptors. *J Neurosci* 23(9):3944–3952
26. Hall DA, Láinez MJ, Newman CW, Sanchez TG, Egler M, Tennigkeit F, Langguth B et al (2011) Treatment options for subjective tinnitus: self reports from a sample of general practitioners and ENT physicians within Europe and the USA. *BMC Health Serv Res* 11(1):302
27. Hasson D, Theorell T, Wallén MB, Leineweber C, Canlon B (2011) Stress and prevalence of hearing problems in the Swedish working population. *BMC Public Health* 11(1):130
28. Hallam RS, McKenna L, Shurlock L (2004) Tinnitus impairs cognitive efficiency. *Int J Audiol* 43(4):218–226
29. Hallam RS, Rachman S, Hinchcliffe R (1984) Psychological aspects of tinnitus. In: Rachman S (ed) *Contributions to medical psychology*, vol 3. Pergamon press, Oxford
30. Handscomb L, Hall DA, Shorter GW, Hoare DJ (2016) Online data collection to evaluate a theoretical cognitive model of tinnitus. *Am J Audiol* 25(3S):313–317

31. Hébert S, Canlon B, Hasson D (2012) Emotional exhaustion as a predictor of tinnitus. *Psychother Psychosom* 81(5):324–326
32. Henry JA, Dennis KC, Schechter MA (2005) General review of Tinnitus: prevalence, mechanisms, effects, and management. *J Speech Lang Hear Res* 48(5):1204–1235
33. Hesser H, Pereswetoff-Morath CE, Andersson G (2009) Consequences of controlling background sounds: the effect of experiential avoidance on tinnitus interference. *Rehabil Psychol* 54(4):381
34. Hoare DJ, Gander PE, Collins L, Smith S, Hall DA (2012) Management of tinnitus in English NHS Audiology Departments: an evaluation of current practice. *J Eval Clin Pract* 18(2):326–334
35. Jastreboff PJ, Jastreboff MM (2000) Tinnitus Retraining Therapy (TRT) as a method for treatment of tinnitus and hyperacusis patients. *J Am Acad Audiol* 11(3):162–177
36. Jastreboff PJ, Hazell JW (1993) A neurophysiological approach to tinnitus: clinical implications. *Br J Audiol* 27(1):7–17
37. Jastreboff PJ (1990) Phantom auditory perception (tinnitus): mechanisms of generation and perception. *Neurosci Res* 8(4):221–254
38. Jastreboff PJ, Brennan JF, Coleman JK, Sasaki CT (1988) Phantom auditory sensation in rats: an animal model for tinnitus. *Behav Neurosci* 102(6):811–822
39. Jastreboff PJ, Brennan JF, Sasaki CT (1988) An animal model for tinnitus. *Laryngoscope* 98(3):280–286
40. Jastreboff PJ (2007) Tinnitus retraining therapy. *Prog Brain Res* 166:415–423
41. Kleinstäuber M, Jasper K, Schweda I, Hiller W, Andersson G, Weise C (2013) The role of fear-avoidance cognitions and behaviors in patients with chronic tinnitus. *Cogn Behav Ther* 42(2):84–99
42. Krog NH, Engdahl BO, Tams K (2010) The association between tinnitus and mental health in a general population sample: results from the HUNT Study. *J Psychosom Res* 69(3):289–298
43. Langers DR, de Kleine E, van Dijk P (2012) Tinnitus does not require macroscopic tonotopic map reorganization. *Front Syst Neurosci* 6:2. <https://doi.org/10.3389/fnsys.2012.00002>
44. Langguth B, Landgrebe M, Kleinjung T, Sand GP, Hajak G (2011) Tinnitus and depression. *World J Biol Psychiatry* 12(7):489–500
45. Langguth B, Elgoyhen AB (2012) Current pharmacological treatments for tinnitus. *Expert Opin Pharmacother* 13(17):2495–2509
46. Leaver AM, Seydell-Greenwald A, Rauschecker JP (2016) Auditory-limbic interactions in chronic tinnitus: challenges for neuroimaging research. *Hear Res* 334:49–57
47. Leaver AM, Seydell-Greenwald A, Turesky TK, Morgan S, Kim HJ, Rauschecker JP (2012) Cortico-limbic morphology separates tinnitus from tinnitus distress. *Front Syst Neurosci* 6:21
48. LePage EL (1995) A model for cochlear origin of subjective tinnitus: excitatory drift in the operating point of inner hair cells. *Mechanisms of tinnitus vol 48*. Allyn & Bacon, London
49. Liberman MC, Dodds LW (1987) Acute ultrastructural changes in acoustic trauma: serial-section reconstruction of stereocilia and cuticular plates. *Hear Res* 26(1):45–64
50. Liberman MC (2017) Noise-induced and age-related hearing loss: new perspectives and potential therapies. *F1000Res* 6:927
51. Liberman MC, Kujawa SG (2017) Cochlear synaptopathy in acquired sensorineural hearing loss: manifestations and mechanisms. *Hear Res* 349:138–147
52. Llinás R, Urbano FJ, Leznik E, Ramírez RR, van Marle HJ (2005) Rhythmic and dysrhythmic thalamocortical dynamics: GABA systems and the edge effect. *Trends Neurosci* 28(6):325–333
53. Llinás RR, Ribary U, Jeanmonod D, Kronberg E, Mitra PP (1999) Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci USA* 96(26):15222–15227
54. Liu C, Pan J, Li S, Zhao Y, Wu LY, Berkman CE, Xian M et al (2011) Capture and visualization of hydrogen sulfide by a fluorescent probe. *Angew Chem Int Ed Engl* 50(44):10327–10329
55. Mazurek B, Szczepek AJ, Hebert S (2015) Stress and tinnitus. *HNO* 63(4):258–265
56. McFerran D, Hoare DJ, Carr S, Ray J, Stockdale D (2018) Tinnitus services in the United Kingdom: a survey of patient experiences. *BMC Health Serv Res* 18(1):110
57. McKenna L, Handscomb L, Hoare DJ, Hall DA (2014) A scientific cognitive-behavioral model of tinnitus: novel conceptualizations of tinnitus distress. *Front Neurol* 5:196
58. McKenna L, Hallam RS, Hinchliffe R (1991) The prevalence of psychological disturbance in neuro-otology outpatients. *Clin Otolaryngol* 16(5):452–456
59. McCormack A, Edmondson-Jones M, Somerset S, Hall D (2016) A systematic review of the reporting of tinnitus prevalence and severity. *Hear Res* 337:70–79
60. Mohamad N, Hoare DJ, Hall DA (2016) The consequences of tinnitus and tinnitus severity on cognition: a review of the behavioural evidence. *Hear Res* 332:199–209
61. Noreña AJ (2015) Revisiting the cochlear and central mechanisms of tinnitus and therapeutic approaches. *Audiol Neurotol* 20(Suppl. 1):53–59
62. Noreña AJ, Farley BJ (2013) Tinnitus-related neural activity: theories of generation, propagation, and centralization. *Hear Res* 295:161–171
63. Patuzzi R (2002) Outer hair cells, EP regulation and tinnitus. In: Patuzzi R (ed) *Proceedings of the Seventh International Tinnitus Seminars*. The University of Western Australia, Crawley, pp 16–24
64. Rauschecker JP, Leaver AM, Mühlau M (2010) Tuning out the noise: limbic-auditory interactions in tinnitus. *Neuron* 66(6):819–826
65. Rossiter S, Stevens C, Walker G (2006) Tinnitus and its effect on working memory and attention. *J Speech Lang Hear Res* 49(1):150–160
66. Ruel J, Chabbert C, Nouvian R, Bendris R, Eybalin M, Leger CL, Bourin J, Mersel M, Puel JL (2008) Salicylate enables cochlear arachidonic-acid-sensitive NMDA receptor responses. *J Neurosci* 28(29):7313–7323
67. Seydel C, Haupt H, Olze H, Szczepek AJ, Mazurek B (2013) Gender and chronic tinnitus: differences in tinnitus-related distress depend on age and duration of tinnitus. *Ear Hear* 34(5):661–672
68. Shargorodsky J, Curhan GC, Farwell WR (2010) Prevalence and characteristics of tinnitus among US adults. *Am J Med* 123(8):711–718
69. Shore SE (2005) Multisensory integration in the dorsal cochlear nucleus: unit responses to acoustic and trigeminal ganglion stimulation. *Eur J Neurosci* 21(12):3334–3348
70. Shore SE, El Kashan H, Lu J (2003) Effects of trigeminal ganglion stimulation on unit activity of ventral cochlear nucleus neurons. *Neuroscience* 119(4):1085–1101
71. Shore SE, Roberts LE, Langguth B (2016) Maladaptive plasticity in tinnitus—triggers, mechanisms and treatment. *Nat Rev Neurol* 12(3):150
72. Stevens C, Walker G, Boyer M, Gallagher M (2007) Severe tinnitus and its effect on selective and divided attention. *Int J Audiol* 46(5):208–216
73. Valero MD, Burton JA, Hauser SN, Hackett TA, Ramachandran R, Liberman MC (2017) Noise-induced cochlear synaptopathy in rhesus monkeys (*Macaca mulatta*). *Hear Res* 353:213–223
74. Vlaeyen JW, Linton SJ (2000) Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 85(3):317–332
75. Wu C, Stefanescu RA, Martel DT, Shore SE (2016) Tinnitus: maladaptive auditory–somatosensory plasticity. *Hear Res* 334:20–29
76. Yew KS (2014) Diagnostic approach to patients with tinnitus. *Am Fam Physician* 89(2):106–113
77. Zirke N, Goebel G, Mazurek B (2010) Tinnitus und psychische Komorbiditäten. *HNO* 58(7):726–732
78. Brouwers MC, Kho ME, Brouman GP, Burgers JS, Cluzeau F, Feder G, Littlejohns P et al (2010) AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ* 182(18):E839–E842
79. Fuller TE, Haider HF, Kikidis D, Lapira A, Mazurek B, Noreña A, Brueggemann PG et al (2017) Different teams, same conclusions? A systematic review of existing clinical guidelines for the assessment and treatment of tinnitus in adults. *Front Psychol* 8:206
80. Maes IH, Cima RF, Anteunis LJ, Scheijen DJ, Baguley DM, El Rafea A, Joore MA et al (2014) Cost-effectiveness of specialized treatment based on cognitive behavioral therapy versus usual care for tinnitus. *Otol Neurotol* 35(5):787–795
81. Maes IH, Cima RF, Vlaeyen JW, Anteunis LJ, Joore MA (2013) Tinnitus: a cost study. *Ear Hear* 34(4):508–514
82. Stockdale D, McFerran D, Brazier P, Pritchard C, Kay T, Dowrick C, Hoare DJ (2017) An economic evaluation of the healthcare cost of tinnitus management in the UK. *BMC Health Serv Res* 17(1):577
83. Ahmad N, Seidman M (2004) Tinnitus in the older adult. *Drugs Aging* 21(5):297–305
84. Cima RFF, Andersson G, Schmidt CJ, Henry JA (2014) Cognitive-behavioral treatments or Tinnitus: a review of the literature. *J Am Acad Audiol* 25(1):29–61
85. Coles RRA, Lutman ME, Axelsson A, Hazell JWP (1991) Tinnitus severity gradings: cross sectional studies. Fourth International Tinnitus Seminar, Bourdeaux, 27–30. August, pp 453–455
86. Fackrell K, Hall DA, Barry J, Hoare DJ (2018) Performance of the Tinnitus Functional Index as a diagnostic instrument in a UK clinical population. *Hear Res* 358:74–85
87. Hall DA, Haider H, Szczepek AJ, Lau P, Rabau S, Jones-Diette J, Fuller T et al (2016) Systematic review of outcome domains and instruments used in clinical trials of tinnitus treatments in adults. *Trials* 17(1):270
88. Hallam RS (1996) *Manual of the tinnitus questionnaire (TQ)*. Psychological Corporation, London
89. Kamalski DM, Hoekstra CE, Zanten BGV, Grolman W, Rovers MM (2010) Measuring disease-specific health-related quality of life to evaluate treatment outcomes in tinnitus patients: a systematic review. *Otolaryngol Head Neck Surg* 143(2):181–185
90. Kennedy V, Wilson C, Stephens D (2004) Quality of life and tinnitus. *Audiol Med* 2:29–40
91. Kuk FK, Tyler RS, Russell D, Jordan H (1990) The psychometric properties of a tinnitus handicap questionnaire. *Ear Hear* 11(6):434–445

92. Meikle MB, Griest SE, Stewart BJ, Press LS (1995) Measuring the negative impact of tinnitus: a brief severity index. In: *Abstr Assoc Res Otolaryngol*, vol 167
93. Meikle MB, Henry JA, Griest SE, Stewart BJ, Abrams HB, McArdle R, Folmer RL et al (2012) The tinnitus functional index: development of a new clinical measure for chronic, intrusive tinnitus. *Ear Hear* 33(2):153–176
94. Newman CW, Jacobson GP, Spitzer JB (1996) Development of the tinnitus handicap inventory. *Arch Otolaryngol Neck Surg* 122(2):143–148
95. Wilson PH, Henry J, Bowen M, Haralambous G (1991) Tinnitus reaction questionnaire: psychometric properties of a measure of distress associated with tinnitus. *J Speech Lang Hear Res* 34(1):197–201
96. Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta Psychiatr Scand* 67(6):361–370
97. Arda HN, Tuncel U, Akdogan O, Ozluoglu LN (2003) The role of zinc in the treatment of tinnitus. *Otol Neurotol* 24(1):86–89
98. Arlinger S (2003) Negative consequences of uncorrected hearing loss—a review. *Int J Audiol* 42:2517–2520
99. Association of the Scientific Medical Societies, German S3 guideline 017/064: Chronic tinnitus [AWMF-Register Nr. 017/064 Klasse: S3 Chronischer Tinnitus]. 2015: Germany.
100. Baldo P, Doree C, Lazzarini R, Molin P, McFerran DJ (2006) Antidepressants for patients with tinnitus. *Cochrane Database Syst Rev* 2006(4):CD003853
101. Benninger D, Pal N, Stephan M, Herrmann F, Maire R (2014) Transcranial direct current stimulation for the treatment of chronic tinnitus: a randomized, double-blind, sham controlled study (P2. 283). *Baillieres Clin Neurol* 82(10 Supplement):P2–P283
102. Cavalcanti K, Brasil-Neto JP, Allam N, Boechat-Barros R (2015) A double-blind, placebo-controlled study of the effects of daily tDCS sessions targeting the dorsolateral prefrontal cortex on tinnitus handicap inventory and visual analog scale scores. *Brain Stimul* 8(5):978–980
103. Coelho CB, Tyler R, Hansen M (2007) Zinc as a possible treatment for tinnitus. *Prog Brain Res* 166:279–285
104. Coelho C, Witt SA, Ji H, Hansen MR, Gantz B, Tyler R (2013) Zinc to treat tinnitus in the elderly: a randomized placebo controlled crossover trial. *Otol Neurotol* 34(6):1146–1154
105. Davis A (2011) National survey of hearing and communication
106. Davis PB, Paki B, Hanley PJ (2007) Neuromonic tinnitus treatment: third clinical trial. *Ear Hear* 28(2):242–259
107. Davis PB, Wilde RA, Steed LG, Hanley PJ (2008) Treatment of tinnitus with a customized acoustic neural stimulus: a controlled clinical study. *Ear Nose Throat J* 87(6):330
108. De Ridder D, Vanneste S, Engineer ND, Kilgard MP (2014) Safety and efficacy of vagus nerve stimulation paired with tones for the treatment of tinnitus: a case series. *Neuromodulation* 17(2):170–179
109. Del Bo L, Ambrosetti U (2007) Hearing aids for the treatment of tinnitus. *Prog Brain Res* 166:341–345
110. Dobson K (ed) (2010) *Handbook of cognitive behavioral therapies*, 3rd edn. Guilford, New York
111. Ferguson MA, Kitterick PT, Chong LY, Edmondson-Jones M, Barker F, Hoare DJ (2017) Hearing aids for mild to moderate hearing loss in adults. *Cochrane Database Syst Rev* 2017(9):CD12023
112. Fregni F, Marcondes R, Boggio PS, Marcolin MA, Rigonatti SP, Sanchez TEEA, Pascual-Leone A et al (2006) Transient tinnitus suppression induced by repetitive transcranial magnetic stimulation and transcranial direct current stimulation. *Eur J Neurol* 13(9):996–1001
113. Grewal R, Spielmann PM, Jones SEM, Hussain SSM (2014) Clinical efficacy of tinnitus retraining therapy and cognitive behavioural therapy in the treatment of subjective tinnitus: a systematic review. *J Laryngol Otol* 128(12):1028–1033
114. Hanley PJ, Davis PB (2008) Treatment of tinnitus with a customized, dynamic acoustic neural stimulus: underlying principles and clinical efficacy. *Trends Amplif* 12:210–222
115. Henry JA, Schechter MA, Zaugg TL, Griest S, Jastreboff PJ, Vernon JA, Stewart BJ et al (2006) Clinical trial to compare tinnitus masking and tinnitus retraining therapy. *Acta Otolaryngol* 126(sup556):64–69
116. Henry JA, Frederick M, Sell S, Griest S, Abrams H (2015) Validation of a novel combination hearing aid and tinnitus therapy device. *Ear Hear* 36(1):42–52
117. Henry JA, McMillan G, Dann S, Bennett K, Griest S, Theodoroff S, Saunders G et al (2017) Tinnitus management: randomized controlled trial comparing extended-wear hearing aids, conventional hearing aids, and combination instruments. *J Am Acad Audiol* 28(6):546–561
118. Hesse G (2015) Neueste Behandlungsansätze bei chronischem Tinnitus. *HNO* 63(4):283–290
119. Hesse G (2016) Evidence and evidence gaps in tinnitus therapy. *GMS Current Topics in Otorhinolaryngology*. *Head Neck Surg* 15:Doc04
120. Hesser H, Weise C, Westin VZ, Andersson G (2011) A systematic review and meta-analysis of randomized controlled trials of cognitive-behavioral therapy for tinnitus distress. *Clin Psychol Rev* 31(4):545–553
121. Hilton M, Stuart E (2004) Ginkgo biloba for tinnitus. *Cochrane Database Syst Rev* 2004(2):CD003852
122. Hoare DJ, Hall DA (2011) Clinical guidelines and practice: a commentary on the complexity of tinnitus management. *Eval Health Prof* 34(4):413–420
123. Hoare DJ, Adjajian P, Sereda M (2016) Electrical stimulation of the ear, head, cranial nerve, or cortex for the treatment of tinnitus: a scoping review. *Neural Plast* 2016:5130503
124. Hoare DJ, Edmondson-Jones M, Sereda M, Akeroyd MA, Hall D (2014) Amplification with hearing aids for patients with tinnitus and co-existing hearing loss. *Cochrane Libr* 2014(1):CD010151
125. Hoare DJ, Kowalkowski VL, Kang S, Hall DA (2011) Systematic review and meta-analyses of randomized controlled trials examining tinnitus management. *Laryngoscope* 121(7):1555–1564
126. Hobson J, Chisholm E, El Refaie A (2012) Sound therapy (masking) in the management of tinnitus in adults. *Cochrane Libr* 2012(11):CD006371
127. Hofmann S, Asnaani A, Vonk IJ, Sawyer A, Fang A (2012) The efficacy of cognitive behavioral therapy: a review of meta-analyses. *Cognit Ther Res* 36(5):427–440
128. Humes LE, Rogers SE, Quigley TM, Main AK, Kinney DL, Herring C (2017) The effects of service-delivery model and purchase price on hearing-aid outcomes in older adults: a randomized double-blind placebo-controlled clinical trial. *Am J Audiol* 26(1):53–79
129. Hyvärinen P, Yrttiaho S, Lehtimäki J, Ilmonemi RJ, Mäkitie A, Ylikoski J, Mäkelä JP, Aarnisalo AA (2015) Transcutaneous vagus nerve stimulation modulates tinnitus-related beta- and gamma-band activity. *Ear Hear* 36(3):E76–E85
130. Jastreboff PJ (2011) Tinnitus retraining therapy. In: Möller AR, Kleinjung T, Langguth B, Ridder D (eds) *Textbook of Tinnitus*. Springer, New York, pp 575–596
131. Jufas NE, Wood R (2015) The use of benzodiazepines for tinnitus: systematic review. *J Laryngol Otol* 129(S3):S14–S22
132. Kim JI, Choi JY, Lee DH, Choi TY, Lee MS, Ernst E (2012) Acupuncture for the treatment of tinnitus: a systematic review of randomized clinical trials. *BMC Complement Altern Med* 12(1):97
133. Kompis M, Pelizzone M, Dillier N, Allum J, DeMin N, Senn P (2012) Tinnitus before and 6 months after cochlear implantation. *Audiol Neurotol* 17(3):161–168
134. Kreuzer PM, Landgrebe M, Resch M, Husser O, Schecklmann M, Geisreiter F, Langguth B (2014) Feasibility, safety and efficacy of transcutaneous vagus nerve stimulation in chronic tinnitus: an open pilot study. *Brain Stimul* 7(5):740–747
135. Langguth B (2015) Treatment of tinnitus. *Curr Opin Otolaryngol Head Neck Surg* 23(5):361–368
136. Lehtimäki J, Hyvärinen P, Ylikoski M, Bergholm M, Mäkelä JP, Aarnisalo A, Ylikoski J et al (2013) Transcutaneous vagus nerve stimulation in tinnitus: a pilot study. *Acta Otolaryngol* 133(4):378–382
137. Li SA, Bao L, Chrostowski M (2016) Investigating the effects of a personalized, spectrally altered music-based sound therapy on treating Tinnitus: a blinded, randomized controlled trial. *Audiol Neurotol* 21(5):296–304
138. Lin FR et al (2011) Hearing loss and incident dementia. *Arch Neurol* 68(2):214–220
139. Martinez-Devesa P, Perera R, Theodoulou M, Waddell A (2010) Cognitive behavioural therapy for tinnitus. *Cochrane Database Syst Rev* 2010(1):CD005233
140. Matthews L (2013) Hearing loss, tinnitus and mental health. A literature review. *Action on Hearing Loss*, London, UK
141. McKenna L, Marks EM, Hallsworth CA, Schaeffer R (2017) Mindfulness-based cognitive therapy as a treatment for chronic tinnitus: a randomized controlled trial. *Psychother Psychosom* 86(6):351–361
142. Megwalu U, Piccirillo JF, Finnell J (2005) The effects of melatonin on Tinnitus and sleep. *Otolaryngol Head Neck Surg* 133(2):P89
143. Meng Z, Liu S, Zheng Y, Phillips JS (2011) Repetitive transcranial magnetic stimulation for tinnitus. *Cochrane Libr* 2011(10):CD007946
144. Nguyen MF, Bonnefoy M, Adrait A, Gueugnon M, Petitot C, Collet L, Perrot X et al (2017) Efficacy of hearing aids on the cognitive status of patients with Alzheimer's disease and hearing loss: A multicenter controlled randomized trial. *J Alzheimers Dis* 58(1):123–137
145. Newman CW, Sandridge SA (2012) A comparison of benefit and economic value between two sound therapy tinnitus management options. *J Am Acad Audiol* 23(2):126–138
146. Noreña AJ (2011) An integrative model of tinnitus based on a central gain controlling neural sensitivity. *Neurosci Biobehav Rev* 35(5):1089–1109
147. Nyenhuis N, Goltz D, Kröner-Herwig B (2013) A systematic review and meta-analysis on the efficacy of self-help interventions in Tinnitus. *Cogn Behav Ther* 42(2):159–169
148. Okamoto H, Stracke H, Stoll W, Pantev C (2010) Listening to tailor-made notched music reduces

- tinnitus loudness and tinnitus-related auditory cortex activity. *Proc Natl Acad Sci USA* 107(3):1207–1210
149. Pal N, Maire R, Stephan MA, Herrmann FR, Benninger DH (2015) Transcranial direct current stimulation for the treatment of chronic tinnitus: a randomized controlled study. *Brain Stimul* 8(6):1101–1107
 150. Pantev C, Okamoto H, Teismann H (2012) Tinnitus: the dark side of the auditory cortex plasticity. *Ann NY Acad Sci* 1252(1):253–258
 151. Parazzini M, Del Bo L, Jastreboff M, Tognola G, Ravazzani P (2011) Open ear hearing aids in tinnitus therapy: an efficacy comparison with sound generators. *Int J Audiol* 50(8):548–553
 152. Parra LC, Pearlmutter BA (2007) Illusory percepts from auditory adaptation. *J Acoust Soc Am* 121(3):1632–1641
 153. Phillips JS, McFerran D (2010) Tinnitus retraining therapy (TRT) for tinnitus. *Cochrane Libr* 2010(3):CD007330
 154. Pirodda A, Raimondi MC, Ferri GG (2010) Exploring the reasons why melatonin can improve tinnitus. *Med Hypotheses* 75(2):190–191
 155. Posadzki P, Watson L, Ernst E (2013) Herb-drug interactions: an overview of systematic reviews. *Br J Clin Pharmacol* 75(3):603–618
 156. Quaranta N, Wagstaff S, Baguley DM (2004) Tinnitus and cochlear implantation. *Int J Audiol* 43(5):245–251
 157. Ramakers GG, van Zon A, Stegeman I, Grolman W (2015) The effect of cochlear implantation on tinnitus in patients with bilateral hearing loss: a systematic review. *Laryngoscope* 125(11):2584–2592
 158. Reato D, Rahman A, Bikson M, Parra LC (2013) Effects of weak transcranial alternating current stimulation on brain activity—a review of known mechanisms from animal studies. *Front Hum Neurosci* 7:687
 159. Rejali D, Sivakumar A, Balaji N (2004) Ginkgo biloba does not benefit patients with tinnitus: a randomized placebo-controlled double-blind trial and meta-analysis of randomized trials. *Clin Otolaryngol Allied Sci* 29(3):226–231
 160. Rosenberg SI, Silverstein H, Rowan PT, Olds MJ (1998) Effect of melatonin on tinnitus. *Laryngoscope* 108(3):305–310
 161. Sanchez TG, Medeiros IR, Levy CP, Ramalho JR, Bento RF (2005) Tinnitus in normally hearing patients: clinical aspects and repercussions. *Braz J Otorhinolaryngol* 71(4):427–431
 162. Schaeffe R, McAlpine D (2011) Tinnitus with a normal audiogram: physiological evidence for hidden hearing loss and computational model. *J Neurosci* 31(38):13452–13457
 163. Shekhawat GS, Searchfield GD, Stinear CM (2014) Randomized trial of transcranial direct current stimulation and hearing aids for tinnitus management. *Neurorehabil Neural Repair* 28(5):410–419
 164. Shin JW, Lee HK (2016) Tinnitus retraining therapy. *Hanyang Med Rev* 36(2):120–124
 165. Soleimani R, Jalali MM, Hasandokht T (2016) Therapeutic impact of repetitive transcranial magnetic stimulation (rTMS) on tinnitus: a systematic review and meta-analysis. *Eur Arch Otorhinolaryngol* 273(7):1663–1675
 166. Song JJ, Vanneste S, Van de Heyning P, De Ridder D (2012) Transcranial direct current stimulation in tinnitus patients: a systemic review and meta-analysis. *Sci World J* 2012:427941
 167. Stein A, Wunderlich R, Lau P, Engell A, Wollbrink A, Shaykevich A, Pantev C et al (2016) Clinical trial on tonal tinnitus with tailor-made notched music training. *BMC Neurol* 16(1):38
 168. Topak M, Sahin-Yilmaz A, Ozdoganoglu T, Yilmaz HB, Ozbay M, Kulekci M (2009) Intratympanic methylprednisolone injections for subjective tinnitus. *J Laryngol Otol* 123(11):1221–1225
 169. Tunkel DE, Bauer CA, Sun GH, Rosenfeld RM, Chandrasekhar SS, Cunningham ER Jr, Henry JA et al (2014) Clinical practice guideline: tinnitus. *Otolaryngol Head Neck Surg* 151:S1–S40
 170. Tyler RS, Noble W, Coelho CB, Ji H (2012) Tinnitus retraining therapy: mixing point and total masking are equally effective. *Ear Hear* 33(5):588–594
 171. Tyler R, Cacace A, Stocking C, Tarver B, Engineer N, Martin J, Burrell C et al (2017) Vagus nerve stimulation paired with tones for the treatment of tinnitus: a prospective randomized double-blind controlled pilot study in humans. *Sci Rep* 7(1):11960
 172. Vanneste S, Song JJ, De Ridder D (2013) Tinnitus and musical hallucinosis: the same but more. *Neuroimage* 82:373–383
 173. von Boetticher A (2011) Ginkgo biloba extract in the treatment of tinnitus: a systematic review. *Neuropsychiatr Dis Treat* 7:441
 174. Von Korff M, Moore JC (2001) Stepped care for back pain: activating approaches for primary care. *Ann Intern Med* 134(9Pt 2):911–917
 175. Wegger M, Ovesen T, Larsen DG (2017) Acoustic coordinated reset neuromodulation: a systematic review of a novel therapy for Tinnitus. *Front Neurol* 8:36
 176. Wilson BS, Tucci DL, Merson MH, O'Donoghue GM (2017) Global hearing health care: new findings and perspectives. *Lancet* 390(10111):2503–2515
 177. Yang S, Weiner BD, Zhang LS, Cho SJ, Bao S (2011) Homeostatic plasticity drives tinnitus perception in an animal model. *Proc Natl Acad Sci USA* 108(36):14974–14979
 178. Ylikoski J, Lehtimäki J, Pirvola U, Mäkitie A, Aarnisalo A, Hyvärinen P, Ylikoski M (2017) Non-invasive vagus nerve stimulation reduces sympathetic preponderance in patients with tinnitus. *Acta Otolaryngol* 137(4):426–431
 179. Altissimi G, Salvati M, Turchetta R, Orlando MP, Greco A, De Vincentiis M, Cianfrone G et al (2016) When alarm bells ring: emergency tinnitus. *Eur Rev Med Pharmacol Sci* 20(14):2955–2973
 180. Axelsson A, Prasher D (2000) Tinnitus induced by occupational and leisure noise. *Noise Health* 2(8):47
 181. Barlow J, Wright C, Sheasby J, Turner A, Hainsworth J (2002) Self-management approaches for people with chronic conditions: a review. *Patient Educ Couns* 48(2):177–187
 182. Brodney S, Fiwler FJ, Wexler R, Bowen M (2016) Shared decision-making in clinical practice: examples of successful implementation. *Eur J Pers Cent Healthc* 4(4):656–659
 183. Crönlein T, Langguth B, Pregler M, Kreuzer PM, Wetter TC, Schecklmann M (2016) Insomnia in patients with chronic tinnitus: cognitive and emotional distress as moderator variables. *J Psychosom Res* 83:65–68
 184. Dibb B, Yardley L (2006) How does social comparison within a self-help group influence adjustment to chronic illness? A longitudinal study. *Soc Sci Med* 63(6):1602–1613
 185. Eysenbach G, Powell J, Englesakis M, Rizo C, Stern A (2004) Health related virtual communities and electronic support groups: systematic review of the effects of online peer to peer interactions. *BMJ* 328(7449):1166
 186. Fackrell K, Fearnley C, Hoare DJ, Sereda M (2015) Hyperacusis questionnaire as a tool for measuring hypersensitivity to sound in a tinnitus research population. *Biomed Res Int* 2015:290425
 187. Gander PE, Hoare DJ, Collins L, Smith S, Hall DA (2011) Tinnitus referral pathways within the National Health Service in England: a survey of their perceived effectiveness among audiology staff. *BMC Health Serv Res* 11(1):162
 188. Greenwell K, Sereda M, Coulson N, El Refaie A, Hoare DJ (2016) A systematic review of techniques and effects of self-help interventions for tinnitus: Application of taxonomies from health psychology. *Int J Audiol* 55(sup3):S79–S89
 189. Hallberg LR, Erlandsson SI (1993) Tinnitus characteristics in tinnitus complainers and noncomplainers. *Br J Audiol* 27(1):19–27
 190. Hayes SC, Luoma JB, Bond FW, Masuda A, Lillis J (2006) Acceptance and commitment therapy: model, processes and outcomes. *Behav Res Ther* 44(1):1–25
 191. Henry JL, Wilson PH (2002) Tinnitus: a self-management guide for the ringing in your ears. Allyn & Bacon, Boston, US
 192. Henry JA, Griest S, Zaugg TL, Thielman E, Kaelin C, Galvez G, Carlson KF (2015) Tinnitus and hearing survey: a screening tool to differentiate bothersome tinnitus from hearing difficulties. *Am J Audiol* 24(1):66–77
 193. Hesser H, Weise C, Rief W, Andersson G (2011) The effect of waiting: a meta-analysis of wait-list control groups in trials for tinnitus distress. *J Psychosom Res* 70(4):378–384
 194. Hoare DJ, Ainscough E, Smith S, Greenwell K (2017) A qualitative analysis of threads in online support groups for tinnitus. *J Hear Sci* 7(2):103
 195. Hong O, Kerr MJ, Poling GL, Dhar S (2013) Understanding and preventing noise-induced hearing loss. *Dis Mon* 59(4):110–118
 196. Kabat-Zinn J (1982) An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: Theoretical considerations and preliminary results. *Gen Hosp Psychiatry* 4(1):33–47
 197. Kreuzer PM, Goetz M, Holl M, Schecklmann M, Landgrebe M, Staudinger S, Langguth B (2012) Mindfulness- and body-psychotherapy-based group treatment of chronic tinnitus: a randomized controlled pilot study. *BMC Complement Altern Med* 12(1):235
 198. Meltzer JB, Herzfeld M (2014) Tinnitus, hyperacusis, and misophonia toolbox. *Semin Hear* 35(2):121–130
 199. Sereda M, Hoare DJ (2015) Self-help interventions for Tinnitus. In: Baguley D, Fagelson M (eds) Tinnitus: clinical and research perspectives, vol 227. Pleural Publishing, San Diego
 200. Malouff JM, Noble W, Schutte NS, Bhullar N (2010) The effectiveness of bibliotherapy in alleviating tinnitus-related distress. *J Psychosom Res* 68(3):245–251
 201. McKenna L, Baguley D, McFerran D (2010) Living with tinnitus and hyperacusis. SPCK, London, UK
 202. McMullan M (2006) Patients using the Internet to obtain health information: how this affects the patient-health professional relationship. *Patient Educ Couns* 63(1):24–28
 203. Newman CW, Sandridge SA (2005) Incorporating group and individual sessions into a tinnitus management clinic. In: Tinnitus treatment: clinical protocols. Thieme, New York, pp 187–197
 204. Nondahl DM, Cruickshanks KJ, Wiley TL, Klein R, Klein BE, Tweed TS (2002) Prevalence and 5-year

- incidence of tinnitus among older adults: the epidemiology of hearing loss study. *J Am Acad Audiol* 13(6):323–331
205. Paglialonga A, Tognola G, Pincirolfi F (2015) Apps for hearing healthcare. In: MIE, pp 666–668
 206. Philippot P, Neff F, Clauw L, Romrée M, Segal Z (2012) A randomized controlled trial of mindfulness-based cognitive therapy for treating tinnitus. *Clin Psychol Psychother* 19(5):411–419
 207. Phillips JS, McFerran DJ, Hall DA, Hoare DJ (2018) The natural history of subjective tinnitus in adults: a systematic review and meta-analysis of no-intervention periods in controlled trials. *Laryngoscope* 128(1):217–227
 208. Pryce H, Culhane BA, Swift S, Claesen B, Shaw R, Hall A (2017) Shared decision making in the Tinnitus clinic—what are patient preferences for treatment? *J Hear Sci* 7(2):121
 209. Sereda M, Hoare DJ, Nicholson R, Smith S, Hall DA (2015) Consensus on hearing aid candidature and fitting for mild hearing loss, with and without tinnitus: Delphi review. *Ear Hear* 36(4):417–429
 210. Sereda M, Stockdale D, Newton K, Smith S (2017) Mobile applications for management of tinnitus. *J Hear Sci* 7(2):124
 211. Shekhawat GS, Searchfield GD, Stinear CM (2013) Role of hearing aids in tinnitus intervention: a scoping review. *J Am Acad Audiol* 24(8):747–762
 212. Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, Trevena L (2017) Decision aids for people facing health treatment or screening decisions. *Cochrane Libr* 2017(4):CD001431
 213. St. Claire LS, Stothart G, McKenna L, Rogers PJ (2010) Caffeine abstinence: an ineffective and potentially distressing tinnitus therapy. *Int J Audiol* 49(1):24–29
 214. Weise C, Hesser H, Andersson G, Nyenhuis N, Zastrutski S, Kröner-Herwig B, Jäger B (2013) The role of catastrophizing in recent onset tinnitus: its nature and association with tinnitus distress and medical utilization. *Int J Audiol* 52(3):177–188
 215. Weise C, Kleinstäuber M, Andersson G (2016) Internet-delivered cognitive-behavior therapy for tinnitus: a randomized controlled trial. *Psychosom Med* 78(4):501–510
 216. Kircher ML, Standring RT, Leonetti JP (2008) Neuroradiologic assessment of pulsatile tinnitus. *Otolaryngol Head Neck Surg* 139(2):144
 217. Sismanis A (1998) Pulsatile Tinnitus A 15-year experience. *Am J Otol* 19(4):1998
 218. Sismanis A (2003) Pulsatile Tinnitus. *Otolaryngol Clin North Am* 36(2):389–402
 219. Pegge SA, Steens SC, Kunst HP, Meijer FJ (2017) Pulsatile tinnitus: differential diagnosis and radiological work-up. *Curr Radiol Rep* 5(1):5
 220. Hofmann E, Behr R, Neumann-Haefelin T, Schwager K (2013) Pulsatile tinnitus: imaging and differential diagnosis. *Dtsch Arztebl Int* 110(26):451
 221. Abel MD, Levine RA (2004) Muscle contractions and auditory perception in tinnitus patients and nonclinical subjects. *Cranio* 22:181–191
 222. Cacace AT (2003) Expanding the biological basis of tinnitus: crossmodal origins and the role of neuroplasticity. *Hear Res* 175:112–132
 223. Cacace AT, Cousins JP, Parnes SM, McFarland DJ, Semenov D, Holmes T et al (1999) Cutaneous-evoked tinnitus. II. Review of neuroanatomical, physiological and functional imaging studies. *Audiol Neurootol* 4:258–268
 224. Cacace AT, Cousins JP, Parnes SM, McFarland DJ, Semenov D, Holmes T et al (1999) Cutaneous-evoked tinnitus. I. Phenomenology, psychophysical and functional imaging. *Audiol Neurootol* 4:247–268
 225. Haider HF, Hoare DJ, Costa RF, Potgieter I, Kikidis D, Lapira A, Paço JC et al (2017) Pathophysiology, diagnosis and treatment of somatosensory tinnitus: a scoping review. *Front Neurosci* 11:207
 226. Herraiz C (2008) Assessing the cause of tinnitus for therapeutic options. *Expert Opin Med Diagn* 2(10):1183–1196
 227. Koehler SD, Shore SE (2013) Stimulus timing-dependent plasticity in dorsal cochlear nucleus is altered in tinnitus. *J Neurosci* 33:19647–19656
 228. Latifpour DH, Grenner J, Sjødahl C (2009) The effect of a new treatment based on somatosensory stimulation in a group of patients with somatically related tinnitus. *Int Tinnitus J* 15:94
 229. Levine RA, Abel M, Cheng H (2003) CNS somatosensory-auditory interactions elicit or modulate tinnitus. *Exp Brain Res* 153:643–648
 230. Levine RA, Nam EC, Oron Y, Melcher JR (2007) Evidence for a tinnitus subgroup responsive to somatosensory based treatment modalities. *Prog Brain Res* 166:195–207
 231. Levine RA, Nam EC, Melcher J (2008) Somatosensory pulsatile tinnitus syndrome: somatic testing identifies a pulsatile tinnitus subtype that implicates the somatosensory system. *Trends Amplif* 2:242–253
 232. Lockwood AH, Wack DS, Burkard RF, Coad ML, Reyes SA, Arnold SA et al (2001) The functional anatomy of gaze-evoked tinnitus and sustained lateral gaze. *Baillieres Clin Neurol* 56:472–480
 233. Oostendorp RA, Bakker I, Elvers H, Mikolajewska E, Michiels S, De Hertogh W et al (2016) Cervicogenic somatosensory tinnitus: an indication for manual therapy? Part 1: theoretical concept. *Man Ther* 23:120–123
 234. Rocha CAB, Sanchez TG, Tesseroli de Siqueira JT (2008) Myofascial trigger point: a possible way of modulating tinnitus. *Audiol Neurootol* 13:153–160
 235. Sanchez TG, Akemi M (2008) Modulating tinnitus with visual, muscular, and tactile stimulation. *Semin Hear* 29:350–360
 236. Sanchez TG, Guerra GCY, Lorenzi MC, Brandão AL, Bento RF (2002) The influence of voluntary muscle contractions upon the onset and modulation of tinnitus. *Audiol Neurootol* 7:370–375
 237. Sanchez TG, da Silva Lima A, Brandão AL, Lorenzi MC, Bento RF (2007) Somatic modulation of tinnitus: test reliability and results after repetitive muscle contraction training. *Ann Otol Rhinol Laryngol* 116:30–35
 238. Sanchez TG, Rocha CB (2011) Diagnosis and management of somatosensory tinnitus: review article. *Clinics* 66:1089–1094
 239. Simmons R, Dambra C, Lobarinas E, Stocking C, Salvi R (2008) Head, neck, and eye movements that modulate tinnitus. *Semin Hear* 29:361–370
 240. Vielsmeier V, Kleinjung T, Strutz J, Bürgers R, Kreuzer PM, Langguth B (2011) Tinnitus with temporomandibular joint disorders a specific entity of tinnitus patients? *Otolaryngol Head Neck Surg* 145:748–752
 241. Won JY, Yoo S, Lee SK, Choi HK, Yakunina N, Le Q et al (2013) Prevalence and factors associated with neck and jaw muscle modulation of tinnitus. *Audiol Neurootol* 18:261–273
 242. Beck AT (1976) Cognitive therapy and the emotional disorders. International Universities Press, Oxford
 243. Bohlmeijer E, Prenger R, Taal E, Cuijpers P (2010) The effects of mindfulness-based stress reduction therapy on mental health of adults with a chronic medical disease: a meta-analysis. *J Psychosom Res* 68(6):539–544
 244. Ellis A, Grieger R (1977) Handbook of rational-emotive therapy. Springer, New York
 245. Fjorback LO, Arendt M, Ørnbøl E, Fink P, Walach H (2011) Mindfulness-based stress reduction and mindfulness-based cognitive therapy—a systematic review of randomized controlled trials. *Acta Psychiatr Scand* 124(2):102–119
 246. Hallam RS, Jakes SC, Hinchcliffe R (1988) Cognitive variables in tinnitus annoyance. *Br J Clin Psychol* 27(3):213–222
 247. Hayes SC, Strosahl KD, Wilson KG (1999) Acceptance and commitment therapy: an experiential approach to behavior change. Guilford, New York
 248. Hayes SC, Wilson KG (1994) Acceptance and commitment therapy: altering the verbal support for experiential avoidance. *Behav Anal* 17(2):289–303
 249. Hofmann SG, Sawyer AT, Fang A (2010) The empirical status of the “new wave” of cognitive behavioral therapy. *Psychiatr Clin North Am* 33(3):701–710
 250. Kauth MR, Sullivan G, Blevins D, Cully JA, Landes RD, Said Q, Teasdale TA (2010) Employing external facilitation to implement cognitive behavioral therapy in VA clinics: a pilot study. *Implement Sci* 5:75
 251. Ludwig DS, Kabat-Zinn J (2008) Mindfulness in medicine. *JAMA* 300(11):1350–1352
 252. Scott B, Lindberg P, Lyttkens L, Melin L (1985) Psychological treatment of tinnitus. An experimental group study. *Scand Audiol* 14(4):223–230
 253. Shapiro SL, Brown KW, Thoresen C, Plante TG (2011) The moderation of Mindfulness-based stress reduction effects by trait mindfulness: results from a randomized controlled trial. *J Clin Psychol* 67(3):267–277
 254. Sweetow RW (1986) Cognitive aspects of tinnitus patient management. *Ear Hear* 7(6):390–396
 255. Teasdale JD, Scott J, Moore RG, Hayhurst H, Pope M, Paykel ES (2001) How does cognitive therapy prevent relapse in residual depression? Evidence from a controlled trial. *J Consult Clin Psychol* 69(3):347–357
 256. Volders S, den Hollander M, de Peuter S, Vlaeyen JWS (2011) Protocolaire behandeling van patiënten met chronische lage rugpijn. Exposure-in-vivo therapie Protocolaire behandelingsenvoor volwassenen met psychische klachten, vol 2. Boom, Amsterdam, pp 169–196
 257. Wilson PH (2006) Classical conditioning as the basis for the effective treatment of tinnitus-related distress. *ORL J Otorhinolaryngol Relat Spec* 68(1):6–13
 258. Hall DA, Domingo SZ, Hamdache LZ, Manchiaiah V, Thammaiah S, Evans C, Wong L et al (2017) A good practice guideline for translating questionnaires in ENT and Audiology. *J Hear Sci* 7(2):146