

A New Bidirectional Associative Neutrosophic Cognitive Dynamical System approaches to study Youth Violence

A. Victor Devadoss
Head & Associate Professor
PG & Research Department of mathematics
Loyola College, Chennai-34, India

A. Felix
Ph.D. Research scholar
PG & Research Department of mathematics
Loyola College, Chennai-34, India

ABSTRACT

BAM and NCMs have been applied in many fields to show the relationship between the essential components. In this paper we introduce a new Neutrosophic Bimodal called Bidirectional Associative Neutrosophic Cognitive Maps (BANCMs) to analyze the problem of youth violence since enormous growth of violent escalation among youth in every country and culture is alarming. This new model can act simultaneously to give the solution in more accurately. This article organized as follows, section one describes the problem of our study. Section two recalls the basic concepts of BAM and NCM. Section three gives the new modal BANCM. Section four, the adaptation of the problem using BANCM and final section derived conclusion, suggestion and scope for future study.

Keywords: BAM, NCM, BANCM, fixed point, limit cycle, bihidden pattern.

1. INTRODUCTION

Cognitive maps were initially introduced by Robert Axelrod [1] in 1976. He used this modal to study decision making in social and political systems. Bart Kosco enhanced the power of cognitive maps considering fuzzy values for the concepts of the cognitive map and fuzzy degrees of interrelationships between concepts [2]. In 2003, NCM was introduced by W.B.Vasanhakandasamy and Florentin Samaranche to study the aspects of migrant laborers living with HIV/AIDS [7]. The BAM modal was introduced by Bart Kosco in 1988 and modified in the year 2001[3]. Praveen prakash has introduced a bimodal called Fuzzy Cognitive Relational Maps (FCRM) in the year 2010. Thirusangu et.al 2012 introduced a Bimodal Called Bidirectional Associative Fuzzy Cognitive maps (BACM) [6]. In this paper we introduce a new BANCM modal by combining BAM and NCM since to connect the attributes there is an indeterminate occurs. This model work simultaneously to study the problem of youth violence. Violence among young people is one of the most visible forms of violence. Both fatal and non-fatal assaults involving young people contribute greatly to the burden of premature death, injury and disability. Youth violence deeply harms not only its victims, but also their families, friends and communities. An average of 565 children, adolescents and young adults between the ages of 10 and 29 years die each day as a result of interpersonal violence across the world says the World Report on Violence and Health,2002. A large number of these are from the South-East Asia Region. And also enormous growth of violent escalation among youth in every country and culture is alarming. With this background of information the study was conducted to assess the causes and effect of youth violence. Moreover the data is an unsupervised one obtained from survey as well as interviews. We have got two sets of attributes by the two different

experts, in which one relates to the BAM dynamical system and other with a NCM dynamical system and if we wish to see the stage-by-stage effect of the attributes on the system, we need a dynamical system. Hence we introduce a dynamical bisystem

2. THEORITICAL BACKGROUND

2.1 BAM MODEL

A group of neurons forms a field. Neural networks contain many fields of neurons. F_x denotes a neuron field which contains n neurons and F_y denotes a neuron field which contains p neurons.

Neuronal Dynamical Systems: The neuronal dynamical system is described by a system of first order differential equations that govern the time evaluation of the neuronal activations or membrane potentials.

$$\dot{X}_i = g_i(X, Y, \dots), \dot{Y}_j = h_j(X, Y, \dots)$$

where x_i and y_j denote respectively the activation time function of the i^{th} neuron in F_x and the j^{th} neuron in F_y . The over dot denotes time differentiation, g_i and h_j are functions of X, Y etc., where $X(t) = (x_1(t), \dots, x_n(t))$, $Y(t) = (y_1(t), \dots, y_p(t))$

Define the state of the neuronal dynamical system at time t . Additive bivalent Models describe asynchronous and stochastic behavior. At each moment each neuron can randomly decide whether to change state, or whether to omit a new signal given its current activation. The BAM is a non-adaptive, additive, bivalent neural network.

2.1.1 Bivalent Additive BAM

In neural literature, the discrete version of the earlier equations is often referred to as the Bidirectional Associative Memories or BAMs. A discrete additive BAM with threshold signal functions, arbitrary thresholds and inputs, an arbitrary but a constant synaptic connection matrix M and discrete time steps K are defined by the equations.

$$x_i^{k+1} = \sum_j^p S_j(y_j^k) m_{ij} + I_i$$

$$y_j^{k+1} = \sum_i^n S_i(x_i^k) m_{ij} + I_j$$

Where, $m_{ij} \in M$ S_i and S_j are the signal functions. They represent binary or bipolar threshold functions. For arbitrary real-valued thresholds $U=(U_1, U_2, \dots, U_n)$ for F_x neurons and $V=(V_1, V_2, \dots, V_p)$ for F_y neurons. The threshold binary signal functions corresponds neurons.

2.1.2 Synaptic connection Matrices

Let us suppose that the field F_x with n neurons is synaptically connected to the field F_y with p neurons. Let m_{ij} be a synapse where the axon from the i^{th} neuron in F terminates, m_{ij} can be positive, negative or zero. The synaptic matrix M is a $n \times p$ matrix of real numbers whose entries are the synaptic efficacies m_{ij} . The matrix M describes the forward projections from the neuronal field F_x to the neuronal field F_y . Similarly, M^T , a $p \times n$ synaptic matrix and describes the backward projections F_y to F_x .

2.1.3 Unidirectional Networks

These kinds of networks occur when a neuron synoptically interconnects to itself. The matrix N is $n \times n$ square matrix.

Bidirectional Networks.

A network is said to be a bidirectional network if $M = N^T$ and $N = M^T$

2.1.4 Bidirectional Associative Memories

When the activation dynamics of the neuronal fields F_x and F_y lead to the overall stable behavior, the bi-directional networks are called as Bi-directional Associative Memories or BAM. A unidirectional network also defines a BAM if M is symmetric i.e. $M = M$. In the next section, we proceed on to give more details about this BAM.

2.1.5 Additive Activation Models

An additive activation model is defined by a system of $n + p$ coupled first-order differential equations that interconnects the fields F_x and F_y through the constant synaptic matrices M and N described earlier. $S_i(x_i)$ and $S_j(y_j)$ denote respectively the signal function of the i^{th} neuron in the field F_x and the signal function of the j^{th} neuron in the field F_y . Discrete additive activation models correspond to neurons with threshold signal functions. The neurons can assume only two values **ON** and **OFF**. **ON** represents the signal value $+1$ and **OFF** represents 0 or -1 (-1 when the representation is bipolar). The bipolar version of these equations yield the signal value -1 when $x_i < V_i$ or $y_j < V_j$.

$$\begin{aligned} \dot{x} &= -A_i x_i + \sum_j^p S_j(y_j^k) m_{ji} + I_i \\ \dot{y} &= -A_j y_j + \sum_i^n S_i(x_i^k) m_{ij} + I_j \end{aligned}$$

The bivalent signal functions allow us to model complex asynchronous state-change patterns. At any moment different neurons can decide whether to compare their activation to their threshold. At each moment any of the 2^n subsets of F_x neurons or the 2^p subsets of the F_y neurons can decide to change state. Each neuron may randomly decide whether to check the threshold conditions in the equations given above.

At each moment each neuron defines a random variable that can assume the value **ON** ($+1$) or **OFF** (0 or -1). The network is often assumed to be deterministic and state changes are synchronous i.e. an entire field of neurons is updated at a time. In case of simple asynchrony only one neuron makes a state change decision at a time. When the subsets represent the entire fields F_x and F_y synchronous state change results.

In a real life problem the entries of the constant synaptic matrix M depends upon the investigator's feelings. The synaptic matrix is given a weight age according to their feelings. If $x = F_x$ and $y = F_y$ the forward projections from F_x to F_y is defined by the matrix M : $\{P(x_i, x_j)\} = M$, $1 < i < n$, $1 < j < p$.

The backward projection is defined by the Matrix M^T : $\{F(y_j, x_i)\} = (m_{ij}) = M^T$, $1 < i < n$, $1 < j < p$.

2.1.6 Bidirectional Stability

All BAM state changes lead a fixed-point stability. This property holds for synchronous as well as asynchronous state changes.

A BAM system (F_x, F_y, M) is bidirectionally stable if all inputs converge to fixed point equilibrium. Bidirectional stability is a dynamic equilibrium. The same signal information flows back and forth in a bidirectional fixed point.

Let us suppose that A denotes a binary n -vector and B denotes a binary p -vector. Let A be initial input to the BAM system. Then the BAM equilibrates a bi directional fixed point (A_i, B_i) as

$$\begin{aligned} A &\rightarrow M \rightarrow B \\ A' &\leftarrow M^T \leftarrow B \\ A' &\rightarrow M \rightarrow B' \\ A'' &\leftarrow M^T \leftarrow B' \\ &\dots \\ A_f &\rightarrow M \rightarrow B_f \\ A_f &\leftarrow M^T \leftarrow B_f \end{aligned}$$

Where A' , A'' ... and B' , B'' ... represents intermediate or transient signal state vectors between A and A_f , B and B_f respectively. The fixed point of a bidirectional system is time dependent. The fixed point for the initial input vectors can be attained at different times which are illustrated later. Based on the synaptic matrix M which is developed by the investigators feelings, the time at which bi directional stability is attained also varies accordingly

2.3 NEUTROSOPHIC COGNITIVE MAPS

Fuzzy Cognitive Maps (FCMs) are more applicable when the data in the first Place is an unsupervised one. The FCMs work on the opinion of experts. FCMs model the world as a collection of classes and causal relations between classes.

2.3.1 Definition

When the nodes of the NCM are fuzzy sets then they are called as fuzzy nodes.

2.3.2 Definition

NCMs with edge weights or causalities from the set $\{-1, 0, 1, I\}$ are called simple NCMs

2.3.3 Definition

Let C_i and C_j denote the two nodes of the NCM. The directed edge from C_i to C_j denote the causality of C_i on C_j called connections. Every edge in the NCM is weighted with a number in the set $\{-1, 0, 1, I\}$. Let e_{ij} be the weight of the directed edge $C_i C_j$, $e_{ij} \in \{-1, 0, 1, I\}$. $e_{ij} = 0$ if C_i does not have any effect on C_j , $e_{ij} = 1$ if increase (or decrease) in C_i causes increase (or decreases) in C_j . $e_{ij} = -1$ if increase (or decrease) in C_i causes decrease (or increase) in C_j . $e_{ij} = I$ if the relation or effect of C_i on C_j is an indeterminate.

2.3.4 Definition

Let C_1, C_2, \dots, C_n be nodes of a NCM. Let the neutrosophic matrix $N(E)$ be defined as $N(E) = (e_{ij})$ where e_{ij} is the weight of

the directed edge $C_i C_j$, where $e_{ij} \in \{.1, 0, 1, I\}$. $N(E)$ is called the neutrosophic adjacency matrix of the NCM.

2.3.5 Definition

Let C_1, C_2, \dots, C_n be the nodes of the NCM. Let $A = \{a_1, a_2, \dots, a_n\}$, where $a_i \in \{0, 1, I\}$. A is called the instantaneous state neutrosophic vector and it denotes the ON-OFF-INDETERMINATE state position of the node at an instant

$a_i = 0$ if a_i is OFF (no effect)

$a_i = 1$ if a_i is ON (has effect)

$a_i = I$ if a_i is INDETERMINATE (effect cannot be determined) for $i = 1, 2, \dots, n$

2.3.6 Definition

Let C_1, C_2, \dots, C_n be the nodes of the FCM. Let $C_1 C_2, C_2 C_3, \dots, C_i C_j$ be the edges of the NCM. Then the edges form a directed cycle. An NCM is said to be cyclic if it possesses a directed cyclic. An NCM is said to be acyclic if it does not possess any directed cycle.

2.3.7 Definition

An NCM with cycles is said to have a feedback. When there is a feedback in the NCM i.e. when the causal relations flow through a cycle in a revolutionary manner the NCM is called a dynamical system.

2.3.8 Definition

Let $C_1 C_2, C_2 C_3, \dots, C_{n-1} C_n$ be cycle, when C_i is switched on and if the causality flow through the edges of a cycle and if it again causes C_i , we say that the dynamical system goes round and round. This is true for any node C_i , for $i = 1, 2, \dots, n$ the equilibrium state for this dynamical system is called the hidden pattern.

2.3.9 Definition

If the equilibrium state of a dynamical system is a unique state vector, then it is called a fixed point. Consider the NCM with C_1, C_2, \dots, C_n as nodes. For example let us start the dynamical system by switching on C_1 Let us assume that the NCM settles down with C_1 and C_n on, i.e. the state vector remain as $(1, 0, \dots, 1)$ this neutrosophic state vector $(1, 0, \dots, 0, 1)$ is called the fixed point.

2.3.10 Definition

If the NCM settles with a neutrosophic state vector repeating in the form $A_1 \rightarrow A_2 \rightarrow \dots \rightarrow A_1$ then this equilibrium is called a limit cycle of the NCM

3. The New BANCM Bimodal

3.1 Definition

Let $A = A_1 \cup A_2$ where A_1 and A_2 are non-empty sets; with $A_1 A_2$ and $A_1 A_2$ then we call A as a biset.

Let $A = \{(2\ 5\ 1\ 0\ 1\ 4), (1\ 9\ 3\ 5\ 7\ 0), \dots\} \cup \{(2\ 2\ 1\ 3), (5\ 0\ 2\ 9)\}$ = $A_1 \cup A_2$, clearly A is a biset.

3.2 Definition

Let $V_1 = (v_1, v_2, \dots, v_n)$, $V_2 = (v_1', v_2', v_3', \dots, v_m')$ be two vectors of length n and m respectively. Then $V = V_1 \cup V_2$ is a neutrosophic bivector.

Example:

Let $V = V_1 \cup V_2 = (6\ 5\ 2I\ 1\ 3I) \cup (3\ 4I\ 5\ 1)$, V is a neutrosophic bivector.

3.3 Definition:

A matrix E is said to be a Neutrosophic bimatrix. If $E = E_1 \cup E_2$ where E_1 and E_2 are two different matrices.

$$E = \begin{bmatrix} 4 & I & 2 \\ 2I & 6 & 3 \\ 0 & 7 & 5I \end{bmatrix} \cup \begin{bmatrix} 1 & I & 4I \\ 2I & 6 & 7I \\ 0 & 7 & 1I \end{bmatrix}$$

3.4 Definition

Let $G = G_1 \cup G_2$ where G_1 and G_2 are two distinct graphs then we call G as a bigraph.

3.5 Definition

Let $E = E_1 \cup E_2$ be a neutrosophic bimatrix. Then the bitranspose of the bimatrix E is defined as $E^t = (E_1 \cup E_2)^t = E_1^t \cup E_2^t$

3.6 Definition

A Bidirectional Associative Neutrosophic Cognitive Maps (BANCM) is a directed special bigraph with concepts like policies, events, etc as nodes and causalities as edges. It represents causal relationship between concepts. In a BANCM we call the pair of associated nodes as binodes. If the order of the bimatrix associated with BANCM is a $p \times m$ matrix and a $n \times n$ square matrix then the binodes are bivectors of length (p, n) or (m, n) .

3.7 Definition

Consider the binodes biconcepts $\{X_1, \dots, X_p\}$ and $\{Y_1, \dots, Y_m\}$ of the BAM and $\{C_1, C_2, \dots, C_n\}$ of the NCM of the BANCM bimodal. Suppose the directed graph is drawn using the edge biweight $e_{ij}^1 = [-5, 5]$; $e_{ij}^2 = [-1, 1]$, I . The bimatrix $E = E_1 \cup E_2$ is defined by $e_{ks}^1 \cup e_{ij}^2$ where e_{ks}^1 is the directed edge of $X_k Y_s$ and e_{ij}^2 is the weight of the directed edge $C_i C_j$. $E = E_1 \cup E_2$ is called adjacency bimatrix of the new BANCM bimodal.

3.8 Definition

The new BANCMs with edge biweight $\{1, 0, -1, I\}$ are called simple BANCMs. Let $\{(X_1, \dots, X_m), (Y_1, \dots, Y_n)\} \cup \{C_1, \dots, C_p\}$ be the binodes of an BANCM. $V = V_1 \cup V_2 = (x_1, \dots, x_m)$ (or (y_1, \dots, y_n)) $\cup (z_1, \dots, z_p)$, where $x_i, y_j, z_k \in \{0, 1, I\}$; $1 \leq i \leq m$, $1 \leq j \leq n$ and $1 \leq k \leq p$. V is called instantaneous state bivector and it denotes the ON-OFF-INDETERMINATE position of the node at an instant.

$x_j = 0, y_i = 0, z_k = 0$ if x_j, y_i, z_k are OFF,

$x_j = 1, y_i = 1, z_k = 1$ if x_j, y_i, z_k are ON

$x_j = I, y_i = I, z_k = I$ if x_j, y_i, z_k are Indeterminate for $1 \leq j \leq m, 1 \leq i \leq n$ and $1 \leq k \leq p$

3.9 Definition

Let $\{(X_1, \dots, X_m), (Y_1, \dots, Y_n)\} \cup \{C_1, \dots, C_p\}$ be the binodes of an BANCM. Let $X_k Y_s \cup C_i C_j$ be the bidedges of the BANCMs; $1 \leq k \leq m, 1 \leq s \leq n, 1 \leq i, j \leq p, (i \neq j)$. Then the bidedges form a directed bicycle. A BANCM is said to be bicyclic if it possesses a directed bicycle. An BANCM is said to be a bicyclic if it does not possess any directed bicycle.

3.10 Definition

A BAFCM with bicycles is said to have a feedback. When there is a feed back in an BAFCM, i.e., when the casual relations flow through a cycle in a revolutionary way, the BAFCM is called a dynamical bisystem.

3.11 Definition

If the BAFCM settles down with a bistate, bivector repeating in the form $B_1 \rightarrow B_2 \rightarrow \dots \rightarrow B_j \rightarrow B_1$ (or $D_1 \rightarrow D_2 \rightarrow \dots \rightarrow D_k \rightarrow D_1$) \cup $A_1 \rightarrow A_2 \rightarrow \dots \rightarrow A_1$ then this equilibrium is called a limit bicycle.

3.12 Definition

The biedges $e_{ij} = (e_{ks}^1) \cup (e_{ij}^2)$ take the values in fuzzy causal biinterval $[-1,1] \cup [-1,1] \cup I$.

- i) $e_{ij} = 0$ indicates no causality between the binodes.
 - ii) $e_{ij} > 0$ implies that both $e_{ij}^2 > 0$ and $e_{ks}^1 > 0$; implies increase in the binodes $C_i \cup X_k$ (or Y_s); implies increase in the binodes $C_j \cup Y_s$ (or X_k).
 - iii) $e_{ij} < 0$ implies that both $e_{ij}^2 < 0$ and $e_{ks}^1 < 0$; similarly decrease in the binodes $C_i \cup X_k$ (or Y_s); implies decrease in the binodes $C_j \cup Y_s$ (or X_k).
 - iv) $e_{ij} = I$ indicates Indeterminate between the binodes
- However, unlike the NCM and BAM model we can have the following possibilities other than that of $e_{ij} = 0, e_{ij} > 0$ and $e_{ij} < 0$.

- i) $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ can be such that $(e_{ij}^2) = 0$ and $(e_{ks}^1) > 0$. No relation in one binode and an increase in other node.
- ii) $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ we can have $(e_{ij}^2) = 0$ and $(e_{ks}^1) < 0$. No causality in the NCM node and decreasing relation in the BAM mode.
- iii) $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ we can have $(e_{ij}^2) \leq 0$ and $(e_{ks}^1) > 0$
- iv) In $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ we can have $(e_{ij}^2) < 0$ and $(e_{ks}^1) = 0$
- v) In $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ we can have $(e_{ij}^2) > 0$ and $(e_{ks}^1) = 0$
- vi) In $e_{ij} = (e_{ij}^2) \cup (e_{ks}^1)$ we can have $(e_{ij}^2) > 0$ and $(e_{ks}^1) < 0$.

Thus in the case of BANCM we can have 9 possibilities where as in NCMs or BAMs we have only 4 possibilities. Thus the extra 6 possibilities can help in making the Solution of the problem more sensitive or accurate.

4. ADAPTATION OF THE PROBLEM

Youth violence is a devastating public health problem worldwide. According to the World Health Organization, over 540 adolescents and young adults die every day from interpersonal violence. Annually, anywhere from 3.5 to 7.5 million young people experience injuries from violence requiring hospital treatment. Indeed, the health consequences of violence are severe, including death, permanent physical disabilities, high costs of medical care and rehabilitation, and immeasurable grief and suffering.

To analyze the causes and effect of youth violence, we have interviewed and collected a data from 100 youth in and around Chennai. After that some of the data was transformed in to a BAM as 8 main attributes of causes of violence and 7 attributes as effect of violence. We list out briefly the main attributes given below

Cause of youth violence:

- A₁- Poor family functioning
- A₂-Academic failure / dropping out of school

- A₃-Delinquent peers / gang membership
- A₄-Poverty in the community
- A₅-Castesam / inequality
- A₆-Influences of mass media
- A₇-Opportunities are denied
- A₈-Prizes hike in the basic needs

Effects of youth violence:

- B₁- Audit for drugs and alcohol
- B₂- Damage public property/ burning down Public transport
- B₃- Involving in other forms of anti-social behavior such as terrorism, robbery etc.
- B₄- Murdering / fake encounter
- B₅-Long drawn strike
- B₆-Run away from home
- B₇-Increasing aggressive behavior

Also through another expert the linguistic questionnaire was transformed in to NCM. In which 8 main attributes of youth violence is chosen, which are given below

- C₁- Parental criminality / Domestic violence
- C₂- Truancy and dropping out of school
- C₃- Ill treatment by teacher
- C₄- Bad Company
- C₅- Poverty and seeks self-respect
- C₆- Drug alcohol and tobacco use
- C₇-Violence on mass media / violent video games
- C₈- Increasing aggressive behavior.

As the data is an unsupervised one and involves lot of uncertainties we are justified in using BANCM to analyses the problem of youth violence.

$$\begin{bmatrix} 1 & 0 & 3 & -2 & -3 & 5 & 4 \\ 2 & -1 & 1 & 0 & -3 & 2 & 3 \\ 3 & 1 & 4 & 2 & 1 & 0 & -4 \\ 0 & -3 & 3 & 0 & -2 & -3 & -2 \\ -1 & 2 & 5 & 3 & 2 & -2 & 0 \\ 0 & -4 & 2 & 1 & -3 & 0 & 4 \\ -2 & 0 & 3 & 1 & 0 & -2 & 3 \\ -4 & 3 & 2 & 0 & 4 & 0 & -1 \end{bmatrix} \cup \begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$E = E_1 \cup E_2$$

Let the initial vector X_1 in BANCM

$$(3 -2 -4 -2 0 -1 0 -5) \cup (1 0 0 0 0 0 0 0)$$

Therefore $X_1 = (1 0 0 0 0 0 0 0) \cup (1 0 0 0 0 0 0 0)$

The effect X_1 on the dynamical system E

$$\begin{aligned} X_1 E &= (1 0 0 0 0 0 0 0) E_1 \cup (1 0 0 0 0 0 0 0) E_2 \\ &= (1 0 3 -2 -3 5 4) \cup (0 1 0 1 0 1 0 1) \\ &\hookrightarrow (1 0 1 0 0 1 1) \cup (1 1 0 1 0 1 0 1) \\ &\Rightarrow (1 0 1 0 0 1 1) E_1^T \cup (1 1 0 1 0 1 0 1) E_2 \\ &= (13 5 3 -2 2 6 2 -3) \cup (0 2 1 3 2 1 1 1) \\ &\hookrightarrow (1 1 1 0 1 1 1 0) \cup (1 1 1 1 1 1 1 1) = X_2 \\ X_2 E &= (1 1 1 0 1 1 1 0) E_1 \cup (1 1 1 1 1 1 1 1) E_2 \\ &= (3 -2 18 4 -6 0 8) \cup (1 2 1 1 1 1 1 1) \\ &\hookrightarrow (1 0 1 1 0 0 1) \cup (1 1 1 1 1 1 1 1) \\ &\Rightarrow (1 0 1 1 0 0 1) E_1^T \cup (1 1 1 1 1 1 1 1) E_2 \\ &= (6 6 5 -2 5 7 3 -3) \cup (1 2 1 1 1 1 1 1) \\ &\hookrightarrow (1 1 1 0 1 1 1 0) \cup (1 1 1 1 1 1 1 1) = X_3 \\ X_3 E &= (1 1 1 0 1 1 1 0) E_1 \cup (1 1 1 1 1 1 1 1) E_2 \\ &= (3 -2 18 4 -6 0 10) \cup (1 2 1 1 1 1 1 1) \\ &\hookrightarrow (1 0 1 1 0 0 1) \cup (1 1 1 1 1 1 1 1) \end{aligned}$$

Now $X_2 = X_3$. Hence the limit points is $((1\ 1\ 1\ 0\ 1\ 1\ 1\ 0), (1\ 0\ 1\ 1\ 0\ 0\ 1)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$

The set of all limit points corresponding to the different input vectors

No	Input Vector	Limit points
1	(10000000)	$((11101110), (101100\ 1)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$
2	(01000000)	$((11101100), (1011011)) \cup (1\ 1\ 1\ 1\ 1\ 0\ 1\ 1)$
3	(00100000)	$((11101001), (1111111)) \cup (1\ 1\ 1\ 1\ 1\ 0\ 1\ 1)$
4	(00010000)	$((11111111), (11101001)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$
5	(00001000)	$((00101011), (0111100)) \cup (1\ 1\ 0\ 1\ 1\ 1\ 1\ 1)$
6	(00000100)	$((11111111), (11101001)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$
7	(00000010)	$((11111111), (11101001)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$
8	(00000001)	$((11111111), (11101001)) \cup (1\ 1\ 1\ 1\ 1\ 1\ 1\ 1)$

5. Conclusion

We analyzed the causes of youth violence using BANCM, the set of all limit points are highlighting that $A_1, A_2, A_3, A_5, A_6, A_8, B_1, B_3, B_4, B_6, B_7, C_1, C_2, C_4, C_5, C_7$ and C_8 are the main causes and effects of youth violence. i.e., The effect of Poor family functioning, Academic failure / dropping out of school, Delinquent peers / gang membership, Castesam / inequality, Influences of mass media and Prizes hike in the basic needs, Parental criminality / Domestic violence, Poverty and seeks self-respect and Violence on mass media / violent video games are leads to Audit for drugs and alcohol, Involving in other forms of anti-social behavior such as terrorism, robbery etc., Murdering / fake encounter, Run away from home and Increasing aggressive behavior but Ill treatment by teacher is an indeterminate position.

Parent's responsibility to protect their children from violence

1. Parents have to talk to their teen, and listen - show caring and concern know where teens go, what they do, and whom they're with.
2. We Include teens in family activities, and be home during at least one of these times: when our teen wakes up, when he or she comes home from school, at dinner time, or when our teen goes to bed.
3. Be consistent and firm - but not harsh - in our discipline
4. If applicable, help our teen or other family members get help for drug or alcohol problems.
5. Monitor and control our teen's exposure to violence in the media, including television, movies, video games, music, etc.
6. Parents should get more involvement in their community and their child's school; talk to school administrators about violence or bullying that occurs at school.
7. Parents should be a good example of non-violent ways to resolve conflicts

Future Direction

The violence due to aggressive behavior and family condition are correlated. The broken families are the fertile nurseries for the anti-social behavior among the children. Family plays a vital role in formation of mental and psychic abilities of children. Family is placed at the centre of great struggle between good and evil, between life and death, between love and all that is opposed to love. Hence parents have a greater role in shaping educating the next generation in to a society in to a peaceful society. We planned to study the behavior of parents in influencing the children in to violent behavior. .

REFERENCES

- [1] Axelrod, R. (1976). *Structure of decision: The cognitive maps of political elites*. Princeton, NJ: Princeton University Press.
- [2] Kosco, B. (1986), "Fuzzy Cognitive Maps" International journal of man-machine studies, January, 62-75
- [3] Kosco, B., (1997) "Neural Networks and Fuzzy Systems: A Dynamical Systems Approach to Machine Intelligence", Prentice Hall of India,
- [4] Lian-Hua Chen "Violent Scene Detection in Movies: World Scientific Journal.
- [5] Strauss M.A , 2007 "Conflict Tactics Scales. In: NA. Jackson, editor Encyclopedia of Domestic violence",.
- [6] Thirusangu et.al. (2012) "A new bidirectional associative fuzzy cognitive dynamical system". Indian Journal of Science and Technology,.
- [7] Vasantha Kandasamy W.B and Smarandache Florentin (2004) "Analysis of social aspects of migrant laborers living with HIV/AIDS using Fuzzy Theory and Neutrosophic Cognitive Maps", Xiquan, Phoenix.
- [8] Vasantha Kandasamy W.B and Smarandache Florentin, "Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps", Xiquan, Phoenix (2003)
- [9] Varma, D. Chandra, PS, Thomas, T. Carey, MP. (2007) "Intimate partner violence and sexual coercion among pregnant women in India: Relationship with depression and post traumatic stress disorder". J Affect Disord; 102:227-35.
- [10] Victor Devadoss, A. Felix, A. (2012), "Fuzzy Clustering of the Domestic Violence for the Degree of Suicide thought based on Married Women Perception". International Journal of Engineering Research and Application.
- [11] Victor Devadoss, A. Clement Joe Anand, M. Felix, A. (2012); "A Study on the Impact Violent Video-Games playing among Children in Chennai using Neutrosophic Cognitive Maps". International Journal of Scientific & Engineering Research.
- [12] Zimmermann H.J (2011), "Fuzzy Set Theory and its application", Fourth Edition Springer.