A NEW ESTIMATE OF THE DIFFERENCE AMONG QUASI-ARITHMETIC MEANS

PAWEŁ PASTECZKA

Abstract. In the 1960s Cargo and Shisha proved some majorizations for the distance among quasi-arithmetic means (defined as $f^{-1}(\sum_{i=1}^{n} w_i f(a_i))$ for any continuous, strictly monotone function $f: I \to \mathbb{R}$, where *I* is an interval, and (a_1, \ldots, a_n) is a vector with entries in *I*, (w_1, \ldots, w_n) is a sequence of corresponding weights $w_i > 0$, $\sum w_i = 1$).

Nearly thirty years later, in 1991, Páles presented an iff condition for a sequence of quasi-arithmetic means to converge to another QA mean. It was closely related with the three-parameter operator (f(x) - f(y))/(f(x) - f(z)).

The author presented recently an estimate for the distance among such quasi-arithmetic means whose underlying functions satisfy some smoothness conditions. Used was the operator $f \mapsto f''/f'$ introduced in the 1940s by Mikusiński and Łojasiewicz. It is natural to look for similar estimate(s) in the case of the underlying functions *not* being smooth. For instance, by the way of using Páles' operator. This is done in the present note. Moreover, the result strengthens author's earlier estimates.

Mathematics subject classification (2010): 26E60, 26D15, 26D07. *Keywords and phrases:* Quasi-arithmetic means, metric, Arrow-Pratt index.

REFERENCES

- [1] G. T. CARGO AND O. SHISHA, On comparable means, Pacific J. Math. 14 (3), 1964, 1053–1058.
- [2] G. T. CARGO AND O. SHISHA, A metric space connected with generalized means, J. Approx. Theory 2 (2), 1969, 207–222.
- [3] B. DE FINETTI, Sul concetto di media, G. Ist. It. Att. 2 (3), 1931, 369–396.
- [4] A. KOLMOGOROFF, Sur la notion de la moyenne, Rend. Accad. dei Lincei 12 (6), 1930, 388–391.
- [5] J. G. MIKUSIŃSKI, Sur les moyennes de la forme $\psi^{-1}[\Sigma q \psi(x)]$, Studia Math. 10 (1), 1948, 90–96.
- [6] M. NAGUMO, Über eine Klasse der Mittelwerte, Japan. J. Math. 7 (1), 1930, 71-79.
- [7] M. NOSARZEWSKA, On uniform convergence in some classes of functions, Fund. Math. 39 (1), 1952, 38–52.
- [8] ZS. PÁLES, Report of Meeting 15th International Conference on Functional Equations and Inequalities, Ann. Univ. Paedagog. Crac. Stud. Math. XII (1), 2013, 121–122.
- [9] ZS. PÁLES, On the convergence of Means, J. Math. Anal. Appl. 156 (1), 1991, 52-60.
- [10] P. PASTECZKA, When is a Family of Generalized Means a Scale?, Real Anal. Exchange 38 (1), 2013, 193–210.
- [11] G. PÓLYA, Wohlbekannt ist folgender Satz von Dini, Arxiv der Mathematic und Phisik **28** (1), 1920, 174.

