

A New Family of q -analogue of Genocchi Numbers and Polynomials of Higher Order

SERKAN ARACI* AND MEHMET ACIKGOZ

*University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,
27310 Gaziantep, Turkey*

e-mail: mtsrkn@hotmail.com and acikgoz@gantep.edu.tr

JONG JIN SEO

Department of Applied Mathematics, Pukyong National University, Busan, 608-737, Republic of Korea

e-mail: seo2011@pknu.ac.kr

ABSTRACT. In the present paper, we introduce the new generalization of q -Genocchi polynomials and numbers of higher order. Also, we give some interesting identities. Finally, by applying q -Mellin transformation to the generating function for q -Genocchi polynomials of higher order put we define novel q -Hurwitz-Zeta type function which is an interpolation for this polynomials at negative integers.

1. Introduction

Recently, the Bernoulli, Euler and Genocchi polynomials and their families have been studied by many mathematicians for a long time (for details, see [1-22]). These polynomials possess a number of interesting properties not only in Analytic numbers theory and Complex analysis, but also in Mathematical physics, q -analysis, p -adic analysis and other areas of mathematics.

Throughout this work, we assume that $q \in \mathbb{C}$ with $|q| < 1$. The q -integer of x is defined by $[x]_q = \frac{1-q^x}{1-q}$ and we note that $\lim_{q \rightarrow 1} [x]_q = x$. The q -derivative is also defined by F. H. Jackson as follows:

$$(1.1) \quad D_q f(x) = \frac{d_q f(x)}{d_q x} = \frac{f(x) - f(qx)}{(1-q)x}.$$

* Corresponding Author.

Received December 24, 2012; accepted April 04, 2013.

2010 Mathematics Subject Classification: 11S80, 11B68.

Key words and phrases: Genocchi numbers and polynomials, q -Genocchi numbers and polynomials of higher order, q -Mellin transformation, q -Hurwitz-zeta function, q -Gamma function, q -Exponential function.

Taking $f(x) = x^n$ in (1.1), it becomes as follows:

$$D_q x^n = \frac{x^n - (qx)^n}{(1-q)x} = [n]_q x^{n-1} \text{ and } \left(\frac{d}{d_q x}\right)^n f(x) = [n]_q!$$

where $[n]_q! = [n]_q [n-1]_q \cdots [1]_q$. Now, we give definitions of two kinds of q -exponential functions as follows:

For any $z \in \mathbb{C}$ with $|z| < 1$,

$$(1.2) \quad e_q(z) = \sum_{l=0}^{\infty} \frac{z^l}{[l]_q!} \text{ and } E_q(z) = \sum_{l=0}^{\infty} q^{\binom{l}{2}} \frac{z^l}{[l]_q!}.$$

By (1.2), it is not difficult to show that $[l]_{\frac{1}{q}}! = q^{-\binom{l}{2}} [l]_q!$. Then, we have the following

$$(1.3) \quad e_{\frac{1}{q}}(z) = E_q(z).$$

For the q -commuting variables x and y such that $yx = qxy$, we know that

$$(1.4) \quad e_q(x+y) = e_q(x) e_q(y).$$

The q -integral was defined by Jackson as follows:

$$(1.5) \quad \int_0^x f(\xi) d_q \xi = (1-q)x \sum_{l=0}^{\infty} f(q^l x) q^l$$

provided that the series on the right hand side converges absolutely.

In particular, if $f(\xi) = \xi^n$, then we have

$$(1.6) \quad \int_0^x \xi^n d_q \xi = \frac{1}{[n+1]_q} x^{n+1}.$$

The definitions of q -integral and q -derivative imply the following formula:

$$(1.7) \quad D_q \left(\int_0^x f(t) d_q t \right) = f(x)$$

and

$$(1.8) \quad D_q (f(x)g(x)) = f(x)D_q(g(x)) + g(qx)D_q(f(x)).$$

For more information of Eqs. (1-8), one can refer to [23-27].

The ordinary Euler polynomials are defined via the following generating function:

$$e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} = \frac{2}{e^t + 1} e^{xt}, \quad |t| < \pi$$

with the usual convention about replacing $E^n(x)$ by $E_n(x)$ (see [1-22]).

In [16], the new q -generalization of Euler polynomials was introduced by T. Kim as follows:

$$\sum_{j=0}^{\infty} \frac{z^j}{[j]_q!} E_{j,q}(x) = \frac{[2]_q}{e_q(z) + 1} e_q(xz).$$

By using the above generating function, Kim gave some interesting properties for new q -generalization of Euler numbers and polynomials. We note that these polynomials are useful to study in the theory of special functions. By the same motivation, in the next section, we shall introduce the generating function of q -Genocchi numbers and polynomials of higher order. Additionally, we shall give their applications.

2. On The New q -Genocchi Numbers and Polynomials of Higher Order

In this section, we introduce the generating function for q -Genocchi numbers of higher order due to Kim's method in [16]. Thus, we now start as follows:

$$(1.1) \quad S_q(z : \alpha) = \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} G_{n,q}^{(\alpha)} = \left(\frac{[2]_q z}{e_q(z) + 1} \right)^\alpha.$$

Here $G_{n,q}^{(\alpha)}$ are called the q -Genocchi numbers of higher order. By using q -derivative operator, we compute as follows:

$$(1.2) \quad S_q\left(\frac{d}{d_q x} : \alpha\right) x^k = \sum_{n=0}^{\infty} \frac{1}{[n]_q!} G_{n,q}^{(\alpha)} \left(\frac{d}{d_q x}\right)^n x^k = \sum_{n=0}^k \binom{k}{n}_q G_{n,q}^{(\alpha)} x^{k-n} = G_{k,q}^{(\alpha)}(x).$$

where $G_{k,q}^{(\alpha)}(x)$ are called q -Genocchi polynomials of higher order and

$$\binom{k}{n}_q = \frac{[k]_q [k-1]_q \cdots [k-n+1]_q}{[n]_q!}.$$

Similarly, by (1.2), we develop as follows:

$$\begin{aligned} S_q\left(\frac{d}{d_q x} : \alpha\right) e_q(tx) &= \sum_{j=0}^{\infty} \frac{G_{j,q}^{(\alpha)}}{[j]_q!} \left(\frac{d}{d_q x}\right)^j \sum_{k=0}^{\infty} \frac{x^k}{[k]_q!} t^k \\ &= \sum_{j=0}^{\infty} \frac{t^j}{[j]_q!} G_{j,q}^{(\alpha)}(x) \\ &= S_q(x, t : \alpha). \end{aligned}$$

From this point of view, we can also consider the q -Genocchi polynomials of higher order in the form:

$$(1.3) \quad \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} G_{n,q}^{(\alpha)}(x) = \left(\frac{[2]_q z}{e_q(z) + 1} \right)^\alpha e_q(zx).$$

As $q \rightarrow 1$ and $\alpha = 1$ in Eq. (1.3), we easily reach the following

$$\lim_{q \rightarrow 1} G_{n,q}^{(1)}(x) = G_n(x)$$

in which $G_n(x)$ are known ordinary Genocchi polynomials (for details, see [14], [18], [26], [2], [4], [5]).

By (1.1) and (1.3), we readily see that

$$\sum_{j=0}^{\infty} \frac{z^j}{[j]_q!} G_{j,q}^{(\alpha)}(x) = \sum_{j=0}^{\infty} \left(\sum_{n=0}^j \binom{j}{n}_q x^{j-n} G_{n,q}^{(\alpha)} \right) \frac{z^j}{[j]_q!}.$$

Matching the coefficients of $\frac{z^j}{[j]_q!}$ in the both sides of the above equation, then we obtain the following theorem.

Theorem 2.1. For any $j \in \mathbb{N}$, we have

$$G_{j,q}^{(\alpha)}(x) = \sum_{n=0}^j \binom{j}{n}_q x^{j-n} G_{n,q}^{(\alpha)}.$$

By applying q -derivative operator to (1.3), then we see that

$$\sum_{n=1}^{\infty} \frac{z^n}{[n]_q!} \left\{ \frac{d}{d_q x} G_{n,q}^{(\alpha)}(x) \right\} = z \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} G_{n,q}^{(\alpha)}(x).$$

By comparing the coefficients of z^n on the both sides of the above equation, we arrive at the following theorem.

Theorem 2.2. For any $n \in \mathbb{N}^* = \mathbb{N} \cup \{0\}$, we have

$$\frac{d}{d_q x} G_{n,q}^{(\alpha)}(x) = [n]_q G_{n-1,q}^{(\alpha)}(x).$$

For q -commuting variables x and y ($yx = qxy$), we note that

$$\begin{aligned} \sum_{l=0}^{\infty} \frac{z^l}{[l]_q!} G_{l,q}^{(\alpha)}(x+y) &= \left(\frac{[2]_q z}{e_q(z) + 1} \right)^\alpha e_q(z(x+y)) \\ &= e_q(zy) \left(e_q(zx) \left(\frac{[2]_q z}{e_q(z) + 1} \right)^\alpha \right) \\ &= \sum_{l=0}^{\infty} \left(\sum_{j=0}^l \binom{l}{j}_q y^{l-j} G_{j,q}^{(\alpha)}(x) \right) \frac{z^l}{[l]_q!}. \end{aligned}$$

As a result, we procure the following theorem.

Theorem 2.3. For any $n \in \mathbb{N}^*$, we have

$$G_{n,q}^{(\alpha)}(x+y) = \sum_{j=0}^n \binom{n}{j}_q y^{n-j} G_{n,q}^{(\alpha)}(x).$$

By expression (1.3), we compute as follows:

$$\begin{aligned} (1.4) \quad & \sum_{l=0}^{\infty} \frac{z^l}{[l]_q!} G_{l,q}^{(\alpha+\beta)}(x) \\ &= \left[\left(\frac{[2]_q z}{e_q(z)+1} \right)^\alpha \right] \left[\left(\frac{[2]_q z}{e_q(z)+1} \right)^\beta e_q(zx) \right] \\ &= \left[\sum_{j=0}^{\infty} \frac{z^j}{[j]_q!} G_{j,q}^{(\alpha)} \right] \left[\sum_{k=0}^{\infty} \frac{z^k}{[k]_q!} G_{k,q}^{(\beta)}(x) \right] \end{aligned}$$

by using Cauchy product on the above equation, we derive that

$$(1.5) \quad \sum_{l=0}^{\infty} \frac{z^l}{[l]_q!} \left(\sum_{n=0}^l \binom{l}{n}_q G_{n,q}^{(\alpha)} G_{l-n,q}^{(\beta)}(x) \right).$$

Comparing the coefficients of Eqs. (1.4) and (1.5), we get the following theorem.

Theorem 2.4. For any $n \in \mathbb{N}^*$, we have

$$G_{n,q}^{(\alpha+\beta)}(x) = \sum_{k=0}^n \binom{n}{k}_q G_{k,q}^{(\alpha)} G_{n-k,q}^{(\beta)}(x).$$

Jackson defined the q -analogue of the Gamma function by

$$(1.6) \quad \Gamma_q(x) = \frac{(q; q)_\infty}{(q^x; q)_\infty} (1-q)^{1-x}, \quad x \neq 0, -1, -2, \dots.$$

which have the following properties:

$$\Gamma_q(x+1) = [x]_q \Gamma_q(x), \quad \Gamma_q(1) = 1 \quad \text{and} \quad \lim_{q \rightarrow 1^-} \Gamma_q(x) = \Gamma(x), \quad \Re(x) > 0.$$

It has the q -integral representation as follows:

$$(1.7) \quad \Gamma_q(s) = \int_0^{\frac{1}{1-q}} t^{s-1} E_q(-qt) d_q t.$$

When $\frac{\log(1-q)}{\log q} \in \mathbb{Z}$, becomes

$$(1.8) \quad \Gamma_q(s) = \int_0^\infty t^{s-1} E_q(-qt) d_q t.$$

The q -Mellin transformation of a suitable function f on $\mathbb{R}_{q,+}$ is defined by

$$(1.9) \quad M_q(f)(s) = \int_0^\infty t^{s-1} f(t) d_q t$$

(for details of Eqs. (14-17), see [20], [10], [15]).

In [27], the novel q -differential operator was defined by Rubin as follows:

$$(1.10) \quad \partial_q(f)(x) = \frac{f(q^{-1}x) + f(-q^{-1}x) - f(qx) + f(-qx) - 2f(-x)}{2(1-q)x}.$$

By (1.10), we note that

$$(1.11) \quad \lim_{q \rightarrow 1} \partial_q(f)(x) = f'(x).$$

By applying Rubin's q -differential operator to the generating function of q -Genocchi numbers and polynomials of higher order, we compute as follows:

$$\begin{aligned} & \sum_{n=0}^\infty \frac{z^n}{[n]_q!} \partial_q G_{n,q}^{(\alpha)}(x) = \partial_q \left\{ \left(\frac{[2]_q z}{e_q(z) + 1} \right)^\alpha e_q(xz) \right\} \\ & = \sum_{n=0}^\infty \left(\frac{1}{2(1-q)} \sum_{l=0}^n \binom{n}{l}_q \left\{ \begin{matrix} q^{-l} + (-1)^l q^{-l} - q^l \\ + (-1)^l q^l + 2(-1)^l \end{matrix} \right\} x^{l-1} G_{n-l,q}^{(\alpha)} \right) \frac{z^n}{[n]_q!}. \end{aligned}$$

By comparing the coefficients of $\frac{z^n}{[n]_q!}$ on the both sides of the above equation. Then, we state the following theorem.

Theorem 2.5. Let $T_q(l) = q^{-l} + (-1)^l q^{-l} - q^l + (-1)^l q^l + 2(-1)^l$, then we get

$$(1.12) \quad \partial_q G_{n,q}^{(\alpha)}(x) = \frac{1}{2(1-q)} \sum_{l=0}^n \binom{n}{l}_q T_q(l) x^{l-1} G_{n-l,q}^{(\alpha)}.$$

By (1.12), we readily derive the following

$$\begin{aligned} \partial_q G_{n,q}^{(\alpha)}(x) &= \frac{1}{(1-q)} \sum_{l=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2l}_q \{q^{-2l} + 1\} G_{n-2l,q}^{(\alpha)} \\ &+ \frac{1}{(q-1)} \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2l+1}_q \{q^{2l+1} + 1\} G_{n-1-2l,q}^{(\alpha)}. \end{aligned}$$

Here $[\cdot]$ is Gauss' symbol. Consequently, we state the following theorem.

Theorem 2.6. For any $n \in \mathbb{N}^*$, we have

$$\begin{aligned} \partial_q G_{n,q}^{(\alpha)}(x) &= \frac{1}{(1-q)} \sum_{l=0}^{[\frac{n}{2}]} \binom{n}{2l}_q \{q^{-2l} + 1\} G_{n-2l,q}^{(\alpha)} \\ &+ \frac{1}{(q-1)} \sum_{l=0}^{[\frac{n-1}{2}]} \binom{n}{2l+1}_q \{q^{2l+1} + 1\} G_{n-1-2l,q}^{(\alpha)}. \end{aligned}$$

From (1.12) and Theorem 2.6, we conclude as follows:

Corollary 2.7. The following identity

$$\begin{aligned} &\frac{1}{2(1-q)} \sum_{l=0}^n \binom{n}{l}_q T_q(l) x^{l-1} G_{n-l,q}^{(\alpha)} \\ &= \frac{1}{(1-q)} \sum_{l=0}^{[\frac{n}{2}]} \binom{n}{2l}_q \{q^{-2l} + 1\} G_{n-2l,q}^{(\alpha)} \\ &+ \frac{1}{(q-1)} \sum_{l=0}^{[\frac{n-1}{2}]} \binom{n}{2l+1}_q \{q^{2l+1} + 1\} G_{n-1-2l,q}^{(\alpha)} \end{aligned}$$

is true.

By (1.12), we have the following corollary.

Corollary 2.8. For any $n \in \mathbb{N}^*$, we get

$$\lim_{q \rightarrow 1} \partial_q G_{n,q}^{(\alpha)}(x) = n G_{n-1}^{(\alpha)}(x)$$

where $G_n^{(\alpha)}(x)$ are known Genocchi polynomials of higher order defined by the following generating function:

$$\sum_{n=0}^{\infty} G_n^{(\alpha)}(x) \frac{t^n}{n!} = \left(\frac{2t}{e^t + 1} \right)^\alpha e^{xt}.$$

We now give a q -analogue of D. Milićić's Lemma (see [22, page 1, Lemma 1.2.1]).

Lemma 2.9. Let $a_{n,q}$, $n \in \mathbb{N}^*$ be complex numbers such that $\sum_{n=0}^{\infty} |a_{n,q}|$ converges. Let

$$\lambda = \{-n \mid n \in \mathbb{N}^* \text{ and } a_{n,q} \neq 0\}.$$

Then,

$$g_q(z) = \sum_{n=0}^{\infty} \frac{a_{n,q}}{[z+n]_q}$$

converges absolutely for $z \in \mathbb{C} - \lambda$ and uniformly on bounded subsets of $\mathbb{C} - \lambda$. The q -function is a meromorphic function on complex plane with simple poles at the points in λ and $\text{Re } s(g_q, -n)$ for any $-n \in \lambda$.

Proof. By using similar method in lecture notes of D. Milićić's in [22], the proof of Lemma can be easily done. So, we omit it.

We now want to indicate that Γ extends to a meromorphic function by using Lemma 2.9. That is, we discover the following

$$\Gamma_q(z) = \int_0^{\infty} t^{z-1} E_q(-qt) d_q t = \int_0^1 t^{z-1} E_q(-qt) d_q t + \int_1^{\infty} t^{z-1} E_q(-qt) d_q t.$$

Then, the second integral converges for any complex z and represents an entire function. On the other hand, the q -exponential function is entire, and we have

$$\begin{aligned} \int_0^1 t^{z-1} E_q(-qt) d_q t &= \int_0^1 t^{z-1} \left\{ \sum_{j=0}^{\infty} \frac{(-1)^j q^{\binom{j+1}{2}}}{[j]_q!} t^j \right\} d_q t \\ &= \sum_{j=0}^{\infty} \frac{(-1)^j q^{\binom{j+1}{2}}}{[j]_q!} \left\{ \int_0^1 t^{z+j-1} d_q t \right\} \\ &= \sum_{j=0}^{\infty} \frac{(-1)^j q^{\binom{j+1}{2}}}{[j]_q!} \frac{1}{[z+j]_q} \end{aligned}$$

for any $z \in \mathbb{C}$. Now also, we can write as follows:

$$\Gamma_q(z) = \int_1^{\infty} t^{z-1} E_q(-qt) d_q t + \sum_{j=0}^{\infty} \frac{(-1)^j q^{\binom{j+1}{2}}}{[j]_q!} \frac{1}{[z+j]_q}$$

for any z in the right half plane. From Lemma 2.9, the right hand-side of the above identity defines a meromorphic function on the complex plane with simple poles at $z = -j, j \in \mathbb{N}^*$. Then, we have the following theorem.

Theorem 2.10. For any $j \in \mathbb{N}^*$, we derive the following

$$(1.13) \quad \text{Res}(\Gamma_q, -j) = \frac{(-1)^j q^{\frac{j(j+1)}{2}}}{[j]_q!}.$$

As $q \rightarrow 1$ into (1.13), we easily derive that

$$\lim_{q \rightarrow 1} \text{Res}(\Gamma_q, -j) = \frac{(-1)^j}{j!}$$

which is residue of Euler's Gamma function (see [25]).

Now also, by applying q -Mellin Transformation to generating function of q -Genocchi polynomials of higher order, then we compute as follows:

For q -commuting variables x and y ($yx = qxy$),

$$\begin{aligned} \mathfrak{S}_q(z, x : \alpha) &= \frac{1}{\Gamma_{\frac{1}{q}}(z)} \int_0^\infty t^{z-\alpha-1} \{(-1)^\alpha S_q(x, -t : \alpha)\} d_{\frac{1}{q}}t \\ &= \sum_{l_1, l_2, \dots, l_\alpha=0}^\infty (-1)^{l_1+l_2+\dots+l_\alpha} \left\{ \frac{1}{\Gamma_{\frac{1}{q}}(z)} \int_0^\infty t^{z-1} E_{\frac{1}{q}} \left(-\frac{t}{q} \left(qx + q \sum_{k=1}^\alpha l_k \right) \right) d_{\frac{1}{q}}t \right\} \\ &= [2]_q^\alpha \sum_{l_1, l_2, \dots, l_\alpha=0}^\infty \frac{q^{-z} (-1)^{l_1+l_2+\dots+l_\alpha}}{(l_1 + l_2 + \dots + l_\alpha + x)^z}. \end{aligned}$$

So, we now introduce definition of q -Hurwitz-Zeta type function as follows:

Definition 2.11. For any $z \in \mathbb{C}$, we define

$$\mathfrak{S}_q(z, x : \alpha) = [2]_q^\alpha \sum_{l_1, l_2, \dots, l_\alpha=0}^\infty \frac{q^{-z} (-1)^{l_1+l_2+\dots+l_\alpha}}{(l_1 + l_2 + \dots + l_\alpha + x)^z}.$$

By the above definition, we derive interpolation functions for q -Genocchi polynomials of higher order at negative integers.

Theorem 2.12. The following equality holds true:

$$\mathfrak{S}_q(-n, x : \alpha) = \frac{q^{-n} G_{n+\alpha, q}^{(\alpha)}(x)}{[\alpha]_q! \binom{n+\alpha}{\alpha}_q}.$$

References

- [1] S. Araci, Novel identities involving Genocchi numbers and polynomials arising from applications of umbral calculus, *Applied Mathematics and Computation*, **233**(2014), 599–607.
- [2] M. Acikgoz, S. Araci and I. N. Cangul, A note on the modified q -Bernstein polynomials for functions of several variables and their reflections on q -Volkenborn integration, *Applied Mathematics and Computation*, **218**(3)(2011), 707–712.
- [3] S. Araci, M. Acikgoz, E. Şen, On the extended Kim's p -adic q -deformed fermionic integrals in the p -adic integer ring, *Journal of Number Theory*, **133**(2013), 3348–3361.
- [4] S. Araci, M. Acikgoz and K. H. Park, A note on the q -analogue of Kim's p -adic log gamma type functions associated with q -extension of Genocchi and Euler numbers with weight α , *Bull. Korean Math. Soc.*, **50**(2)(2013), 583–588.

- [5] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p -adic q -integral representation on \mathbb{Z}_p associated with weighted q -Bernstein and q -Genocchi polynomials, *Abstract and Applied Analysis*, Volume 2011, Article ID 649248, 10 pages.
- [6] S. Araci, M. Acikgoz and A. Gürsul, Analytic continuation of weighted q -Genocchi numbers and polynomials, *Commun. Korean Math. Soc.*, **28(3)**(2013), 457–462.
- [7] S. Araci and M. Acikgoz, A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials, *Advanced Studies in Contemporary Mathematics*, **22(3)**(2012), 399–406.
- [8] S. Araci, M. Acikgoz, H. Jolany, J. J. Seo, A unified generating function of the q -Genocchi polynomials with their interpolation functions, *Proc. Jangjeon Math. Soc.*, **15(2)**(2012), 227–233.
- [9] S. Araci, J. J. Seo and D. Erdal, New construction weighted (h, q) -Genocchi numbers and polynomials related to Zeta-type functions, *Discrete Dynamics in Nature and Society*, **2011**(2011), Article ID 487490, 7 pages.
- [10] K. Brahim, and R. Guanés, Some applications of the q -Mellin Transform, *Tamsui Oxford Journal of Mathematical Sciences*, **26(3)**(2010), 335–343.
- [11] Y. He and C. Wang, Some formulae of products of the Apostol-Bernoulli and Apostol-Euler polynomials, *Discrete Dynamics in Nature and Society*, vol. 2012, Article ID 927953, 11 pages, 2012.
- [12] F. H. Jackson, On q -definite integrals, *The Quarterly Journal of Pure and Applied Mathematics*, **41**(1910), 193–2036.
- [13] L. C. Jang, The q -analogue of twisted Lerch type Euler Zeta functions, *Bull. Korean Math. Soc.*, **47(6)**(2010), 1181–1188.
- [14] H. Jolany and H. Sharifi, Some results for Apostol-Genocchi Polynomials of higher order, *Bulletin of the Malaysian Mathematical Sciences Society*, **36(2)**(2013).
- [15] V. G. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, New York, 2002.
- [16] T. Kim, q -Generalized Euler numbers and polynomials, *Russ. J. Math. Phys.*, **13(3)**(2006), pp. 293–308
- [17] T. Kim, Some identities on the q -Euler polynomials of higher order and q -stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p , *Russian J. Math. Phys.*, **16**(2009), 484–491.
- [18] T. Kim, On the q -extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.*, **326**(2007), 1458–1465.
- [19] T. Kim, On the analogs of Euler numbers and polynomials associated with p -adic q -integral on \mathbb{Z}_p at $q = -1$, *J. Math. Anal. Appl.*, **331** (2007), 779–792.

- [20] T. Kim, Some Identities on the integral representation of the product of several q -Bernstein-type polynomials, *Abstract and Applied Analysis*, Volume 2011, Article ID 634675, 11 pages.
- [21] T. Kim, S. H. Lee, H. H. Han and C. S. Ryoo, On the values of the weighted q -Zeta and L -functions, *Discrete Dynamics in Nature and Society*, Volume 2011, Article ID 476381, 7 pp.
- [22] T. Kim, Euler numbers and polynomials associated with Zeta functions, *Abstract and Applied Analysis*, Volume 2008, Article ID 581582, 11 pages.
- [23] T. Kim, Power series and asymptotic series associated with the q -analogue of the two-variable p -adic L -function, *Russian Journal of Mathematical Physics*, **12(2)**(2005), 186–196.
- [24] T. Kim, On the q -extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.*, **326**(2007), 1458–1465.
- [25] D. Milićić, Notes on the Riemann's Zeta function, <http://www.math.utah.edu/~milicic/zeta.pdf>.
- [26] S-H. Rim, J-H. Jin, E-J. Moon and S-J. Lee, On multiple interpolation functions of the q -Genocchi polynomials, *Journal of Inequalities and Applications*, Volume 2010, Article ID 351419, 13 pages.
- [27] Richard L. Rubin, A q^2 -analogue operator for q^2 -analogue Fourier Analysis, *J. Math. Anal. App.*, **212**(1997), 571-582.