

A NEW FORM OF THE GENERALIZED COMPLETE ELLIPTIC INTEGRALS*

SHINGO TAKEUCHI

Dedicated to Professor Noboru Okazawa on the occasion of his 70th birthday

Abstract

Generalized trigonometric functions are applied to Legendre's form of complete elliptic integrals, and a new form of the generalized complete elliptic integrals of the Borweins is presented. According to the form, it can be easily shown that these integrals have similar properties to the classical ones. In particular, it is possible to establish a computation formula of the generalized π in terms of the arithmetic-geometric mean, in the classical way as the Gauss-Legendre algorithm for π by Brent and Salamin. Moreover, an elementary alternative proof of Ramanujan's cubic transformation is also given.

1. Introduction

Complete elliptic integrals of the first kind and of the second kind

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} = \int_0^1 \frac{dt}{\sqrt{(1 - t^2)(1 - k^2 t^2)}},$$
$$E(k) = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta = \int_0^1 \sqrt{\frac{1 - k^2 t^2}{1 - t^2}} dt$$

are classical integrals which have helped us, for instance, to evaluate the length of curves and to express exact solutions of differential equations.

In this paper we give a generalization of complete elliptic integrals as an application of generalized trigonometric functions. For this, we need the

2010 *Mathematics Subject Classification.* 33E05, 33C75, 11Z05.

Key words and phrases. Generalized trigonometric functions; Generalized complete elliptic integrals; Ramanujan's cubic transformation; Arithmetic-geometric mean; Gauss-Legendre's algorithm; p -Laplacian.

*This work was supported by JSPS KAKENHI Grant Number 24540218.

Received April 1, 2015; revised July 30, 2015.

generalized sine function $\sin_p \theta$ and the generalized π denoted by π_p , where $\sin_p \theta$ is the inverse function of

$$\sin_p^{-1} \theta := \int_0^\theta \frac{dt}{(1-t^p)^{1/p}}, \quad 0 \leq \theta \leq 1,$$

and π_p is the number defined by

$$\pi_p := 2 \sin_p^{-1} 1 = 2 \int_0^1 \frac{dt}{(1-t^p)^{1/p}} = \frac{2\pi}{p \sin \frac{\pi}{p}}.$$

Clearly, $\sin_2 \theta = \sin \theta$ and $\pi_2 = \pi$. These two appear in the eigenvalue problem of one-dimensional p -Laplacian:

$$-(|u'|^{p-2}u')' = \lambda|u|^{p-2}u, \quad u(0) = u(1) = 0.$$

Indeed, the eigenvalues are given as $\lambda_n = (p-1)(n\pi_p)^p$, $n = 1, 2, 3, \dots$, and the corresponding eigenfunction to λ_n is $u_n(x) = \sin_p(n\pi_p x)$ for each n . There are a lot of literature on generalized trigonometric functions and related functions. See [9, 10, 14, 16, 18, 22, 23, 24, 25, 28] for general properties as functions; [13, 14, 15, 22, 26, 29] for applications to differential equations involving p -Laplacian; [5, 10, 11, 16, 17, 22, 30] for basis properties for sequences of these functions.

Now, applying $\sin_p \theta$ and π_p to the complete elliptic functions, we define the *complete p -elliptic integrals of the first kind* $K_p(k)$ and of the *second kind* $E_p(k)$: for $p \in (1, \infty)$ and $k \in [0, 1)$

$$(1.1) \quad K_p(k) := \int_0^{\pi_p/2} \frac{d\theta}{(1-k^p \sin_p^p \theta)^{1-1/p}} = \int_0^1 \frac{dt}{(1-t^p)^{1/p}(1-k^p t^p)^{1-1/p}},$$

$$(1.2) \quad E_p(k) := \int_0^{\pi_p/2} (1-k^p \sin_p^p \theta)^{1/p} d\theta = \int_0^1 \left(\frac{1-k^p t^p}{1-t^p} \right)^{1/p} dt.$$

Here, each second equality of the definitions is obtained by setting $\sin_p \theta = t$. It is easy to see that for $p = 2$ these integrals are equivalent to the classical complete elliptic integrals $K(k)$ and $E(k)$.

It is worth pointing out that the Borweins [7, Section 5.5] define the *generalized complete elliptic integrals of the first and of the second kind* by

$$(1.3) \quad \mathbf{K}_s(k) := \frac{\pi}{2} F\left(\frac{1}{2} - s, \frac{1}{2} + s; 1; k^2\right),$$

$$(1.4) \quad \mathbf{E}_s(k) := \frac{\pi}{2} F\left(-\frac{1}{2} - s, \frac{1}{2} + s; 1; k^2\right)$$

for $|s| < 1/2$ and $0 \leq k < 1$, where $F(a, b; c; x)$ denotes the Gaussian hypergeometric function (see Section 2 for the definition). Note that $\mathbf{K}_0(k) = K(k)$

and $E_0(k) = E(k)$. According to Euler's integral representation (see [2, Theorem 2.2.1] or [33, p. 293]), we have

$$K_s(k) = \frac{\cos \pi s}{2s+1} \int_0^1 \frac{dt}{(1-t^{2/(2s+1)})^{(2s+1)/2} (1-k^2 t^{2/(2s+1)})^{1-(2s+1)/2}},$$

$$E_s(k) = \frac{\cos \pi s}{2s+1} \int_0^1 \left(\frac{1-k^2 t^{2/(2s+1)}}{1-t^{2/(2s+1)}} \right)^{(2s+1)/2} dt.$$

Thus

$$K_s(k) = \frac{\pi}{\pi_p} K_p(k^{2/p}), \quad E_s(k) = \frac{\pi}{\pi_p} E_p(k^{2/p}),$$

where $p = 2/(2s+1)$. To show this, one may use Proposition 2.8 below instead of integral representations. Anyway, we emphasize that the complete p -elliptic integrals (1.1) and (1.2) give representations of generalized complete elliptic integrals in Legendre's form with generalized trigonometric functions. The advantage of using the complete p -elliptic integrals lies in the fact that it is possible to prove formulas of the generalized complete elliptic integrals simply as well as that of the classical complete elliptic integrals. For example, we have known the following Legendre relation between $K(k)$ and $E(k)$ (see [2, 7, 20, 33]).

$$(1.5) \quad K'(k)E(k) + K(k)E'(k) - K(k)K'(k) = \frac{\pi}{2},$$

where $k' := \sqrt{1-k^2}$, $K'(k) := K(k')$ and $E'(k) := E(k')$. For this we can show the following relation between $K_p(k)$ and $E_p(k)$.

THEOREM 1.1. For $k \in (0, 1)$

$$(1.6) \quad K'_p(k)E_p(k) + K_p(k)E'_p(k) - K_p(k)K'_p(k) = \frac{\pi_p}{2},$$

where $k' := (1-k^p)^{1/p}$, $K'_p(k) := K_p(k')$ and $E'_p(k) := E_p(k')$.

In fact, it is known that $K_s(k)$ and $E_s(k)$ also satisfy the similar relation (2.17) below to (1.6), which follows from Elliott's identity (2.18) below. In contrast to this, our approach with generalized trigonometric functions seems to be more elementary and self-contained.

As an application of complete p -elliptic integrals, we establish a computation formula of π_p . Let us first mention the case $p = 2$. In that case, consider the sequences $\{a_n\}$ and $\{b_n\}$ satisfying $a_0 = a$, $b_0 = b$, where $a \geq b > 0$, and

$$a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n}, \quad n = 0, 1, 2, \dots$$

It is easily checked that these sequences converge to a common limit, the *arithmetic-geometric mean of a and b*, denoted by $M(a, b)$. It is well-known that on May 30th in 1799 Gauss discovered a celebrated relation between $M(a, b)$ and $K(k)$ (precisely, a relation between $M(1, \sqrt{2})$ and the lemniscate integral):

$$(1.7) \quad K(k) = \frac{\pi}{2} \frac{1}{M(1, k')},$$

where $k' = \sqrt{1 - k^2}$. Combining (1.5) and (1.7) with $k = k' = 1/\sqrt{2}$, Brent [8] and Salamin [27] independently established the following formula (see also [2, 7] for the proof).

$$(1.8) \quad \pi = \frac{2M\left(1, \frac{1}{\sqrt{2}}\right)^2}{1 - \sum_{n=0}^{\infty} 2^n (a_n^2 - b_n^2)},$$

where the initial data of $\{a_n\}$ and $\{b_n\}$ are $a = 1$ and $b = 1/\sqrt{2}$. This is known as a fundamental formula to the Gauss-Legendre algorithm, or the Brent-Salamin algorithm, for computing the value of π .

Owing to (1.8), it is natural to try to establish a computation formula of π_p . In addition we are interested in finding such an elementary way of its construction as Brent and Salamin.

In the present paper, we give the formula only for the case $p = 3$. We prepare notation for stating results. Let $a \geq b > 0$, and assume that $\{a_n\}$ and $\{b_n\}$ are sequences satisfying $a_0 = a$, $b_0 = b$ and

$$(1.9) \quad a_{n+1} = \frac{a_n + 2b_n}{3}, \quad b_{n+1} = \sqrt[3]{\frac{(a_n^2 + a_n b_n + b_n^2)b_n}{3}}, \quad n = 0, 1, 2, \dots$$

It is easy to see that both the sequences converge to the same limit as $n \rightarrow \infty$, denoted by $M_3(a, b)$. Then, we obtain

THEOREM 1.2. *Let $0 \leq k < 1$. Then*

$$K_3(k) = \frac{\pi_3}{2} \frac{1}{M_3(1, k')},$$

where $k' = \sqrt[3]{1 - k^3}$.

Actually, Theorem 1.2 is identical to the result of the Borweins [6, Theorem 2.1 (b)] (with some trivial typos). In either proof, it is essential to show Ramanujan's cubic transformation (Lemma 3.1 below). We will give an alternative proof for this by more elementary calculation with properties of $K_3(k)$.

By Theorems 1.1 and 1.2 we obtain the following formula of π_3 .

THEOREM 1.3. *Let $a = 1$ and $b = 1/\sqrt[3]{2}$. Then*

$$\pi_3 = \frac{2M_3\left(1, \frac{1}{\sqrt[3]{2}}\right)^2}{1 - 2 \sum_{n=1}^{\infty} 3^n (a_n + c_n) c_n},$$

where $\{a_n\}$ and $\{b_n\}$ are the sequences (1.9) and $c_n := \sqrt[3]{a_n^3 - b_n^3}$.

By the theory of theta functions, the Borweins [6, Section 3] give three iterations for π . One of them is obtained from the same sequences as (1.9):

$$(1.10) \quad \pi = \frac{3M_3\left(1, \left(\frac{\sqrt{3}-1}{2}\right)'\right)^2}{1 - \sum_{n=0}^{\infty} 3^{n+1} (a_n^2 - a_{n+1}^2)}.$$

However, note that the initial data $a = 1$ and $b = ((\sqrt{3}-1)/2)'$ of (1.10) is different from our initial data $a = 1$ and $b = 1/\sqrt[3]{2}$.

It is a simple matter to obtain other formulas for π_3 if we combine $\pi_3 = 4\sqrt{3}\pi/9 = 2.418\dots$ with (1.8) or (1.10). The former converges quadratically to π_3 and the latter does cubically. On the other hand, our formula in Theorem 1.3 converges cubically to π_3 (Table 1). However, we are not interested in such trivial formulas obtained from those of π , and it is not our purpose to study the speed of convergence and we will not develop this point here.

	25 digits	Error
q_1	2.418399152309345558425031	2.9449×10^{-12}
q_2	2.418399152312290467458771	4.0425×10^{-40}
q_3	2.418399152312290467458771	1.0367×10^{-124}
q_4	2.418399152312290467458771	1.8728×10^{-379}

Table 1. Convergence of q_m to π_3 , where $q_m := \frac{2a_{m+1}^2}{1 - 2 \sum_{n=1}^m 3^n (a_n + c_n) c_n}$.

This paper is organized as follows. In Section 2 we have compiled some basic facts of complete p -elliptic integrals. In particular we show Legendre's relation for $K_p(k)$ and $E_p(k)$ (Theorem 1.1) and observe relationship between the complete p -elliptic integrals and the Gaussian hypergeometric functions (Theorem 1.2). Section 3 establishes a computation formula of π_p with $p = 3$ as an application of complete p -elliptic integrals (Theorem 1.3). In particular, we give

an elementary proof of Ramanujan’s cubic transformation by using our representation of integrals (Lemma 3.1).

2. Complete p -elliptic integrals

In this section, we present some basic properties of complete p -elliptic integrals $K_p(k)$ and $E_p(k)$.

Let $1 < p < \infty$. We repeat the definition of complete p -elliptic integrals of the first kind $K_p(k)$ and of the second kind $E_p(k)$: for $k \in [0, 1)$,

$$(1.1) \quad K_p(k) := \int_0^{\pi_p/2} \frac{d\theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} = \int_0^1 \frac{dt}{(1 - t^p)^{1/p}(1 - k^p t^p)^{1-1/p}},$$

$$(1.2) \quad E_p(k) := \int_0^{\pi_p/2} (1 - k^p \sin_p^p \theta)^{1/p} d\theta = \int_0^1 \left(\frac{1 - k^p t^p}{1 - t^p} \right)^{1/p} dt,$$

where $\sin_p \theta$ and π_p have been defined in the Introduction. As is traditional, we will use the notation $k' := (1 - k^p)^{1/p}$. The variable k is often called the *modulus*, and k' is the *complementary modulus*. The *complementary integrals* $K'_p(k)$ and $E'_p(k)$ are defined by $K'_p(k) := K_p(k')$ and $E'_p(k) := E_p(k')$.

Let $\cos_p \theta := (1 - \sin_p^p \theta)^{1/p}$. The following formulas will be frequently used:

$$\begin{aligned} \sin_p^p \theta + \cos_p^p \theta &= 1, \\ \frac{d}{d\theta}(\sin_p \theta) &= \cos_p \theta, \quad \frac{d}{d\theta}(\cos_p^{p-1} \theta) = -(p - 1) \sin_p^{p-1} \theta. \end{aligned}$$

If $p = 2$ then $\sin_p \theta$, $\cos_p \theta$ and π_p coincide with the usual $\sin \theta$, $\cos \theta$ and π , respectively, so that these properties above are familiar.

The functions $K_p(k)$ and $E_p(k)$ satisfy a system of differential equations.

PROPOSITION 2.1.

$$\frac{dE_p}{dk} = \frac{E_p - K_p}{k}, \quad \frac{dK_p}{dk} = \frac{E_p - (k')^p K_p}{k(k')^p}.$$

Proof. Differentiating $E_p(k)$ we have

$$\begin{aligned} \frac{dE_p}{dk} &= \int_0^{\pi_p/2} \frac{d}{dk} (1 - k^p \sin_p^p \theta)^{1/p} d\theta \\ &= \int_0^{\pi_p/2} \frac{-k^{p-1} \sin_p^p \theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} d\theta \\ &= \frac{1}{k} \int_0^{\pi_p/2} \frac{1 - k^p \sin_p^p \theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} d\theta - \frac{1}{k} \int_0^{\pi_p/2} \frac{d\theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} \\ &= \frac{1}{k} (E_p - K_p). \end{aligned}$$

Next, for $K_p(k)$

$$(2.1) \quad \frac{dK_p}{dk} = \int_0^{\pi_p/2} \frac{(p-1)k^{p-1} \sin_p^p \theta}{(1-k^p \sin_p^p \theta)^{2-1/p}} d\theta.$$

Here we see that

$$\begin{aligned} & \frac{d}{d\theta} \left(\frac{-\cos_p^{p-1} \theta}{(1-k^p \sin_p^p \theta)^{1-1/p}} \right) \\ &= \frac{(p-1) \sin_p^{p-1} \theta (1-k^p \sin_p^p \theta) - (p-1)k^p \sin_p^{p-1} \theta \cos_p^p \theta}{(1-k^p \sin_p^p \theta)^{2-1/p}} \\ &= \frac{(p-1)(k')^p \sin_p^{p-1} \theta}{(1-k^p \sin_p^p \theta)^{2-1/p}}, \end{aligned}$$

so that we use integration by parts as

$$\begin{aligned} \frac{dK_p}{dk} &= \int_0^{\pi_p/2} \frac{k^{p-1}}{(k')^p} \frac{d}{d\theta} \left(\frac{-\cos_p^{p-1} \theta}{(1-k^p \sin_p^p \theta)^{1-1/p}} \right) \sin_p \theta d\theta \\ &= \frac{k^{p-1}}{(k')^p} \left[\frac{-\cos_p^{p-1} \theta \sin_p \theta}{(1-k^p \sin_p^p \theta)^{1-1/p}} \right]_0^{\pi_p/2} + \frac{k^{p-1}}{(k')^p} \int_0^{\pi_p/2} \frac{\cos_p^p \theta}{(1-k^p \sin_p^p \theta)^{1-1/p}} d\theta \\ &= \frac{k^{p-1}}{(k')^p} \int_0^{\pi_p/2} \frac{1}{k^p} \cdot \frac{1-k^p \sin_p^p \theta - (1-k^p)}{(1-k^p \sin_p^p \theta)^{1-1/p}} d\theta \\ &= \frac{1}{k(k')^p} (E_p - (k')^p K_p). \end{aligned}$$

This completes the proof. □

Proposition 2.1 now yields Theorem 1.1.

Proof of Theorem 1.1. We will differentiate the left-hand side of (1.6) and apply Proposition 2.1. As $dk'/dk = -(k/k')^{p-1}$ we have

$$(2.2) \quad \frac{dK'_p}{dk} = \frac{k^p K'_p - E'_p}{k(k')^p}, \quad \frac{dE'_p}{dk} = k^{p-1} \frac{K'_p - E'_p}{(k')^p}.$$

Hence a direct computation shows that

$$\begin{aligned}
 & \frac{d}{dk} (K'_p E_p + K_p E'_p - K_p K'_p) \\
 &= \frac{k^p K'_p - E'_p}{k(k')^p} \cdot E_p + K'_p \cdot \frac{E_p - K_p}{k} + \frac{E_p - (k')^p K_p}{k(k')^p} \cdot E'_p \\
 & \quad + K_p \cdot k^{p-1} \frac{K'_p - E'_p}{(k')^p} - \frac{E_p - (k')^p K_p}{k(k')^p} \cdot K'_p - K_p \cdot \frac{k^p K'_p - E'_p}{k(k')^p} \\
 &= E'_p E_p \left(-\frac{1}{k(k')^p} + \frac{1}{k(k')^p} \right) + (K'_p E_p - K_p E'_p) \left(\frac{k^{p-1}}{(k')^p} + \frac{1}{k} - \frac{1}{k(k')^p} \right) \\
 & \quad + K'_p K_p \left(-\frac{1}{k} + \frac{k^{p-1}}{(k')^p} + \frac{1}{k} - \frac{k^{p-1}}{(k')^p} \right) \\
 &= 0.
 \end{aligned}$$

Therefore the left-hand side of (1.6) is a constant C .

We will evaluate C as follows. It is easy to see that $\lim_{k \rightarrow +0} K_p E'_p = \pi_p/2$. Moreover, since

$$\begin{aligned}
 |(K_p - E_p)K'_p| &= \int_0^{\pi_p/2} \left(\frac{1}{(1 - k^p \sin_p^p \theta)^{1-1/p}} - (1 - k^p \sin_p^p \theta)^{1/p} \right) d\theta \\
 & \quad \cdot \int_0^{\pi_p/2} \frac{d\theta}{(1 - (k')^p \sin_p^p \theta)^{1-1/p}} \\
 &= \int_0^{\pi_p/2} \frac{k^p \sin_p^p \theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} d\theta \cdot \int_0^{\pi_p/2} \frac{d\theta}{(\cos_p^p \theta + k^p \sin_p^p \theta)^{1-1/p}} \\
 &\leq k^p K_p(k) \cdot \frac{1}{k^{p-1}} \frac{\pi_p}{2} \\
 &= k K_p(k) \frac{\pi_p}{2},
 \end{aligned}$$

we obtain $\lim_{k \rightarrow +0} (K_p - E_p)K'_p = 0$. Thus, letting $k \rightarrow +0$ in the left-hand side of (1.6), we conclude that $C = \pi_p/2$. □

PROPOSITION 2.2. $K_p(k)$ and $K'_p(k)$ satisfy

$$\frac{d}{dk} \left(k(k')^p \frac{dy}{dk} \right) = (p - 1)k^{p-1}y,$$

that is

$$k(1 - k^p) \frac{d^2y}{dk^2} + (1 - (p + 1)k^p) \frac{dy}{dk} - (p - 1)k^{p-1}y = 0.$$

Moreover $E_p(k)$ and $E'_p(k) - K'_p(k)$ satisfy

$$(k')^p \frac{d}{dk} \left(k \frac{dy}{dk} \right) = -k^{p-1}y,$$

that is

$$k(1-k^p) \frac{d^2y}{dk^2} + (1-k^p) \frac{dy}{dk} + k^{p-1}y = 0.$$

Proof. Let us first give proofs for K_p and E_p . Repeated application of Proposition 2.1 and $d(k')^p/dk = -pk^{p-1}$ enables us to see that

$$\begin{aligned} \frac{d}{dk} \left(k(k')^p \frac{dK_p}{dk} \right) &= \frac{d}{dk} (E_p - (k')^p K_p) \\ &= \frac{1}{k} (E_p - K_p) - \left(-pk^{p-1} K_p + \frac{1}{k} (E_p - (k')^p K_p) \right) \\ &= (p-1)k^{p-1} K_p \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dk} \left(k \frac{dE_p}{dk} \right) &= \frac{d}{dk} (E_p - K_p) \\ &= \frac{1}{k} (E_p - K_p) - \frac{1}{k(k')^p} (E_p - (k')^p K_p) \\ &= -\frac{k^{p-1}}{(k')^p} E_p. \end{aligned}$$

Similarly, it follows easily from (2.2) that K'_p satisfies

$$\frac{d}{dk} \left(k(k')^p \frac{dK'_p}{dk} \right) = (p-1)k^{p-1} K'_p.$$

Set $H_p(k) := E_p(k) - K_p(k)$. Using (2.2) repeatedly, we have $dH'_p/dk = E'_p/k$ and

$$\frac{d}{dk} \left(k \frac{dH'_p}{dk} \right) = \frac{dE'_p}{dk} = -\frac{k^{p-1}}{(k')^p} H'_p.$$

The proof is complete. \square

Define $K_p^*(k)$ and $E_p^*(k)$ as conjugates for $K_p(k)$ and $E_p(k)$ respectively: for $k \in [0, 1)$

$$(2.3) \quad K_p^*(k) := \int_0^{\pi_p/2} \frac{d\theta}{(1-k^p \sin_p^p \theta)^{1/p}} = \int_0^1 \frac{dt}{(1-t^p)^{1/p} (1-k^p t^p)^{1/p}},$$

$$(2.4) \quad E_p^*(k) := \int_0^{\pi_p/2} (1-k^p \sin_p^p \theta)^{1-1/p} d\theta = \int_0^1 \frac{(1-k^p t^p)^{1-1/p}}{(1-t^p)^{1/p}} dt.$$

It is clear that $K_2^*(k) = K_2(k) = K(k)$ and $E_2^*(k) = E_2(k) = E(k)$. The integral $K_p^*(k)$ appears in the study [29] for bifurcation problems of p -Laplacian. In [32], $K_p^*(k)$ with t^p replaced by t^2 is applied to the planar p -elastic problem.

Here are some elementary relations between the integrals we have introduced. To state the relations, it is convenient to use the notation $i_p := e^{i\pi/p}$ and for a nonnegative number ℓ

$$(2.5) \quad K_p^*(i_p \ell) := \int_0^{\pi_p/2} \frac{d\theta}{(1 + \ell^p \sin_p^p \theta)^{1/p}} = \int_0^1 \frac{dt}{(1 - t^p)^{1/p}(1 + \ell^p t^p)^{1/p}},$$

$$(2.6) \quad E_p^*(i_p \ell) := \int_0^{\pi_p/2} (1 + \ell^p \sin_p^p \theta)^{1-1/p} d\theta = \int_0^1 \frac{(1 + \ell^p t^p)^{1-1/p}}{(1 - t^p)^{1/p}} dt.$$

PROPOSITION 2.3. *Let $0 \leq k < 1$. Then*

$$(2.7) \quad K_{p^*}(k^{p-1}) = (p - 1) \frac{1}{k'} K_p^* \left(i_p \frac{k}{k'} \right),$$

$$(2.8) \quad K_{p^*}(k^{p-1}) = (p - 1) K_p(k),$$

$$(2.9) \quad E_{p^*}(k^{p-1}) = (p - 1)(k')^{p-1} E_p^* \left(i_p \frac{k}{k'} \right),$$

$$(2.10) \quad E_{p^*}(k^{p-1}) = E_p(k) + (p - 2)(k')^p K_p(k),$$

where $p^* := p/(p - 1)$.

Proof. Let us first show (2.7) and (2.9). Setting $1 - t^{p^*} = u^p$ in each integral, we have

$$(2.11) \quad \begin{aligned} K_{p^*}(k^{p-1}) &= \int_0^1 \frac{dt}{(1 - t^{p^*})^{1/p^*} (1 - k^p t^{p^*})^{1-1/p^*}} \\ &= (p - 1) \int_0^1 \frac{du}{(1 - u^p)^{1/p} (1 - k^p + k^p u^p)^{1/p}} \\ &= (p - 1) \frac{1}{k'} K_p^* \left(i_p \frac{k}{k'} \right) \end{aligned}$$

and

$$(2.12) \quad \begin{aligned} E_{p^*}(k^{p-1}) &= \int_0^1 \left(\frac{1 - k^p t^{p^*}}{1 - t^{p^*}} \right)^{1/p^*} dt \\ &= (p - 1) \int_0^1 \frac{(1 - k^p + k^p u^p)^{1-1/p}}{(1 - u^p)^{1/p}} du \\ &= (p - 1)(k')^{p-1} E_p^* \left(i_p \frac{k}{k'} \right). \end{aligned}$$

Next we will prove (2.8). Changing the variable as

$$u^p = \frac{(1 - k^p)t^p}{1 - k^p t^p},$$

in (2.11) we obtain

$$\begin{aligned} K_{p^*}(k^{p-1}) &= (p-1) \int_0^1 \frac{\frac{(1 - k^p)^{1/p}}{(1 - k^p t^p)^{1+1/p}}}{\left(\frac{1 - t^p}{1 - k^p t^p}\right)^{1/p} \left(\frac{1 - k^p}{1 - k^p t^p}\right)^{1/p}} dt \\ &= (p-1) \int_0^1 \frac{dt}{(1 - t^p)^{1/p} (1 - k^p t^p)^{1-1/p}} \\ &= (p-1) K_p(k). \end{aligned}$$

To deduce (2.10), we make use of the same change of variable above to (2.12).

$$\begin{aligned} E_{p^*}(k^{p-1}) &= (p-1) \int_0^1 \frac{\left(\frac{1 - k^p}{1 - k^p t^p}\right)^{1-1/p}}{\left(\frac{1 - t^p}{1 - k^p t^p}\right)^{1/p}} \frac{(1 - k^p)^{1/p}}{(1 - k^p t^p)^{1+1/p}} dt \\ &= (p-1)(k')^p \int_0^1 \frac{dt}{(1 - t^p)^{1/p} (1 - k^p t^p)^{2-1/p}}. \end{aligned}$$

In the last integral, after setting $t = \sin_p \theta$, using (2.1) and Proposition 2.1 we have

$$\begin{aligned} \int_0^1 \frac{dt}{(1 - t^p)^{1/p} (1 - k^p t^p)^{2-1/p}} &= \int_0^{\pi_p/2} \frac{d\theta}{(1 - k^p \sin_p^p \theta)^{2-1/p}} \\ &= \int_0^{\pi_p/2} \frac{d\theta}{(1 - k^p \sin_p^p \theta)^{1-1/p}} \\ &\quad + \int_0^{\pi_p/2} \frac{k^p \sin_p^p \theta}{(1 - k^p \sin_p^p \theta)^{2-1/p}} d\theta \\ &= K_p + \frac{k}{p-1} \frac{dK_p}{dk} \\ &= K_p + \frac{1}{(p-1)(k')^p} (E_p - (k')^p K_p). \end{aligned}$$

Thus

$$\begin{aligned} E_{p^*}(k^{p-1}) &= (p-1)(k')^p \left(K_p + \frac{1}{(p-1)(k')^p} (E_p - (k')^p K_p) \right) \\ &= E_p + (p-2)(k')^p K_p. \end{aligned}$$

Therefore we conclude (2.10). □

From Proposition 2.3 it immediately follows

COROLLARY 2.4. *Let $0 \leq k < 1$. Then*

$$\begin{aligned} K_p(k) &= \frac{1}{k'} K_p^* \left(i_p \frac{k}{k'} \right), \\ E_p(k) &= (k')^{p-1} \left((p-1) E_p^* \left(i_p \frac{k}{k'} \right) - (p-2) K_p^* \left(i_p \frac{k}{k'} \right) \right). \end{aligned}$$

For $p = 2$, the identities of Corollary 2.4 are equivalent to:

$$K(k) = \frac{1}{k'} K \left(i \frac{k}{k'} \right), \quad E(k) = k' E \left(i \frac{k}{k'} \right),$$

which can be found in [21, Table 4, p. 319].

The next corollary means that $pK_p(\ell^{1/p})$ and $E_p(\ell^{1/p}) - (1-\ell)K_p(\ell^{1/p})$ have duality properties with respect to p .

COROLLARY 2.5. *Let $0 \leq \ell < 1$. Then*

$$(2.13) \quad p^* K_{p^*}(\ell^{1/p^*}) = p K_p(\ell^{1/p}),$$

$$(2.14) \quad E_{p^*}(\ell^{1/p^*}) - (1-\ell) K_{p^*}(\ell^{1/p^*}) = E_p(\ell^{1/p}) - (1-\ell) K_p(\ell^{1/p}).$$

Proof. Putting $k = \ell^{1/p}$ in (2.8) and multiplying it by p^* , we obtain (2.13) at once. To deduce (2.14), putting $k = \ell^{1/p}$ in (2.10) and using (2.13) we have

$$\begin{aligned} &E_{p^*}(\ell^{1/p^*}) - (1-\ell) K_{p^*}(\ell^{1/p^*}) \\ &= E_p(\ell^{1/p}) + (p-2)(1-\ell) K_p(\ell^{1/p}) - (1-\ell)(p-1) K_p(\ell^{1/p}) \\ &= E_p(\ell^{1/p}) - (1-\ell) K_p(\ell^{1/p}). \end{aligned}$$

Therefore we conclude (2.14). □

Letting $\ell = 0$ in (2.13) we have $p^* \pi_{p^*} = p \pi_p$, which was indicated in [10, 22].

PROPOSITION 2.6.

$$K_p\left(\frac{1}{\sqrt[p]{2}}\right) = \frac{\sqrt[p]{2}\Gamma\left(\frac{1}{2p}\right)^2}{4p\Gamma\left(\frac{1}{p}\right)\cos\frac{\pi}{2p}},$$

$$E_p\left(\frac{1}{\sqrt[p]{2}}\right) = \frac{\sqrt[p]{2}}{8p\Gamma\left(\frac{1}{p}\right)}\left(\frac{\Gamma\left(\frac{1}{2p}\right)^2}{\cos\frac{\pi}{2p}} + \frac{2p\Gamma\left(\frac{1}{2p} + \frac{1}{2}\right)^2}{\sin\frac{\pi}{2p}}\right),$$

where Γ is the gamma function.

Proof. Let $k = 1/\sqrt[p]{2}$, then $k' = 1/\sqrt[p]{2}$. By Corollary 2.4 we have

$$K_p\left(\frac{1}{\sqrt[p]{2}}\right) = \sqrt[p]{2}K_p^*(i_p) = \sqrt[p]{2}\int_0^1 \frac{dt}{(1-t^{2p})^{1/p}}.$$

Putting $t^{2p} = x$, we obtain

$$\int_0^1 \frac{dt}{(1-t^{2p})^{1/p}} = \frac{1}{2p}B\left(\frac{1}{2p}, 1 - \frac{1}{p}\right) = \frac{\Gamma\left(\frac{1}{2p}\right)\Gamma\left(1 - \frac{1}{p}\right)}{2p\Gamma\left(1 - \frac{1}{2p}\right)},$$

where B is the beta function. Since

$$(2.15) \quad \Gamma\left(1 - \frac{1}{p}\right) = \frac{\pi}{\Gamma\left(\frac{1}{p}\right)\sin\frac{\pi}{p}} = \frac{\pi}{2\Gamma\left(\frac{1}{p}\right)\sin\frac{\pi}{2p}\cos\frac{\pi}{2p}},$$

$$(2.16) \quad \Gamma\left(1 - \frac{1}{2p}\right) = \frac{\pi}{\Gamma\left(\frac{1}{2p}\right)\sin\frac{\pi}{2p}},$$

the first formula in the proposition follows.

Similarly, Corollary 2.4 yields

$$E_p\left(\frac{1}{\sqrt[p]{2}}\right) = \frac{1}{\sqrt[p]{2}}((p-1)E_p^*(i_p) - (p-2)K_p^*(i_p)).$$

Since

$$E_p^*(i_p) = \int_0^1 \frac{1+t^p}{(1-t^{2p})^{1/p}} dt = K_p^*(i_p) + \int_0^1 \frac{t^p}{(1-t^{2p})^{1/p}} dt,$$

we have

$$\begin{aligned} E_p\left(\frac{1}{\sqrt[p]{2}}\right) &= \frac{1}{\sqrt[p]{2}} \left(K_p^*(i_p) + (p-1) \int_0^1 \frac{t^p}{(1-t^{2p})^{1/p}} \right) \\ &= \frac{1}{2p \sqrt[p]{2}} \left(\frac{\Gamma\left(\frac{1}{2p}\right)\Gamma\left(1-\frac{1}{p}\right)}{\Gamma\left(1-\frac{1}{2p}\right)} + (p-1) \frac{\Gamma\left(\frac{1}{2p}+\frac{1}{2}\right)\Gamma\left(1-\frac{1}{p}\right)}{\Gamma\left(\frac{3}{2}-\frac{1}{2p}\right)} \right). \end{aligned}$$

By (2.15), (2.16) and

$$\Gamma\left(\frac{3}{2}-\frac{1}{2p}\right) = \frac{\pi}{2p^* \Gamma\left(\frac{1}{2p}+\frac{1}{2}\right) \cos \frac{\pi}{2p}},$$

the second formula in the proposition follows. □

For $p = 2$ Proposition 2.3 gives that

$$K\left(\frac{1}{\sqrt{2}}\right) = \frac{\Gamma\left(\frac{1}{4}\right)^2}{4\sqrt{\pi}}, \quad E\left(\frac{1}{\sqrt{2}}\right) = \frac{\Gamma\left(\frac{1}{4}\right)^2 + 4\Gamma\left(\frac{3}{4}\right)^2}{8\sqrt{\pi}},$$

which can be found in [7, Theorem 1.7]. See also [21, Section 13.8] and [33, Section 22.8].

On account of Proposition 2.6, it is possible to show $C = \pi_p/2$ in the proof of Theorem 1.1 in another way. Indeed, letting $k = 1/\sqrt[p]{2}$ we have

$$\begin{aligned} C &= 2K_p\left(\frac{1}{\sqrt[p]{2}}\right)E_p\left(\frac{1}{\sqrt[p]{2}}\right) - K_p\left(\frac{1}{\sqrt[p]{2}}\right)^2 \\ &= \frac{(\sqrt[p]{2})^2 \Gamma\left(\frac{1}{2p}\right)^2}{16p^2 \Gamma\left(\frac{1}{p}\right)^2 \cos \frac{\pi}{2p}} \left(\frac{\Gamma\left(\frac{1}{2p}\right)^2}{\cos \frac{\pi}{2p}} + \frac{2p \Gamma\left(\frac{1}{2p}+\frac{1}{2}\right)^2}{\sin \frac{\pi}{2p}} \right) - \frac{(\sqrt[p]{2})^2 \Gamma\left(\frac{1}{2p}\right)^4}{16p^2 \Gamma\left(\frac{1}{p}\right)^2 \cos^2 \frac{\pi}{2p}} \\ &= \frac{(\sqrt[p]{2})^2 \Gamma\left(\frac{1}{2p}\right)^2 \Gamma\left(\frac{1}{2p}+\frac{1}{2}\right)^2}{4p \Gamma\left(\frac{1}{p}\right)^2 \sin \frac{\pi}{p}} \\ &= \frac{2^{2/p-2}}{p \sin \frac{\pi}{p}} \left(\frac{\sqrt{\pi}}{2^{1/p-1}} \right)^2 = \frac{\pi_p}{2}, \end{aligned}$$

where we have used the duplication formula

$$\Gamma(2z) = \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right)$$

with $z = 1/(2p)$.

The remainder of this section will be devoted to the study of relation between complete p -elliptic integrals and hypergeometric series.

For a real number a and a natural number n , we define

$$(a)_n := \frac{\Gamma(a+n)}{\Gamma(a)} = (a+n-1)(a+n-2) \cdots (a+1)a.$$

We adopt the convention that $(a)_0 := 1$. For $|x| < 1$ the series

$$F(a, b; c; x) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}$$

is called a *Gaussian hypergeometric series*. See [2, 7, 20, 33] for more details.

LEMMA 2.7. For $n = 0, 1, 2, \dots$

$$\int_0^{\pi_p/2} \sin_p^{pn} \theta \, d\theta = \frac{\pi_p}{2} \frac{\left(\frac{1}{p}\right)_n}{n!}.$$

Proof. Letting $\sin_p^p \theta = t$, we have

$$\int_0^{\pi_p/2} \sin_p^{pn} \theta \, d\theta = \frac{1}{p} \int_0^1 t^{n+1/p-1} (1-t)^{-1/p} dt = \frac{1}{p} B\left(n + \frac{1}{p}, 1 - \frac{1}{p}\right).$$

Moreover,

$$\begin{aligned} \frac{1}{p} B\left(n + \frac{1}{p}, 1 - \frac{1}{p}\right) &= \frac{1}{p} B\left(\frac{1}{p}, 1 - \frac{1}{p}\right) \frac{B\left(n + \frac{1}{p}, 1 - \frac{1}{p}\right)}{B\left(\frac{1}{p}, 1 - \frac{1}{p}\right)} \\ &= \frac{\pi_p}{2} \frac{\Gamma\left(n + \frac{1}{p}\right)}{\Gamma\left(\frac{1}{p}\right) \Gamma(n+1)} \\ &= \frac{\pi_p}{2} \frac{\left(\frac{1}{p}\right)_n}{n!}, \end{aligned}$$

and the lemma follows. □

PROPOSITION 2.8. For $k \in (0, 1)$

$$K_p(k) = \frac{\pi_p}{2} F\left(\frac{1}{p}, 1 - \frac{1}{p}; 1; k^p\right),$$

$$E_p(k) = \frac{\pi_p}{2} F\left(\frac{1}{p}, -\frac{1}{p}; 1; k^p\right),$$

$$K_p^*(k) = \frac{\pi_p}{2} F\left(\frac{1}{p}, \frac{1}{p}; 1; k^p\right),$$

$$E_p^*(k) = \frac{\pi_p}{2} F\left(\frac{1}{p}, \frac{1}{p} - 1; 1; k^p\right).$$

Proof. Binomial series expansion gives

$$K_p(k) = \int_0^{\pi_p/2} (1 - k^p \sin_p^p \theta)^{1/p-1} d\theta = \sum_{n=0}^{\infty} (-1)^n \binom{\frac{1}{p}-1}{n} k^{pn} \int_0^{\pi_p/2} \sin_p^{pn} \theta d\theta.$$

Here, using Lemma 2.7 and the fact

$$(-1)^n \binom{\frac{1}{p}-1}{n} = (-1)^n \frac{\left(\frac{1}{p}-1\right)\left(\frac{1}{p}-2\right)\cdots\left(\frac{1}{p}-n\right)}{n!} = \frac{\left(1-\frac{1}{p}\right)_n}{(1)_n},$$

we see that

$$K_p(k) = \frac{\pi_p}{2} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{p}\right)_n \left(1-\frac{1}{p}\right)_n k^{pn}}{(1)_n n!} = \frac{\pi_p}{2} F\left(\frac{1}{p}, 1 - \frac{1}{p}; 1; k^p\right).$$

The other cases are similar and we left to the reader. □

A hypergeometric series $F(a, b; c; x)$ satisfies the hypergeometric differential equation

$$x(1-x) \frac{d^2y}{dx^2} + (c - (a+b+1)x) \frac{dy}{dx} - aby = 0.$$

From the fact and Proposition 2.8 we can also prove Proposition 2.2.

As mentioned in the Introduction, the Borweins [7, Section 5.5] define the generalized complete elliptic integrals of the first kind $K_s(k)$ and of the second kind $E_s(k)$ by (1.3) and (1.4) respectively. They indicate that these functions satisfy

$$(2.17) \quad K'_s(k)E_s(k) + K_s(k)E'_s(k) - K_s(k)K'_s(k) = \frac{\pi \cos \pi s}{2(1+2s)},$$

where $K'_s(k) := K_s(k')$, $E'_s(k) := E_s(k')$ and $k' := \sqrt{1-k^2}$. Letting $s = 1/p - 1/2$ in this equality we can also prove Theorem 1.1.

They obtained (2.17), relying on the following identity of hypergeometric functions with $a = -b = c = s$:

$$\begin{aligned}
 (2.18) \quad & F\left(\frac{1}{2} + a, -\frac{1}{2} - c; a + b + 1; x\right) F\left(\frac{1}{2} - a, \frac{1}{2} + c; b + c + 1; 1 - x\right) \\
 & + F\left(\frac{1}{2} + a, \frac{1}{2} - c; a + b + 1; x\right) F\left(-\frac{1}{2} - a, \frac{1}{2} + c; b + c + 1; 1 - x\right) \\
 & - F\left(\frac{1}{2} + a, \frac{1}{2} - c; a + b + 1; x\right) F\left(\frac{1}{2} - a, \frac{1}{2} + c; b + c + 1; 1 - x\right) \\
 & = \frac{\Gamma(a + b + 1)\Gamma(b + c + 1)}{\Gamma(a + b + c + \frac{3}{2})\Gamma(b + \frac{1}{2})},
 \end{aligned}$$

which was given by Elliott [19] (see also [1], [2, Theorem 3.2.8] and [20, (13) p. 85]). In contrast to this, our approach to Theorem 1.1 is more self-contained.

Finally in this section, we refer the reader to [1] for generalized elliptic integrals in geometric function theory and the relationship with hypergeometric functions. In [1] they also define a generalized Jacobian elliptic function related to $K_s(k)$. The author [30] produces generalized Jacobian elliptic functions with two parameters p and q related to $K_p(k)$, but as real functions (cf. [29]).

3. Application

In this section, we will apply the complete p -elliptic integrals (1.1) and (1.2) to compute π_p , and prove Theorem 1.3. For the very special case of π_p with $p = 3$, we are able to obtain a computation formula like (1.8) for π by Brent [8] and Salamin [27].

Let $a \geq b > 0$. Consider the sequences $\{a_n\}$ and $\{b_n\}$ satisfying $a_0 = a$, $b_0 = b$ and

$$a_{n+1} = \frac{a_n + 2b_n}{3}, \quad b_{n+1} = \sqrt[3]{\frac{(a_n^2 + a_n b_n + b_n^2)b_n}{3}}, \quad n = 0, 1, 2, \dots$$

It is easy to see that $a_n \geq b_n$ for any n , $\{a_n\}$ is decreasing and $\{b_n\}$ is increasing. Hence each sequence converges to a limit as $n \rightarrow \infty$. Moreover, since

$$(3.1) \quad a_{n+1} - b_{n+1} \leq \frac{a_n + 2b_n}{3} - b_n = \frac{1}{3}(a_n - b_n),$$

these limits are same. We will denote by $M_3(a, b)$ the common limit for a and b .

To show Theorem 1.3, the following identity by Ramanujan for the hypergeometric function $F(1/3, 2/3; 1; x)$ is extremely important.

LEMMA 3.1 (Ramanujan’s cubic transformation). For $0 < k \leq 1$

$$(3.2) \quad F\left(\frac{1}{3}, \frac{2}{3}; 1; 1 - k^3\right) = \frac{3}{1 + 2k} F\left(\frac{1}{3}, \frac{2}{3}; 1; \left(\frac{1 - k}{1 + 2k}\right)^3\right).$$

Proof. This identity has been proved by, for instance, the Borweins [6], Berndt et al. [4, Corollary 2.4] or [3, Corollary 2.4 and (2.25)], and Chan [12], though Ramanujan did not leave his proof. We will present an alternative proof with elementary calculation.

Since Proposition 2.8 yields

$$K_3(k) = \frac{\pi_3}{2} F\left(\frac{1}{3}, \frac{2}{3}; 1; k^3\right),$$

so that (3.2) is equivalent to

$$(3.3) \quad K_3(k') = \frac{3}{1 + 2k} K_3\left(\frac{1 - k}{1 + 2k}\right).$$

We have known from Proposition 2.2 that $K_3(k')$ satisfies

$$(3.4) \quad \frac{d}{dk} \left(k(k')^3 \frac{dy}{dk} \right) = 2k^2 y.$$

To show (3.3) we will verify that the function of right-hand side of (3.3) also satisfies (3.4). Now we let

$$f(k) = \frac{3}{1 + 2k} K_3\left(\frac{1 - k}{1 + 2k}\right).$$

Applying Proposition 2.1 we have

$$(3.5) \quad \frac{df(k)}{dk} = \frac{f(k)}{1 - k} - \frac{(1 + 2k)E_3(\ell)}{k(k')^3},$$

where $\ell = (1 - k)/(1 + 2k)$. Thus, differentiating both sides of

$$k(k')^3 \frac{df(k)}{dk} = (1 + k + k^2)kf(k) - (1 + 2k)E_3(\ell)$$

gives

$$(3.6) \quad \frac{d}{dk} \left(k(k')^3 \frac{df(k)}{dk} \right) = (1 + 2k + 3k^2)f(k) + (1 + k + k^2)k \frac{df(k)}{dk} - \left(2E_3(\ell) + (1 + 2k) \frac{dE_3(\ell)}{dk} \right).$$

Here, by Proposition 2.1

$$\frac{dE_3(\ell)}{d\ell} = \frac{f(k)}{1-k} - \frac{3E_3(\ell)}{(1-k)(1+2k)}.$$

Applying this and (3.5) to (3.6), we see that the right-hand side of (3.6) is equal to $2k^2f(k)$. This shows that $f(k)$ also satisfies (3.4) as $K_3(k')$ does.

The equation (3.4) has a regular singular point at $k=1$ and the roots of the associated indicial equation are both 0. Thus, it follows from the theory of ordinary differential equations (see for instance [33, §10.32]) that the functions $K_3(k')$ and $f(k)$, which agree at $k=1$, must be equal. This concludes the lemma. \square

PROPOSITION 3.2. *Let $0 \leq k < 1$, then*

- (i) $K_3(k) = \frac{1}{1+2k} K_3\left(\frac{\sqrt[3]{9(1+k+k^2)k}}{1+2k}\right),$
- (ii) $K_3(k) = \frac{3}{1+2k'} K_3\left(\frac{1-k'}{1+2k'}\right),$
- (iii) $E_3(k) = \frac{1+2k}{3} E_3\left(\frac{\sqrt[3]{9(1+k+k^2)k}}{1+2k}\right) + \frac{(1-k)(2+k)}{3} K_3(k),$
- (iv) $E_3(k) = (1+2k')E_3\left(\frac{1-k'}{1+2k'}\right) - k'(1+k')K_3(k),$

where $k' = \sqrt[3]{1-k^3}$.

Proof. It is obvious that (ii) is equivalent to (3.3), hence to Lemma 3.1. Therefore we will show (i), (iv) and (iii) in this order.

(i) Setting in (ii)

$$\frac{1-k'}{1+2k'} = \ell,$$

we get $0 \leq \ell < 1$ and

$$k' = \frac{1-\ell}{1+2\ell}, \quad k = \frac{\sqrt[3]{9(1+\ell+\ell^2)\ell}}{1+2\ell}.$$

Then (ii) is equivalent to

$$K_3\left(\frac{\sqrt[3]{9(1+\ell+\ell^2)\ell}}{1+2\ell}\right) = (1+2\ell)K_3(\ell).$$

Replacing ℓ by k , we obtain (i).

(iv) Let ℓ be the number above, then

$$\frac{d\ell}{dk} = \frac{3k^2}{(1+2k')^2(k')^2}.$$

It follows from (ii) that $(1 + 2k')K_3(k) = 3K_3(\ell)$. Differentiating both sides in k , we have

$$\begin{aligned} & -2\left(\frac{k}{k'}\right)^2 K_3(k) + \frac{1 + 2k'}{k(k')^3} E_3(k) - \frac{1 + 2k'}{k} K_3(k) \\ &= \frac{9k^2}{(1 + 2k')^2(k')^2} \cdot \frac{1}{\ell(\ell')^3} E_3(\ell) - \frac{9k^2}{(1 + 2k')^2(k')^2\ell} K_3(\ell). \end{aligned}$$

Applying

$$\ell = \frac{1 - k'}{1 + 2k'}, \quad \ell' = \frac{\sqrt[3]{9(1 + k' + (k')^2)k'}}{1 + 2k'}$$

and (ii), we see that the right-hand side is written as

$$\frac{(1 + 2k')^2}{k(k')^3} E_3(\ell) - \frac{3k^2}{(1 - k')(k')^2} K_3(k).$$

Thus we have

$$\frac{1 + 2k'}{k(k')^3} E_3(k) = \frac{(1 + 2k')^2}{k(k')^3} E_3(\ell) - \frac{(1 + 2k')(1 + k')}{k(k')^2} K_3(k).$$

Multiplying this by $k(k')^3/(1 + 2k')$, we obtain (iv).

(iii) It is obvious that (iv) can be written in ℓ , that is,

$$E_3\left(\frac{\sqrt[3]{9(1 + \ell + \ell^2)\ell}}{1 + 2\ell}\right) = \frac{3}{1 + 2\ell} E_3(\ell) - \frac{(1 - \ell)(2 + \ell)}{(1 + 2\ell)^2} K_3\left(\frac{\sqrt[3]{9(1 + \ell + \ell^2)\ell}}{1 + 2\ell}\right).$$

From (i) we have (iii). The proof is complete. □

Let us introduce auxiliary functions

$$\begin{aligned} I_p(a, b) &:= \int_0^{\pi_p/2} \frac{d\theta}{(a^p \cos_p^p \theta + b^p \sin_p^p \theta)^{1-1/p}}, \\ J_p(a, b) &:= \int_0^{\pi_p/2} (a^p \cos_p^p \theta + b^p \sin_p^p \theta)^{1/p} d\theta. \end{aligned}$$

It is easy to check that $K_p(k)$, $K'_p(k)$, $E_p(k)$ and $E'_p(k)$ are written as

$$\begin{aligned} K_p(k) &= I_p(1, k'), & K'_p(k) &= I_p(1, k), \\ E_p(k) &= J_p(1, k'), & E'_p(k) &= J_p(1, k). \end{aligned}$$

In the remainder of this section we assume $p = 3$. We will write $I_3(a, b)$ and $J_3(a, b)$ simply $I(a, b)$ and $J(a, b)$ respectively when no confusion can arise. The next lemma is crucial to show Theorem 1.2.

LEMMA 3.3. For $a \geq b > 0$,

$$aI(a, b) = \frac{a+2b}{3} I\left(\frac{a+2b}{3}, \sqrt[3]{\frac{(a^2+ab+b^2)b}{3}}\right).$$

Proof. From Proposition 3.2 (ii) (with k replaced by k') we get

$$\begin{aligned} aI(a, b) &= \frac{1}{a} K'_3\left(\frac{b}{a}\right) \\ &= \frac{3}{a+2b} K_3\left(\frac{a-b}{a+2b}\right) \\ &= \frac{3}{a+2b} I\left(1, \frac{\sqrt[3]{9(a^2+ab+b^2)b}}{a+2b}\right) \\ &= \frac{a+2b}{3} I\left(\frac{a+2b}{3}, \sqrt[3]{\frac{(a^2+ab+b^2)b}{3}}\right). \end{aligned}$$

This proves the lemma. □

Lemma 3.3 implies that $\{a_n I(a_n, b_n)\}$ is a constant sequence:

$$(3.7) \quad aI(a, b) = a_1 I(a_1, b_1) = a_2 I(a_2, b_2) = \cdots = a_n I(a_n, b_n) = \cdots.$$

Letting $n \rightarrow \infty$ in (3.7) we have

$$aI(a, b) = M_3(a, b) I(M_3(a, b), M_3(a, b)) = \frac{1}{M_3(a, b)} I(1, 1),$$

hence,

PROPOSITION 3.4. For $a \geq b > 0$

$$aI(a, b) = \frac{\pi_3}{2} \frac{1}{M_3(a, b)}.$$

From above, Theorem 1.2 immediately follows.

Proof of Theorem 1.2. Put $a = 1$ and $b = k'$ in Proposition 3.4. □

Let $I_n := I(a_n, b_n)$, $J_n := J(a_n, b_n)$, then

LEMMA 3.5. For $a \geq b > 0$

$$3J_{n+1} - J_n = a_n b_n (a_n + b_n) I_n, \quad n = 0, 1, 2, \dots$$

Proof. Set $\kappa_n := \sqrt[3]{1 - (b_n/a_n)^3}$. We see at once that

$$(3.8) \quad I_n = \frac{1}{a_n^2} K_3(\kappa_n), \quad J_n = a_n E_3(\kappa_n), \quad n = 0, 1, 2, \dots$$

Now, letting $k = \kappa_n$ in Proposition 3.2 (iv), we have

$$E_3(\kappa_n) = (1 + 2\kappa'_n) E_3\left(\frac{1 - \kappa'_n}{1 + 2\kappa'_n}\right) - \kappa'_n (1 + \kappa'_n) K_3(\kappa_n).$$

It is easily seen that $\kappa'_n = b_n/a_n$ and $\kappa_{n+1} = (1 - \kappa'_n)/(1 + 2\kappa'_n)$. Thus

$$E_3(\kappa_n) = \frac{a_n + 2b_n}{a_n} E_3(\kappa_{n+1}) - \frac{b_n}{a_n} \left(1 + \frac{b_n}{a_n}\right) K_3(\kappa_n).$$

Multiplying this by a_n and using $a_n + 2b_n = 3a_{n+1}$ we obtain

$$a_n E_3(\kappa_n) = 3a_{n+1} E_3(\kappa_{n+1}) - b_n \left(1 + \frac{b_n}{a_n}\right) K_3(\kappa_n).$$

From (3.8) we accomplished the proof. □

PROPOSITION 3.6. *Let $a \geq b > 0$, then*

$$J(a, b) = \left(a^3 - a \sum_{n=1}^{\infty} 3^n (a_n + c_n) c_n \right) I(a, b),$$

where $c_n := \sqrt[3]{a_n^3 - b_n^3}$.

Proof. We denote $I(a, b)$ and $J(a, b)$ briefly by I and J respectively. Lemma 3.3 gives $a_n I_n = aI$ for any n . By Lemma 3.5 and $c_{n+1} = (a_n - b_n)/3$, we obtain

$$\begin{aligned} 3(J_{n+1} - aa_{n+1}^2 I) - (J_n - aa_n^2 I) &= (ab_n(a_n + b_n) - 3aa_{n+1}^2 + aa_n^2) I \\ &= \frac{a}{3} (2a_n^2 - a_n b_n - b_n^2) I \\ &= \frac{a}{3} (2a_n + b_n)(a_n - b_n) I \\ &= 3a(a_{n+1} + c_{n+1})c_{n+1} I. \end{aligned}$$

Multiplying this by 3^n and summing both sides from $n = 0$ to $n = m - 1$, we obtain

$$(3.9) \quad 3^m (J_m - aa_m^2 I) - (J - a^3 I) = a \left(\sum_{n=1}^m 3^n (a_n + c_n) c_n \right) I.$$

On the other hand, since $aI = a_m I_m$, we have

$$\begin{aligned} 3^m(J_m - aa_m^2 I) &= 3^m \int_0^{\pi_3/2} \frac{a_m^3 \cos^3 \theta + b_m^3 \sin^3 \theta - a_m^3}{(a_m^3 \cos^3 \theta + b_m^3 \sin^3 \theta)^{2/3}} d\theta \\ &= 3^m c_m^3 \int_0^{\pi_3/2} \frac{-\sin^3 \theta}{(a_m^3 \cos^3 \theta + b_m^3 \sin^3 \theta)^{2/3}} d\theta. \end{aligned}$$

By (3.1) we get

$$0 \leq 3^m c_m^3 \leq \frac{1}{9m} (a - b)^3,$$

which means $\lim_{m \rightarrow \infty} 3^m(J_m - aa_m^2 I) = 0$. Therefore, as $m \rightarrow \infty$ in (3.9) the proposition follows. \square

Now we are in a position to show Theorem 1.3.

Proof of Theorem 1.3. Let $k = 1/\sqrt[3]{2}$ in Theorem 1.1, then

$$(3.10) \quad 2K_3\left(\frac{1}{\sqrt[3]{2}}\right)E_3\left(\frac{1}{\sqrt[3]{2}}\right) - K_3\left(\frac{1}{\sqrt[3]{2}}\right)^2 = \frac{\pi_3}{2}.$$

Letting $a = 1$ and $b = 1/\sqrt[3]{2}$ in Proposition 3.6 we get

$$E_3\left(\frac{1}{\sqrt[3]{2}}\right) = \left(1 - \sum_{n=1}^{\infty} 3^n (a_n + c_n) c_n\right) K_3\left(\frac{1}{\sqrt[3]{2}}\right),$$

where $c_n = \sqrt[3]{a_n^3 - b_n^3}$. Substituting this to (3.10), we have

$$\left(2\left(1 - \sum_{n=1}^{\infty} 3^n (a_n + c_n) c_n\right) - 1\right) K_3\left(\frac{1}{\sqrt[3]{2}}\right)^2 = \frac{\pi_3}{2}.$$

Finally, applying Theorem 1.2 with $k = 1/\sqrt[3]{2}$ to this, we obtain

$$\left(1 - 2 \sum_{n=1}^{\infty} 3^n (a_n + c_n) c_n\right) \frac{\pi_3^2}{4M_3\left(1, \frac{1}{\sqrt[3]{2}}\right)^2} = \frac{\pi_3}{2}.$$

This leads the result. \square

Remark 3.7. In Theorem 1.2, we proved the identity

$$K_3(k) = \frac{\pi_3}{2} \frac{1}{M_3(1, k')}.$$

From the fact and Corollary 2.4 we can formally deduce

$$K_3^*(k) = \frac{\pi_3}{2} \frac{1}{M_3(k', 1)},$$

where $M_3(a, b)$ for $0 < a \leq b$ is also defined by (1.9) in the same way as that for $a \geq b > 0$. This means

$$F\left(\frac{1}{3}, \frac{1}{3}; 1; 1 - k^3\right) = \sqrt[3]{\frac{3}{1 + k + k^2}} F\left(\frac{1}{3}, \frac{1}{3}; 1; \frac{(1 - k)^3}{9(1 + k + k^2)}\right),$$

which is reported in [31].

REFERENCES

- [1] G. D. ANDERSON, S. L. QIU, M. K. VAMANAMURTHY AND M. VUORINEN, Generalized elliptic integrals and modular equations, *Pacific J. Math.* **192** (2000), 1–37.
- [2] G. ANDREWS, R. ASKEY AND R. ROY, *Special functions, Encyclopedia of mathematics and its applications* **71**, Cambridge University Press, Cambridge, 1999.
- [3] B. C. BERNDT, *Ramanujan's notebooks, Part V*, Springer-Verlag, New York, 1998.
- [4] B. C. BERNDT, S. BHARGAVA AND F. G. GARVAN, Ramanujan's theories of elliptic functions to alternative bases, *Trans. Amer. Math. Soc.* **347** (1995), 4163–4244.
- [5] P. BINDING, L. BOULTON, J. ČEPIČKA, P. DRÁBEK AND P. GIRG, Basis properties of eigenfunctions of the p -Laplacian, *Proc. Amer. Math. Soc.* **134** (2006), 3487–3494.
- [6] J. M. BORWEIN AND P. B. BORWEIN, A cubic counterpart of Jacobi's identity and the AGM, *Trans. Amer. Math. Soc.* **323** (1991), 691–701.
- [7] J. M. BORWEIN AND P. B. BORWEIN, *Pi and the AGM, A study in analytic number theory and computational complexity*, reprint of the 1987 original, Canadian Mathematical Society series of monographs and advanced texts **4**, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1998.
- [8] R. P. BRENT, Fast multiple-precision evaluation of elementary functions, *J. Assoc. Comput. Mach.* **23** (1976), 242–251.
- [9] F. D. BURGONE, Generalized trigonometric functions, *Math. Comp.* **18** (1964), 314–316.
- [10] P. J. BUSHSELL AND D. E. EDMUNDS, Remarks on generalized trigonometric functions, *Rocky Mountain J. Math.* **42** (2012), 25–57.
- [11] L. BOULTON AND G. LORD, Approximation properties of the q -sine bases, *Proc. R. Soc. Lond., Ser. A. Math. Phys. Eng. Sci.* **467** (2011), 2690–2711.
- [12] H. H. CHAN, On Ramanujan's cubic transformation formula for ${}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; z\right)$, *Math. Proc. Cambridge Philos. Soc.* **124** (1998), 193–204.
- [13] M. DEL PINO, M. ELGUETA AND R. MANÁSEVICH, A homotopic deformation along p of a Leray-Schauder degree result and existence for $(|u'|^{p-2}u')' + f(t, u) = 0$, $u(0) = u(T) = 0$, $p > 1$, *J. Differential Equations* **80** (1989), 1–13.
- [14] O. DOŠLÝ AND P. ŘEHÁK, *Half-linear differential equations*, North-Holland mathematics studies **202**, Elsevier Science B.V., Amsterdam, 2005.
- [15] P. DRÁBEK AND R. MANÁSEVICH, On the closed solution to some nonhomogeneous eigenvalue problems with p -Laplacian, *Differential Integral Equations* **12** (1999), 773–788.
- [16] D. E. EDMUNDS, P. GURKA AND J. LANG, Properties of generalized trigonometric functions, *J. Approx. Theory* **164** (2012), 47–56.

- [17] D. E. EDMUNDS, P. GURKA AND J. LANG, Basis properties of generalized trigonometric functions, *J. Math. Anal. Appl.* **420** (2014), 1680–1692.
- [18] A. ELBERT, A half-linear second order differential equation, *Qualitative theory of differential equations*, I, II, Szeged, 1979, 153–180, *Colloq. Math. Soc. Janos Bolyai* **30**, North-Holland, Amsterdam-New York, 1981.
- [19] E. B. ELLIOTT, A formula including Legendre's $EK' + KE' - KK' = \frac{1}{2}\pi$, *Messenger Math.* **33** (1903/1904), 31–32.
- [20] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER AND F. G. TRICOMI, *Higher transcendental functions, I*, Robert E. Krieger Publishing Co., Inc., Melbourne, Fla., 1981. Based on notes left by Harry Bateman. With a preface by Mina Rees. With a foreword by E. C. Watson. Reprint of the 1953 original.
- [21] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER AND F. G. TRICOMI, *Higher transcendental functions, II*, Robert E. Krieger Publishing Co., Inc., Melbourne, Fla., 1981. Based on notes left by Harry Bateman. Reprint of the 1953 original.
- [22] J. LANG AND D. E. EDMUNDS, Eigenvalues, embeddings and generalised trigonometric functions, *Lecture notes in mathematics* **2016**, Springer, Heidelberg, 2011.
- [23] P. LINDQVIST, Some remarkable sine and cosine functions, *Ricerche Mat.* **44** (1995), 269–290.
- [24] P. LINDQVIST AND J. PEETRE, p -arclength of the q -circle, *The Mathematics Student* **72** (2003), 139–145.
- [25] P. LINDQVIST AND J. PEETRE, Comments on Erik Lundberg's 1879 thesis, Especially on the work of Göran Dillner and his influence on Lundberg, *Memorie dell'Institutio Lombardo (Classe di Scienze Matem. Nat.)* **31**, 2004.
- [26] Y. NAITO, Uniqueness of positive solutions of quasilinear differential equations, *Differential Integral Equations* **8** (1995), 1813–1822.
- [27] E. SALAMIN, Computation of π using arithmetic-geometric mean, *Math. Comp.* **30** (1976), 565–570.
- [28] D. SHELUPSKY, A generalization of the trigonometric functions, *Amer. Math. Monthly* **66** (1959), 879–884.
- [29] S. TAKEUCHI, Generalized Jacobian elliptic functions and their application to bifurcation problems associated with p -Laplacian, *J. Math. Anal. Appl.* **385** (2012), 24–35.
- [30] S. TAKEUCHI, The basis property of generalized Jacobian elliptic functions, *Commun. Pure Appl. Anal.* **13** (2014), 2675–2692.
- [31] M.-K. WANG, Y.-M. CHU AND Y.-P. JIANG, Ramanujan's cubic transformation and generalized modular equation, *Sci. China Math.* **58** (2015), 2387–2404.
- [32] K. WATANABE, Planar p -elastic curves and related generalized complete elliptic integrals, *Kodai Math. J.* **37** (2014), 453–474.
- [33] E. T. WHITTAKER AND G. N. WATSON, *A course of modern analysis, An introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions*, reprint of the fourth edition, 1927, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1996.

Shingo Takeuchi
 DEPARTMENT OF MATHEMATICAL SCIENCES
 SHIBAURA INSTITUTE OF TECHNOLOGY
 307 FUKASAKU, MINUMA-KU
 SAITAMA-SHI, SAITAMA 337-8570
 JAPAN
 E-mail: shingo@shibaura-it.ac.jp