



A new giant *Vellozia* (Velloziaceae) from Minas Gerais, Brazil with comments on the *V. compacta* complex and conservation

RUY JOSÉ VÁLKA ALVES^{1*}, ALESSANDRA RIBEIRO GUIMARÃES², CLAUDIA DE MORAES REZENDE³, LAURA DI SPIRITO BRAGA⁴ & NÍLBER GONÇALVES DA SILVA²

¹Departamento de Botânica, Museu Nacional, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista s. no., São Cristóvão, 20940-040, Rio de Janeiro, RJ, Brasil,

²Pós-Graduação em Ciências Biológicas (Botânica), Museu Nacional, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista s. no., São Cristóvão, 20940-040, Rio de Janeiro, RJ, Brasil

³Laboratório de Análise de Aromas, Departamento de Química Orgânica, Instituto de Química, Universidade Federal do Rio de Janeiro, 21945-970 Rio de Janeiro, RJ, Brasil,

⁴Graduação em Ciências Biológicas, Universidade Federal de Minas Gerais, Brasil

*author for correspondence, ruyvalka@mn.ufjr.br

Abstract

A new giant *Vellozia* species from the Ouro Branco range, Minas Gerais, is described and compared morphologically to the other dracenoid species. Leaf-anatomical characters and a unique suite of chemical constituents which help to separate the new species from *V. compacta*, sympatric in the range, are provided. Furthermore we discuss the circumscription of *V. compacta* and argue that this binomial is applied to a species complex in need of further taxonomic study.

Resumo

Uma nova espécie gigante de *Vellozia* da Serra do Ouro Branco, Minas Gerais é descrita e morfologicamente comparada às demais espécies dracenóides. São descritos caracteres de anatomia foliar e um conjunto único de constituintes químicos que ajudam a separar a espécie nova de *V. compacta* que é simpátrica na Serra. Adicionalmente discutimos a circunscrição de *V. compacta*, argumentando que esse binômio é aplicado a um complexo de espécies que necessita estudos taxonômicos mais detalhados.

Key words: Pandanales, endemism, Neotropics, monocotyledons, dracenoid habit

Introduction

The Velloziaceae are a relatively small, amphiatlantic family with about 250 species (Mello-Silva 2005). Despite many recent studies the delimitation of genera and many species of Velloziaceae remains controversial [e.g. *Nanuza* Smith & Ayensu (1976: 38): Alves (2002) vs. Mello–Silva *et al.* (2011)]. While some species of *Vellozia* can be easily distinguished by morphology, many require complementary leaf anatomic and chemical studies. Several species complexes occur in *Vellozia* which are in need of more thorough investigations (e.g. Barbosa *et al.* 2012, Mello–Silva 2010). One such complex is *V. compacta* Martius (1829: 293), a group of tall shrubby species dubbed “dracenoid” by Mello–Silva & Menezes (1999). The new species described herein based on morphology, anatomy and chemistry is one of the tallest in the genus and grows in the Ouro Branco mountain range which represents the southernmost limit of distribution of the *V. compacta* species complex (*viz.* Lousada *et al.* 2013).

Material and Methods

Plant material was repeatedly collected between 1992 and 2007 in the Ouro Branco range, Minas Gerais. Between 1989 and 2006, vegetation surveys were conducted annually by the first author, mostly with students, in several mountain

Acknowledgements

During this research, the team was supported by grants of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) to R.J.V. Alves, C.M. Rezende and L.D.S. Braga, and scholarships of the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) to A.R. Guimarães and N.G. Silva. We thank Prof. Nanuza Luiza de Menezes for discussing the anatomy during the elaboration of the manuscript and Ricardo Vieira and Martin Callmänder for their valuable reviews.

References

- Alves, R.J.V. (1994) Morphological age determination and longevity in some *Vellozia* populations in Brazil. *Folia Geobotanica et Phytotaxonomica* 29: 55–59.
- Alves, R.J.V. (2002) Two new species of *Nanuza* (Velloziaceae) from Brazil. *Novon* 12: 12–17.
<http://dx.doi.org/10.2307/3393230>
- Ayensu, E.S. (1974) Leaf anatomy and systematics of New World Velloziaceae. *Smithsonian Contributions to Botany* 15: 1–172.
<http://dx.doi.org/10.5479/si.0081024x.15>
- Barbosa, A.R., Fiorini, C.F., Silva-Pereira, V., Melo-Silva, R. & Borba, E.L. (2012) Geographical genetic structuring and phenotypic variation in the *Vellozia hirsuta* (Velloziaceae) ochlopecies complex. *American Journal of Botany* 99: 1477–1488.
<http://dx.doi.org/10.3732/ajb.1200070>
- Barnes, R.A., Pereira, A.L., Scofield, T.C.V., Braz-Filho, R. and Pinto, A.C. (1984) A new triterpene from *Vellozia compacta*. *Chemical and Pharmaceutical Bulletin* 32: 3674–3677.
<http://dx.doi.org/10.1248/cpb.32.3674>
- Bukatsch, F. (1972) Bemerkungen zur Doppelfärbung Astrablau-Safranin. *Mikrokosmos* 61: 255.
- Carmo, F.F. & Jacobi, C.M. (2013) A vegetação de canga no Quadrilátero Ferrífero, Minas Gerais: caracterização e contexto fitogeográfico. *Rodriguésia* 64: 527–541.
<http://dx.doi.org/10.1590/s2175-78602013000300005>
- Cronk, Q.C.B. (1998) The ochlopecies concept. In: Huxley, C.R. Lock, J.M. & Cutler, D.E (eds.) *Chorology, Taxonomy and Ecology of the Floras of Africa and Madagascar*. Royal Botanic Gardens, Kew, pp. 155–170.
- Giulietti, A.M. & Pirani, J.R. (1988) Patterns of geographic distribution of some plant species from the Espinhaço Range, Minas Gerais and Bahia, Brazil. In: Vanzolini, P.E. & Heyer, W. R. (eds) *Proceedings of a workshop on Neotropical distribution patterns*, Academia Brasileira de Ciências, Rio de Janeiro, pp. 39–69 .
- Giulietti, A.M., Pirani, J.R. & Harley, R.M. (1997) Espinhaço Range Region, Eastern Brazil. In: Davis, S.D., Heywood, V.H., Herrera-MacBryde, O., Villa-Lobos, J. & Hamilton, A.C. (eds.) *Centres of plant diversity – A Guide and Strategy for their Conservation*. World Wildlife Fund & World Conservation Union, Gland, pp. 397–404.
- Goethart, J.W.C. & Henrard, J.J.T. (1937) Tijdschrift voor de Systematiek en de Geografie der Planten. *Blumea* 2: 374.
- Gray, A. (1870) Botanical contributions 1. Notes on Labiatae. *Proceedings of the American Academy of Arts and Sciences* 8: 369.
- IUCN (2012) *IUCN Red List Categories and Criteria, Version 3.1*, 2nd edition. IUCN Species Survival Commission, Gland & Cambridge.
Available at: http://jr.iucnredlist.org/documents/redlist_cats_crit_en.pdf.
- Jacobi, C.M. & Carmo, F.F. (eds.) (2012) *Diversidade florística nas Cangas do Quadrilátero Ferrífero*. IDM, Belo Horizonte.
- Jacobi, C.M., Carmo, F.F., Vincent, R.C. & Stehmann, J.R. (2007) Plant communities on ironstone outcrops: a diverse and endangered Brazilian ecosystem. *Biodiversity and Conservation* 16: 2185–2200.
<http://dx.doi.org/10.1007/s10531-007-9156-8>
- Johansen, D.A. (1940) *Plant Microtechnique*. McGraw, Hill Book Company, New York, 523 pp.
- Linnaeus, C. (1767) *Systema naturæ per regna tria naturæ, secundum classes, ordines, genera, species, cum characteribus & differentiis*. Tomus II, Holmiæ. (Salvius), 735 pp.
<http://dx.doi.org/10.5962/bhl.title.7378>
- Lousada, J.M., Lovato, M.B. & Borba, E.L. (2013) High genetic divergence and low genetic variability in disjunct populations of the endemic *Vellozia compacta* (Velloziaceae) occurring in two edaphic environments of Brazilian campos rupestres. *Brazilian Journal of Botany* 36: 45–53.
<http://dx.doi.org/10.1007/s40415-013-0001-x>
- Martius, K.F.P. (1829) *Vellosia compacta*. In: Roemer, J.J. & Schultes, J. (eds.) *Systema Vegetabilium* ed. 15 bis, 7(1): 293.

- Martius, K.F.P. (1847) *Flora Brasiliensis* 3(1): 78.
- Mello-Silva, R. (2000) Partial cladistic analysis of *Vellozia* and characters for the phylogeny of Velloziaceae. In: Wilson, K.L. & Morrison, D.A. (eds.) *Monocots: Systematics and Evolution*. CSIRO, Melbourne, pp. 505–522.
- Mello-Silva, R. (2004) Novitates Velloziacearum florum phanerogamicarum Sancti Pauli. *Revista Brasileira de Botânica* 27: 453–462.
<http://dx.doi.org/10.1590/s0100-84042004000300006>
- Mello-Silva, R. (2005) Morphological analysis, phylogenies and classification in Velloziaceae. *Botanical Journal of the Linnean Society* 148: 157–173.
<http://dx.doi.org/10.1111/j.1095-8339.2005.00399.x>
- Mello-Silva, R. & Menezes, N.L. (1999) Two New Brazilian Velloziaceae, *Vellozia auriculata* and *Vellozia gigantea*, and a Key to the Related Dracenoid Species of *Vellozia*. *Novon* 9: 536–541.
<http://dx.doi.org/10.2307/3392159>
- Mello-Silva, R., Santos, D.Y.A.C., Salatino, M.L.F., Motta, L.B., Cattai, M.B., Sasaki, D., Lovo, J., Pita P.B., Rocini, C., Rodrigues, C.D.N., Zarrei, M. & Chase, M.W. (2011) Five vicarious genera from Gondwana: the Velloziaceae as shown by molecules and morphology. *Annals of Botany* 108: 87–102.
<http://dx.doi.org/10.1093/aob/mcr107>
- Mikan, J.C. (1822) *Delectus Florae et Faunae Brasiliensis* part 2: tab. 7.
<http://dx.doi.org/10.5962/bhl.title.63994>
- Morrison III, W.R., Lohr, J.L., Duchon, P., Wilches, R., Trujillo, D., Mair, M. & Renner, S.S. (2009) The impact of taxonomic change on conservation: Does it kill, can it save, or is it just irrelevant? *Biological Conservation* 142: 3201–3206.
<http://dx.doi.org/10.1016/j.biocon.2009.07.019>
- Pinto, A.C. (1980) Um novo diterpeno de *Vellozia compacta*. *Anais da Academia Brasileira de Ciências* 52: 473–475.
- Pinto, A.C. (1985). *Constituintes químicos de Vellozia compacta Martius & Schultes f.* Thesis. Instituto de Química, Universidade Federal do Rio de Janeiro.
- Pinto, A.C. & Borges, C. (1983) Six diterpenes from *Vellozia compacta*. *Phytochemistry* 22: 2011.
[http://dx.doi.org/10.1016/0031-9422\(83\)80034-x](http://dx.doi.org/10.1016/0031-9422(83)80034-x)
- Pinto, A.C., Silva, A.J.R., Mayer, L.M., & Braz F.R. (1979) Chemistry of South American Velloziaceae. Part VI: Compactone, a new diterpenoid from *Vellozia compacta*. *Phytochemistry* 18: 2036.
- Pinto, A.C., Epifanio, R. de, A. & Pizzolatti, M. (1992) Diterpenoids from *Vellozia declinans*. *Phytochemistry* 31: 4241–4243.
[http://dx.doi.org/10.1016/0031-9422\(92\)80451-j](http://dx.doi.org/10.1016/0031-9422(92)80451-j)
- Pinto, A.C., Furtado, V.L.R. & Gonzaga, L. (1990) Terpenoid constituents from *Vellozia epidendroides*. *Química Nova* 13: 64.
- Pinto, A.C., Antunes, O.A.C., Rezende, C.M. & Correia, C.R.D. (1997) Separation of Acidic Components of *Vellozia flavicans* by Silica Gel/Potassium Hydroxide Chromatography. *Phytochemical Analysis* 8: 14–17.
[http://dx.doi.org/10.1002/\(sici\)1099-1565\(199701\)8:1<14::aid-pca326>3.3.co;2-y](http://dx.doi.org/10.1002/(sici)1099-1565(199701)8:1<14::aid-pca326>3.3.co;2-y)
- Riehl, C.A.S. & Pinto, A.C. (2000) A cleistanthane diterpene lactone from *Vellozia compacta*. *Phytochemistry* 53: 917–919.
[http://dx.doi.org/10.1016/s0031-9422\(99\)00579-8](http://dx.doi.org/10.1016/s0031-9422(99)00579-8)
- Rodrigues, R.L. (2010) *Fungos endofíticos associados à Vellozia compacta Mart. ex Schult. F. (Velloziaceae) presente em afloramentos rochosos nos estados de Minas Gerais e Tocantins*. Dissertation, Universidade Federal de Ouro Preto. Available at: <http://www.biomias.ufop.br/biomias/dissertacoes/Rogério%20Leonardo%20Rodrigues.pdf>.
- Silva, G.C. da, Valente, L.M.M., Patitucci, M.L., Pinto, A.C. & Menezes, N.L. (2001) Diterpenóides com esqueleto cleistanano de *Vellozia* aff. *carunculares* Martius ex Seubert (Velloziaceae). *Química Nova* 24: 619–625.
<http://dx.doi.org/10.1590/s0100-40422001000500009>
- Smith, L.B. (1962) A synopsis of American Velloziaceae. *Contributions from the United States National Herbarium* 35: 251–294.
- Smith, L.B. & Ayensu, E.S. (1976) A revision of American Velloziaceae. *Smithsonian Contributions to Botany* 30: 1–172.
<http://dx.doi.org/10.5479/si.0081024x.30>
- SpeciesLink 2014. Available at: <http://www.splink.org.br/>.
- Spix, J.B. von & Martius, K.F.P. von (1824) *Travels in Brazil, in the years 1817-1820*. London: Longman, Hurst, Rees, Orme, Brown, and Green. Vol. 1.
- Touche, E.M.G., Lopez, E.G., Reyes, A.P., Sanchez, H., Honecker, F. & Achenbach, H. (1997) Parryin, a diterpene with a tricyclic 6-7-5-ring system from *Salvia parryi*. *Phytochemistry* 45: 387–390.
[http://dx.doi.org/10.1016/s0031-9422\(96\)00807-2](http://dx.doi.org/10.1016/s0031-9422(96)00807-2)
- Vincent, R.C. & Meguro, M. (2008) Influence of soil properties on the abundance of plant species in ferruginous rocky soils vegetation, southeastern Brazil. *Revista Brasileira de Botânica* 31: 377–388.
<http://dx.doi.org/10.1590/s0100-84042008000300002>