

A New Hilbert-type Inequality with the Integral in Whole Plane

ZITIAN XIE*

Department of Mathematics, Zhaoqing University, Zhaoqing, Guangdong 526061, China

e-mail: gdzqxzt@163.com

ZHENG ZENG

Department of Mathematics, Shaoguan University, Shaoguan, Guangdong 512005, China

e-mail: zz@sgu.edu.cn

ABSTRACT. In this paper, by estimating the weight function, we give a new Hilbert-type inequality with the integral in whole plane. As its applications, we consider the equivalent and a particular result.

1. Introduction

If $f(x), g(x) \geq 0$, such that $0 < \int_0^\infty f^2(x)dx < \infty$ and $0 < \int_0^\infty g^2(x)dx < \infty$, then [1]

$$(1.1) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(x)}{x+y} dx dy < \pi \left\{ \int_0^\infty f^2(x)dx \int_0^\infty g^2(x)dx \right\}^{1/2},$$

where the constant factor π is the best possible. Inequality (1.1) is well-known as Hilbert's integral inequality, which has been extended by Hardy-Riesz:

If $p > 1, \frac{1}{p} + \frac{1}{q} = 1, f(x), g(x) \geq 0$, such that $0 < \int_0^\infty f^p(x)dx < \infty$ and $0 < \int_0^\infty g^q(x)dx < \infty$, then we have the following Hardy-Hilbert's integral inequality:

$$(1.2) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \frac{\pi}{\sin(\pi/p)} \left\{ \int_0^\infty f^p(x)dx \right\}^{1/p} \left\{ \int_0^\infty g^q(x)dx \right\}^{1/q},$$

where the constant factor $\frac{\pi}{\sin(\pi/p)}$ also is the best possible.

* Corresponding Author.

Received February 1, 2011; accepted September 23, 2011.

2010 Mathematics Subject Classification: 26D15.

Key words and phrases: Hilbert-type integral inequality, weight function, Hölder's inequality, equivalent form.

Hilbert's inequality is important in analysis and its applications. It attracted some attention in the recent years. Actually, inequalities (1.1) and (1.2) have many generalizations and variants. (1.1) has been strengthened by Yang and others. (including double series inequalities) [2-18].

In 2008 Xie and Zitian Xie and Zheng Zeng gave a new Hilbert-type Inequality [2] as follows:

If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $f(x), g(x) \geq 0$ such that $0 < \int_0^\infty x^{-1-p/2} f^p(x) dx < \infty$ and $0 < \int_0^\infty x^{-1-q/2} g^q(x) dx < \infty$, then

$$(1.3) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(y)}{(x+a^2y)(x+b^2y)(x+c^2y)} dx dy < K \left\{ \int_0^\infty x^{-1-p/2} f^p(x) dx \right\}^{1/p} \left\{ \int_0^\infty x^{-1-q/2} g^q(x) dx \right\}^{1/q},$$

where the constant factor $K = \frac{\pi}{(a+b)(a+c)(b+c)}$ is the best possible.

In this paper, by obtaining the weight function, we give a new Hilbert-type inequality with the integral in whole plane.

In the following, we always suppose that $a = \cos \theta$, $\theta \in (0, \frac{\pi}{2})$; $p > 1$, $1/p + 1/q = 1$; $m + n = 2$, $m, n \in (0, 2)$.

2. Some lemmas

Lemma 2.1. If $k_1 := \int_0^\infty \frac{dt}{(t^2+2at+1)t^{1-n}}$, $k_2 := \int_0^\infty \frac{dt}{(t^2-2at+1)t^{1-n}}$, and $k := \int_{-\infty}^\infty \frac{dt}{(t^2+2at+1)|t|^{1-n}}$, then

$$(2.1) \quad k_1 = \begin{cases} \pi \csc \theta \csc(n\pi) \sin((1-n)\theta), & \text{if } n \neq 1, \\ \theta \csc \theta, & \text{if } n = 1 \end{cases}$$

$$k_2 = \begin{cases} \pi \csc \theta \csc(n\pi) \sin((1-n)(\pi-\theta)), & \text{if } n \neq 1, \\ (\pi-\theta) \csc \theta, & \text{if } n = 1 \end{cases}$$

and

$$(2.2) \quad k = k_1 + k_2 = \begin{cases} 2\pi \csc \theta \csc(n\pi) \sin \frac{(1-n)\pi}{2} \cos \frac{(1-n)(\pi-\theta)}{2}, & \text{if } n \neq 1, \\ \pi \csc \theta, & \text{if } n = 1, \end{cases}$$

$$= \begin{cases} \pi \csc \theta \sec \frac{n\pi}{2} \cos \frac{(1-n)(\pi-\theta)}{2}, & \text{if } n \neq 1, \\ \pi \csc \theta, & \text{if } n = 1 \end{cases}.$$

Proof. Let $f(z) = \frac{1}{(1+2bz+z^2)z^\alpha} = \frac{1}{(z-z_1)(z-z_2)z^\alpha}$, $\alpha \in (-1, 1)$, then

$$k^* := \int_0^\infty \frac{dt}{(t^2+2bt+1)t^\alpha} = \frac{2\pi i}{1-e^{-2\alpha\pi i}} [Res(f, z_1) + Res(f, z_2)];$$

if $b = \cos \vartheta$ ($0 < \vartheta < \pi$), and $z_1 = -e^{i\vartheta}$, $z_2 = -e^{-i\vartheta}$, then

$$k^* = \frac{2\pi i}{1-e^{-2\alpha\pi i}} \left[\frac{1}{(-2i \sin \vartheta)(-e^{i\vartheta})^\alpha} + \frac{1}{(2i \sin \vartheta)(-e^{-i\vartheta})^\alpha} \right] = \pi \csc \vartheta \csc(\alpha\pi) \sin(\alpha\vartheta),$$

setting $\alpha = 1 - n$, $\vartheta = \theta$ and $\vartheta = \pi - \theta$, we have (2.1).

On the other hand,

$$k = \int_0^\infty \frac{dt}{(t^2 + 2at + 1)t^{1-n}} + \int_{-\infty}^0 \frac{dt}{(t^2 + 2at + 1)(-t)^{1-n}} = k_1 + k_2$$

$$= \begin{cases} 2\pi \csc \theta \csc(n\pi) \sin \frac{(1-n)\pi}{2} \cos \frac{(1-n)(\pi-\theta)}{2}, & \text{if } n \neq 1, \\ \pi \csc \theta, & \text{if } n = 1. \end{cases}$$

We obtain (2.2). □

Lemma 2.2. Define the weight functions as follow:

$$w(x) := \int_{-\infty}^\infty \frac{|x|^m dy}{(x^2 + 2axy + y^2)|y|^{1-n}}, \quad \tilde{w}(y) := \int_{-\infty}^\infty \frac{|y|^n dx}{(x^2 + 2axy + y^2)|x|^{1-m}}.$$

Then

$$(2.3) \quad w(x) = \tilde{w}(y) = k.$$

Proof. We only prove that $w(x) = k$ for $x \in (-\infty, 0)$.

$$w(x) = \int_{-\infty}^0 \frac{|x|^m dy}{(x^2 + 2axy + y^2)|y|^{1-n}} + \int_0^\infty \frac{|x|^m dy}{(x^2 + 2axy + y^2)|y|^{1-n}} := w_1 + w_2,$$

setting $y = tx$, then $w_1 = \int_{-\infty}^0 \frac{(-x)^{2-n} dy}{(x^2 + 2axy + y^2)(-y)^{1-n}} = \int_0^\infty \frac{dt}{(t^2 + 2at + 1)t^{1-n}} = k_1$.

Similarly, setting $y = -tx$, $w_2 = \int_0^\infty \frac{(-x)^{2-n} dy}{(x^2 + 2axy + y^2)y^{1-n}} = \int_0^\infty \frac{dt}{(t^2 - 2at + 1)t^{1-n}} = k_2$, and $w(x) = k$.

Easily $\tilde{w}(x) = k$, using lemma 2.1, the lemma is proved. □

Lemma 2.3. For $\varepsilon > 0$, and $(n - \max\{\frac{2\varepsilon}{p}, \frac{2\varepsilon}{q}\}) \in (0, 2)$, define both functions, \tilde{f} and \tilde{g} , as follow:

$$(2.4) \quad \tilde{f}(x) = \begin{cases} x^{m-1-2\varepsilon/p}, & \text{if } x \in (1, \infty), \\ 0, & \text{if } x \in [-1, 1], \\ (-x)^{m-1-2\varepsilon/p}, & \text{if } x \in (-\infty, -1) \end{cases} ;$$

$$\tilde{g}(x) = \begin{cases} x^{n-1-2\varepsilon/q}, & \text{if } x \in (1, \infty), \\ 0, & \text{if } x \in [-1, 1], \\ (-x)^{n-1-2\varepsilon/q}, & \text{if } x \in (-\infty, -1) \end{cases} .$$

Then

$$(2.5) \quad I(\varepsilon) := \varepsilon \left\{ \int_{-\infty}^\infty |x|^{p(1-m)-1} \tilde{f}^p(x) dx \right\}^{1/p} \left\{ \int_{-\infty}^\infty |x|^{q(1-n)-1} \tilde{g}^q(x) dx \right\}^{1/q} = 1;$$

$$(2.6) \quad \tilde{I}(\varepsilon) := \varepsilon \int_{-\infty}^\infty \int_{-\infty}^\infty \frac{\tilde{f}(x)\tilde{g}(y)}{x^2 + 2axy + y^2} dx dy = k + o(1) \quad (\varepsilon \rightarrow 0^+).$$

Proof. Easily,

$$I(\varepsilon) = 2\varepsilon \left\{ \int_1^\infty x^{-1} x^{-2\varepsilon} dx \right\}^{1/p} \left\{ \int_1^\infty x^{-1} x^{-2\varepsilon} dx \right\}^{1/q} = 1.$$

Let $y = -Y$, using $\tilde{f}(-x) = \tilde{f}(x)$, $\tilde{g}(-x) = \tilde{g}(x)$, and

$$\tilde{f}(-x) \int_{-\infty}^\infty \frac{\tilde{g}(y) dy}{x^2 - 2axy + y^2} = \tilde{f}(x) \int_{-\infty}^\infty \frac{\tilde{g}(Y) dY}{x^2 + 2axY + Y^2}$$

we have that $\tilde{f}(x) \int_{-\infty}^\infty \frac{\tilde{g}(y) dy}{x^2 + 2axy + y^2}$ is an even function on x , then

$$\begin{aligned} \tilde{I}(\varepsilon) &= 2\varepsilon \int_0^\infty \tilde{f}(x) \left(\int_{-\infty}^\infty \frac{\tilde{g}(y)}{x^2 + 2axy + y^2} dy \right) dx \\ &= 2\varepsilon \left[\int_1^\infty x^{m-1-\frac{2\varepsilon}{p}} \left(\int_{-\infty}^{-1} \frac{(-y)^{n-1-\frac{2\varepsilon}{q}}}{x^2 + 2axy + y^2} dy \right) dx \right. \\ &\quad \left. + \int_1^\infty x^{m-1-\frac{2\varepsilon}{p}} \left(\int_1^\infty \frac{y^{n-1-\frac{2\varepsilon}{q}}}{x^2 + 2axy + y^2} dy \right) dx \right] \\ &:= I_1 + I_2. \end{aligned}$$

Setting $y = tx$ then

$$\begin{aligned} I_1 &= 2\varepsilon \left[\int_1^\infty x^{m-1-\frac{2\varepsilon}{p}} \left(\int_1^\infty \frac{y^{n-1-\frac{2\varepsilon}{q}}}{x^2 - 2axy + y^2} dy \right) dx \right] \\ &= 2\varepsilon \left[\int_1^\infty x^{-1-2\varepsilon} \left(\int_{\frac{1}{x}}^\infty \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt \right) dx \right] \\ &= 2\varepsilon \left[\int_1^\infty x^{-1-2\varepsilon} \left(\int_1^\infty \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt \right) dx \right. \\ &\quad \left. + \int_1^\infty x^{-1-2\varepsilon} \left(\int_{\frac{1}{x}}^1 \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt \right) dx \right] \\ &= \int_1^\infty \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt + 2\varepsilon \int_0^1 \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} \left(\int_{\frac{1}{t}}^\infty x^{-1-2\varepsilon} dx \right) dt \\ &= \int_1^\infty \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt + \int_0^1 \frac{t^{n-1+\frac{2\varepsilon}{p}}}{t^2 - 2at + 1} dt \\ &= \int_0^\infty \frac{t^{n-1-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} dt + \int_0^1 \frac{t^{\frac{2\varepsilon}{p}} - t^{-\frac{2\varepsilon}{q}}}{t^2 - 2at + 1} t^{n-1} dt \\ &= \pi \csc \theta \csc \left(\left(n - \frac{2\varepsilon}{q} \right) \pi \right) \sin \left[\left(1 - \left(n - \frac{2\varepsilon}{q} \right) \right) (\pi - \theta) \right] + \eta(\varepsilon). \end{aligned}$$

there $\lim_{\varepsilon \rightarrow 0^+} \eta(\varepsilon) = 0$, and we have $I_1 \rightarrow k_2$ ($\varepsilon \rightarrow 0^+$).

Similarly $I_2 \rightarrow k_1$ ($\varepsilon \rightarrow 0^+$). The lemma is proved. □

Lemma 2.4. *We have*

$$(2.7) \quad \begin{aligned} J &:= \int_{-\infty}^{\infty} |y|^{np-1} \left(\int_{-\infty}^{\infty} \frac{f(x)}{x^2 + 2axy + y^2} dx \right)^p dy \\ &\leq k^p \int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx \end{aligned}$$

Proof. By lemma 2.2, we find

$$\begin{aligned} &\left(\int_{-\infty}^{\infty} \frac{f(x)}{x^2 + 2axy + y^2} dx \right)^p \\ &= \left[\int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \left(\frac{|x|^{(1-m)/q}}{|y|^{(1-n)/p}} f(x) \right) \left(\frac{|y|^{(1-n)/p}}{|x|^{(1-m)/q}} \right) dx \right]^p \\ &\leq \int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \frac{|x|^{(1-m)(p-1)}}{|y|^{1-n}} f^p(x) dx \\ &\quad \left(\int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \frac{|y|^{(1-n)(q-1)}}{|x|^{1-m}} dx \right)^{p-1} \\ &= k^{p-1} |y|^{-np+1} \int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \frac{|x|^{(1-m)(p-1)}}{|y|^{1-n}} f^p(x) dx, \\ J &\leq k^{p-1} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \frac{|x|^{(1-m)(p-1)}}{|y|^{1-n}} f^p(x) dx \right] dy \\ &= k^{p-1} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{x^2 + 2axy + y^2} \frac{|x|^{(1-m)(p-1)}}{|y|^{1-n}} dy \right] f^p(x) dx \\ &= k^p \int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx. \end{aligned}$$

3. Main results

Theorem 3.1. *If both functions, $f(x)$ and $g(x)$ are nonnegative measurable functions, and satisfy*

$0 < \int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx < \infty$ and $0 < \int_{-\infty}^{\infty} |x|^{q(1-n)-1} g^q(x) dx < \infty$, then

$$(3.1) \quad \begin{aligned} I^* &:= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(x)g(y)}{x^2 + 2axy + y^2} dx dy \\ &< k \left(\int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-n)-1} g^q(x) dx \right)^{1/q}, \end{aligned}$$

and

$$(3.2) \quad \begin{aligned} J &= \int_{-\infty}^{\infty} |y|^{p(1-m)-1} \left(\int_{-\infty}^{\infty} \frac{f(x)}{x^2 + 2axy + y^2} dx \right)^p dy \\ &< k^p \int_{-\infty}^{\infty} |x|^{p(1-n)-1} f^p(x) dx. \end{aligned}$$

Inequalities (3.1) and (3.2) are equivalent, and where the constant factors k and k^p are the best possible.

Proof. If (2.7) takes the form of equality for some $y \in (-\infty, 0) \cup (0, \infty)$, then there exists constants M and N , such that they are not all zero, and

$$M \frac{|x|^{(1-m)(p-1)}}{|y|^{1-n}} f^p(x) = N \frac{|y|^{(1-n)(q-1)}}{|x|^{1-m}} \quad \text{a.e. in } (-\infty, \infty),$$

Hence, there exists a constant C , such that

$$M|x|^{(1-m)p} f^p(x) = N|y|^{(1-n)q} = C \quad \text{a.e. in } (-\infty, \infty).$$

We claim that $M = 0$. In fact, if $M \neq 0$, then $|x|^{p(1-m)-1} f^p(x) = \frac{C}{M|x|}$ a.e. in $(-\infty, \infty)$ which contradicts the fact that $0 < \int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx < \infty$. In the same way, we claim that $N = 0$. This is too a contradiction and hence by (2.7), we have (3.2).

By Hölder's inequality with weight and (3.2), we have,

$$(3.3) \quad \begin{aligned} I^* &= \int_{-\infty}^{\infty} \left[|y|^{n-1+\frac{1}{q}} \int_{-\infty}^{\infty} \frac{f(x)}{x^2 + 2axy + y^2} dx \right] \left[|y|^{1-n-\frac{1}{q}} g(y) \right] dy \\ &\leq (J)^{\frac{1}{p}} \left(\int_{-\infty}^{\infty} |y|^{q(1-n)-1} g^q(y) dy \right)^{1/q}. \end{aligned}$$

Using (3.2), we have (3.1).

Setting $g(y) = |y|^{pn-1} \left(\int_{-\infty}^{\infty} \frac{f(x)}{x^2 + 2axy + y^2} dx \right)^{p-1}$, then $J = \int_{-\infty}^{\infty} |y|^{q(1-n)-1} g^q(y) dy$, by (2.7) we have $J < \infty$. If $J = 0$ then (3.2) is proved; if $0 < J < \infty$, by (3.1), we obtain

$$\begin{aligned} 0 &< \int_{-\infty}^{\infty} |y|^{q(1-n)-1} g^q(y) dy = J = I^* \\ &< k \left(\int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-n)-1} g^q(x) dx \right)^{1/q}, \\ \left(\int_{-\infty}^{\infty} |x|^{q(1-m)-1} g^q(x) dx \right)^{1/p} &= J^{1/p} < k \left(\int_{-\infty}^{\infty} |x|^{p(1-n)-1} f^p(x) dx \right)^{1/p}. \end{aligned}$$

Inequalities (3.1) and (3.2) are equivalent.

If the constant factor k in (3.1) is not the best possible, then there exists a positive h (with $h < k$), such that

$$(3.4) \quad \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(x)g(y)}{x^2 + 2axy + y^2} dx dy < h \left(\int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-n)-1} g^q(x) dx \right)^{1/q}.$$

For $\varepsilon > 0$, by (3.4), using lemma 2.3, we have

$$k + o(1) < \varepsilon h \left(\int_{-\infty}^{\infty} |x|^{-1} \tilde{f}^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{-1} \tilde{g}^q(x) dx \right)^{1/q} = h.$$

Hence we find, $k + o(1) < h$. For $\varepsilon \rightarrow 0^+$, it follows that $k \leq h$, which contradicts the fact that $h < k$. Hence the constant k in (3.1) is the best possible.

Thus we complete the prove of the theorem. □

Remark. For $a = \frac{1}{2}$, $(\theta = \frac{\pi}{3}), n \neq 1$, in (3.1), we have the following particular result:

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(x)g(y)}{x^2 + xy + y^2} dx dy < \frac{2}{\sqrt{3}} \pi \sec \frac{n\pi}{2} \cos \frac{(1-n)\pi}{3} \left(\int_{-\infty}^{\infty} |x|^{p(1-m)-1} f^p(x) dx \right)^{\frac{1}{p}} \left(\int_{-\infty}^{\infty} |x|^{q(1-n)-1} g^q(x) dx \right)^{\frac{1}{q}}.$$

References

- [1] Hardy G. H., Littlewood J. E. and Pólya G., *Inequalities*, Cambridge University Press, Cambridge, 1952.
- [2] Zitian Xie and Zeng Zheng, *A Hilbert-type integral inequality whose kernel is a homogeneous form of degree -3* , J. Math. Appl., **339**(2008), 324-331.
- [3] Bicheng Yang, *On the the norm of an intergal operator and applications*, J. Math. Anal. Appl., **321**(2006), 182-192.
- [4] Zitian Xie and Zheng Zeng, *A Hilbert-type integral inequality with a non-homogeneous form and a best constant factor*, Advances and applications in mathematical sciences, **3**(1)(2010), 61-71.
- [5] Xie Zitian, *A New Hilbert-type integral inequality with the homogeneous kernel of real number-degree*, Journal of Jishou University(Natturnal Science Edition), **32**(4)(2011), 26-30.
- [6] Bicheng Yang, *A Hilbert-type with a mixed kemel and extensions*, Journal of Sichuan Normal University(Natural Science), **31**(3)(2008), 281-284.

- [7] Zheng Zeng and Zitian Xie, *A new Hilbert-type integral inequality with a best constant factor*, Journal of South China Normal University (Natural Science Edition), **3**(2010), 31-33
- [8] Zheng Zeng and Zitian Xie, *On a new Hilbert-type integral inequality with the the integral in whole plane*, Journal of Inequalities and Applications vol. 2010, Article ID 256796, 8 pages, 2010. doi:10.1155/2010/256796.
- [9] Zitian Xie, Bicheng Yang, Zheng Zeng, *A New Hilbert-type integral inequality with the homogeneous kernel of real number-degree*, Journal of Jilin University(Science Edition), **48**(6)(2010), 941-945.
- [10] Zitian Xie and Benlu Fu, *A new Hilbert-type integral inequality with the best constant factor*, J. Wuhan Univ. (Nat. Sci. Ed), **55**(6)(2009), 637-640.
- [11] Dongmei Xin, *On a Hilbert-type integral inequality*, Kyungpook Mathematical Journal, **49**(2009), 393-401.
- [12] Zitian Xie and Xingdong Liu, *A new Hilbert-type integral inequality and its reverse*, Journal of Henan University (Science Edition), **39**(1)(2009), 10-13.
- [13] Zheng Zeng and Zitian Xie, *A new Hilbert-type integral inequality with a best constant factor*, Journal of South China Normal University (Natural Science Edition), **3**(2010), 31-33.
- [14] Zitian Xie and Bicheng Yang, *A new Hilbert-type inequality with some parameters and its reverse*, Kyungpook Mathematical Journal, **48**(1)(2008), 93-100.
- [15] Xie Zitian, Zeng Zheng, *A new Hilberttype integral inequality with the homogeneous kernel of degree 0*, Journal of Southwest University (Natturnal Science Edition), **23**(8)(2011), 137-141.
- [16] Xie Zitian, Zeng Zheng, *A Hilbert-type integral inequality with the homogeneous kernel of real number-degree and its equivalent*, Journal of Zhejiang University (Science Edition).
- [17] Bicheng Yang, *On a relation to Hilbert's integral inequality and a Hilbert-type inequality*, Kyungpook Mathematical Journal, **49**(2009), 563-572.
- [18] Zitian Xie, Zheng Zeng, *The Hilbert-type integral inequality with the system kernel of $-\lambda$ degree homogeneous form*, Kyungpook Mathematical Journal, **50**(2)(2010), 297-306.