



# A new numerical technique for local fractional diffusion equation in fractal heat transfer

Xiao-Jun Yang<sup>a,b</sup>, J. A. Tenreiro Machado<sup>c</sup>, Dumitru Baleanu<sup>d,e</sup>, Feng Gao<sup>a,b,\*</sup>

<sup>a</sup>School of Mechanics and Civil Engineering, China University of Mining and Technology, Xuzhou 221116, China.

<sup>b</sup>State Key Laboratory for Geomechanics and Deep Underground Engineering, School of Mechanics and Civil Engineering, China University of Mining and Technology, Xuzhou 221116, China.

<sup>c</sup>Institute of Engineering, Polytechnic of Porto, Department of Electrical Engineering, Rua Dr. Antonio Bernardino de Almeida, 4249-015 Porto, Portugal.

<sup>d</sup>Department of Mathematics, Cankaya University, Ogretmenler Cad. 14, Balgat-06530, Ankara Turkey.

<sup>e</sup>Institute of Space Sciences, Magurele-Bucharest, Romania.

Communicated by A. Atangana

---

## Abstract

In this paper, a new numerical approach, embedding the differential transform (DT) and Laplace transform (LT), is firstly proposed. It is considered in the local fractional derivative operator for obtaining the non-differential solution for diffusion equation in fractal heat transfer. ©2016 All rights reserved.

*Keywords:* Numerical solution, diffusion equation, differential transform, Laplace transform, fractal heat transfer, local fractional derivative.

*2010 MSC:* 76R50, 26A33, 44A10, 28A80.

---

## 1. Introduction

Fractional calculus and its generalizations started to be considered by many researchers as an optimal tool to describe the dynamics of complex systems [3, 4, 11, 16, 24, 26, 30]. Ordinary differential equations (ODEs) and partial differential equations (PDEs) within the derivative of non-integer orders were used to describe complex phenomena in the time-space sense of differentiability [1, 12, 23, 27, 28, 33] and non-differentiability [7, 17, 37, 38]. Several approaches were developed to find the numerical and analytical solutions for fractional ordinary differential equations (FODEs) and fractional partial differential equations

---

\*Corresponding author

Email address: [jsppw@sohu.com](mailto:jsppw@sohu.com) (Feng Gao)

(FPDEs), namely the methods of Mellin and Laplace transforms [5], implicit local radial basis function (ILRBF) [19], finite difference (FD) [32], homotopy perturbation (HP)[18], homotopy analysis (HA) [10], heat-balance integral (HBI) [15], variational iteration (VI) [14], generalized differential transform (GDT) [29], and others [9, 22, 25, 31]. There are distinct strategies for solving the local fractional partial differential equations [13, 20, 34, 35], such as the local fractional perspectives to differential transform (DT) [8], variation iteration method (VIM) [36], Fourier transform (FT) [39], Laplace transform (LT) [40], Laplace variation iteration method (LVIM) [21], and functional method (FM) [6].

In this paper, we consider the diffusion equation in fractal heat transfer with local fractional derivative [34, 35, 37, 38]

$$\frac{\partial^\alpha \Omega(x, t)}{\partial t^\alpha} - \frac{\partial^{2\alpha} \Omega(x, t)}{\partial x^{2\alpha}} = 0, \quad (1.1)$$

where the local fractional derivative partial derivative operator is defined as [8, 34]

$$\frac{\partial^\alpha \Omega(x, y)}{\partial x^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha (\Omega(x, y) - \Omega(x_0, y))}{(x - x_0)^\alpha},$$

with

$$\Delta^\alpha (\Omega(x, y) - \Omega(x_0, y)) \cong \Gamma(1 + \alpha) \Delta (\Omega(x, y) - \Omega(x_0, y)).$$

The classical Laplace differential transform was presented in [2]. However, the Laplace differential transform via local fractional differential operator is not addressed. The aim of the manuscript is to point out the local fractional Laplace differential transform to solve the diffusion equation in fractal heat transfer with local fractional derivative.

The structure of the manuscript is as follows. In Section 2, we introduce the fundamental theory of Laplace and differential transforms via local fractional derivative. In Section 3, we analyze the Laplace differential transform method. In Section 4, we give the numerical solution for diffusion equation in fractal heat transfer. In Section 5, we discuss the error analysis. Finally, in Section 6 we outlines the main conclusions.

## 2. Fundamental tools

The local fractional derivative of  $\Omega(x)$  is defined as follows [34–36, 40]:

$$\frac{\partial^\alpha \Omega(x)}{\partial x^\alpha} \Big|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Delta^\alpha (\Omega(x) - \Omega(x_0))}{(x - x_0)^\alpha},$$

where

$$\Delta^\alpha (\Omega(x) - \Omega(x_0)) \cong \Gamma(1 + \alpha) \Delta (\Omega(x) - \Omega(x_0)).$$

The local fractional Laplace transform of  $\Omega(\tau)$  can be formulated as follows [21, 36]:

$$\tilde{Y}_\alpha \{\Omega(\tau)\} = \Omega_y^{\tilde{Y}, \alpha}(y) = \frac{1}{\Gamma(1 + \alpha)} \int_0^\infty E_\alpha(-y^\alpha \tau^\alpha) \Omega(\tau) (d\tau)^\alpha, \quad 0 < \alpha \leq 1,$$

where the local fractional integral operator of  $\theta(x)$  is defined as follows [6, 21, 34, 36, 40]:

$${}_a I_b^{(\alpha)} \theta(\tau) = \frac{1}{\Gamma(1 + \alpha)} \int_a^b \theta(\tau) (d\tau)^\alpha = \frac{1}{\Gamma(1 + \alpha)} \lim_{\Delta\tau \rightarrow 0} \sum_{j=0}^{N-1} \theta(\tau) (\Delta\tau)^\alpha,$$

where  $\Delta\tau = t_{j+1} - t_j$ ,  $j = 0, \dots, N - 1$ ,  $t_0 = a$ , and  $t_N = b$ .

The inverse local fractional Laplace transform is defined as [6, 21, 36]:

$$\Omega(\tau) = \tilde{Y}_\alpha^{-1} \left\{ \Omega_y^{\tilde{Y}, \alpha}(y) \right\} = \frac{1}{(2\pi)^\alpha} \int_{\beta - i\infty}^{\beta + i\infty} E_\alpha(y^\alpha \tau^\alpha) \Omega_y^{\tilde{Y}, \alpha}(y) (dy)^\alpha,$$

where  $y^\alpha = \beta^\alpha + i^\alpha \infty^\alpha$  and  $Re(y^\alpha) = \beta^\alpha$ .

The properties applied in this paper are given below [21, 36]:

$$\begin{aligned} \tilde{Y}_\alpha \{a\Omega_1(\tau) + b\Omega_2(\tau)\} &= a\tilde{Y}_\alpha \{\Omega_1(\tau)\} + b\tilde{Y}_\alpha \{\Omega_2(\tau)\}, \\ \tilde{Y}_\alpha \{\Omega^{(n\alpha)}(\tau)\} &= y^{n\alpha} \tilde{Y}_\alpha \{\Omega(\tau)\} - \sum_{k=1}^n y^{(k-1)\alpha} \Omega^{(n-k)\alpha}(0), \\ \tilde{Y}_\alpha \{E_\alpha(x^\alpha)\} &= \frac{1}{y^\alpha - 1}, \\ \tilde{Y}_\alpha \left\{ \frac{\tau^{k\alpha}}{\Gamma(1+k\alpha)} \right\} &= \frac{1}{y^{(k+1)\alpha}}. \end{aligned}$$

If  $g(\tau)$  is a local fractional analytic function [8] in the domain  $\Omega$ , then the differential transform (the revised differential transform) via local fractional derivative becomes

$$G(\kappa) = \frac{1}{\Gamma(1+\kappa\alpha)} \left. \frac{d^{\kappa\alpha} g(\tau)}{d\tau^{\kappa\alpha}} \right|_{\tau=\tau_0}, \quad \tau, \tau_0 \in \Omega,$$

where  $\kappa = 0, 1, \dots, n, 0 < \alpha \leq 1$ , and the inverse transform of  $G(\kappa)$  in the domain  $\Omega$  via local fractional derivative is given by:

$$g(\tau) = \sum_{\kappa=0}^{\infty} G(\kappa) (\tau - \tau_0)^{\kappa\alpha}, \quad \kappa \in K. \tag{2.1}$$

The function  $G(\kappa)$  is the non-differentiable spectrum of  $g(\tau)$ . The properties of differential transform via local fractional derivative are listed in Table 1.

Table 1: The fundamental operations of the local fractional DTM (see[8]).

Original functions	Transformed functions
$s(\tau) = g(\tau) + h(\tau)$	$S(\kappa) = G(\kappa) + H(\kappa)$
$s(\tau) = ag(\tau)$	$S(\kappa) = aG(\kappa), \quad a \text{ is a constant}$
$s(\tau) = g(\tau)h(\tau)$	$S(\kappa) = \sum_{l=0}^{\kappa} G(l)H(\kappa-l)$
$s(\tau) = E_\alpha(\tau^\alpha)$	$S(\kappa) = \frac{1}{\Gamma(1+\kappa\alpha)}$
$s(\tau) = D^{n\alpha}g(\tau)$	$S(\kappa) = \frac{\Gamma(1+(\kappa+n)\alpha)}{\Gamma(1+\kappa\alpha)} G(\kappa+n)$

### 3. The method of the Laplace differential transform via local fractional derivative

In order to present the Laplace differential transform method via local fractional derivative, we consider the initial-boundary conditions of Eq. (1.1) as

$$\begin{aligned} \Omega(0, t) &= \varphi_1(t), \\ \frac{\partial^\alpha}{\partial x^\alpha} \Omega(0, t) &= \varphi_2(t), \\ \Omega(x, 0) &= \phi(x). \end{aligned}$$

We take the local fractional Laplace transform of Eq. (1.1) with respect to  $t$ , namely,

$$y^\alpha \Omega(x, y) - \phi(x) - \frac{\partial^{2\alpha} \Omega(x, y)}{\partial x^{2\alpha}} = 0,$$

and the initial conditions are presented as follows:

$$\begin{aligned} \frac{\partial^\alpha}{\partial x^\alpha} \Omega(0, y) &= \varphi_2(y), \\ \Omega(0, y) &= \varphi_1(y). \end{aligned} \tag{3.1}$$

Using the LFDT to Eq. (1.1) with respect to  $x$ , we have

$$\Omega(x, y) = \sum_{\kappa=0}^{\infty} \Omega(\kappa, y)x^{\kappa\alpha}, \quad \kappa \in K,$$

so that

$$y^\alpha \Omega(\kappa, y) - \phi(\kappa) - \frac{\Gamma(1 + (\kappa + 2)\alpha)}{\Gamma(1 + \kappa\alpha)} \Omega(\kappa + 2, y) = 0.$$

Therefore, after taking the inverse local fractional Laplace transform, we obtain

$$\Omega(x, t) = \tilde{Y}_\alpha^{-1} \{ \Omega(x, y) \} = \tilde{Y}_\alpha^{-1} \left\{ \sum_{\kappa=0}^{\infty} \Omega(\kappa, y)x^{\kappa\alpha} \right\} = \sum_{\kappa=0}^{\infty} \Omega(\kappa, t)x^{\kappa\alpha},$$

which, using the inverse local fractional Laplace transform of Eq. (2.1), leads to the solution of Eq. (1.1).

#### 4. Application to diffusion equation in fractal heat transfer

In this section we consider a typical application to diffusion equation in fractal heat transfer. We now suggest

$$\varphi_1(t) = E_\alpha(t^\alpha), \quad \varphi_2(t) = E_\alpha(t^\alpha), \quad \phi(x) = E_\alpha(x^\alpha).$$

From (3.1), we obtain

$$y^\alpha \Omega(\kappa, y) - \frac{1}{\Gamma(1 + \kappa\alpha)} - \frac{\Gamma(1 + (\kappa + 2)\alpha)}{\Gamma(1 + \kappa\alpha)} \Omega(\kappa + 2, y) = 0, \quad \kappa = 0, 1, 2, \dots, \tag{4.1}$$

where

$$\frac{\partial^\alpha}{\partial x^\alpha} \Omega(0, y) = \frac{1}{y^\alpha - 1}, \quad \Omega(0, y) = \frac{1}{y^\alpha - 1}. \tag{4.2}$$

Submitting Eq. (4.2) into the iterative relationship (4.1) leads to the simulation results for the components  $\Omega(\kappa, y)$  listed in Table 2.

Table 2: The simulation values for the components  $\Omega(\kappa, y)$ .

$\Omega(0, y)$	$\Omega(1, y)$	$\Omega(2, y)$	$\Omega(3, y)$	$\Omega(4, y)$
$\frac{1}{y^\alpha - 1}$	$\frac{1}{y^\alpha - 1} \frac{1}{\Gamma(1 + \alpha)}$	$\frac{1}{y^\alpha - 1} \frac{1}{\Gamma(1 + 2\alpha)}$	$\frac{1}{y^\alpha - 1} \frac{1}{\Gamma(1 + 3\alpha)}$	$\frac{1}{y^\alpha - 1} \frac{1}{\Gamma(1 + 4\alpha)}$

Therefore, we obtain

$$\Omega(\kappa, y) = \sum_{\kappa=0}^4 \frac{1}{y^\alpha - 1} \frac{1}{\Gamma(1 + \kappa\alpha)}.$$

Making use of the inverse local fractional Laplace transform, it yields:

$$\Omega(\kappa, t) = E_\alpha(t^\alpha) \sum_{\kappa=0}^4 \frac{1}{\Gamma(1 + \kappa\alpha)}. \tag{4.3}$$

Adopting the inverse local fractional Laplace transform to Eq. (4.3), the simulation value of the solution becomes:

$$\Omega(\kappa, t) = E_\alpha(t^\alpha) \sum_{\kappa=0}^4 \frac{x^{\kappa\alpha}}{\Gamma(1 + \kappa\alpha)}. \tag{4.4}$$

The plot of Eq. (4.4) is shown in Figure 1 when  $\alpha = \ln 2 / \ln 3$ .




Figure 1: The simulation value of Eq. (4.4) when  $\alpha = \ln 2 / \ln 3$ .

### 5. Error analysis

In this section we present the error analysis of the local fractional Laplace differential transform. Taking the successively iterative process, we have

$$\Omega(x, t) = E_\alpha(t^\alpha) \lim_{n \rightarrow \infty} \sum_{\kappa=0}^n \frac{x^{\kappa\alpha}}{\Gamma(1 + \kappa\alpha)}, \tag{5.1}$$

so that the exact solution of Eq. (1.1) takes the form:

$$\Omega^*(x, t) = E_\alpha(t^\alpha) \lim_{n \rightarrow \infty} \sum_{\kappa=0}^n \frac{x^{\kappa\alpha}}{\Gamma(1 + \kappa\alpha)} = E_\alpha(t^\alpha) E_\alpha(x^\alpha). \tag{5.2}$$

From Eqs. (5.1) and (5.2) the error value can be written as:

$$\mathfrak{R}(x, t) = |\Omega^*(x, t) - \Omega(x, t)| = E_\alpha(t^\alpha) \sum_{\kappa=n+1}^{\infty} \frac{x^{\kappa\alpha}}{\Gamma(1 + \kappa\alpha)}.$$

The comparison between the numerical solution when  $n = 4$  and exact solution of Eq. (1.1) is displayed in Figure 2. The components of the term  $\tilde{Y}_\alpha^{-1}\{\Omega(\kappa, y)\}$ , when  $\kappa = 0, 1, 2, 3, 4$ , are listed in Table 3. The plots of the components of the term  $\tilde{Y}_\alpha^{-1}\{\Omega(\kappa, y)\}$ , when  $\kappa = 0, 1, 2, 3, 4$ , are illustrated in Figure 3. The numerical solutions of Eq. (1.1), when  $n = 0, 1, 2, 3, 4$ , are listed in Table 4. The plots of numerical solutions of Eq. (1.1), when  $n = 0, 1, 2, 3, 4$ , are demonstrated in Figure 4.

Table 3: The components of the term  $\tilde{Y}_\alpha^{-1}\{\Omega(\kappa, y)\}$  when  $\kappa = 0, 1, 2, 3, 4$ .

$\kappa = 0$	$\kappa = 1$	$\kappa = 2$	$\kappa = 3$	$\kappa = 4$
$E_\delta(t^\delta)$	$\frac{E_\alpha(t^\alpha)x^\alpha}{\Gamma(1+\alpha)}$	$\frac{E_\alpha(t^\alpha)x^{2\alpha}}{\Gamma(1+2\alpha)}$	$\frac{E_\alpha(t^\alpha)x^{3\alpha}}{\Gamma(1+3\alpha)}$	$\frac{E_\alpha(t^\alpha)x^{4\alpha}}{\Gamma(1+4\alpha)}$

Table 4: The numerical solutions of Eq.(1.1) when  $n = 0, 1, 2, 3, 4$ .

$n = 0$	$n = 1$	$n = 2$	$n = 3$	$n = 4$
$E_\delta(t^\delta)$	$\sum_{\kappa=0}^1 \frac{E_\alpha(t^\alpha)x^{\kappa\alpha}}{\Gamma(1+\kappa\alpha)}$	$\sum_{\kappa=0}^2 \frac{E_\alpha(t^\alpha)x^{\kappa\alpha}}{\Gamma(1+\kappa\alpha)}$	$\sum_{\kappa=0}^3 \frac{E_\alpha(t^\alpha)x^{\kappa\alpha}}{\Gamma(1+\kappa\alpha)}$	$\sum_{\kappa=0}^4 \frac{E_\alpha(t^\alpha)x^{\kappa\alpha}}{\Gamma(1+\kappa\alpha)}$




Figure 2: The comparison between numerical solution and exact solution of Eq. (1.1).




Figure 3: The components of the term  $\tilde{Y}_\alpha^{-1}\{\Omega(\kappa, y)\}$  when  $\kappa = 0, 1, 2, 3, 4$ .




Figure 4: The numerical solutions of Eq.(1.1) when  $n = 0, 1, 2, 3, 4$ .

## 6. Conclusions

The local fractional Laplace differential transform was firstly proposed based upon local fractional calculus (LFC). We adopted the method to solve the diffusion equation in fractal heat transfer in the presence of the local fractional derivative. The obtained result strongly illustrates that the method is easy, simple, efficient and accurate for a class of local fractional partial differential equations.

## Acknowledgment

This work is supported by the State Key Research Development Program of China (Grant No. 2016YFC0600705).

## References

- [1] R. Almeida, A. B. Malinowska, D. F. M. Torres, *A fractional calculus of variations for multiple integrals with application to vibrating string*, J. Math. Phys., **51** (2010), 12 pages. 1
- [2] M. Alquran, K. Al-Khaled, M. Ali, A. Ta'any, *The combined Laplace transform-differential transform method for solving linear non-homogeneous PDEs*, J. Math. Comput. Sci., **2** (2012), 690–701. 1

- [3] D. Baleanu, K. Diethelm, E. Scalas, J. J. Trujillo, *Models and numerical methods*, World Sci., **3** (2012), 10–16. 1
- [4] D. Baleanu, J. A. T. Machado, A. C. J. Luo, *Fractional dynamics and control*, Springer, Berlin, (2012). 1
- [5] A. H. Bhrawy, E. H. Doha, D. Baleanu, S. S. Ezz-Eldien, *A spectral tau algorithm based on Jacobi operational matrix for numerical solution of time fractional diffusion-wave equations*, J. Comput. Phys., **293** (2015), 142–156. 1
- [6] Y. Cao, W.-G. Ma, L.-C. Ma, *Local fractional functional method for solving diffusion equations on Cantor sets*, Abstr. Appl. Anal., **2014** (2014), 6 pages. 1, 2
- [7] A. Carpinteri, P. Cornetti, *A fractional calculus approach to the description of stress and strain localization in fractal media*, Chaos Solitons Fractals, **13** (2002), 85–94. 1
- [8] C. Cattani, H. M. Srivastava, X.-J. Yang, *Fractional dynamics*, De Gruyter Open, Berlin, (2015). 1, 1, 2, 1
- [9] K. Diethelm, N. J. Ford, A. D. Freed, Y. Luchko, *Algorithms for the fractional calculus: a selection of numerical methods*, Comput. Methods Appl. Mech. Engrg., **194** (2005), 743–773. 1
- [10] A. K. Golmankhaneh, A. K. Golmankhaneh, D. Baleanu, *On nonlinear fractional Klein–Gordon equation*, Signal Process., **91** (2011), 446–451. 1
- [11] R. Gorenflo, F. Mainardi, *Fractional calculus: integral and differential equations of fractional order*, Fractals and fractional calculus in continuum mechanics, Udine, (1996), 223–276, CISM Courses and Lectures, Springer, Vienna, (1997). 1
- [12] R. Gorenflo, F. Mainardi, D. Moretti, G. Pagnini, P. Paradisi, *Fractional diffusion: probability distributions and random walk models*, Non extensive thermodynamics and physical applications, Villasimius, (2001), Phys. A, **305** (2002), 106–112. 1
- [13] J.-H. He, *A tutorial review on fractal spacetime and fractional calculus*, Internat. J. Theoret. Phys., **53** (2014), 3698–3718. 1
- [14] J. Hristov, *Approximate solutions to fractional subdiffusion equations*, Eur. Phys. J. Spec. Top., **193** (2013), 229–243. 1
- [15] H. Jafari, S. Seifi, *Solving a system of nonlinear fractional partial differential equations using homotopy analysis method*, Commun. Nonlinear Sci. Numer. Simul., **14** (2009), 1962–1969. 1
- [16] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*, North-Holland Mathematics Studies, Elsevier Science B.V., Amsterdam, (1997). 1
- [17] K. M. Kolwankar, A. D. Gangal, *Local fractional Fokker-Planck equation*, Phys. Rev. Lett., **80** (1998), 214–217. 1
- [18] C. P. Li, F. H. Zeng, *The finite difference methods for fractional ordinary differential equations*, Numer. Funct. Anal. Optim., **34** (2013), 149–179. 1
- [19] F. Liu, V. V. Anh, I. Turner, P. Zhuang, *Time fractional advection-dispersion equation*, J. Appl. Math. Comput., **13** (2003), 233–245. 1
- [20] H.-Y. Liu, J.-H. He, Z.-B. Li, *Fractional calculus for nanoscale flow and heat transfer*, Internat. J. Numer. Methods Heat Fluid Flow, **24** (2014), 1227–1250. 1
- [21] C.-F. Liu, S.-S. Kong, S.-J. Yuan, *Reconstructive schemes for variational iteration method within Yang-Laplace transform with application to fractal heat conduction problem*, Therm. Sci., **17** (2013), 715–721. 1, 2
- [22] Y. Luchko, R. Gorenflo, *An operational method for solving fractional differential equations with the Caputo derivatives*, Acta Math. Vietnam., **24** (1999), 207–233. 1
- [23] Y. F. Luchko, H. M. Srivastava, *The exact solution of certain differential equations of fractional order by using operational calculus*, Comput. Math. Appl., **29** (1995), 73–85. 1
- [24] C. I. Muresan, C. Ionescu, S. Folea, R. De Keyser, *Fractional order control of unstable processes: the magnetic levitation study case*, Nonlinear Dyn., **80** (2014), 1761–1772. 1
- [25] Z. Odibat, S. Momani, *A generalized differential transform method for linear partial differential equations of fractional order*, Appl. Math. Lett., **21** (2008), 194–199. 1
- [26] M. D. Ortigueira, *Fractional calculus for scientists and engineers*, Lecture Notes in Electrical Engineering, Springer, Dordrecht, (2011). 1
- [27] I. Podlubny, A. Chechkin, T. Skovranek, Y. Q. Chen, B. M. Vinagre Jara, *Matrix approach to discrete fractional calculus. II. Partial fractional differential equations*, J. Comput. Phys., **228** (2009), 3137–3153. 1
- [28] Y. A. Rossikhin, M. V. Shitikova, *Application of fractional calculus for dynamic problems of solid mechanics: novel trends and recent results*, Appl. Mech. Rev., **63** (2010), 52 pages. 1
- [29] M. G. Sakar, F. Erdogan, A. Yildirim, *Variational iteration method for the time-fractional Fornberg-Whitham equation*, Comput. Math. Appl., **63** (2012), 1382–1388. 1
- [30] S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional integrals and derivatives*, Theory and applications, Edited and with a foreword by S. M. Nikolskiĭ, Translated from the 1987 Russian original, Revised by the authors, Gordon and Breach Science Publishers, Yverdon, (1993). 1
- [31] N. T. Shawagfeh, *Analytical approximate solutions for nonlinear fractional differential equations*, Appl. Math. Comput., **131** (2002), 517–529. 1
- [32] S. Wei, W. Chen, Y.-C. Hon, *Implicit local radial basis function method for solving two-dimensional time fractional diffusion equations*, Therm. Sci., **19** (2015), 59–67. 1
- [33] B. J. West, P. Grigolini, R. Metzler T. F. Nonnenmacher, *Fractional diffusion and Lévy stable processes*, Phys.

- Rev. E, **55** (1997), 99–106. 1
- [34] X.-J. Yang, *Advanced local fractional calculus and its applications*, World Sci., New York, (2012). 1, 1, 2
- [35] X.-J. Yang, D. Baleanu, H. M. Srivastava, *Local fractional similarity solution for the diffusion equation defined on Cantor sets*, Appl. Math. Lett., **47** (2015), 54–60. 1
- [36] X.-J. Yang, D. Baleanu, H. M. Srivastava, *Local fractional integral transforms and their applications*, Elsevier/Academic Press, Amsterdam, (2016). 1, 2
- [37] X.-J. Yang, J. T. Machado, J. Hristov, *Nonlinear dynamics for local fractional Burgers' equation arising in fractal flow*, Nonlinear Dyn., **84** (2015), 3–7. 1
- [38] X.-J. Yang, H. M. Srivastava, *An asymptotic perturbation solution for a linear oscillator of free damped vibrations in fractal medium described by local fractional derivatives*, Commun. Nonlinear Sci. Numer. Simul., **29** (2015), 499–504. 1
- [39] A.-M. Yang, Y.-Z. Zhang, Y. Long, *The Yang-Fourier transforms to heat-conduction in a semi-infinite fractal bar*, Therm. Sci., **17** (2013), 707–713. 1
- [40] Y.-Z. Zhang, *Initial boundary value problem for fractal heat equation in the semi-infinite region by Yang-Laplace transform*, Therm. Sci., **18** (2014), 677–681. 1, 2