

A New Old Class of Maximal Monotone Operators

M. Marques Alves*

IMPA, Estrada Dona Castorina 110, 22460-320 Rio de Janeiro, Brazil
maicon@impa.br

B. F. Svaiter†

IMPA, Estrada Dona Castorina 110, 22460-320 Rio de Janeiro, Brazil
benar@impa.br

Dedicated to Stephen Simons on the occasion of his 70th birthday.

Received: April 29, 2008

Revised manuscript received: June 5, 2008

In a recent paper in Journal of Convex Analysis the authors studied, in non-reflexive Banach spaces, a class of maximal monotone operators, characterized by the existence of a function in Fitzpatrick's family of the operator which conjugate is above the duality product. This property was used to prove that such operators satisfies a strict version of Brøndsted-Rockafellar property.

In this work we will prove that if a single Fitzpatrick function of a maximal monotone operator has a conjugate above the duality product, then all Fitzpatrick function of the operator have a conjugate above the duality product. As a consequence, the family of maximal monotone operators with this property is just the class NI, previously defined and studied by Simons.

We will also prove that an auxiliary condition used by the authors to prove the strict Brøndsted-Rockafellar property is equivalent to the assumption of the conjugate of the Fitzpatrick function to majorize the duality product.

Keywords: 47H05, 49J52, 47N10

2000 Mathematics Subject Classification: Maximal monotone operators, Brøndsted-Rockafellar property, non-reflexive Banach spaces, Fitzpatrick functions

1. Introduction

Let X be a real Banach space. We use the notation X^* for the topological dual of X and $\langle \cdot, \cdot \rangle$ for the duality product in $X \times X^*$:

$$\langle x, x^* \rangle = x^*(x).$$

In $X \times X^*$ we use the norm-topology. Whenever necessary, we will identify X with its image under the canonical injection of X into X^{**} . To simplify the notation, from now

*Partially supported by Brazilian CNPq scholarship 140525/2005-0.

†Partially supported by CNPq grants 300755/2005-8, 475647/2006-8 and by PRONEX-Optimization.

on π and π_* stand for the duality product in $X \times X^*$ and $X^* \times X^{**}$ respectively:

$$\begin{aligned} \pi : X \times X^* &\rightarrow \mathbb{R}, & \pi_* : X^* \times X^{**} &\rightarrow \mathbb{R} \\ \pi(x, x^*) &= \langle x, x^* \rangle, & \pi_*(x^*, x^{**}) &= \langle x^*, x^{**} \rangle. \end{aligned} \quad (1)$$

The *indicator function* of $A \subset X$ is $\delta_A : X \rightarrow \bar{\mathbb{R}}$,

$$\delta_A(x) := \begin{cases} 0, & x \in A \\ \infty, & \text{otherwise.} \end{cases}$$

For $f : X \rightarrow \bar{\mathbb{R}}$, the lower semicontinuous convex closure of f is $\text{cl conv } f : X \rightarrow \bar{\mathbb{R}}$, the largest lower semicontinuous convex function majorized by f . The conjugate of f is $f^* : X^* \rightarrow \bar{\mathbb{R}}$,

$$f^*(x^*) = \sup_{x \in X} \langle x, x^* \rangle - f(x).$$

It is trivial to check that $f^* = (\text{cl conv } f)^*$.

A point-to-set operator $T : X \rightrightarrows X^*$ is a relation on $X \times X^*$:

$$T \subset X \times X^*$$

and $x^* \in T(x)$ means $(x, x^*) \in T$. An operator $T : X \rightrightarrows X^*$ is *monotone* if

$$\langle x - y, x^* - y^* \rangle \geq 0, \quad \forall (x, x^*), (y, y^*) \in T.$$

and it is *maximal monotone* if it is monotone and maximal (with respect to the inclusion) in the family of monotone operators of X to X^* . Maximal monotone operators in Banach spaces arise, for example, in the study of PDE's, equilibrium problems and calculus of variations.

Given a maximal monotone operator $T : X \rightrightarrows X^*$, Fitzpatrick defined [15] the family \mathcal{F}_T as those convex, lower semicontinuous functions in $X \times X^*$ which are bounded below by the duality product and coincide with it at T :

$$\mathcal{F}_T = \left\{ h \in \bar{\mathbb{R}}^{X \times X^*} \left| \begin{array}{l} h \text{ is convex and lower semicontinuous} \\ \langle x, x^* \rangle \leq h(x, x^*), \quad \forall (x, x^*) \in X \times X^* \\ (x, x^*) \in T \Rightarrow h(x, x^*) = \langle x, x^* \rangle \end{array} \right. \right\}. \quad (2)$$

Fitzpatrick found an explicit formula for the minimal element of \mathcal{F}_T , from now on *Fitzpatrick function* of T , $\varphi_T : X \times X^* \rightarrow \bar{\mathbb{R}}$

$$\varphi_T(x, x^*) = \sup_{(y, y^*) \in T} \langle x, y^* \rangle + \langle y, x^* \rangle - \langle y^*, y \rangle. \quad (3)$$

Moreover, he also proved that if $h \in \mathcal{F}_T$ then h represents T in the following sense:

$$(x, x^*) \in T \iff h(x, x^*) = \langle x, x^* \rangle.$$

Note that

$$\varphi_T(x, x^*) = (\pi + \delta_T)^*(x^*, x).$$

The supremum of the Fitzpatrick family is the \mathcal{S} -function, originally defined and studied by Burachik and Svaiter in [12, Eq. (35)], [11, Eq. (29)]

$$\mathcal{S}_T : X \times X^* \rightarrow \bar{\mathbb{R}}, \quad \mathcal{S}_T = \text{cl conv}(\pi + \delta_T). \tag{4}$$

Beside being the point-wise supremum of the Fitzpatrick Family, the \mathcal{S} -function also belongs to the Fitzpatrick family [12, 11]:

$$\mathcal{S}_T = \sup_{h \in \mathcal{F}_T} h, \quad \mathcal{S}_T \in \mathcal{F}_T. \tag{5}$$

Some authors [4, 25, 6] attribute the \mathcal{S} -function to [18] although [18] was submitted after the publication of [12]. Moreover, the content of [12], and specifically the \mathcal{S} -function, was presented on Erice workshop on July 2001, by R. S. Burachik [9]. A list of the talks of this congress, which includes [20], is available on the [www](http://www.polyu.edu.hk/~ama/events/conference/EriceItaly-OCA2001/Abstract.html)¹. It shall also be noted that [11], the preprint of [12], was published (and available on [www](http://www.polyu.edu.hk/~ama/events/conference/EriceItaly-OCA2001/Abstract.html)) at IMPA preprint server in August 2001.

Burachik and Svaiter defined [12], for $h : X \times X^* \rightarrow \bar{\mathbb{R}}$,

$$\mathcal{J}h : X \times X^* \rightarrow \bar{\mathbb{R}}, \quad \mathcal{J}h(x, x^*) = h^*(x^*, x)$$

and proved that if T is maximal monotone, then \mathcal{J} maps \mathcal{F}_T into itself and $\mathcal{J} \mathcal{S}_T = \varphi_T$:

$$\mathcal{S}_T^*(x^*, x) = \varphi_T(x, x^*).$$

Note that any $h \in \mathcal{F}_T$ satisfies the condition below:

$$\begin{aligned} h(x, x^*) &\geq \langle x, x^* \rangle \\ h^*(x^*, x) &\geq \langle x, x^* \rangle \end{aligned} \quad \forall (x, x^*) \in X \times X^*. \tag{6}$$

What about the converse? Burachik and Svaiter proved in [13, Theorem 3.1] that if a lower semicontinuous convex function h satisfies (6) in a reflexive Banach space, then h represents a maximal monotone operator and h belongs to the Fitzpatrick family of this operator. This result has been used for ensuring maximal monotonicity in reflexive Banach spaces [23, 18, 19, 5, 7, 1, 2, 3, 6, 22].

For the case of a non-reflexive Banach space, Marques-Alves and Svaiter proved [16] that if h is a convex lower semicontinuous function in $X \times X^*$ and

$$\begin{aligned} h(x, x^*) &\geq \langle x, x^* \rangle, & \forall (x, x^*) \in X \times X^* \\ h^*(x^*, x^{**}) &\geq \langle x^*, x^{**} \rangle, & \forall (x^*, x^{**}) \in X^* \times X^{**} \end{aligned} \tag{7}$$

then again h and $\mathcal{J}h$ represent a maximal monotone operator and belong to Fitzpatrick family of this operator. Moreover, the operator T satisfies a strict version of the Brøndsted-Rockafellar property. In particular, Marques-Alves and Svaiter proved that if T is maximal monotone and *one* function in Fitzpatrick family of T satisfies (7), then T satisfies a strict Brøndsted-Rockafellar property. The case of h convex (but not lower semicontinuous) and satisfying (7) was also examined in [16]. The “strong” version of Brøndsted-Rockafellar property for maximal monotone operators in reflexive Banach

¹<http://www.polyu.edu.hk/~ama/events/conference/EriceItaly-OCA2001/Abstract.html>

spaces was established by Torralba [24] and independently rediscovered in Hilbert spaces and in reflexive Banach spaces in [14, 10] respectively.

A maximal monotone operator $T : X \rightrightarrows X^*$ satisfies the Brøndsted-Rockafellar property whenever, for any $x \in X$, $x^* \in X^*$ and $\varepsilon \geq 0$, the condition

$$\langle x - y, x^* - y^* \rangle \geq -\varepsilon, \quad \forall (y, y^*) \in T \tag{8}$$

implies that for any $\lambda > 0$ there exists $x_\lambda \in X$, $x_\lambda^* \in X^*$ such that

$$(x_\lambda, x_\lambda^*) \in T, \quad \|x_\lambda - x\| \leq \lambda, \quad \|x_\lambda^* - x^*\| \leq \frac{\varepsilon}{\lambda}.$$

This condition is an extension for maximal monotone operators of a property of the sub-differential, proved by Brøndsted and Rockafellar in the seminal paper [8]. For Marques Alves and Svaiter, a maximal monotone operator has the strict Brøndsted-Rockafellar property [16] whenever, for any $x \in X$, $x^* \in X^*$ and $\varepsilon \geq 0$, the condition (8) implies that for any $\lambda > 0$ and $\eta > \varepsilon$ there exists $x_\lambda \in X$, $x_\lambda^* \in X^*$ such that

$$(x_\lambda, x_\lambda^*) \in T, \quad \|x_\lambda - x\| < \lambda, \quad \|x_\lambda^* - x^*\| < \frac{\eta}{\lambda}.$$

Martínez-Legaz and Svaiter [17] defined (with a different notation), for $h : X \times X^* \rightarrow \bar{\mathbb{R}}$ and $(x_0, x_0^*) \in X \times X^*$

$$\begin{aligned} h_{(x_0, x_0^*)} : X \times X^* &\rightarrow \bar{\mathbb{R}}, \\ h_{(x_0, x_0^*)}(x, x^*) &:= h(x + x_0, x^* + x_0^*) - [\langle x, x_0^* \rangle + \langle x_0, x^* \rangle + \langle x_0, x_0^* \rangle]. \end{aligned} \tag{9}$$

The operation $h \mapsto h_{(x_0, x_0^*)}$ preserves many properties of h , as convexity, lower semicontinuity and can be seen as the action of the group $(X \times X^*, +)$ on $\bar{\mathbb{R}}^{X \times X^*}$, because

$$(h_{(x_0, x_0^*)})_{(x_1, x_1^*)} = h_{(x_0+x_1, x_0^*+x_1^*)}.$$

Moreover

$$(h_{(x_0, x_0^*)})^* = (h^*)_{(x_0^*, x_0)},$$

where the rightmost x_0 is identified with its image under the canonical injection of X into X^{**} . Therefore,

1. $h \geq \pi \iff h_{(x_0, x_0^*)} \geq \pi$,
2. $(h_{(x_0, x_0^*)})^* \geq \pi_* \iff (h^*)_{(x_0^*, x_0)} \geq \pi_*$,

and finally,

$$h \in \mathcal{F}_T \iff h_{(x_0, x_0^*)} \in \mathcal{F}_{T - \{(x_0, x_0^*)\}}.$$

Marques-Alves and Svaiter work [16] was heavily based on these nice properties of the map $h \mapsto h_{(x_0, x_0^*)}$. These authors also used the fact that if h satisfies condition (7), then it also satisfies the following *auxiliary condition*:

$$\inf_{(x, x^*) \in X \times X^*} h_{(x_0, x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0, \quad \forall (x_0, x_0^*) \in X \times X^*. \tag{10}$$

A possible generalization of [16] would be to require only the auxiliary condition (10) for one Fitzpatrick function of a maximal monotone operator T and then conclude that

this operator satisfies the strict Brøndsted-Rockafellar property. Unfortunately, condition (10) is not more general than condition (7), as we will prove.

The class of operators studied in [16] is the class of maximal Monotone operators for which there exists a function in Fitzpatrick family with a conjugate Above the duality product. So, for the time being, we will call these operators type MA. We will also prove that MA condition is equivalent to NI condition. A maximal monotone $T : X \rightrightarrows X^*$ is type (NI) [21] if

$$\inf_{(y,y^*) \in T} \langle x^{**} - y, x^* - y^* \rangle \leq 0, \quad \forall (x^*, x^{**}) \in X^* \times X^{**}.$$

For proving this equivalence we will show that if *some* $h \in \mathcal{F}_T$ satisfies condition (7), then *all* functions in Fitzpatrick family of T satisfy condition (7). Observe again that, for a function in Fitzpatrick family, $h \geq \pi$ holds by definition.

The main results of this work are the two theorems below:

Theorem 1.1. *Let X be a real Banach space and h be a convex function on $X \times X^*$. Then h satisfies the condition [16, Eq. (4)]*

$$h \geq \pi \quad h^* \geq \pi_* \tag{11}$$

if, and only if, h satisfies the auxiliary condition [16, Eq. immediately below Eq. (29)],

$$\inf_{(x,x^*) \in X \times X^*} h_{(x_0,x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0, \quad \forall (x_0, x_0^*) \in X \times X^*. \tag{12}$$

Theorem 1.2. *Let X be a real Banach space and $T : X \rightrightarrows X^*$. The following conditions are equivalent:*

1. T is type MA, that is, T is maximal monotone and there exists some $h \in \mathcal{F}_T$ such that $h^* \geq \pi_*$ (and $h \geq \pi$),
2. T is maximal monotone and all $h \in \mathcal{F}_T$, satisfies the condition $h^* \geq \pi_*$ (and $h \geq \pi$),
3. T is maximal monotone and some $h \in \mathcal{F}_T$ satisfies the condition

$$\inf_{(x,x^*) \in X \times X^*} h_{(x_0,x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0, \quad \forall (x_0, x_0^*) \in X \times X^*.$$

4. T is maximal monotone and all $h \in \mathcal{F}_T$ satisfies the condition

$$\inf_{(x,x^*) \in X \times X^*} h_{(x_0,x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0, \quad \forall (x_0, x_0^*) \in X \times X^*.$$

5. T is type NI:

$$\inf_{(y,y^*) \in T} \langle x^{**} - y, x^* - y^* \rangle \leq 0, \quad \forall (x^*, x^{**}) \in X^* \times X^{**}.$$

where π and π_* are the duality products in $X \times X^*$ and $X^* \times X^{**}$, as described in (1).

2. Proof of the main results

Proof of Theorem 1.1. Let $\bar{h} := \text{cl conv } h$. As h is convex,

$$\bar{h}(x, x^*) = \lim_{(y, y^*) \rightarrow (x, x^*)} \inf h(y, y^*),$$

and, for any $(x_0, x_0^*) \in X \times X^*$,

$$\bar{h}_{(x_0, x_0^*)}(x, x^*) = \lim_{(y, y^*) \rightarrow (x, x^*)} \inf h_{(x_0, x_0^*)}(y, y^*).$$

As the duality product is continuous and $(\text{cl conv } h)^* = h^*$, condition (11) holds for h if, and only if, it holds for \bar{h} . As the norms are continuous (this is indeed trivial), condition (12) holds for h if, and only if, it holds for \bar{h} . So, it suffices to prove the theorem for the case where h is lower semicontinuous, and we assume it from now on in this proof.

For the sake of completeness, we discuss the implication (11) \Rightarrow (12). Take $(x_0, x_0^*) \in X \times X^*$. If condition (11) holds for h , then it holds for $h_{(x_0, x_0^*)}$ and using [16, Theorem 3.1, Eq. (12)] we conclude that condition (12) holds.

For proving the implication (12) \Rightarrow (11), first note that, for any $(z, z^*) \in X \times X^*$,

$$h_{(z, z^*)}(0, 0) \geq \inf_{(x, x^*)} h_{(z, z^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2.$$

Therefore, using also (12) we obtain

$$h(z, z^*) - \langle z, z^* \rangle = h_{(z, z^*)}(0, 0) \geq 0.$$

Since (z, z^*) is an arbitrary element of $X \times X^*$ we conclude that $h \geq \pi$.

For proving that, under assumption (12), $h^* \geq \pi_*$, take some $(y^*, y^{**}) \in X^* \times X^{**}$. First, use Fenchel-Young inequality to conclude that for any $(x, x^*), (z, z^*) \in X \times X^*$,

$$h_{(z, z^*)}(x, x^*) \geq \langle x, y^* - z^* \rangle + \langle x^*, y^{**} - z \rangle - (h_{(z, z^*)})^*(y^* - z^*, y^{**} - z).$$

As $(h_{(z, z^*)})^* = (h^*)_{(z^*, z)}$,

$$\begin{aligned} (h_{(z, z^*)})^*(y^* - z^*, y^{**} - z) &= h^*(y^*, y^{**}) - \langle z, y^* - z^* \rangle - \langle z^*, y^{**} - z \rangle - \langle z, z^* \rangle \\ &= h^*(y^*, y^{**}) - \langle y^*, y^{**} \rangle + \langle y^* - z^*, y^{**} - z \rangle. \end{aligned}$$

Combining the two above equations we obtain

$$\begin{aligned} &h_{(z, z^*)}(x, x^*) \\ &\geq \langle x, y^* - z^* \rangle + \langle x^*, y^{**} - z \rangle - \langle y^* - z^*, y^{**} - z \rangle + \langle y^*, y^{**} \rangle - h^*(y^*, y^{**}). \end{aligned}$$

Adding $(1/2)\|x\|^2 + (1/2)\|x^*\|^2$ in both sides of the above inequality we have

$$\begin{aligned} h_{(z, z^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 &\geq \langle x, y^* - z^* \rangle + \langle x^*, y^{**} - z \rangle + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 \\ &\quad - \langle y^* - z^*, y^{**} - z \rangle + \langle y^*, y^{**} \rangle - h^*(y^*, y^{**}). \end{aligned}$$

Note that

$$\langle x, y^* - z^* \rangle + \frac{1}{2}\|x\|^2 \geq -\frac{1}{2}\|y^* - z^*\|^2, \quad \langle x^*, y^{**} - z \rangle + \frac{1}{2}\|x^*\|^2 \geq -\frac{1}{2}\|y^{**} - z\|^2.$$

Therefore, for any $(x, x^*), (z, z^*) \in X \times X^*$,

$$\begin{aligned} & h_{(z, z^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 \\ & \geq -\frac{1}{2}\|y^* - z^*\|^2 - \frac{1}{2}\|y^{**} - z\|^2 - \langle y^* - z^*, y^{**} - z \rangle + \langle y^*, y^{**} \rangle - h^*(y^*, y^{**}). \end{aligned}$$

Using now assumption (12) we conclude that the infimum, for $(x, x^*) \in X \times X^*$, at the left hand side of the above inequality is 0. Therefore, taking the infimum on $(x, x^*) \in X \times X^*$ at the left hand side of the above inequality and rearranging the resulting inequality we have

$$h^*(y^*, y^{**}) - \langle y^*, y^{**} \rangle \geq -\frac{1}{2}\|y^* - z^*\|^2 - \frac{1}{2}\|y^{**} - z\|^2 - \langle y^* - z^*, y^{**} - z \rangle.$$

Note that

$$\sup_{z^* \in X^*} -\langle y^* - z^*, y^{**} - z \rangle - \frac{1}{2}\|y^* - z^*\|^2 = \frac{1}{2}\|y^{**} - z\|^2.$$

Hence, taking the sup over $z^* \in X^*$ at the right hand side of the previous inequality we obtain

$$h^*(y^*, y^{**}) - \langle y^*, y^{**} \rangle \geq 0$$

and condition (11) holds. □

Proof of Theorem 1.2. First use Theorem 1.1 to conclude that item 1. and 3. are equivalent. The same theorem also shows that items 2. and 4. are equivalent.

Now assume that item 3. holds, that is, for some $h \in \mathcal{F}_T$,

$$\inf_{(x, x^*) \in X \times X^*} h_{(x_0, x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0, \quad \forall (x_0, x_0^*) \in X \times X^*.$$

Take $g \in \mathcal{F}_T$, and $(x_0, x_0^*) \in X \times X^*$. First observe that, for any $(x, x^*) \in X \times X^*$, $g_{(x_0, x_0^*)}(x, x^*) \geq \langle x, x^* \rangle$ and

$$g_{(x_0, x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 \geq \langle x, x^* \rangle + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 \geq 0.$$

Therefore,

$$\inf_{(x, x^*) \in X \times X^*} g_{(x_0, x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 \geq 0. \tag{13}$$

Note that the square of the norm is coercive, that is, it grows faster than any scalar multiple of the norm, as the norm tends to infinity. Therefore, there exists $M > 0$ such that

$$\left\{ (x, x^*) \in X \times X^* \mid h_{(x_0, x_0^*)}(x, x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 < 1 \right\} \subset B_{X \times X^*}(0, M),$$

where

$$B_{X \times X^*}(0, M) = \left\{ (x, x^*) \in X \times X^* \mid \sqrt{\|x\|^2 + \|x^*\|^2} < M \right\}.$$

For any $\varepsilon > 0$, there exists (\tilde{x}, \tilde{x}^*) such that

$$\min \{1, \varepsilon^2\} > h_{(x_0, x_0^*)}(\tilde{x}, \tilde{x}^*) + \frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2.$$

Therefore

$$\begin{aligned} \varepsilon^2 > h_{(x_0, x_0^*)}(\tilde{x}, \tilde{x}^*) + \frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2 \geq h_{(x_0, x_0^*)}(\tilde{x}, \tilde{x}^*) - \langle \tilde{x}, \tilde{x}^* \rangle \geq 0, \\ M^2 \geq \|\tilde{x}\|^2 + \|\tilde{x}^*\|^2. \end{aligned} \tag{14}$$

In particular,

$$\varepsilon^2 > h_{(x_0, x_0^*)}(\tilde{x}, \tilde{x}^*) - \langle \tilde{x}, \tilde{x}^* \rangle.$$

Using now the fact that operators type MA satisfies the strict Brøndsted-Rockafellar property [16, Theorem 3.4] we conclude that there exists (\bar{x}, \bar{x}^*) such that

$$h_{(x_0, x_0^*)}(\bar{x}, \bar{x}^*) = \langle \bar{x}, \bar{x}^* \rangle, \quad \|\tilde{x} - \bar{x}\| < \varepsilon, \quad \|\tilde{x}^* - \bar{x}^*\| < \varepsilon. \tag{15}$$

Therefore,

$$h(\bar{x} + x_0, \bar{x}^* + x_0^*) - \langle \bar{x} + x_0, \bar{x}^* + x_0^* \rangle = h_{(x_0, x_0^*)}(\bar{x}, \bar{x}^*) - \langle \bar{x}, \bar{x}^* \rangle = 0,$$

and $(\bar{x} + x_0, \bar{x}^* + x_0^*) \in T$. As $g \in \mathcal{F}_T$,

$$g(\bar{x} + x_0, \bar{x}^* + x_0^*) = \langle \bar{x} + x_0, \bar{x}^* + x_0^* \rangle,$$

i.e.,

$$g_{(x_0, x_0^*)}(\bar{x}, \bar{x}^*) = \langle \bar{x}, \bar{x}^* \rangle. \tag{16}$$

Using the first line of (14) we have

$$\varepsilon^2 > h_{(x_0, x_0^*)}(\tilde{x}, \tilde{x}^*) + \left[\frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2 + \langle \tilde{x}, \tilde{x}^* \rangle \right] - \langle \tilde{x}, \tilde{x}^* \rangle \geq \frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2 + \langle \tilde{x}, \tilde{x}^* \rangle.$$

Therefore,

$$\varepsilon^2 > \frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2 + \langle \tilde{x}, \tilde{x}^* \rangle. \tag{17}$$

Direct use of (15) gives

$$\begin{aligned} \langle \bar{x}, \bar{x}^* \rangle &= \langle \tilde{x}, \tilde{x}^* \rangle + \langle \bar{x} - \tilde{x}, \tilde{x}^* \rangle + \langle \tilde{x}, \bar{x}^* - \tilde{x}^* \rangle + \langle \bar{x} - \tilde{x}, \bar{x}^* - \tilde{x}^* \rangle \\ &\leq \langle \tilde{x}, \tilde{x}^* \rangle + \|\bar{x} - \tilde{x}\| \|\tilde{x}^*\| + \|\tilde{x}\| \|\bar{x}^* - \tilde{x}^*\| + \|\bar{x} - \tilde{x}\| \|\bar{x}^* - \tilde{x}^*\| \\ &\leq \langle \tilde{x}, \tilde{x}^* \rangle + \varepsilon[\|\tilde{x}^*\| + \|\tilde{x}\|] + \varepsilon^2 \end{aligned}$$

and

$$\begin{aligned} \|\bar{x}\|^2 + \|\bar{x}^*\|^2 &\leq (\|\tilde{x}\| + \|\bar{x} - \tilde{x}\|)^2 + (\|\tilde{x}^*\| + \|\bar{x}^* - \tilde{x}^*\|)^2 \\ &\leq \|\tilde{x}\|^2 + \|\tilde{x}^*\|^2 + 2\varepsilon[\|\tilde{x}\| + \|\tilde{x}^*\|] + 2\varepsilon^2. \end{aligned}$$

Combining the two above equations with (16) we obtain

$$g_{(x_0, x_0^*)}(\bar{x}, \bar{x}^*) + \frac{1}{2}\|\bar{x}\|^2 + \frac{1}{2}\|\bar{x}^*\|^2 \leq \langle \tilde{x}, \tilde{x}^* \rangle + \frac{1}{2}\|\tilde{x}\|^2 + \frac{1}{2}\|\tilde{x}^*\|^2 + 2\varepsilon[\|\tilde{x}\| + \|\tilde{x}^*\|] + 2\varepsilon^2$$

Using now (17) and the second line of (14) we conclude that

$$g_{(x_0, x_0^*)}(\bar{x}, \bar{x}^*) + \frac{1}{2}\|\bar{x}\|^2 + \frac{1}{2}\|\bar{x}^*\|^2 \leq 2\varepsilon M\sqrt{2} + 3\varepsilon^2.$$

As ε is an arbitrary strictly positive number, using also (13) we conclude that

$$\inf_{(x,x^*) \in X \times X^*} g_{(x_0,x_0^*)}(x,x^*) + \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x^*\|^2 = 0.$$

Altogether, we conclude that if item 3. holds then item 4. holds. The converse item 4. \Rightarrow item 3. is trivial to verify. Hence item 3. and item 4. are equivalent. As item 1. is equivalent to 3. and item 2. is equivalent to 4., we conclude that items 1., 2., 3. and 4. are equivalent.

Now we will deal with item 5. First suppose that item 2. holds. Since, from (5), $\mathcal{S}_T \in \mathcal{F}_T$

$$(\mathcal{S}_T)^* \geq \pi_*.$$

As has already been observed, for any proper function h it holds that $(\text{cl conv } h)^* = h^*$. Therefore, using (4)

$$(\mathcal{S}_T)^* = (\pi + \delta_T)^* \geq \pi_*,$$

that is

$$\sup_{(y,y^*) \in T} \langle y, x^* \rangle + \langle y^*, x^{**} \rangle - \langle y, y^* \rangle \geq \langle x^*, x^{**} \rangle, \quad \forall (x^*, x^{**}) \in X^* \times X^{**}. \quad (18)$$

After some algebraic manipulations we conclude that (18) is equivalent to

$$\inf_{(y,y^*) \in T} \langle x^{**} - y, x^* - y^* \rangle \leq 0, \quad \forall (x^*, x^{**}) \in X^* \times X^{**},$$

that is, T is type (NI). If item 5. holds, by the same reasoning we conclude that (18) holds and therefore $(\mathcal{S}_T)^* \geq \pi_*$. As $\mathcal{S}_T \in \mathcal{F}_T$, we conclude that item 5. \Rightarrow item 1. As has been proved previously item 1. \Rightarrow item 2. \square

Acknowledgements. We thanks the anonymous referees for the suggestions and criticism which improved this work.

References

[1] S. Bartz, H. H. Bauschke, J. M. Borwein, S. Reich, X. Wang: Fitzpatrick functions, cyclic monotonicity and Rockafellar’s antiderivative, *Nonlinear Anal., Theory Methods Appl.* 66A(5) (2007) 1198–1223.
 [2] H. H. Bauschke, J. M. Borwein, X. Wang: Fitzpatrick functions and continuous linear monotone operators, *SIAM J. Optim.* 18(3) (2007) 789–809.
 [3] H. H. Bauschke, X. Wang: A convex-analytical approach to extension results for n -cyclically monotone operators, *Set-Valued Anal.* 15(3) (2007) 297–306.
 [4] J. M. Borwein: Maximal monotonicity via convex analysis, *J. Convex Analysis* 13(3&4) (2006) 561–586.
 [5] J. M. Borwein: Maximality of sums of two maximal monotone operators, *Proc. Amer. Math. Soc.* 134(10) (2006) 2951–2955.
 [6] J. M. Borwein: Maximality of sums of two maximal monotone operators in general Banach space, *Proc. Amer. Math. Soc.* 135(12) (2007) 3917–3924.

- [7] R. I. Boţ, S.-M. Grad, G. Wanka: Maximal monotonicity for the precomposition with a linear operator, *SIAM J. Optim.* 17(4) (2006) 1239–1252.
- [8] A. Brøndsted, R. T. Rockafellar: On the subdifferentiability of convex functions, *Proc. Amer. Math. Soc.* 16 (1965) 605–611.
- [9] R. S. Burachik: Maximal monotone operators, convex functions and a special family of enlargements, in: *International Workshop on Optimization and Control with Applications (Erice, 2001)*, Short Talk.
- [10] R. S. Burachik, B. F. Svaiter: ϵ -Enlargements of maximal monotone operators in Banach spaces, *Set-Valued Anal.* 7(2) (1999) 117–132.
- [11] R. S. Burachik, B. F. Svaiter: Maximal monotone operators, convex functions and a special family of enlargements, Technical Report A094, IMPA (August 2001), available at: http://www.preprint.impa.br/Shadows/SERIE_A/2001/94.html.
- [12] R. S. Burachik, B. F. Svaiter: Maximal monotone operators, convex functions and a special family of enlargements, *Set-Valued Anal.* 10(4) (2002) 297–316.
- [13] R. S. Burachik, B. F. Svaiter: Maximal monotonicity, conjugation and the duality product, *Proc. Amer. Math. Soc.* 131(8) (2003) 2379–2383.
- [14] R. S. Burachik, C. A. Sagastizábal, B. F. Svaiter: ϵ -Enlargements of maximal monotone operators: theory and applications, in: *Reformulation: Nonsmooth, Piecewise Smooth, Semismooth and Smoothing Methods (Lausanne, 1997)*, M. Fukushima et al. (ed.), *Appl. Optim.* 22, Kluwer, Dordrecht (1999) 25–43.
- [15] S. Fitzpatrick: Representing monotone operators by convex functions, in: *Workshop / Miniconference on Functional Analysis and Optimization (Canberra, 1988)*, *Proc. Centre Math. Anal. Austral. Nat. Univ.* 20, Austral. Nat. Univ., Canberra (1988) 59–65.
- [16] M. Marques Alves, B. F. Svaiter: Brøndsted-Rockafellar property and maximality of monotone operators representable by convex functions in non-reflexive Banach spaces, *J. Convex Analysis* 15(4) (2008) 345–348.
- [17] J.-E. Martínez-Legaz, B. F. Svaiter: Monotone operators representable by l.s.c. convex functions, *Set-Valued Anal.* 13(1) (2005) 21–46.
- [18] J.-P. Penot: The relevance of convex analysis for the study of monotonicity, *Nonlinear Anal., Theory Methods Appl.* 58A(7–8) (2004) 855–871.
- [19] J.-P. Penot, C. Zălinescu: On the convergence of maximal monotone operators, *Proc. Amer. Math. Soc.* 134(7) (2006) 1937–1946.
- [20] J.-P. Penot: The use of nonsmooth analysis and of duality methods for the study of hamilton-jacobi equations, in: *International Workshop on Optimization and Control with Applications (Erice, 2001)*, Short Talk.
- [21] S. Simons: The range of a monotone operator, *J. Math. Anal. Appl.* 199(1) (1996) 176–201.
- [22] S. Simons: Positive sets and monotone sets, *J. Convex Analysis* 14(2) (2007) 297–317.
- [23] S. Simons, C. Zălinescu: A new proof for Rockafellar’s characterization of maximal monotone operators, *Proc. Amer. Math. Soc.* 132(10) (2004) 2969–2972.
- [24] D. Torralba: *Convergence Épigraphique et Changements d’Échelle en Analyse Variationnelle et Optimisation*, Ph.D. Thesis, Université Montpellier II, Montpellier (1996).
- [25] M. D. Voisei: The sum and chain rules for maximal monotone operators, *Set-Valued Anal.* 16(4) (2008) 461–476.