

A NOTE ON THE DIAMETER OF A GRAPH

J. A. Bondy

The distance $d(x, y)$ between vertices x, y of a graph G is the length of the shortest path from x to y in G . The diameter $\delta(G)$ of G is the maximum distance between any pair of vertices in G . i. e. $\delta(G) = \max_{x \in G} \max_{y \in G} d(x, y)$. In this note we obtain an upper bound

for $\delta(G)$ in terms of the numbers of vertices and edges in G . Using this bound it is then shown that for any complement-connected graph G with N vertices

$$\delta(G) + \delta(\bar{G}) \leq N + 1$$

where \bar{G} is the complement of G .

THEOREM. Let δ be the diameter of an undirected connected graph G . If G has N vertices $\{x_i\}_1^N$ and E edges then

$$2\delta - 3 - \left(\frac{\delta^2 - \delta - 4}{N}\right) \leq \frac{N^2 - 2E}{N} .$$

Proof. Let $x_1, x_2, \dots, x_{\delta+1}$ be a diametral path. If $m > \delta + 1$, x_m can be joined to at most three vertices of this path. For otherwise, suppose x_m is joined to x_i and to x_{i+k} ($k > 2$). Then $x_1, x_2, \dots, x_i, x_m, x_{i+k}, \dots, x_{\delta+1}$ is a path of length $\delta - k + 2 < \delta$, contradicting the supposition that $x_1, x_2, \dots, x_{\delta+1}$ is a diametral path.

Hence $E \leq \delta$ (the diametral path)

+ $3(N - \delta - 1)$ (the above-mentioned connections)

+ $\frac{1}{2}(N - \delta - 1)(N - \delta - 2)$ (x_m joined to x_n for $m, n > \delta + 1, m \neq n$)

i. e. $2\delta - 3 - \left(\frac{\delta^2 - \delta - 4}{N}\right) \leq \frac{N^2 - 2E}{N}$ as stated.

Note. This upper bound is best possible in the following sense. Given N, E there exists a graph G with N vertices, E edges and diameter δ such that

$$(1) \quad 2\delta - 3 - \left(\frac{\delta^2 - \delta - 4}{N}\right) \leq \frac{N^2 - 2E}{N} < 2(\delta + 1) - 3 - \left(\frac{(\delta + 1)^2 - (\delta + 1) - 4}{N}\right).$$

To construct such a G , let G' be the graph on N vertices $\{x_i\}_1^N$ obtained by taking a path $x_1, x_2, \dots, x_{\delta+2}$ and joining x_m ($\forall m \geq \delta+3$) to x_n ($\forall n \geq \delta, n \neq m$). Then G' has diameter $\delta+1$ and E' edges, where $E' = \delta + 1 + 3(N - \delta - 2) + \frac{1}{2}(N - \delta - 2)(N - \delta - 3)$. The inequalities (1) imply that $0 < E - E' \leq N - \delta$. Let $E - E' = k$. Then G is obtained from G' by adding the following k edges: if $k < N - \delta$ join $x_{\delta-1}$ to $x_{\delta+r}$, $r = 2, \dots, k + 1$; if $k = N - \delta$ join $x_{\delta-1}$ to $x_{\delta+r}$, $r = 2, \dots, N - \delta$, and x_δ to $x_{\delta+2}$.

COROLLARY 1. In an undirected connected graph,

$$\delta < 1 + \frac{N^2 - 2E}{N}$$

Proof. If $\delta = 1$, $2E = N(N-1)$ and therefore $1 + \frac{N^2 - 2E}{N} = 2 > \delta = 1$. If $\delta = 2$, $2E < N(N-1)$ and hence $1 + \frac{N^2 - 2E}{N} > 2 = \delta$. When $\delta \geq 3$, $\delta^2 - \delta - 4 > 0$. Therefore, on using the trivial bound $\delta \leq N - 1$, the Theorem gives

$$\begin{aligned} \frac{N^2 - 2E}{N} &> 2\delta - 3 - \left(\frac{\delta^2 - \delta - 4}{\delta + 1}\right) \\ &= 2\delta - 3 - (\delta - 2) + \frac{2}{\delta + 1} > \delta - 1. \end{aligned}$$

COROLLARY 2. If both G and its complement \bar{G} are connected graphs,

$$\delta(G) + \delta(\bar{G}) \leq N + 1.$$

Proof. From Corollary 1

$$\delta(G) < 1 + \frac{N^2 - 2E}{N}.$$

Similarly $\delta(\bar{G}) < 1 + \frac{N^2 - 2\bar{E}}{N}$ where $E + \bar{E} = \frac{1}{2}N(N-1)$.

Therefore $\delta(G) + \delta(\bar{G}) < N + 3$
i.e.

$$(2) \quad \delta(G) + \delta(\bar{G}) \leq N + 2.$$

If $\delta(G) \leq 2$, then, since $\delta(\bar{G}) \leq N - 1$, we have immediately that $\delta(G) + \delta(\bar{G}) \leq N + 1$.

If $\delta(G) = 3$, then $\delta(\bar{G}) \leq N - 2$. For there is only one graph \bar{G} with $\delta(\bar{G}) = N - 1$ (the simple path) and for this graph $\delta(G) = 2$. Hence in this case $\delta(G) + \delta(\bar{G}) \leq N + 1$. By symmetry Corollary 2 also holds if $\delta(\bar{G}) \leq 3$.

We now assume that $\delta(G) > 3$, $\delta(\bar{G}) > 3$.

By the Theorem

$$2\delta - 3 \leq \frac{N^2 - 2E}{N} + \left(\frac{\delta^2 - \delta - 4}{N} \right)$$

$$2\bar{\delta} - 3 \leq \frac{N^2 - 2\bar{E}}{N} + \left(\frac{\bar{\delta}^2 - \bar{\delta} - 4}{N} \right)$$

where $\delta \equiv \delta(G)$, $\bar{\delta} \equiv \delta(\bar{G})$.

Therefore $2(\delta + \bar{\delta}) - 6 \leq N + 1 + \left(\frac{\delta^2 + \bar{\delta}^2 - \delta - \bar{\delta} - 8}{\delta + \bar{\delta} - 2} \right)$

by addition and (2).

Hence $2(\delta + \bar{\delta}) - 6 \leq N + 1 + \delta + \bar{\delta} - 5 - \frac{2(\delta - 3)(\bar{\delta} - 3)}{\delta + \bar{\delta} - 2}$.

i. e. $\delta + \bar{\delta} \leq N + 2 - \frac{2(\delta - 3)(\bar{\delta} - 3)}{\delta + \bar{\delta} - 2} < N + 2$.

Therefore $\delta + \bar{\delta} \leq N + 1$.

Note added in proof: The author has recently noticed that these results are essentially contained in Lemma 1.1 and Lemma 3 of [1]. In fact Lemma 3 of [1] implies that, apart from the case $\delta(G) = \delta(\bar{G}) = 3$, $\delta(G) + \delta(\bar{G}) = \max \{ \delta(G) + 2, \delta(\bar{G}) + 2 \}$.

REFERENCE

1. J. Bosák, A. Rosa, and Š. Znam, On decomposition of complete graphs into factors with given diameters. Proceedings of the Colloquium on Theory of Graphs, held at Tihany, Hungary, September 1966, edited by P. Erdős and G. Katona. (Academic Press, New York, 1968).

Mathematical Institute, Oxford