


ARTICLE

Open Access

# A novel antibody-TCR (AbTCR) platform combines Fab-based antigen recognition with gamma/delta-TCR signaling to facilitate T-cell cytotoxicity with low cytokine release

Yiyang Xu<sup>1</sup>, Zhiyuan Yang<sup>1</sup>, Lucas H. Horan<sup>1</sup>, Pengbo Zhang<sup>1</sup>, Lianxing Liu<sup>1</sup>, Bryan Zimdahl<sup>1</sup>, Shon Green<sup>1</sup>, Jingwei Lu<sup>1</sup>, Javier F. Morales<sup>1</sup>, David M. Barrett<sup>2</sup>, Stephan A. Grupp<sup>2</sup>, Vivien W. Chan<sup>1</sup> , Hong Liu<sup>1</sup> and Cheng Liu<sup>1</sup>

## Abstract

The clinical use of genetically modified T-cell therapies has led to unprecedented response rates in leukemia and lymphoma patients treated with anti-CD19 chimeric antigen receptor (CAR)-T. Despite this clinical success, FDA-approved T-cell therapies are currently limited to B-cell malignancies, and challenges remain with managing cytokine-related toxicities. We have designed a novel antibody-T-cell receptor (AbTCR) platform where we combined the Fab domain of an antibody with the  $\gamma$  and  $\delta$  chains of the TCR as the effector domain. We demonstrate the ability of anti-CD19-AbTCR-T cells to trigger antigen-specific cytokine production, degranulation, and killing of CD19-positive cancer cells in vitro and in xenograft mouse models. By using the same anti-CD19 binding moiety on an AbTCR compared to a CAR platform, we demonstrate that AbTCR activates cytotoxic T-cell responses with a similar dose-response as CD28/CD3 $\zeta$  CAR, yet does so with less cytokine release and results in T cells with a less exhausted phenotype. Moreover, in comparative studies with the clinically validated CD137 (4-1BB)-based CAR, CTL019, our anti-CD19-AbTCR shows less cytokine release and comparable tumor inhibition in a patient-derived xenograft leukemia model.

## Introduction

The remarkable efficacy of anti-CD19-chimeric antigen receptor (CAR)-T cell therapy in both B-cell acute lymphoblastic leukemia (B-ALL) and lymphomas<sup>1,2</sup> has demonstrated the clinical importance of genetically modified T-cells as a cancer therapy, and simultaneously exemplified Eshhar's original vision to make a chimeric cell that combines the antibody specificity of a B-cell with the cytotoxic properties of a T cell<sup>3</sup>. The first chimeric

receptor design from Eshhar's group replaced the antigen recognition variable regions of the alpha ( $\alpha$ ) and beta ( $\beta$ ) TCR chains with the variable regions of an anti-SP6 antibody<sup>3</sup>. While they were able to demonstrate antigen specific T-cell activation through this chimeric antibody-TCR receptor, there were technical hurdles with the mispairing with the T-cell's endogenous  $\alpha$  and  $\beta$  TCR chains and having to express two synthetic molecules in the same cell. The group subsequently addressed these problems by engineering a single chain molecule that fused an antibody in scFv format onto the Immunoreceptor Tyrosine-based Activation Motifs (ITAM)-containing domain of CD3 $\zeta$ <sup>4</sup>. The efficient single-chain design has demonstrated clinical efficacy as the backbone for the majority of CAR-T therapies to date.

Correspondence: Cheng Liu ([cheng.liu@eurekainc.com](mailto:cheng.liu@eurekainc.com))

<sup>1</sup>Eureka Therapeutics, Inc, Emeryville, CA 94608, USA

<sup>2</sup>Division of Oncology, Department of Pediatrics, Children's Hospital of Philadelphia and Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA 19104, USA

These authors contributed equally: Yiyang Xu, Zhiyuan Yang, Lucas H. Horan, Pengbo Zhang

© The Author(s) 2018



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

However, the direct fusion of antigen recognition to cellular activation domains creates a synthetic activation signal that likely differs from the cellular activation signal propagated from an endogenous TCR-CD3 complex.

T cells are molecularly defined by TCRs present on their cell surface. The TCR contributes to tumor immune surveillance<sup>5</sup> by enabling T cells to recognize abnormal cells and triggering a cascade of signaling events that lead to T-cell activation and subsequent cancer cell lysis. In the majority of T cells, the TCR consists of an  $\alpha$  chain and a  $\beta$  chain, whereas in 1–5% of T cells the TCR consists of a gamma ( $\gamma$ ) and a delta ( $\delta$ ) chain<sup>6</sup>. The extracellular regions of the  $\alpha\beta$  chains (or the  $\gamma\delta$  chains) are responsible for antigen recognition and engagement. Antigen binding stimulates downstream signaling through the multimeric CD3 complex that associates with the intracellular domains of the  $\alpha\beta$  (or  $\gamma\delta$ ) chains as three dimers ( $\epsilon\gamma$ ,  $\epsilon\delta$ ,  $\zeta\zeta$ )<sup>7</sup>. The entire CD3 complex contains 10 ITAMs which feed into a network of phosphorylation pathways that create the T-cell activation signal<sup>7</sup>. We hypothesized that by replacing the antigen recognition domain of a  $\gamma\delta$ TCR with an antibody-derived Fab fragment, we could create a synthetic receptor that uses endogenous TCR signaling pathways while having the flexibility to target either a peptide-MHC complex with a TCR-mimic (TCRm) antibody, or an extracellular antigen with a conventional antibody.

TCR-T cell therapy is another active field of research. While it has shown clinical response<sup>8</sup>, TCR-T therapies has been predominantly limited to targets that are MHC (major histocompatibility complex)-restricted. TCRm antibodies that recognize peptide-MHC complexes<sup>9</sup> have allowed direct functional comparisons between single-chain CAR activation and activation through the endogenous signaling pathways used by TCRs with a matched antigen-recognition motif<sup>10–12</sup>. Head-to-head comparisons demonstrate that activation through the TCR leads to a T cell with more potent anti-tumor cytotoxicity and notably in one study, higher antigen sensitivity with less cytokine release<sup>10</sup>. These data suggest there may be therapeutic advantages to an engineered T-cell therapy that uses a cellular activation mechanism that more closely resembles the activation signal propagated from the endogenous TCR.

In this study, we describe the design, characterization, and preclinical validation of our two-chained antibody-TCR (AbTCR). Unlike previous designs that were built on  $\alpha\beta$  chains of the TCR, our AbTCR platform avoids mispairing with the T cell's endogenous  $\alpha\beta$ TCR by using the transmembrane and intracellular domains from the  $\gamma\delta$ TCR. To functionally characterize the new platform, we developed a human anti-CD19 antibody (ET190L1), generated both ET190L1-AbTCR-T cells and ET190L1-CAR-T cells (employing CD28 and CD3 $\zeta$ ) and compared


phenotypes between the T cells before and after antigen stimulation. Finally, we show both in vitro and with tumor xenograft models that ET190L1-AbTCR-T cells maintain comparable anti-tumor potency to ET190L1-CAR and CD137-based CTL019 T cells, yet T-cell activation through ET190L1-AbTCR results in lower concentrations of inflammatory cytokines.

## Results

### The antibody-TCR (AbTCR) forms a multimeric T cell signaling molecule with the endogenous CD3 complex

Our synthetic receptor design is an AbTCR that fuses an antigen-binding domain from an antibody with the C-terminal signaling domain of a  $\gamma\delta$ TCR. More specifically, the antigen-binding domain of our AbTCR is an antibody Fab fragment, and the effector, or cellular-activation domain of our AbTCR consists of portions of the  $\delta$  and  $\gamma$  chains of a  $\gamma\delta$ TCR (Fig. 1a). In the AbTCR receptor, the heavy chain domain of the Fab fragment is fused to a portion of the  $\delta$  chain of a  $\gamma\delta$  TCR. Whereas, the light chain domain of the Fab fragment is fused to a portion of the  $\gamma$  chain of a  $\gamma\delta$  TCR. We named this  $\gamma\delta$ -based synthetic receptor the ARTEMIS™ AbTCR. Similar to the exogenously-expressed TCR in TCR-T cells, our AbTCR engages the endogenous CD3 complex to initiate T-cell activation. However, since the  $\gamma\delta$ TCR chains do not bind with the TCRs in  $\alpha\beta$  T-cells<sup>13,14</sup>, AbTCR avoids the mispairing challenge of traditional  $\alpha\beta$ TCR-based synthetic receptors<sup>15,16</sup>. In addition, by employing a Fab as the antigen-binding domain, the AbTCR can be used to target either peptide-MHC complexes or cell surface antigens by using TCR-mimic or conventional antibodies respectively. Overall, the AbTCR design couples the antigen-binding utility of an antibody to the endogenous TCR activation pathways.

CAR-T therapy has been extensively studied in the clinical setting against CD19<sup>+</sup> tumors. Thus, throughout this study we use a human anti-CD19 antibody (ET190L1, Eureka Therapeutics, Supplementary Fig. S1) to evaluate the AbTCR platform in both in vitro and in vivo CD19<sup>+</sup> tumor models. The ET190L1 antibody was isolated from a human-derived antibody library and competes for the same epitope as the murine-derived FMC63 antibody (Supplementary Fig. S1b). Using an anti-human Fab antibody, which recognizes the Fab domain of the AbTCR, we observed efficient expression and even distribution of the receptor on the surface of primary T cells (Fig. 1b). Crucial to the AbTCR design is the ability of the receptor to trigger T-cell activation through the association with the endogenous CD3 complex. CD3 is a multi-chain complex composed of a CD3 $\zeta$  homodimer and two other heterodimers formed from CD3 $\delta$ , CD3 $\epsilon$ , CD3 $\gamma$  chains. J.RT3-T3.5 is a T cell line that lacks CD3 $\epsilon$  surface expression because of a mutation in the TCR $\beta$  chain that



prevents TCR $\alpha\beta$ :CD3 complex formation. Fig. 1c shows that the AbTCR interacts with endogenous CD3 molecules and restores CD3 $\epsilon$  surface expression on J.RT3-T3.5 cells. To test if AbTCR associates with the additional members of the CD3 complex, a FLAG-tagged ET190L1-

AbTCR was expressed in primary T cells for biochemical pull-down. All four CD3 chains co-immunoprecipitated with the AbTCR, suggesting AbTCR can form a complex with CD3 that is biochemically similar to the endogenous TCR:CD3 complex (Fig. 1d).

### AbTCR-T cells differ from CAR-T cells prior to antigen engagement

To compare the phenotypes between AbTCR-T cells and CAR-T cells, we engineered a scFv using the same human anti-CD19 binding domain and fused it with a second generation CD28/CD3 $\zeta$  CAR (ET190L1-CAR). This allowed us to evaluate AbTCR-T cells in comparison to a CD28/CD3 $\zeta$ -based CAR platform that is widely used clinically. During T cell manufacturing, ET190L1-AbTCR-T cells expanded with similar growth kinetics as ET190L1-CAR-T cells and yielded T cell populations with similar transduction efficiencies and CD4:CD8 ratios (Figs. 2a, b, Supplementary Fig. S5). Retrospective analysis from published CAR-T clinical studies have found that T cells that are more naïve, less differentiated, and less exhausted correlate with improved efficacy<sup>17,18</sup>. Interestingly, after CD3/CD28 expansion, but before antigen engagement, a greater fraction of ET190L1-AbTCR-T cells displayed a naïve ( $T_N$ ) and stem cell memory ( $T_{SCM}$ ) T cell phenotype compared to ET190L1-CAR-T cells (Figs. 2c, d, Supplementary Figs. S6-7). Taken together with the shifts in increased CD28 and lower granzyme B (GranzB) on CD8<sup>+</sup> AbTCR-T cells, the increased CCR7 expression indicates that T-cells engineered with AbTCR are less differentiated<sup>19</sup> (Fig. 2e). Furthermore, expression of the exhaustion markers PD-1 and TIM-3 were lower on AbTCR-T cells than in CAR-T cells (Fig. 2f). Since both ET190L1-CAR-T cells and ET190L1-AbTCR-T cells express the same antigen-specific Ab variable sequences we were surprised to observe differences prior to antigen engagement. We believe the most likely explanations for this difference is tonic signaling that has been reported for both CD137- and CD28-based CAR-T cells<sup>20,21</sup>.

### AbTCR recapitulates functions mediated by TCR

We next characterized the T-cell phenotypes resulting from activation through the AbTCR. ET190L1-AbTCR-T cells were co-incubated with either CD19<sup>+</sup> Raji cells or Raji cells in which CD19 was knocked out (Raji K/O). Upon engagement with CD19<sup>+</sup> Raji cells, ET190L1-AbTCR-T cells express activation markers CD69 and CD25 (Fig. 3a). The accumulation of CD107a was determined as a measure of cellular degranulation, a prerequisite for T-cell-mediated cytotoxicity. As shown in Fig. 3b and Supplementary Fig. S8, T cells degranulated when ET190L1-AbTCR is stimulated with CD19-positive cells. In addition, intracellular flow cytometric analysis of ET190L1-AbTCR-T cells co-incubated with CD19<sup>+</sup> Raji cells shows that TNF $\alpha$ , IL-2, and IFN $\gamma$  are induced in response to antigen (Supplementary Fig. S2). Importantly, no cytokines were produced when ET190L1-AbTCR-T cells were co-cultured with CD19-negative Raji K/O cells (Supplementary Fig. S2). These data demonstrate the


ability of the ET190L1-AbTCR to trigger T-cell activation in an antigen-dependent manner.

To better characterize the tumor-killing activities of AbTCR-T cells, we set up experiments to directly compare phenotypes of ET190L1-AbTCR-T cells with ET190L1-CAR-T cells. The percentage of AbTCR-positive and CAR-positive T cells were matched by dilutions with un-transduced mock T cells and co-cultured with multiple tumor cell lines. Specific lysis of only CD19-positive tumor lines confirmed the antigen specificity of both ET190L1-CAR-T cells and ET190L1-AbTCR-T cells while demonstrating comparable cellular cytotoxicity and degranulation (Fig. 3c, Supplementary Fig. S8). Specific lysis across a range of E:T ratios also showed comparable T-cell killing at low E:T ratios, further demonstrating the cytotoxic potential of using the AbTCR (Supplementary Fig. S3). Replicative capacity of the therapeutic T cell in leukemia patients has been reported to be a key predictive biomarker for clinical efficacy<sup>22,23</sup>. We used a CFSE-based assay to assess in vitro T-cell proliferation upon antigen stimulation. As shown in Fig. 3d, ET190L1-AbTCR-T cells divided in response to antigen with kinetics comparable to that observed with ET190L1-CAR-T cells.


### AbTCR-T cells present less exhaustion markers and release less cytokines during tumor cell lysis

Despite a slight increase in the expression of the CD69 and CD25 activation markers on tumor stimulated AbTCR-T cells compared to CAR-T cells (Fig. 3a), AbTCR<sup>+</sup> CD4<sup>+</sup> cells expressed substantially lower levels of the PD-1 exhaustion marker than CAR<sup>+</sup> CD4<sup>+</sup> cells, and, in both CD4<sup>+</sup> and CD8<sup>+</sup> AbTCR-T cells, LAG-3 was significantly lower (Fig. 3e). Furthermore, while ET190L1-AbTCR-T cells have comparable cytotoxicity and proliferative potential compared to ET190L1-CAR-T cells (Figs. 3c, d), the AbTCR-T cells release lower levels of inflammatory cytokines including TNF- $\alpha$ , IL-2, IFN- $\gamma$ , and GM-CSF after a 16-h in vitro killing assay (Fig. 4a). Comparisons between TCR-T and CAR-T cells have previously shown that activation through the TCR can comparatively reduce cytokine release while simultaneously increasing antigen sensitivity<sup>10</sup>. Although the CAR construct incorporates a covalently-linked CD28 costimulatory domain, Raji cells express CD80 and CD86 and thus provide CD28 costimulation to both CAR-T cells and AbTCR-T cells. Therefore, we believe the cytokine secretion and exhaustion differences between AbTCR and CAR stem from the utilization of endogenous signaling pathways by the  $\gamma\delta$ TCR effector domain of the AbTCR receptor.

While the promise of AbTCR-T cells to reduce the secretion of several inflammatory cytokines has exciting clinical potential, the discovery that tocilizumab, an anti-IL6R antibody, alleviates cytokine release syndrome (CRS)



**Fig. 2** AbTCR-T cells have a less exhausted surface phenotype and a higher percentage of naïve and stem cell memory T cells compared to CAR-T cells. **a** ET190L1-AbTCR T cells (AbTCR) and ET190L1-CAR-T cells (CAR) were cultured and the number of cells determined at the indicated time points. **b–e** Flow cytometry analysis of ET190L1-AbTCR-T cells and ET190L1-CAR-T cells at day 10 of in vitro expansion prior to antigen stimulation (**b**) Proportions of CD4/CD8 within receptor<sup>+</sup> cells. **c** Frequency of naïve (CCR7<sup>+</sup> CD45RA<sup>+</sup>), central memory (CM; CCR7<sup>+</sup> CD45RA<sup>-</sup>), effector memory (EM; CCR7<sup>-</sup> CD45RA<sup>-</sup>) and effector (E; CCR7<sup>-</sup> CD45RA<sup>+</sup>) T cells within CD8<sup>+</sup> receptor<sup>+</sup> cells. **d** Frequency of stem cell memory (SCM; CCR7<sup>+</sup> CD45RO<sup>-</sup> CD95<sup>+</sup> CD122<sup>+</sup>) T cells within CD8<sup>+</sup> receptor<sup>+</sup> cells. **e** Expression of T cell differentiation markers CD28, CCR7, and granzyme B (GranB). **f** Expression of T cell exhaustion markers PD-1, LAG-3, and TIM-3. \*\**P* < 0.01



**Fig. 3** (See legend on next page.)

(see figure on previous page)

**Fig. 3 AbTCR-T cells mediate antigen-specific responses in vitro.** **a** Expression of activation markers CD69 and CD25 on ET190L1-AbTCR-T cells (AbTCR) and ET190L1-CAR-T cells (CAR) following a 16 h co-incubation with Raji or Raji K/O cells. Raji K/O: Raji cells in which CD19 was knocked out using CRISPR technology. Gated on CD3<sup>+</sup> receptor<sup>+</sup> cells. **b** ET190L1-AbTCR-T cells selectively degranulate in the presence of CD19<sup>+</sup> target cells. Representative flow cytometry plot showing CD107a expression on ET190L1-AbTCR incubated with target cells for 4 h at an E:T of 1:1. Degranulation in AbTCR<sup>+</sup> CD8<sup>+</sup> T cells is shown. **c** ET190L1-AbTCR-T cells selectively kill CD19<sup>+</sup> tumor cells and are comparable to ET190L1-CAR-T cells. T cells were incubated with target cells for 16 h at an E:T ratio of 2:1. Cytotoxicity was measured by LDH release assay ( $n = 3$  technical replicates). Error bars, SEM. **d** ET190L1-AbTCR-T cells and ET190L1-CAR-T cells proliferate at similar rates upon antigen stimulation in vitro. T cells were labeled with CFSE and co-cultured with CD19<sup>+</sup> Raji cells and the degree of T cell proliferation was assessed by flow cytometry. **e** ET190L1-AbTCR CD4<sup>+</sup> T cells express lower levels of exhaustion markers following co-incubation with CD19<sup>+</sup> tumor cells

pathology, singles out IL-6 with particular clinical significance<sup>24–27</sup>. Because the majority of IL-6 is produced by antigen presenting cells<sup>28</sup>, including monocytes, macrophages, and dendritic cells, we performed a co-culture assay to measure IL-6 concentrations. Transwells were used to separate T cells/tumor cells from monocyte-lineage cells. ET190L1-AbTCR-T cells were compared to one of the anti-CD19 CAR-T cells that has been extensively studied in the clinic and recently approved (CTL019; research grade version of FDA-approved Kymriah<sup>™</sup>)<sup>29</sup>. In addition, CTL019 uses CD137 (4-1BB) for costimulation, thus offering an opportunity to compare the AbTCR to a CD137-based CAR. Similar to the observed differences in cytokine release between ET190L1-AbTCR-T cells and ET190L1-CAR-T cells (Fig. 4a), ET190L1-AbTCR-T cells released lower levels of TNF- $\alpha$ , IL-2, IFN- $\gamma$ , GM-CSF compared to CTL019-T cells (Fig. 4b). Importantly, we found that ET190L1-AbTCR-T cells induced monocyte-lineage cells to release substantially less IL-6 than CTL019-T cells (Fig. 4b). To test if the reduced cytokine release has an effect on in vivo anti-tumor activity, we used ET190L1-AbTCR-T cells to treat a patient-derived xenograft (PDX) mouse model of primary B-ALL (CHP105R1, negative for both CD80 and CD86) and observed similar tumor inhibition between mice treated with the ET190L1-AbTCR- and CTL019-T cells (Fig. 5c). Thus, consistent with the Raji in vitro studies above, T cells engineered with AbTCR reduce cytokine release without a loss of anti-tumor activity, in a PDX tumor model that lacks CD80 and CD86 costimulation.

#### AbTCR-T cells have similar anti-tumor activities as CAR-T cells in pre-clinical models of B-cell cancers yet release less cytokines and become less exhausted during tumor clearance


We next tested the in vivo anti-tumor activity of ET190L1-AbTCR-T cells in an established human CD19<sup>+</sup> Raji B-cell lymphoma xenograft model. As shown in Fig. 5a and Supplementary Fig. S4, both ET190L1-AbTCR and ET190L1-CAR treatments resulted in tumor regression and long-lasting tumor rejection. At the time when

mice treated with mock-T cells had to be euthanized, tumor burden was on average  $\sim 1000$  fold less in mice treated with ET190L1-CAR-T cells than in the mock-treated mice and remarkably, on average  $\sim 5300$  fold less in mice treated with ET190L1-AbTCR-T cells than in the mock arm in this experiment (Fig. 5a). The ability of persisting ET190L1-AbTCR-T cells to prevent growth of “newly-introduced” tumor cells was tested by re-injecting mice with tumor cells weeks after the T cells had cleared the initial tumor burden (Fig. 5b). While tumors grew rapidly in control mice, mice treated previously with either AbTCR-T cells or CAR-T cells were resistant to Raji lymphoma re-challenge (Fig. 5b). Separate mice in a similar experiment were resistant to a NALM-6 (CD80 and CD86 negative) B-ALL challenge (Fig. 5b). The data from this NALM-6 re-challenge, along with data shown in Fig. 5c, demonstrate that the in vivo anti-tumor activity of ET190L1-AbTCR-T cells extends to CD19<sup>+</sup> tumor types that do not express the CD80 and CD86 costimulatory ligands.

Analysis of in vivo cytokine release and exhaustion markers on T cells recapitulated in vitro findings. Whereas ET190L1-CAR treatment caused marked elevation of inflammatory cytokines, including IL-2, IL-10, IFN- $\gamma$ , and TNF- $\alpha$ , significantly lower levels of these cytokines were released following ET190L1-AbTCR treatment (Fig. 5d). T-cell collected from peripheral blood 9 days and 15 days post-T-cell dosing also revealed that AbTCR-T cells expressed significantly lower levels of PD-1 than CAR-T cells (Fig. 5e). Collectively, these data show that AbTCR-T cells exhibit potent in vitro and in vivo anti-tumor activity, yet release lower levels of inflammatory cytokines and express lower levels of exhaustion markers than CAR-T cells.

#### Discussion

The AbTCR design fuses the Fab domain of an antibody with the effector domains from the  $\gamma$  and  $\delta$  TCR chains, combining the affinity and specificity of antibody recognition with the tumor cytotoxicity potential of a T cell. Similar to the TCR-T platform, this design enables the AbTCR to associate with the CD3 complex, which allows



**Fig. 4** AbTCR-T cells release less CRS-related cytokines than CAR-T cells, including monocyte-lineage-derived IL-6. **a** ET190L1-AbTCR-T cells (AbTCR) release less cytokines than ET190L1-CAR-T cells (CAR) during in vitro killing assays. T cells were incubated with target cells for 16 h at an E:T ratio of 5:1. Secreted cytokine levels in the supernatant were measured. Data shown are representative of three independent experiments. **b** Cytokine expression after trans-well co-culture. T cells, targets (CD19<sup>+</sup> NALM-6 cells) were placed in plate wells, and monocyte-lineage cells (APCs) were placed in transwell inserts. Supernatants were collected after 18 h of co-cultures






Fig. 5 (See legend on next page.)

(see figure on previous page)

**Fig. 5 Robust tumor rejection and reduced cytokine release with ET190L1-AbTCR-T cells in pre-clinical xenograft tumor models.** **a** ET190L1-AbTCR-T cell therapy shows efficacy in a Raji lymphoma model. Bioluminescent images (left panel) and total flux (right panel) over time of Raji-luc implanted mice intravenously administered with  $5 \times 10^6$  (1) un-transduced donor-matched T cells (Mock), (2) ET190L1-CAR-T cells (CAR), or (3) ET190L1-AbTCR-T cells (AbTCR). Doses were based on number of receptor-positive cells;  $n = 6-8$  mice/group. **b** ET190L1-AbTCR treated mice resist Raji lymphoma and NALM-6 B-ALL tumor re-challenge. ET190L1-AbTCR and ET190L1-CAR-treated mice that showed no detectable Raji lymphoma were re-challenged with Raji cells (left panel) or NALM-6 cells (right panel). As controls, Raji-naïve mice were implanted with Raji cells (left panel) or NALM-6 cells (right panel) following an injection of Mock T cells;  $n = 2-3$  mice/group. **c** ET190L1-AbTCR shows anti-tumor activity in a PDX model of CD80/86 negative primary B-ALL (CHP105R1). Total flux overtime of CHP105R1-bearing mice treated with either un-transduced T cells (Mock), CTL019 T cells, or ET190L1-AbTCR T cells;  $n = 5$  mice/group. **d** ET190L1-AbTCR-T cells release significantly less cytokines than ET190L1-CAR-T cells in vivo. Serum cytokine levels (pg/ml) collected from Raji-bearing mice 24 h after T cell dosing. IL-2, IL-10, IFN- $\gamma$  and TNF- $\alpha$  levels are shown. **e** ET190L1-AbTCR-T cells express significantly less PD-1 levels than ET190L1-CAR-T cells in vivo. PD-1 expression levels (mean fluorescent intensity; MFI) on CAR<sup>+</sup> CD4<sup>+</sup> and AbTCR<sup>+</sup> CD4<sup>+</sup> T cells at select times from Raji-bearing mice after T cell infusion. \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; \*\*\*\* $P < 0.0001$ .

the antigen/AbTCR engagement to trigger endogenous T-cell activation and regulatory pathways. Distinct from most TCR-T platforms, however, is the  $\gamma\delta$ TCR-based intracellular domain and the incorporation of an antibody-binding moiety for target recognition. This enables the extension of the TCR platform to non-MHC-restricted targets, such as CD19.

The molecular design of our AbTCR has distinct structural features at both the antigen recognition and effector domains. Our choice of using  $\gamma\delta$  TCR components alleviates the need to disrupt the endogenous  $\alpha\beta$  TCR chains<sup>15,16</sup> since  $\gamma\delta$  TCR subunits do not pair with endogenous  $\alpha\beta$  TCR subunits<sup>13,14</sup>. Another potential advantage of using  $\gamma\delta$  TCR subunits is that they have a higher affinity for the CD3 complex than  $\alpha\beta$  TCRs<sup>30</sup>, therefore signaling downstream of AbTCR may be enhanced relative to that produced by  $\alpha\beta$  TCRs. In addition, the Fab antibody fragment is generally more soluble than its less stable scFv configuration<sup>31</sup>. Since some scFv domains have been reported to induce CAR aggregation and tonic signaling<sup>20</sup>, the Fab-based AbTCR platform could potentially be an alternative for antibodies that are problematic in their scFv-based CAR format<sup>11,20</sup>. In our work characterizing AbTCR, we chose to conduct our experiments along with two of the most used CAR designs currently in clinical studies. Given that variables such as linkers, hinges and transmembrane domains have been shown to affect CAR-T efficacy, we are investigating whether similar modifications may also affect AbTCR.

To lessen concerns of immunogenic responses that have been observed with murine-derived scFv's, we developed a human anti-CD19 antibody (ET190L1) and characterized T cells expressing ET190L1-AbTCR phenotypically and functionally. While ET190L1-AbTCR-T cells expand at a comparable rate as ET190L1-CAR-T cells (Fig. 2a), the AbTCR-T cells are characterized by a more naïve/stem-cell like and less exhausted surface phenotype (Figs. 2c-f). Since T-cell expansion and persistence have been found to correlate with clinical efficacies of anti-CD19 CAR-T

therapy, it will be of great interest to see if the phenotypes observed with AbTCR-T cells translate in the clinical setting.

The role of costimulatory signals, such as CD28 and CD137 (4-1BB), in T-cell activation is well-established, and a critical aspect of CAR-T development was the incorporation of costimulatory ITAMs into the CAR structure<sup>32</sup>. Several independent groups validated that adding CD28 or CD137 endodomains to the first generation scFv-CD3 $\zeta$  structure improved proliferation and often in vivo efficacy<sup>33</sup>. However, the field's understanding of ex vivo manufacturing of genetically modified T cells has evolved since these studies<sup>34</sup>. Early models of ex vivo expansion focused on tumor-infiltrating lymphocyte models, where T cells are expanded to large numbers with high dose IL-2 and anti-CD3<sup>34</sup>. Many findings of improved costimulatory CAR structure were done with similar, now obsolete, ex vivo culture methods that did not include CD28 costimulation<sup>34</sup>. Current *ex vivo* CAR-T manufacturing includes simultaneous CD3/CD28 stimulation with or without supplemental cytokines such as IL-2, IL-7 or IL-15<sup>35,36</sup>. The addition of costimulation, whether in the CAR structure or during ex vivo manufacturing, is required for CAR-T efficacy. However, it is not precisely clear if the CD28 costimulatory signal is needed during CAR engagement or if CD28 stimulation during T-cell expansion is sufficient. It is worth noting that the clinical efficacy observed with blinatumomab, a CD3/CD19 bispecific antibody that is FDA-approved for B-cell ALL, suggests that engaging T cells without additional costimulation can be sufficient for anti-tumor activity.

When we tested the anti-tumor activity of AbTCR-T cells both in vitro and with tumor xenograft models, AbTCR-T cells maintain comparable anti-tumor potency to ET190L1-CAR and CD137-based CTL019-T cells (Figs. 3c, 5a-c). And yet activation of T-cells through ET190L1-AbTCR results in lower concentrations of inflammatory cytokines and a less exhausted T cell

(Figs. 3e, 4a, b, 5d, e). While the engagement through the AbTCR receptor may form an immune synapse that incorporates endogenous costimulatory molecules, there are no fused costimulatory endodomains on the AbTCR platform. Therefore, it is possible that some of the reduction in cytokine release could be due to an AbTCR-T cell with reduced costimulation signal input compared to a CD28/CD137 second generation CAR that has directly fused costimulation and CD3 $\zeta$  domains. However, we hypothesize that the AbTCR's reduction in cytokine release is primarily attributable to the  $\gamma\delta$ TCR effector domain which utilizes more of the T cell's endogenous feedback mechanisms once sufficient T cell-mediated killing has occurred. This hypothesis is supported from our experiments using CD80<sup>+</sup>/CD86<sup>+</sup> Raji tumor models, which provides comparable CD28 costimulation to both CAR-T and AbTCR-T cells. Figs 3e, 4a, b, and 5d, e demonstrated that AbTCR-T cells release dramatically less inflammatory cytokines and produce less exhaustion markers while lysing Raji cells. Moreover, regardless of costimulation from the tumor cell, anti-CD19 AbTCR-T cells had similar anti-tumor activity compared to CAR-T cells in both Raji (Figs. 3c, 5a, b, Supplementary Fig. S4) and CD80<sup>+</sup>/CD86<sup>+</sup> (Figs. 3c, 5b, c) tumor models.

Two of the major side effects associated with current CAR-T cell therapy are CRS and cerebral edema/neurotoxicity<sup>37</sup>. Current mitigation strategies include patient stratification according to disease type and tumor burden, and close monitoring of patients so that high grade CRS/neurotoxicity can be avoided (typically with the combined use of the IL6R inhibitor tocilizumab and steroids)<sup>38,39</sup>. However, treating high-grade cytokine-related toxicities in patients with high disease burdens often requires intensive care unit-level care, increasing both risk and costs. In this work, we show that anti-CD19 AbTCR-T cells functionally match CD28-based CAR-T cells engineered with the same anti-CD19 binding moiety, as well as CD137-based CTL019 CAR-T cells, but release drastically less cytokines upon killing of target-positive tumors both in vitro and in pre-clinical mouse models. Whether these pre-clinical findings for AbTCR translate into the clinical setting is currently being tested (clinicaltrials.gov, NCT03379493).

## Methods

### Cell lines

Cell lines were cultured according to manufacturer's recommendations. The Raji-luc cell line was purchased from Comparative Biosciences, Inc. (Sunnyvale, CA). NALM-6-luc was a generous gift from Dr. Eric Smith (Memorial Sloan Kettering Cancer Center). K562-CD19 is a stable cell line that was engineered to express CD19. Raji CD19 K/O is a CRISPR knockout cell line that lacks CD19 expression<sup>40,41</sup>. The CHP105R1 primary leukemia cell line

(CD19-positive, CD80 and CD86-negative) was created by implanting primary human B-Cell Precursor ALL samples in NSG mice, transducing the expanded cells ex-vivo with luciferase-encoding lentivirus, and serially passaging the cells. Detailed methods are described<sup>42</sup>. Monocytes used in the co-culture assay are a mixture of monocytes, macrophages, immature dendritic cells and mature dendritic cells derived from human peripheral blood monocytes as previously described<sup>28</sup>.

### Generation of CAR and AbTCR-T cells

The anti-CD19 scFv (ET190L1) was identified from Eureka Therapeutics human E-ALPHA<sup>®</sup> phage display library (see Supplemental Methods). ET190L1-CAR is a Myc-tagged second-generation CAR containing the CD28/CD3 $\zeta$  domains. AbTCR is derived from the human TCR gamma (UniProtKB: locus TRGC1\_HUMAN, accession P0CF51) and delta (UniProtKB: locus TRDC\_HUMAN, accession B7Z8K6) chains. CAR and AbTCR constructs were cloned into a 3rd generation pCDH lentiviral vector (Systems Biosciences) for delivery into T cells. Healthy human donor peripheral blood leukocytes were obtained from Blood Centers of the Pacific.

T cells were isolated using negative selection and stimulated with CD3/CD28 Dynabeads (Thermo Fisher Scientific). After 24 h. activated T cells were transduced with lentivirus at an MOI of 2–5, and cultured in RPMI with 10% FBS, IL-7 (10 ng/ml), and IL-15 (5 ng/ml) as previously described<sup>9</sup>. Transduction efficiency for CAR or AbTCR positive T cells was determined by flow cytometry using an anti-Myc antibody (Cell Signaling) or an anti-F(ab')<sub>2</sub> antibody (Jackson ImmunoResearch), respectively. CTL019 T cells and experimentally comparative AbTCR-T cells were generated as previously described<sup>33</sup>.

The percent of receptor positive T cells were normalized with donor-matched untransduced (Mock) T cells for all experiments directly comparing CAR and AbTCR. All comparative results with ET190L1-AbTCR and ET190L1-CAR were observed with no less than 3 donors.

### Co-immunoprecipitation

J.RT3-T3.5 cells were either mock-transduced or transduced with 3xFLAG-tagged AbTCR construct. 7 days after transduction, cells were lysed with digitonin (0.1%) lysis buffer and an anti-FLAG antibody (Sigma) was used to immunoprecipitate. A Western blot was run on the cell lysates or anti-FLAG immunoprecipitates and detected for CD3 $\delta$ ,  $\epsilon$ ,  $\gamma$  and  $\zeta$  chains. Antibodies for CD3 $\delta$ , CD3 $\epsilon$ , CD3 $\gamma$ , and CD3 $\zeta$  were purchased from BioLegend.

### Cytokine assays

Cytokine release in culture supernatants and mouse serum was measured using the Bio-plex Pro Human

Cytokine 8-plex assay (BioRad). Samples in Fig. 4a were pooled from three replicates and measured with a Bio-plex200 (Luminex). Intracellular cytokine staining of cytokines was conducted after 4 h of co-culturing ET190L1-AbTCR-T cells with target cells at an effector-to-target (E:T) ratio of 2:1 in the presence of 5 mg/ml of Brefeldin A. Trans-well co-culture cytokine assays were conducted as previously described<sup>28</sup>.

### Mouse xenograft tumor model

Animal experiments were conducted at either Children's Hospital of Philadelphia (CHOP) or Lumigenics (Richmond, CA). Female NSG mice aged 6–8 weeks were used. For Raji lymphoma models and tumor re-challenge experiments,  $5 \times 10^5$  tumor cells were intravenously injected in PBS and tumors were measured by total bioluminescent flux using a Xenogen Imaging System (Perkin Elmer). Peripheral blood was collected via the submandibular vein. For the Raji re-challenge mice were re-challenged 5 weeks after T cell infusion. For the NALM-6 re-challenge mice were re-challenged 4 weeks after T cell infusion. Mouse studies with PDX and engineered T cells used in those studies were performed at CHOP as previously described<sup>36,43</sup>.

### Statistical analysis

Statistical analyses were performed using Prism GraphPad software. For studies comparing two groups, we used a Student *t* test. For studies with multiple groups, we used a one-way ANOVA followed by Dunnett test which accounts for multiple comparisons. All analyses were two-tailed.

### Acknowledgements

We acknowledge Ed Lim (Lumigenics) for the management of Raji-luc and NALM-6-luc animal studies. The animal facilities at CHOP for carrying out the CHP105R1 PDX mouse studies. Yoko Nakano, Hongruo Yun, Ziyou Cui, and Lu Jin for technical help with experiments. Timothy Acker for assistance in the writing and review of the manuscript. Pei Wang for review of the manuscript. Julie Storm for her excellent technical assistance.

### Author contributions

Y.X., Z.Y., L.H.H., P.Z., L.L., B.Z., S.G., J.L., D.M.B., and H.L. designed and performed research, and analyzed and interpreted data. J.F.M. performed research. V.W.C., C.L. and S.A.G. designed, analyzed and interpreted data. B.Z., L.H.H., S.G., J.F.M., and V.W.C. wrote the manuscript.

### Conflict of interest

Y.X., Z.Y., L.H.H., P.Z., L.L., B.Z., S.G., J.L., J.F.M., V.W.C., H.L., and C.L. are employees of and have equity ownership and/or stock options in Eureka Therapeutics, Inc. S.A.G. is a scientific advisor to Eureka Therapeutics with stock options in the company. D.M.B. is a consultant to Eureka Therapeutics. Other associations: S.A.G. has received research and/or clinical trial support from Novartis, Servier and Kite. He consults for, has participated in ad boards, or serves on study steering committees or scientific advisory boards for the following companies: Novartis, Adaptimmune, TCR2, Juno, GlaxoSmithKline, Cellectis, Vertex, Roche and Janssen. He is listed on a patent related to toxicity management in cell therapy managed by University of Pennsylvania and CHOP policies.

### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Supplementary Information** accompanies the paper at (<https://doi.org/10.1038/s41421-018-0066-6>).

Received: 1 August 2018 Revised: 27 September 2018 Accepted: 28 September 2018

Published online: 20 November 2018

### References

- Oluwole, O. O. & Davila, M. L. At The Bedside: Clinical review of chimeric antigen receptor (CAR) T cell therapy for B cell malignancies. *J. Leukoc. Biol.* **100**, 1265–1272 (2016).
- Maude, S. L. et al. Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N. Engl. J. Med.* **378**, 439–448 (2018).
- Gross, G., Gorochoy, G., Waks, T. & Eshhar, Z. Generation of effector T cells expressing chimeric T cell receptor with antibody type-specificity. *Transplant. Proc.* **21**, 127–130 (1989).
- Eshhar, Z., Waks, T., Gross, G. & Schindler, D. G. Specific activation and targeting of cytotoxic lymphocytes through chimeric single chains consisting of antibody-binding domains and the gamma or zeta subunits of the immunoglobulin and T-cell receptors. *Proc. Natl. Acad. Sci. USA* **90**, 720–724 (1993).
- Shankaran, V. et al. IFN $\gamma$  and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature* **410**, 1107–1111 (2001).
- Zou, C. et al. gammadelta T cells in cancer immunotherapy. *Oncotarget* **8**, 8900–8909 (2017).
- Guy, C. S. & Vignali, D. A. Organization of proximal signal initiation at the TCR-CD3 complex. *Immunol. Rev.* **232**, 7–21 (2009).
- Johnson, L. A. & June, C. H. Driving gene-engineered T cell immunotherapy of cancer. *Cell Res.* **27**, 38–58 (2017).
- Liu, H. et al. Targeting alpha-fetoprotein (AFP)-MHC complex with CAR T-cell therapy for liver cancer. *Clin. Cancer Res.* **23**, 478–488 (2017).
- Harris, D. T. et al. Comparison of T cell activities mediated by human TCRs and CARs that use the same recognition domains. *J. Immunol.* **200**, 1088–1100 (2018).
- Oren, R. et al. Functional comparison of engineered T cells carrying a native TCR versus TCR-like antibody-based chimeric antigen receptors indicates affinity/avidity thresholds. *J. Immunol.* **193**, 5733–5743 (2014).
- Davenport, A. J. et al. Chimeric antigen receptor T cells form nonclassical and potent immune synapses driving rapid cytotoxicity. *Proc. Natl. Acad. Sci. USA* **115**, E2068–E2076 (2018).
- Koning, F., Maloy, W. L., Cohen, D. & Coligan, J. E. Independent association of T cell receptor beta and gamma chains with CD3 in the same cell. *J. Exp. Med.* **166**, 595–600 (1987).
- Saito, T. et al. Surface expression of only gamma delta and/or alpha beta T cell receptor heterodimers by cells with four (alpha, beta, gamma, delta) functional receptor chains. *J. Exp. Med.* **168**, 1003–1020 (1988).
- Ren, J. et al. Multiplex genome editing to generate universal CAR T cells resistant to PD1 inhibition. *Clin. Cancer Res.* **23**, 2255–2266 (2017).
- Ren, J. et al. A versatile system for rapid multiplex genome-edited CAR T cell generation. *Oncotarget* **8**, 17002–17011 (2017).
- Singh, N., Perazzelli, J., Grupp, S. A. & Barrett, D. M. Early memory phenotypes drive T cell proliferation in patients with pediatric malignancies. *Sci. Transl. Med.* **8**, 320ra323 (2016).
- Fraietta, J. A. et al. Biomarkers of response to anti-CD19 chimeric antigen receptor (CAR) T-cell therapy in patients with chronic lymphocytic leukemia. *Blood* **128**, 57–57 (2016).
- Mahnke, Y. D., Brodie, T. M., Sallusto, F., Roederer, M. & Lugli, E. The who's who of T-cell differentiation: human memory T-cell subsets. *Eur. J. Immunol.* **43**, 2797–2809 (2013).
- Long, A. H. et al. 4-1BB costimulation ameliorates T cell exhaustion induced by tonic signaling of chimeric antigen receptors. *Nat. Med.* **21**, 581–590 (2015).

21. Gomes-Silva, D. et al. Tonic 4-1BB costimulation in chimeric antigen receptors impedes T cell survival and is vector-dependent. *Cell Rep.* **21**, 17–26 (2017).
22. Kalos, M. et al. T cells with chimeric antigen receptors have potent antitumor effects and can establish memory in patients with advanced leukemia. *Sci. Transl. Med.* **3**, 95ra73 (2011).
23. Kalos, M. & June, C. H. Adoptive T cell transfer for cancer immunotherapy in the era of synthetic biology. *Immunity* **39**, 49–60 (2013).
24. Teachey, D. T. et al. Identification of predictive biomarkers for cytokine release syndrome after chimeric antigen receptor T-cell therapy for acute lymphoblastic leukemia. *Cancer Discov.* **6**, 664–679 (2016).
25. Grupp, S. A. et al. Chimeric antigen receptor-modified T cells for acute lymphoid leukemia. *N. Engl. J. Med.* **368**, 1509–1518 (2013).
26. Davila, M. L. et al. Efficacy and toxicity management of 19-28z CAR T cell therapy in B cell acute lymphoblastic leukemia. *Sci. Transl. Med.* **6**, 224ra225 (2014).
27. Lee, D. W. et al. T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet* **385**, 517–528 (2015).
28. Singh, N. et al. Monocyte lineage-derived IL-6 does not affect chimeric antigen receptor T-cell function. *Cytotherapy* **19**, 867–880 (2017).
29. Grupp, S. A. Advances in T-cell therapy for ALL. *Best. Pract. Res. Clin. Haematol.* **27**, 222–228 (2014).
30. Nicolas, L. et al. Human gammadelta T cells express a higher TCR/CD3 complex density than alphabeta T cells. *Clin. Immunol.* **98**, 358–363 (2001).
31. Rothlisberger, D., Honegger, A. & Pluckthun, A. Domain interactions in the Fab fragment: a comparative evaluation of the single-chain Fv and Fab format engineered with variable domains of different stability. *J. Mol. Biol.* **347**, 773–789 (2005).
32. Brentjens, R. J. et al. Genetically targeted T cells eradicate systemic acute lymphoblastic leukemia xenografts. *Clin. Cancer Res.* **13**, 5426–5435 (2007).
33. Milone, M. C. et al. Chimeric receptors containing CD137 signal transduction domains mediate enhanced survival of T cells and increased antileukemic efficacy in vivo. *Mol. Ther.* **17**, 1453–1464 (2009).
34. Klapper, J. A. et al. Single-pass, closed-system rapid expansion of lymphocyte cultures for adoptive cell therapy. *J. Immunol. Methods* **345**, 90–99 (2009).
35. Gardner, R. A. et al. Intent-to-treat leukemia remission by CD19 CAR T cells of defined formulation and dose in children and young adults. *Blood* **129**, 3322–3331 (2017).
36. Barrett, D. M. et al. Relation of clinical culture method to T-cell memory status and efficacy in xenograft models of adoptive immunotherapy. *Cytotherapy* **16**, 619–630 (2014).
37. Gust, J. et al. Endothelial activation and blood-brain barrier disruption in neurotoxicity after adoptive immunotherapy with CD19 CAR-T cells. *Cancer Discov.* **7**, 1404–1419 (2017).
38. Lee, D. W. et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood* **124**, 188–195 (2014).
39. Maude, S. L., Barrett, D., Teachey, D. T. & Grupp, S. A. Managing cytokine release syndrome associated with novel T cell-engaging therapies. *Cancer J.* **20**, 119–122 (2014).
40. Jinek, M. et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821 (2012).
41. Ran, F. A. et al. Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**, 2281–2308 (2013).
42. Sotillo, E. et al. Convergence of acquired mutations and alternative splicing of CD19 enables resistance to CART-19 immunotherapy. *Cancer Discov.* **5**, 1282–1295 (2015).
43. Barrett, D. M. et al. Noninvasive bioluminescent imaging of primary patient acute lymphoblastic leukemia: a strategy for preclinical modeling. *Blood* **118**, e112–e117 (2011).