

A Novel Crown-Ether Stabilized Oxonium Halogenochalcogenate(IV): [H₇O₃(Bis-dibromo-dibenzo-30-crown-10)][Se₂Br₉] \cdot 1.5CH₂Cl₂

Adrienne Hammerschmidt, Illenora Beckmann, Mechtild Lage, and Bernt Krebs

Institut fur Anorganische und Analytische Chemie, Westfalische Wilhelms-Universitat,
Corrensstr. 30, D-48149 Munster, Germany

Reprint requests to Prof. Dr. B. Krebs. Fax: +49 (0)251 8338366. E-mail: krebs@uni-muenster.de

Z. Naturforsch. **59b**, 1438 – 1443 (2004); received August 24, 2004

Dedicated to Professor Hubert Schmidbaur on the occasion of his 70th birthday

The title compound [H₇O₃(bis-dibromo-dibenzo-30-crown-10)][Se₂Br₉] \cdot 1.5CH₂Cl₂ (**1**) was isolated from a solution of SeBr₄ and dibenzo-30-crown-10 in CH₃CN/CH₂Cl₂ containing a small amount of hydrobromic acid. During the reaction the crown ether is brominated by HBr.

The compound crystallizes in the monoclinic space group *P*2₁/*n* with *a* = 17.688(11), *b* = 14.921(6), *c* = 20.521(12) , β = 97.71(5)°, and *Z* = 4. **1** is a novel example in the series of supramolecular halogenochalcogeno acids prepared in our group in which different oxonium cations are stabilized and encapsulated by crown ethers. Especially in this class of superacids complexation by cyclic polyethers offers convenient and variable possibilities for the controlled synthesis of oxonium cations. In the present case the large dibenzo-30-crown-10 ring systems are able to stabilize trinuclear [H₇O₃]⁺ cations within their cavities. Besides the macromolecular cations and some dichloromethane solvent molecules, the crystal structure of **1** contains molecular [Se₂Br₉][−] ions with approximate D_{3h} symmetry, each consisting of two face-sharing SeBr₆ octahedra.

Key words: Crystal Structure, Chalcogenates, Oxonium Ions, Crown Ether

Introduction

The structural properties of halogenochalcogenates(IV) can be related to the cubane structures of the tetrahalides. Adding halides results in a stepwise degradation by nucleophilic attack of the tetrameric structure leading to the tri-, di- and monomeric species [Q₃X₁₃][−], [Q₂X₁₀]^{2−} and [QX₆]^{2−}, all based on QX₆-octahedra as building blocks (Q = Se, Te; X = Cl, Br). In the oligomeric species the octahedra are connected *via* common edges. Their structures and properties are well known [1–3]. The size of the resulting anions mainly depends on the solvent, the concentration of the halides and the size of the counterions, respectively.

The presence of anions that do not fit into this scheme of decomposition indicate complex equilibria in solution. The formation of the monomeric [QX₅][−] anions *e.g.* in the compounds [C₈H₈S₂N][TeCl₅] \cdot 1.5C₄H₈O₂ [4] may be caused by ligand exchange in solution. The compounds [PCl₄]_{*n*}[TeCl₅]_{*n*} [5, 6] and [H₉O₄]_{*n*}[Te₂Cl₉]_{*n*} [7] contain polymeric anionic species. The [TeCl₅]_{*n*}[−] anion consists of TeCl₆

octahedra connected *via* common edges whereas in the [Te₂Cl₉]_{*n*}[−] anion TeCl₆ octahedra are alternatingly connected *via* common edges and corners.

In solvents containing aqueous hydrogen halides the chalcogen tetrahalides form halogeno chalcogeno acids. They can be classified as superacids and are completely dissociated in solutions as well as in the solid state. Thus they should be described as oxonium halogenochalcogenates(IV). In more concentrated solutions of these systems the oxonium ions form clusters such as [H₇O₃]⁺ or [H₉O₄]⁺. So far crystalline compounds such as [H₉O₄]₂[TeBr₆] [3], [H₇O₃]₂[TeI₆] \cdot 2H₂O [8], [H₉O₄][TeCl₄OH] \cdot H₂O [7], [H₉O₄][Te₃Br₁₃] [3] and [H₉O₄]_{*n*}[Te₂Cl₉]_{*n*} [7] have been reported. Organic molecules with electron donating groups can be used to stabilize oxonium ions. For instance in the compounds [H₃O][TeBr₅] \cdot 3C₄H₈O₂ [9] and [H₅O₂][Te₂Cl₉] \cdot 2C₄H₈O₂ [10] the oxonium cations are coordinated by three 1,4-dioxane molecules. If the electron donating group is more basic than H₂O the protonated organic molecules can be isolated as counter ions of the halogenochalcogenates(IV). This

was shown for *e.g.* $[(\text{CH}_3)_2\text{SOH}]_2[\text{TeCl}_6] \cdot 2(\text{CH}_3)_2\text{SO}$ and $[(\text{CH}_3)_2\text{CHC}(\text{NH}_2)(\text{OH})][\text{Te}_3\text{Cl}_3] \cdot (\text{CH}_3)_2\text{CHCN}$ [2, 11].

Utilization of crown ethers is another convenient method to stabilize oxonium cations in halogenochalcogenates. Compounds within this class of materials which have been prepared include $[\text{H}_3\text{O}(\text{dibenzo-18-crown-6})][\text{Te}_2\text{Br}_9]$ [10], $[\text{H}_3\text{O}(\text{dibromobenzo-15-crown-5})]_2[\text{SeBr}_6]$ [17], $[\text{H}_3\text{O}(\text{benzo-18-crown-6})]_2[\text{Te}_2\text{Br}_{10}]$ $[\text{H}_5\text{O}_2(\text{dibenzo-24-crown-8})]_2[\text{Te}_2\text{Br}_{10}]$ [18] and $[\text{H}_5\text{O}_2(\text{bis-dibromo-dibenzo-24-crown-8})]_2[\text{Se}_3\text{Br}_8]$ [19, 20]. They have been synthesized and characterized in our research group.

Here we report the novel compound $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ (**1**) where the oxonium ions are encapsulated by especially large crown ether ligands.

Experimental Section

Synthesis

$[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$: 0.5 g SeBr_4 (1.254 mmol) was dissolved in a mixture of 5 ml $\text{CH}_3\text{CN}/\text{CH}_2\text{Cl}_2$ and HBr (0.07 ml, 48%). The solution is heated and dibenzo-30-crown-10 (12.5 mg, 0.024 mmol) was added. After cooling down to -18°C red crystals suitable for single crystal X-ray diffraction have been obtained by careful recrystallization, *m.p.*: 95°C .

Crystal structure determination

For the data collections single crystals of the compound were measured on a SIEMENS P3 four-circle diffractometer under a nitrogen atmosphere. The space group was determined from systematic absences and intensity statistics. $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ crystallizes in space group $P2_1/n$. The structure solution was achieved *via* direct methods using the SHELXTL PLUS software [12]; full-matrix least-squares refinements were carried out using the SHELXL-93 programme. The starting coordinates of the Se and Br atoms were obtained from the initial E-maps, all other non-hydrogen atoms were derived from subsequent difference Fourier syntheses. Se, Br, Cl, O, and C atoms were refined with anisotropic displacement parameters. All atoms are on general positions of the space group.

Aromatic and aliphatic hydrogen atom positions in the organic part were calculated at idealized coordinates and were refined in a restrained system coupled to the adjacent carbon atoms. The approximate coordinates of the H atoms in the oxonium cations could be obtained from careful evaluation

Table 1. Crystal data and details of structure refinement for $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ (**1**).

Empirical formula	$\text{C}_{29.5}\text{O}_{13}\text{H}_{46}\text{Br}_{13}\text{Cl}_3\text{Se}_2$ (1)
Formula weight	1870.20 g/mol
Temperature	153(2) K
Diffractometer	SIEMENS P3
Wavelength	0.71073
Crystal system	monoclinic
Space group	$P2_1/n$ (no. 14)
Unit cell dimensions	$a = 17.688(11) \text{ \AA}$ $b = 14.921(6) \text{ \AA}$ $c = 20.521(12) \text{ \AA}$ $\beta = 97.71(5)^\circ$
Volume	$5372(1) \text{ \AA}^3$
Z	4
Density (calculated)	2.314 g/cm^3
Crystal size	$0.03 \times 0.04 \times 0.20 \text{ mm}^3$
Theta range for data collection	4.0 to 54.0°
Index ranges	$0 \leq h \leq 18$, $0 \leq k \leq 15$, $-23 \leq l \leq 21$
Reflections collected	6759
Independent reflections	2648
Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	2648 / 0 / 401
Goodness-of-fit on F^2	0.970
Final R indices [$I > 2\sigma(I)$]	$R1 = 0.0782$, $wR2 = 0.1266$
R Indices (all data)	$R1 = 0.2263$, $wR2 = 0.1849$
Largest diff. peak and hole	1.715 and -1.135 e\AA^{-3}

of the final difference Fourier maps. They were fully in accordance with chemical arguments considering the assignment of the H_3O group to O(3W) and considering the hydrogen bridge system to the crown ether oxygen atoms. However, they were not refined independently, but rigidly coupled to O(1W) (as H_2O), O(2W) (as H_2O), and O(3W) (as pyramidal H_3O) with a fixed O-H distance of 0.90 \AA .

Besides the well-defined system of $[\text{Se}_2\text{Br}_9]^-$ anions and supramolecular crown ether-oxonium cations the crystal contains a rather ill-defined region of dichloromethane solvate molecules strongly disordered over two sites. In addition, on both sites the Cl atoms of the CH_2Cl_2 species are disordered internally so that a precise localization was difficult. Even at the data collection temperature of 153 K partial loss of crystal solvent from these disordered sites could not be avoided. So, according to the refinement of the occupational factors, a residual content of 1.5 CH_2Cl_2 per formula unit was assumed throughout the structure determination.

The parameters of the data collection as well as details of the structure solution and refinement are summarized in Table 1.

Crystallographic data for the structure have been deposited with the Cambridge Crystallographic Data Centre, CCDC-247194 for $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$.




Fig. 1. Projection of the unit cell in $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ along $[100]$, hydrogen atoms omitted.




Fig. 2. Projection of the unit cell in $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ along $[010]$, hydrogen atoms omitted.

10)][Se_2Br_9] $\cdot 1.5\text{CH}_2\text{Cl}_2$. Copies of the data can be obtained free of charge on application to The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: int.code+(1223)336-033; e-mail for inquiry: fileserv@ccdc.cam.ac.uk).

Results and Discussion

The crystal structure of $[\text{H}_7\text{O}_3(\text{bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ (**1**) consists of a supramolecular arrangement of molecular $[\text{Se}_2\text{Br}_9]^-$ anions and $[\text{H}_7\text{O}_3(\text{bis-dibromo-dibenzo-30-}$




Fig. 3. Dimeric $[\text{Se}_2\text{Br}_9]^-$ anion in $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5\text{CH}_2\text{Cl}_2$ in an ellipsoidal representation (50% probability).

$\text{crown-10}]^+$ cations. In addition the crystals contain some dichloromethane solvent molecules in a disordered manner. Fig. 1 shows a projection of the unit cell along $[100]$, in Fig. 2 the projection along $[010]$ gives a more detailed view on the cations.

The dinuclear anionic unit is formed by condensation of two axially distorted octahedral SeBr_6 entities *via* a common face of three bromine atoms. The anion has D_{3h} symmetry (see Fig. 3). The average Se-Br bond length in **1** (2.602 Å, for details see Table 2) is comparable to the mean Se-Br bond length in the $[\text{SeBr}_6]^{2-}$ anion [*e.g.* 14] with 2.565 Å. The $\text{Se-Br}_{(\text{term})}$ bond lengths (mean value 2.41 Å) are significantly shorter than the $\text{Se-}\mu_2\text{Br}$ values (mean 2.79 Å). These data are in good accordance with other known Se-Br distances for terminal and bridging μ_2 bonds [1]. Moreover the single distances show a clear correlation between the Se-Br bond lengths *trans* to each other (*e.g.* the longest bridge bond $\text{Se}(2)\text{-Br}(1)$ is *trans* to the shortest terminal bond $\text{Se}(2)\text{-Br}(7)$, Table 2). This phenomenon is well known for hypervalent systems of the present kind and is well in accordance with the three-centre-four-electron model.

The short bonds of the terminal bromine atoms to the selenium atoms result in larger mutual repulsion of the bromine atoms resulting in an expansion of the $\text{Br}_{(\text{term})}\text{-Se-Br}_{(\text{term})}$ bond angles (mean 94.2°) and a compression of the $\mu_2\text{Br-Br-}\mu_2\text{Br}$ bond angles (mean 81.9°). The structure within the terminal SeBr_3 units is similar to the one found in tetrameric $[\text{SeBr}_4]_4$ [16]. In the limiting case of an ionic description the anion in **1** can be regarded as $[(\text{SeBr}_3^+)_2(\text{Br}^-)_3]$. This is analogous to the formulation of the tetrabromide $[(\text{SeBr}_3^+)_4\text{Br}_4]$.

It is highly interesting to observe a number of rather short intermolecular $\text{Br}\cdots\text{Br}$ contacts with distances

Table 2. Bond lengths [Å] and angles [°] with standard deviations for [H₇O₃(Bis-dibromo-dibenzo-30-crown-10)] [Se₂Br₉]·1.5CH₂Cl₂ (1).

Se(1)-Br(1)	2.732(4)	Se(2)-Br(1)	2.902(5)	Br(3)-Se(1)-Br(5)	89.9(2)	Br(7)-Se(2)-Br(8)	94.2(2)
Se(1)-Br(2)	2.810(4)	Se(2)-Br(2)	2.748(4)	Br(3)-Se(1)-Br(6)	174.9(2)	Br(7)-Se(2)-Br(9)	94.2(2)
Se(1)-Br(3)	2.737(4)	Se(2)-Br(3)	2.811(5)	Br(4)-Se(1)-Br(5)	94.5(2)	Br(8)-Se(2)-Br(9)	94.6(2)
Se(1)-Br(4)	2.440(4)	Se(2)-Br(7)	2.372(5)	Br(4)-Se(1)-Br(6)	94.1(2)		
Se(1)-Br(5)	2.410(5)	Se(2)-Br(8)	2.418(5)				
Se(1)-Br(6)	2.433(5)	Se(2)-Br(9)	2.407(5)	H ₇ O ₃ cation:			
Se(1)···Se(2)	3.623(5)	Br(2)···Br(9)	3.698(5)	O(1W)···O(3W)	2.477(22)	O(2W)···O(4)	2.757(21)
Br(1)···Br(2)	3.653(5)	Br(3)···Br(4)	3.639(5)	O(1W)···O(1)	3.076(23)	O(2W)···O(5)	3.087(21)
Br(1)···Br(3)	3.677(5)	Br(3)···Br(5)	3.645(5)	O(1W)···O(2)	2.807(25)	O(2W)···O(6)	2.841(21)
Br(1)···Br(5)	3.705(6)	Br(3)···Br(7)	3.670(5)	O(1W)···O(3)	3.046(24)	O(2W)···O(7)	3.125(22)
Br(1)···Br(8)	3.913(6)	Br(3)···Br(8)	3.721(5)	O(1W)···O(9)	2.872(27)	O(3W)···O(3)	3.453(23)
Br(1)···Br(9)	3.934(6)	Br(4)···Br(5)	3.560(5)	O(1W)···O(10)	3.059(24)	O(3W)···O(8)	2.654(21)
Br(2)···Br(3)	3.695(5)	Br(4)···Br(6)	3.567(5)	O(2W)···O(3W)	2.487(22)	O(3W)···O(9)	3.173(25)
Br(2)···Br(4)	3.737(5)	Br(5)···Br(6)	3.534(6)	O(1W)-O(3W)-O(2W)	119.2(8)	O(3W)-O(2W)-O(4)	113.3(7)
Br(2)···Br(6)	3.796(5)	Br(7)···Br(8)	3.508(6)	O(1W)-O(3W)-O(8)	116.1(8)	O(3W)-O(2W)-O(6)	128.5(8)
Br(2)···Br(7)	3.631(5)	Br(7)···Br(9)	3.501(5)	O(2W)-O(3W)-O(8)	111.8(8)	O(2)-O(1W)-O(9)	154.4(9)
Br(8)···Br(9)	3.543(7)			O(3)-O(3W)-O(8)	173.8(8)	O(3W)-O(1W)-O(9)	72.4(7)
				O(4)-O(2W)-O(6)	107.4(7)	O(3W)-O(1W)-O(2)	132.4(8)
intermolecular distances:				H(1W)-O(1W)	0.900	H(1W)···O(2)	2.154
Br(1)···Br(11a)	3.453(4)	Br(6)···Br(6d)	3.613(7)	H(2W)-O(1W)	0.900	H(2W)···O(9)	2.314
Br(2)···Br(9b)	3.913(5)	Br(3)···Br(10d)	3.857(5)	H(3W)-O(3W)	1.200	H(3W)···O(1W)	1.324
Br(7)···Br(12c)	3.348(4)			H(4W)-O(3W)	1.100	H(4W)···O(2W)	1.418
Se(1)-Br(1)-Se(2)	80.0(2)	Br(5)-Se(1)-Br(6)	93.7(2)	H(5W)-O(3W)	0.930	H(5W)···O(8)	1.750
Se(1)-Br(2)-Se(2)	81.4(2)	Br(1)-Se(2)-Br(2)	80.6(2)	H(6W)-O(2W)	0.900	H(6W)···O(6)	1.994
Se(1)-Br(3)-Se(2)	81.6(2)	Br(1)-Se(2)-Br(3)	80.1(1)	H(7W)-O(2W)	0.900	H(7W)···O(4)	1.870
Br(1)-Se(1)-Br(2)	82.5(2)	Br(1)-Se(2)-Br(7)	166.9(2)	H(1W)-O(1W)-H(2W)	86.7	H(4W)-O(2W)-H(7W)	107.1
Br(1)-Se(1)-Br(3)	84.5(1)	Br(1)-Se(2)-Br(8)	94.3(2)	H(1W)-O(1W)-H(3W)	136.3	H(3W)-O(3W)-H(4W)	104.6
Br(1)-Se(1)-Br(4)	171.0(2)	Br(1)-Se(2)-Br(9)	95.2(2)	H(2W)-O(1W)-H(3W)	113.2	H(3W)-O(3W)-H(5W)	106.5
Br(1)-Se(1)-Br(5)	92.0(2)	Br(2)-Se(2)-Br(3)	83.4(2)	H(4W)-O(2W)-H(6W)	107.4	H(4W)-O(3W)-H(5W)	109.2
Br(1)-Se(1)-Br(6)	91.9(2)	Br(2)-Se(2)-Br(7)	90.1(2)	H(6W)-O(2W)-H(7W)	111.8		
Br(2)-Se(1)-Br(3)	83.6(2)	Br(2)-Se(2)-Br(8)	172.4(2)	O(1W)-H(1W)···O(2)	129.2	O(3W)-H(5W)···O(8)	166.3
Br(2)-Se(1)-Br(4)	90.5(2)	Br(2)-Se(2)-Br(9)	91.4(2)	O(1W)-H(2W)···O(9)	120.2	O(2W)-H(6W)···O(8)	156.3
Br(2)-Se(1)-Br(5)	171.8(2)	Br(3)-Se(2)-Br(7)	89.8(2)	O(3W)-H(3W)···O(1W)	158.5	O(2W)-H(7W)···O(4)	168.4
Br(2)-Se(1)-Br(6)	92.5(1)	Br(3)-Se(2)-Br(8)	90.5(2)	O(3W)-H(4W)···O(2W)	161.3		
Br(3)-Se(1)-Br(4)	89.2(2)	Br(3)-Se(2)-Br(9)	173.5(2)				

Symmetry transformations used to generate equivalent atoms: a) $x+1/2, -y+1/2, z+1/2$; b) $-x+1, -y+2, -z+1$; c) $x+1, y+1, z$; d) $-x+1, -y+1, -z+1$.

well below the formal sum of van der Waals radii for bromine (Table 2). As in a number of other halogenochalcogenates (*e. g.* [1]) this is an indication of significant charge transfer bonding interactions between different molecular units. In **1** the most significant interactions are between the [Se₂Br₉][−] units and Br atoms of the brominated crown ether molecules, the smallest distance of 3.340 Å being between Br(7) and Br(12c), (Table 2).

The hydrophilic cavity within the thirty-membered macrocycle of (bis-dibromo-dibenzo-30-crown-10) is large enough to encapsulate a [H₇O₃]⁺ cation (see Fig. 4). The central oxonium oxygen atom is coordinated pyramidally by two water molecules and a crown ether oxygen. The O···O distances (mean 2.48 Å) for O(1W)···O(3W) and O(2W)···O(3W) (Table 2) are

typical for [H₇O₃]⁺ entities. The distance between the central oxonium oxygen and the crown-ether acceptor oxygen atom O(8) is 2.654 Å. Thus, the [H₇O₃]⁺ entity can be regarded as a distorted [H₃O]⁺ cation surrounded by two water molecules. This description is supported by the qualitative location of the hydrogen atoms in the [H₇O₃]⁺ unit, clearly showing three H atoms in bonding distance to O(3W). Moreover, the O···O hydrogen bond distances from the outer water molecules O(1W) and O(2W) of the [H₇O₃]⁺ unit to crown ether oxygen atoms are normal OH_(water)···O values (2.76 Å, 2.84 Å, 2.81 Å, 2.87 Å, see Table 2) in contrast to the shorter value of 2.65 Å for O(3W)···O(8) typical for OH_(oxonium)···O.

Evidently, the 30-membered crown ether tries to optimize its fold for a favorable encapsulation of the




Fig. 4. $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})]^+$ cation in $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5 \text{CH}_2\text{Cl}_2$.

$[\text{H}_7\text{O}_3]^+$ entity, maximizing the hydrogen bond stabilization. As shown in Fig. 5, the polyether molecule has to twist into a very special conformation. Even this optimization leaves some unfavorable details: The angles at O(1W) sharing three H atoms between its acceptor and donor oxygen atoms exhibit large differences with $154.4(9)^\circ$ (O(2)-O(1W)-O(9)), $72.4(7)^\circ$ (O(3W)-O(1W)-O(9)) and $132.4(8)^\circ$ (O(3W)-O(1W)-O(2)), indicating strongly non-linear hydrogen bonds.

Within the crown ether the individual C-O-C and O-C-C bond angles differ significantly from usual values (114.8° and 108.2°). The mean values however are comparable (114.7° and 105.5°).

Conclusion

The present compound extends the series of known oxonium halogenochalcogenates to an especially large 30-membered cyclic polyether as multidentate




Fig. 5. Illustration of the twisting in the $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})]^+$ cation in $[\text{H}_7\text{O}_3(\text{Bis-dibromo-dibenzo-30-crown-10})][\text{Se}_2\text{Br}_9] \cdot 1.5 \text{CH}_2\text{Cl}_2$.

ligand to the oxonium cation. The modes of crown ether coordination, together with their substitution patterns, offer convenient possibilities for the design of oxonium cation size and configuration. Considering the series of the crown members of this class, *e.g.* $[\text{H}_3\text{O}][\text{TeBr}_5] \cdot 3\text{C}_4\text{H}_8\text{O}_2$ [9] and $[\text{H}_5\text{O}_2][\text{Te}_2\text{Cl}_9] \cdot 2\text{C}_4\text{H}_8\text{O}_2$ [10], $[\text{H}_3\text{O}(\text{dibenzo-18-crown-6})][\text{Te}_2\text{Br}_9]$ [10], $[\text{H}_3\text{O}(\text{dibromo-benzo-15-crown-5})]_2[\text{SeBr}_6]$ [17], $[\text{H}_3\text{O}(\text{benzo-18-crown-6})]_2[\text{Te}_2\text{Br}_{10}]$ [18], and $[\text{H}_5\text{O}_2(\text{dibenzo-24-crown-8})]_2[\text{Te}_2\text{Br}_{10}]$ [18], and $[\text{H}_5\text{O}_2(\text{Bis-dibromo-dibenzo-24-crown-8})]_2[\text{Se}_3\text{Br}_8]$ [19, 20] a size-selective stabilization of larger homologues of the oxonium cations by the larger crown ethers is evident.

Acknowledgements

We would like to thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for substantial support of this work.

- [1] B. Krebs, F.-P. Ahlers, *Adv. Inorg. Chem.* **35**, 235 (1990).
 [2] S. Hasche, O. Reich, I. Beckmann, B. Krebs, *Z. Anorg. Allg. Chem.* **623**, 724 (1997).
 [3] B. Krebs, S. Bonmann, K. Erpenstein, *Z. Naturforsch.* **46b**, 919 (1991).
 [4] W. Hinrichs, D. Mandak, G. Klar, *Cryst. Struct. Commun.* **11**, 1781 (1982).
 [5] B. Krebs, B. Buss, W. Berger, *Z. Anorg. Allg. Chem.* **397**, 1 (1973).
 [6] P.H. Collins, M. Webster, *Acta Crystallogr.* **B28**, 1260 (1972).
 [7] B. Krebs, S. Bonmann, K. Gretenkord, *Z. Naturforsch.* **47b**, 93 (1992).
 [8] D. Katryniok, R. Kniep, D. Mootz, *Z. Anorg. Allg. Chem.* **461**, 96 (1980).

- [9] O. Reich, S. Hasche, K. Büscher, I. Beckmann, B. Krebs, *Z. Anorg. Allg. Chem.* **622**, 1011 (1996).
- [10] O. Reich, S. Hasche, S. Bonmann, B. Krebs, *Z. Anorg. Allg. Chem.* **624**, 411 (1998).
- [11] B. Viossat, P. Khodadad, N. Rodier, *J. Mol. Struct.* **71**, 237 (1981).
- [12] G.M. Sheldrick, SHELXTL PLUS, Siemens Analytical X-ray Instruments, Madison, WI (1990); SHELXL-93, Göttingen (1993).
- [13] I. D. Brown, *Can. J. Chem.* **42**, 2758 (1964).
- [14] W. Abriel, *Z. Naturforsch.* **42b**, 415 (1987).
- [15] E. Nagy-Felsobuki, J.B. Peel, *Chem. Phys.* **45**, 189 (1980).
- [16] P. Born, R. Kniep, D. Mootz, *Z. Anorg. Allg. Chem.* **451**, 12 (1979).
- [17] A. Hammerschmidt, I. Beckmann, M. Läge, B. Krebs, *Z. Anorg. Allg. Chem.* **630**, in press.
- [18] A. Hammerschmidt, S. Bonmann, M. Läge, B. Krebs, *Z. Anorg. Allg. Chem.* **630**, in press.
- [19] B. Krebs, I. Beckmann, S. Bonmann, S. Hasche, O. Reich, *Phosphorus, Sulfur and Silicon* **136–138**, 513 (1998).
- [20] I. Beckmann, Ph. D.Thesis, Münster (1994).