

This article was downloaded by: [University of Miami]

On: 27 November 2012, At: 08:47

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lagb20>

A PARTIAL ORDER ON THE ORTHOGONAL GROUP

Thomas Brady^a & Colum Watt^b

^a School of Mathematical Sciences, Dublin City University, Glasnevin, Dublin, Ireland

^b School of Mathematics, Trinity College, Dublin, Ireland

Version of record first published: 01 Sep 2006.

To cite this article: Thomas Brady & Colum Watt (2002): A PARTIAL ORDER ON THE ORTHOGONAL GROUP, Communications in Algebra, 30:8, 3749-3754

To link to this article: <http://dx.doi.org/10.1081/AGB-120005817>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



COMMUNICATIONS IN ALGEBRA
Vol. 30, No. 8, pp. 3749–3754, 2002

A PARTIAL ORDER ON THE ORTHOGONAL GROUP

Thomas Brady¹ and Colum Watt²

¹School of Mathematical Sciences,
Dublin City University, Glasnevin, Dublin 9, Ireland
E-mail: tom.brady@dcu.ie

²School of Mathematics, Trinity College,
Dublin 2, Ireland
E-mail: colum@maths.tcd.ie

ABSTRACT

We define a natural partial order on the orthogonal group and completely describe the intervals in this partial order. The main technical ingredient is that an orthogonal transformation induces a unique orthogonal transformation on each subspace of the orthogonal complement of its fixed subspace.

Let V be an n -dimensional vector space over a field \mathbf{F} and let $O(V)$ be the orthogonal group of V with respect to a fixed anisotropic symmetric bilinear form $\langle \cdot, \cdot \rangle$. In this note we will define a natural partial order on $O(V)$ and completely describe the intervals in this partial order. The main technical ingredient is that an orthogonal transformation A on V induces a unique orthogonal transformation on each subspace of the orthogonal complement of the fixed subspace of A .

3749

Recall that $A \in O(V)$ if $A : V \rightarrow V$ is linear and satisfies $\langle A(\vec{v}), A(\vec{w}) \rangle = \langle \vec{v}, \vec{w} \rangle$ for all $\vec{v}, \vec{w} \in V$. For standard results on symmetric bilinear forms and their associated orthogonal groups see^[1], but note that we are making the further assumption that the form is anisotropic.

For each $A \in O(V)$, we define two subspaces of V , $F(A) = \ker(A - I)$ and $M(A) = \text{im}(A - I)$, where I is the identity operator on V . We note that $F(A)$ is the $+1$ -eigenspace of A , sometimes called the fixed subspace of A . We will write $V = V_1 \perp V_2$ whenever V is the orthogonal direct sum of subspaces V_1 and V_2 .

Proposition 1. $V = F(A) \perp M(A)$

Proof. Since the dimensions of $F(A)$ and $M(A)$ are complementary and the form is anisotropic, it suffices to show that these subspaces are orthogonal. So let $\vec{x} \in F(A)$ and $\vec{y} \in M(A)$. Then $\vec{x} = A(\vec{x})$ and $\vec{y} = (A - I)\vec{z}$ for some $\vec{z} \in V$. Thus

$$\langle \vec{x}, \vec{y} \rangle = \langle \vec{x}, (A - I)\vec{z} \rangle = \langle \vec{x}, A(\vec{z}) \rangle - \langle \vec{x}, \vec{z} \rangle = \langle A(\vec{x}), A(\vec{z}) \rangle - \langle \vec{x}, \vec{z} \rangle = 0.$$

q.e.d.

We will be concerned with how the dimensions of these subspaces behave when we take products in $O(V)$. For notational convenience we will write $|U|$ for $\dim(U)$.

Proposition 2. $|M(AB)| \leq |M(A)| + |M(B)|$ for $A, B \in O(V)$.

Proof. Using the identities $|U| + |V| = |U + V| + |U \cap V|$, $F(A) \cap F(B) \subseteq F(AB)$ and $F(A) + F(B) \subseteq V$ we find that

$$|F(A)| + |F(B)| = |F(A) + F(B)| + |F(A) \cap F(B)| \leq n + |F(AB)|,$$

from which the result follows.

q.e.d.

This result is proved in a more general setting in^[2] However, from the proof above we see that equality occurs if and only if

$$F(A) \cap F(B) = F(AB) \quad \text{and} \quad F(A) + F(B) = V.$$

Thus, using the identities $[U + V]^\perp = U^\perp \cap V^\perp$ and $U^\perp + V^\perp = [U \cap V]^\perp$ we get the following characterization.

Corollary 1. $|M(AB)| = |M(A)| + |M(B)| \Leftrightarrow M(AB) = M(A) \oplus M(B)$.

Definition 1. We will write $A \leq C$ if $|M(C)| = |M(A)| + |M(A^{-1}C)|$.

Proposition 3. The relation \leq is a partial order on $O(V)$ and satisfies

$$A \leq B \leq C \Rightarrow A^{-1}B \leq A^{-1}C.$$

Proof. Reflexivity is immediate. To establish antisymmetry suppose $A \leq C$ and $C \leq A$. Then

$$|M(C)| = |M(A)| + |M(A^{-1}C)| = |M(C)| + |M(C^{-1}A)| + |M(A^{-1}C)|$$

giving $F(C^{-1}A) = F(A^{-1}C) = V$ or $A = C$.

To establish transitivity, suppose $A \leq B$ and $B \leq C$. Then

$$\begin{aligned} |M(C)| &\leq |M(A)| + |M(A^{-1}C)| \\ &= |M(A)| + |M(A^{-1}BB^{-1}C)| \\ &\leq |M(A)| + |M(A^{-1}B)| + |M(B^{-1}C)| \\ &= |M(A)| + \{|M(B)| - |M(A)|\} + \{|M(C)| - |M(B)|\} \\ &= |M(C)| \end{aligned}$$

So both of the inequalities are actually equalities. The first line gives $A \leq C$ and \leq is transitive. The third line gives the second assertion above. q.e.d.

The association of the subspace $M(A)$ to an element $A \in O(V)$ defines a map M from $O(V)$ to the set of subspaces of V . The next sequence of lemmas shows that the restriction of M to the interval $[I, C] = \{A \in O(V) \mid A \leq C\}$ is a bijection onto the set of subspaces of $M(C)$.

In what follows we fix C and a subspace W of $M(C)$ and we suppose that $A \in O(V)$ satisfies $M(A) = W$. We define U to be the unique subspace of $M(C)$ which satisfies $|U| = |W|$ and $(C - I)U = W$. This is possible since $C - I$ is invertible when restricted to $M(C)$.

Lemma 1. *If $W \subseteq M(C)$ then $V = W^\perp \oplus U$.*

Proof. Since the subspaces have complementary dimensions it suffices to show that their intersection is trivial. So let $\vec{x} \in W^\perp \cap U$. Then $\vec{x} \in W^\perp$ and $(C - I)\vec{x} = \vec{w}$ for some $\vec{w} \in W$. Thus $C\vec{x} = \vec{x} + \vec{w}$, with $\vec{x} \in W^\perp$ and $\vec{w} \in W$ so that

$$\langle \vec{x}, \vec{x} \rangle = \langle C\vec{x}, C\vec{x} \rangle = \langle \vec{x} + \vec{w}, \vec{x} + \vec{w} \rangle = \langle \vec{x}, \vec{x} \rangle + \langle \vec{w}, \vec{w} \rangle.$$

Thus $\vec{w} = \vec{0}$ since $\langle \cdot, \cdot \rangle$ is anisotropic and $\vec{x} = \vec{0}$ since $C - I$ is an isomorphism on $M(C)$. q.e.d.

Lemma 2. *$F(A^{-1}C) \subseteq F(C) \perp U$.*

Proof. Let $\vec{x} \in F(A^{-1}C)$. Then $A^{-1}C\vec{x} = \vec{x}$, which implies $C\vec{x} = A\vec{x}$ and $(C - I)\vec{x} = (A - I)\vec{x}$. Using $V = F(C) \perp M(C)$ we can express \vec{x} uniquely as $\vec{x} = \vec{y} + \vec{z}$ with $\vec{y} \in F(C)$ and $\vec{z} \in M(C)$. Thus

$$(C - I)\vec{z} = (C - I)\vec{x} = (A - I)\vec{x} \in M(A) = W,$$

giving $\vec{z} \in U$. This gives $F(A^{-1}C) \subseteq F(C) + U$ and the orthogonality of the subspaces follows since $U \subseteq M(C)$. q.e.d.

Lemma 3. *If $M(A) = W$ and $A \leq C$ then $F(A^{-1}C) = F(C) \perp U$.*

Proof. Since $A \leq C$ we have $|M(A^{-1}C)| = |M(C)| - |M(A)|$ so that

$$|F(A^{-1}C)| = n - |M(A^{-1}C)| = n - |M(C)| + |W| = |F(C)| + |U|.$$

This dimension calculation can now be combined with Lemma 2. q.e.d.

It is now possible to give a formula for A . If $V = V_1 \oplus V_2$ we define the projection $\text{Proj}_{V_1}^{V_2}$ to be the linear transformation which coincides with the identity on V_1 and with the zero transformation on V_2 .

Lemma 4. *If $A \leq C$ and $M(A) = W$ then $A = I + (C - I)\text{Proj}_U^{W^\perp}$.*

Proof. If $M(A) = W$ then $F(A) = W^\perp$ so that A coincides with I on W^\perp . Since $F(A^{-1}C)$ contains U by Lemma 3, A coincides with C on U . Thus $A - I$ coincides with the zero transformation on W^\perp and with $C - I$ on U , giving $A - I = (C - I)\text{Proj}_U^{W^\perp}$, by Lemma 1. q.e.d.

It is not at all clear from this formula that A is orthogonal. However this is indeed the case.

Lemma 5. $A = I + (C - I)\text{Proj}_U^{W^\perp} \in O(V)$.

Proof. Let $\vec{x}, \vec{y} \in V$ and use Lemma 1 to express $\vec{x} = \vec{x}_1 + \vec{x}_2$, $\vec{y} = \vec{y}_1 + \vec{y}_2$, with $\vec{x}_1, \vec{y}_1 \in U$ and $\vec{x}_2, \vec{y}_2 \in W^\perp$. Then, using the fact that A coincides with I on W^\perp and with C on U ,

$$\begin{aligned} \langle A(\vec{x}), A(\vec{y}) \rangle &= \langle C(\vec{x}_1) + \vec{x}_2, C(\vec{y}_1) + \vec{y}_2 \rangle \\ &= \langle C\vec{x}_1, C\vec{y}_1 \rangle + \langle C\vec{x}_1, \vec{y}_2 \rangle + \langle \vec{x}_2, C\vec{y}_1 \rangle + \langle \vec{x}_2, \vec{y}_2 \rangle \\ &= \langle \vec{x}_1, \vec{y}_1 \rangle + \langle C\vec{x}_1, \vec{y}_2 \rangle + \langle \vec{x}_2, C\vec{y}_1 \rangle + \langle \vec{x}_2, \vec{y}_2 \rangle \\ &= \langle \vec{x}, \vec{y} \rangle + \langle (C - I)\vec{x}_1, \vec{y}_2 \rangle + \langle \vec{x}_2, (C - I)\vec{y}_1 \rangle \\ &= \langle \vec{x}, \vec{y} \rangle, \end{aligned}$$

since both $(C - I)\vec{x}_1$ and $(C - I)\vec{y}_1$ lie in W . q.e.d.

We will call A the transformation induced by C on W . Combining the above lemmas we get the following result.

Theorem 1. *If $C \in O(V)$ and W is a subspace of $M(C)$ then there exists a unique $A \in O(V)$ satisfying $A \leq C$ and $M(A) = W$.*

The induced transformations are familiar objects for two special classes of subspace.

Corollary 2. *If $W \subseteq M(C)$ is an invariant subspace of C , then the induced transformation on W is the restriction of C to W .*

Proof. In this case, $U = W$ and the projection in the formula for A becomes an orthogonal projection. q.e.d.

Corollary 3. *If $\text{char}(\mathbf{F}) \neq 2$ and W is a one dimensional subspace of $M(C)$ then the orthogonal transformation induced by C on W is always the orthogonal reflection in W^\perp .*

Proof. Since W is one-dimensional A must act on W by multiplication by a scalar α . The orthogonality of A forces $\alpha^2 = 1$ and $W = M(A)$ gives $\alpha \neq 1$. q.e.d.

The poset $(O(V), \leq)$ is not a lattice, since distinct elements C_1 and C_2 with $M(C_1) = M(C_2)$ cannot have a common upper bound. However the intervals are lattices and can be easily described.

Theorem 2. *If $A \leq C$ in $O(V)$ and $|M(C)| - |M(A)| = m$ then the interval $[A, C] = \{B \in O(V) \mid A \leq B \leq C\}$ is isomorphic to the lattice of subspaces of \mathbf{F}^m under inclusion.*

Proof. The lattices of subspaces of \mathbf{F}^m under inclusion is isomorphic to the interval $[M(A), M(C)]$ in the lattice of subspaces of V . The function $B \mapsto M(B)$ is a bijection from the interval $[A, C]$ to the latter interval by Theorem 1. This map respects the partial orders by Corollary 1. To see that the inverse map respects the partial orders suppose that $M(A) \subseteq W_1 \subseteq W_2 \subseteq M(C)$. Let B_1, B_2 be the transformations induced on W_1, W_2 respectively by C and let B'_1 be the transformation induced on W_1 by B_2 . Then $B'_1 \leq B_2 \leq C$ gives $B'_1 \leq C$, but $M(B_1) = M(B'_1) = W_1$ so the uniqueness part of Theorem 1 gives $B_1 = B'_1$ and $B_1 \leq B_2$. q.e.d.

Each chain in $M(C)$ thus gives rise to a special factorization of C .

Corollary 4. *If $C \in O(V)$ and $W_1 \subset W_2 \subset \dots \subset W_k = M(C)$ is a chain of subspaces in $M(C)$ then C factors uniquely as a product of k transformations $C = B_1 B_2, \dots, B_k$, with $B_1 B_2, \dots, B_i \leq C$ and $M(B_1 B_2, \dots, B_i) = W_i$.*

Proof. If we define C_i to be the transformation induced by C on W_i then $B_i = (C_{i-1})^{-1} C_i$. q.e.d.

The case where this chain is maximal gives a strong version of the Cartan-Dieudonné theorem.

Corollary 5. *If $\text{char}(\mathbf{F}) \neq 2$, $C \in O(V)$ with $|M(C)| = k$ and $W_1 \subset W_2 \subset \cdots \subset W_k = M(C)$ is a maximal flag in $M(C)$ then C factors uniquely as a product of k reflections, $C = R_1 R_2 \cdots R_k$, with $M(R_1 R_2 \cdots R_i) = W_i$.*

Proof. Here the transformation B_i defined in Corollary 4 satisfies $|M(B_i)| = 1$ so that B_i is a reflection by Corollary 3.

Note 1. Using similar methods one can prove analogs of all the above results in the case of a unitary transformation over a finite-dimensional complex vector space. In this case we deal with complex linear subspaces, the induced transformations are unitary (and hence complex linear) and complex reflections replace the above reflections.

REFERENCES

1. Jacobson N., *Basic Algebra I*, Second Edn.; Freeman: San Francisco, 1985.
2. Scherk P., On the Decomposition of Orthogonalities into Symmetries, Proc. Amer. Math. Soc. **1950**, *1*, 481–491.

Received March 2001