

## ORIGINAL ARTICLE

# A population-based survey of sexual activity, sexual problems and associated help-seeking behavior patterns in mature adults in the United States of America

EO Laumann<sup>1</sup>, DB Glasser<sup>2</sup>, RCS Neves<sup>3</sup> and ED Moreira Jr<sup>3</sup>, for the GSSAB Investigators' Group

<sup>1</sup>Department of Sociology, University of Chicago, Chicago, USA; <sup>2</sup>Pfizer Inc., New York, NY, USA and <sup>3</sup>Gonçalo Moniz Research Center, Oswaldo Cruz Foundation, Salvador, Brazil

**To study sexual activity, the prevalence of sexual dysfunction and related help-seeking behaviors among mature adults in the United States of America, a telephone survey was conducted in 2001–2002. A total of 1491 individuals (742 men/749 women) aged 40–80 years completed the survey. Overall, 79.4% of men and 69.3% of women had engaged in sexual intercourse during the 12 months preceding the interview. Early ejaculation (26.2%) and erectile difficulties (22.5%) were the most common male sexual problems. A lack of sexual interest (33.2%) and lubrication difficulties (21.5%) were the most common female sexual problems. Less than 25% of men and women with a sexual problem had sought help for their sexual problem(s) from a health professional. Many men and women in the United States report continued sexual interest and activity into middle age and beyond. Although a number of sexual problems are highly prevalent, few people seek medical help.** *International Journal of Impotence Research* (2009) 21, 171–178; doi:10.1038/ijir.2009.7; published online 26 February 2009

**Keywords:** epidemiology; health surveys; impotence; prevalence; sex; sexual disorders

## Introduction

The development of convenient and effective oral treatments for male erectile dysfunction has stimulated an increasing level of interest in the sexual functioning of middle-aged and older adults. Over the last decade, many studies have investigated the prevalence of sexual problems among middle-aged and elderly people. Most of these studies have involved the populations of industrially developed nations around the world, particularly in Europe,<sup>1–3</sup> and in the Americas.<sup>4–12</sup> The prevalence of the male sexual problems of erectile dysfunction and early ejaculation and their related risk factors have been investigated most extensively, whereas fewer studies have focused specifically on female sexual dysfunction.<sup>13,14</sup> Moreover, relatively little is known

about the average frequency of sexual activity and the importance of sexual relationships among older men and women, although the few studies that have examined sexuality in mature adults have reported that sexual interest and activity persist well into older age.<sup>15,16</sup>

The published studies of the prevalence and correlates of sexual problems in developed and developing countries have used a variety of different study designs and definitions, which makes valid cross-national comparisons difficult. Moreover, there are currently no related studies reporting how men or women from different cultures attempt to manage or overcome their sexual problems, and there are only a few studies that allow a comparison of sexual behaviors across different countries.<sup>17</sup>

The Global Study of Sexual Attitudes and Behaviors (GSSAB) was a population survey of 27 500 men and women aged 40–80 years in 29 countries representing many world regions.<sup>18–21</sup> Here, we report the results from the respondents in the United States. In addition to estimating the prevalence of several sexual problems in men and women, we sought to investigate the factors associated with these problems and described the help-seeking behavior they elicited in this population.

Correspondence: Dr ED Moreira Jr, Unit of Epidemiology and Biostatistics, Oswaldo Cruz Foundation, Rua Walde-mar Falcão 121, Salvador, Bahia 40.295-001, Brazil.  
E-mail: edson@bahia.fiocruz.br

Received 4 December 2008; revised 4 February 2009; accepted 8 February 2009; published online 26 February 2009

## Methods

Using a random-digit dialed sampling design, computer-assisted telephone interviews were carried out in the United States during 2001 and 2002. Respondents were randomly selected by asking for the man or woman in the household aged between 40 and 80 years of age (participants were interviewed by interviewers of the same gender). Women and men were sampled in approximately equal numbers by design and verbal consent was obtained from all study participants. They were also informed about the following issues: (1) all information obtained would be used in aggregate; (2) responses were voluntary; (3) the confidentiality and the privacy of their responses were protected because no personal identifiers were coded into the interview instruments; (4) no list of respondents was retained; (5) the protocol was approved by an institutional review board; and (6) 'refusers' were not called back in an effort to convert them to participating respondents.

A structured questionnaire requested information concerning general health, demographics, relationships and sexual behaviors, attitudes and beliefs. The subjects were asked if they had engaged in sexual intercourse during the previous 12 months and the presence of sexual dysfunction was assessed by means of two sequential questions. The respondents were first asked whether they had experienced one or more of the sexual problems listed in Table 2 for a period of at least 2 months during the previous year, and those who answered 'Yes' were then asked whether they had experienced the problem 'occasionally', 'sometimes' or 'frequently'.

Logistic regression was used to investigate potential factors associated with a selected sexual dysfunction. In these analyses, the presence of a sexual dysfunction was coded only for those respondents who reported experiencing the problem frequently or periodically, whereas those who indicated that they experienced the problem only occasionally were recoded to indicate no sexual dysfunction.

The subjects who reported that they had experienced a sexual problem were asked whether they had sought help from a number of possible sources. The options included: 'Talked to partner', 'Talked to a medical doctor (other than a psychiatrist)', 'Looked for information anonymously (in books/magazines or on the internet)', 'Talked to family member or friend', 'Taken prescription drugs/devices or talked to pharmacist', 'Talked to psychiatrist or psychologist or marriage counsellor', 'Talked to a clergy person or religious adviser', 'Called a telephone help line', 'Other—please specify'. Respondents could indicate that they had sought help from more than one source.

The subjects with sexual problems who had not consulted a physician were asked why they had not done so, and offered a list of 14 possible reasons

(from which they were to check all that applied). The reasons included attitudes and beliefs regarding the sexual problem and the patient–doctor relationship. All respondents (irrespective of whether they reported any sexual problems) were also asked. 'During a routine office visit or consultation in the past 3 years, has your physician asked you about possible sexual difficulties without you bringing it up first?' (Yes/No) and 'Do you think a doctor should routinely ask patients about their sexual function?' (Yes/No).

The categorization of household income as 'low', 'medium' or 'high' was based on the distribution of income in the United States.

The prevalence of a specific characteristic was calculated by dividing the number of cases by the corresponding population. The denominator for the calculation of the prevalence of a sexual problem was the number of sexually active people (that is, at least one episode of intercourse during the previous 12 months). The prevalence estimates are given with their confidence intervals.

## Results

### *Characteristics of the study population*

Overall, 19377 individuals in the United States were contacted, 2817 of whom were not eligible to participate. Of the 16560 eligible individuals, a total of 1491 individuals (742 men and 749 women) completed the survey, for a response rate of 9.0%. The high attrition rate is, in part, attributable to the protocol which stipulated that in the interest of preserving respondents' anonymity, no call backs were permitted to find better times for interviews or to try to persuade 'refusers' to participate.

Table 1 presents selected characteristics of the study sample. A large proportion of the subjects were married or involved in an ongoing partnership (64.8% of men and 62.0% of women). The majority of the men (61.0%) and women (68.5%) were employed and, overall, about 75% of men and women reported that they were in good or excellent general health.

Approximately 80% of men and 70% of women said that they had had sexual intercourse during the 12 months preceding the interview, whereas about one-third of men (35.4%) and more than one-quarter of women (27.8%) engaged in sexual intercourse regularly (that is, more than once a week).

### *Prevalence of sexual problems*

Early ejaculation was the most common male sexual problem, and was reported by 26.2% of the sexually active men in the United States (approximately half of whom said that they experienced this problem periodically or frequently) (Table 2). Erectile diffi-

**Table 1** Selected characteristics (%) of the study sample, USA, 2001–2002

|   | Men<br>(n = 742) | Women<br>(n = 749) |
|---|------------------|--------------------|
| <i>Age group (years)</i>                      |                  |                    |
| 40–49   | 30.6             | 36.0               |
| 50–59   | 27.2             | 30.4               |
| 60–69   | 24.3             | 19.2               |
| 70–80   | 17.9             | 14.3               |
| <i>Relationship status</i>                    |                  |                    |
| Married or ongoing partnership                | 64.8             | 62.0               |
| Divorced/separated without sex partner        | 17.6             | 17.7               |
| Widowed without sex partner                   | 6.5              | 12.8               |
| Single without sex partner                    | 11.1             | 7.4                |
| Urban/suburban residential setting            | 67.0             | 64.1               |
| <i>Education</i>                              |                  |                    |
| Primary school or less                        | 2.0              | 2.0                |
| Secondary/high school                         | 39.5             | 44.8               |
| At least some college                         | 58.5             | 53.2               |
| <i>Household income</i>                       |                  |                    |
| Low   | 10.9             | 19.1               |
| Medium  | 57.5             | 55.2               |
| High  | 31.6             | 25.7               |
| <i>Current employment status</i>              |                  |                    |
| Employed                                      | 61.0             | 68.5               |
| Unemployed                                    | 6.3              | 8.8                |
| Retired                                       | 32.7             | 22.7               |
| <i>Religion</i>                               |                  |                    |
| Christian/Jew                                 | 89.4             | 93.7               |
| Buddhist or other Asian                       | 0.6              | 0.6                |
| Atheist                                       | 6.7              | 2.2                |
| Other, specified                              | 2.2              | 2.2                |
| Not specified                                 | 1.1              | 1.3                |
| Good to excellent general health <sup>a</sup> | 78.8             | 74.4               |
| Intercourse in the last 12 months             | 79.4             | 69.3               |
| Intercourse more than once a week             | 35.4             | 27.8               |
| <i>Number of sexual problems reported</i>     |                  |                    |
| None  | 49.9             | 42.2               |
| 1   | 27.6             | 24.2               |
| 2   | 12.7             | 15.6               |
| ≥3  | 9.8              | 18.0               |

<sup>a</sup>Self-reported 'good' or 'excellent' general health (vs 'fair' or 'poor').

culty was the second most common male sexual problem in the US sample, reported by 22.5% of sexually active men (12.4% said that they experienced this problem periodically or frequently), followed by a lack of sexual interest, which was reported by 18.1% of sexually active men (8.1% said that they experienced this problem periodically or frequently). The other sexual problems investigated (an inability to reach orgasm, a lack of sexual pleasure and pain during sexual intercourse) were experienced somewhat less frequently, particularly pain during intercourse, which was reported by only 3.1% of sexually active men in the United States.

**Table 2** Prevalence of sexual problems in men and women in the USA by severity, 2001–2002

|                           | % (95% confidence interval) |
|---------------------------|-----------------------------|
| <b>Men</b>                |                             |
| Early ejaculation         | <b>26.2 (23.0, 29.5)</b>    |
| Occasional                | 13.8 (11.4, 16.6)           |
| Periodic                  | 7.0 (5.2, 9.1)              |
| Frequent                  | 4.7 (3.3, 6.6)              |
| Erectile difficulties     | <b>22.5 (19.6, 25.7)</b>    |
| Occasional                | 9.6 (7.6, 12.0)             |
| Periodic                  | 5.9 (4.3, 7.9)              |
| Frequent                  | 6.5 (4.8, 8.5)              |
| Lack of sexual interest   | <b>18.1 (15.4, 21.1)</b>    |
| Occasional                | 8.8 (6.8, 11.1)             |
| Periodic                  | 4.8 (3.4, 6.6)              |
| Frequent                  | 3.3 (2.1, 4.9)              |
| Inability to reach orgasm | <b>12.4 (10.1, 15.0)</b>    |
| Occasional                | 5.8 (4.2, 7.8)              |
| Periodic                  | 2.5 (1.5, 4.0)              |
| Frequent                  | 3.1 (2.0, 4.7)              |
| Sex not pleasurable       | <b>11.2 (9.0, 13.8)</b>     |
| Occasional                | 4.8 (3.3, 6.6)              |
| Periodic                  | 2.5 (1.5, 4.0)              |
| Frequent                  | 1.8 (1.0, 3.1)              |
| Pain during sex           | <b>3.1 (2.0, 4.7)</b>       |
| Occasional                | 2.1 (1.2, 3.4)              |
| Periodic                  | 0.7 (0.2, 1.6)              |
| Frequent                  | 0.3 (0.0, 1.0)              |
| <b>Women</b>              |                             |
| Lack of sexual interest   | <b>33.2 (29.8, 36.7)</b>    |
| Occasional                | 14.7 (12.2, 17.5)           |
| Periodic                  | 6.0 (4.4, 8.0)              |
| Frequent                  | 10.2 (8.1, 12.7)            |
| Lubrication difficulties  | <b>21.5 (18.6, 24.7)</b>    |
| Occasional                | 9.3 (7.3, 11.7)             |
| Periodic                  | 5.5 (3.9, 7.5)              |
| Frequent                  | 5.8 (4.2, 7.8)              |
| Inability to reach orgasm | <b>20.7 (17.8, 23.9)</b>    |
| Occasional                | 8.6 (6.6, 10.9)             |
| Periodic                  | 5.4 (3.8, 7.3)              |
| Frequent                  | 5.7 (4.1, 7.7)              |
| Sex not pleasurable       | <b>19.7 (16.8, 22.8)</b>    |
| Occasional                | 8.8 (6.8, 11.1)             |
| Periodic                  | 5.5 (4.0, 7.5)              |
| Frequent                  | 4.2 (2.9, 6.0)              |
| Pain during sex           | <b>12.7 (10.4, 15.4)</b>    |
| Occasional                | 6.6 (4.9, 8.7)              |
| Periodic                  | 2.9 (1.8, 4.4)              |
| Frequent                  | 2.8 (1.7, 4.2)              |

Based on reports from sexually active respondents. Percentage in the first row of each panel indicates the overall prevalence of the sexual dysfunction, defined as an experience of dysfunction for a period of 2 months or more. The difference between the overall prevalence and the sum of the three levels of severity of sexual dysfunction indicates the proportion that failed to specify the level of severity.

Lack of sexual interest (33.2%) was the most common sexual problem reported by sexually active women in the United States, followed by difficulty becoming adequately lubricated (21.5%), an inability to reach orgasm (20.7%) and a lack of sexual pleasure (19.7%) (Table 2). At least one-half of the women who reported each of these problems said that she experienced it frequently or periodically. The other sexual problem investigated was pain

**Table 3** Factors associated with sexual problems by gender, USA, 2001–2002

|                                   | Men               |                           |                           | Women                     |                           |                           |
|-----------------------------------|-------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
|                                   | Early ejaculation | Lack of sexual interest   | Erectile difficulties     | Inability to reach orgasm | Lack of sexual interest   | Lubrication difficulties  |
| <i>Age (years)</i>                |                   |                           |                           |                           |                           |                           |
| 40–49                             | Referent          | Referent                  | Referent                  | Referent                  | Referent                  | Referent                  |
| 50–59                             | 0.58 (0.30, 1.14) | 0.88 (0.42, 1.83)         | 0.88 (0.42, 1.85)         | <b>2.45 (1.17, 5.11)*</b> | 1.59 (0.95, 2.67)         | 1.77 (0.92, 3.44)         |
| 60–80                             | 0.79 (0.43, 1.44) | 0.81 (0.39, 1.69)         | <b>2.19 (1.17, 4.12)*</b> | 0.62 (0.30, 1.27)         | 1.38 (0.78, 2.44)         | <b>2.56 (1.32, 4.98)†</b> |
| <i>Level of physical activity</i> |                   |                           |                           |                           |                           |                           |
| Average and above                 | Referent          | Referent                  | Referent                  | Referent                  | Referent                  | Referent                  |
| Lower than average                | 0.97 (0.48, 1.94) | <b>2.13 (1.08, 4.23)*</b> | 1.21 (0.64, 2.30)         | 1.84 (0.99, 3.39)         | <b>1.76 (1.04, 2.98)*</b> | 1.28 (0.67, 2.44)         |
| <i>Smoking</i>                    |                   |                           |                           |                           |                           |                           |
| Never                             | Referent          | Referent                  | Referent                  | Referent                  | Referent                  | Referent                  |
| Currently/smoked before           | 1.13 (0.65, 1.95) | 0.83 (0.44, 1.55)         | 1.05 (0.61, 1.80)         | 1.20 (0.70, 2.07)         | 1.29 (0.82, 2.01)         | 0.99 (0.59, 1.65)         |
| <i>Education</i>                  |                   |                           |                           |                           |                           |                           |
| Primary school or less            | Referent          | Referent                  | Referent                  | Referent                  | Referent                  | Referent                  |
| Secondary/some college            | 0.30 (0.84, 1.06) | 1.12 (0.12, 10.44)        | 0.60 (0.15, 2.35)         | 0.43 (0.83, 2.25)         | 0.50 (0.12, 2.09)         | 0.66 (0.24, 1.79)         |
| <i>Household income</i>           |                   |                           |                           |                           |                           |                           |
| Low                               | Referent          | Referent                  | Referent                  | Referent                  | Referent                  | Referent                  |
| Medium and high                   | 0.74 (0.35, 1.55) | 0.64 (0.28, 1.45)         | 0.90 (0.44, 1.85)         | 0.63 (0.33, 1.21)         | 1.18 (0.65, 2.16)         | <b>2.31 (1.07, 5.0)*</b>  |
| <i>Medical conditions</i>         |                   |                           |                           |                           |                           |                           |
| Depression diagnosed              | 1.49 (0.73, 3.04) | <b>3.19 (1.55, 6.59)†</b> | <b>2.61 (1.38, 4.96)†</b> | <b>2.73 (1.53, 4.85)†</b> | <b>2.08 (1.25, 3.48)†</b> | <b>2.42 (1.36, 4.33)†</b> |
| Hypertension diagnosed            | 0.97 (0.54, 1.76) | 0.87 (0.43, 1.75)         | 0.96 (0.55, 1.67)         | 0.84 (0.45, 1.57)         | 0.48 (0.27, 1.83)         | 1.19 (0.68, 2.10)         |
| Diabetes diagnosed                | 0.95 (0.43, 2.07) | 1.18 (0.50, 2.78)         | 1.22 (0.62, 2.40)         | 0.58 (0.23, 1.46)         | 0.97 (0.50, 1.89)         | 0.81 (0.37, 1.79)         |
| Heart disease                     | 0.92 (0.43, 1.98) | 0.91 (0.24, 1.54)         | 1.19 (0.63, 2.27)         | 0.71 (0.29, 1.78)         | 1.32 (0.66, 2.64)         | 1.01 (0.46, 2.23)         |
| Prostate disease                  | 0.86 (0.31, 2.34) | <b>2.56 (1.02, 6.44)*</b> | 1.61 (0.76, 3.42)         |                           |                           |                           |

Odds ratios from logistic regression (and 95% confidence intervals), in these analyses, the presence of a sexual problem included only those respondents who reported ‘sometimes’ or ‘frequently’ having the problem (that is, those who indicated ‘occasionally’ were recoded to indicate no sexual problem). Based on reports from sexually active subjects.

\* $P \leq 0.05$ ; † $P \leq 0.01$ .

during sexual intercourse, which was experienced by 12.7% of sexually active women.

Physical/health, demographic and socioeconomic factors associated with three selected sexual dysfunctions in men and women are summarized in Table 3 (odds ratios (OR) from logistic regression). Older age (age 60–80 years compared with the referent age of 40–49 years) was a significant correlate of erectile difficulties in men (OR 2.19,  $P \leq 0.05$ ) and lubrication difficulties in women (OR 2.56,  $P \leq 0.01$ ), whereas the age range of 50–59 years (the age at which many women experience the menopause) was associated with an inability to reach orgasm in women (OR 2.45 compared with the referent of 40–49 years,  $P \leq 0.05$ ). A lower than average level of physical activity was a significant correlate of lack of sexual interest in both men (OR 2.13,  $P \leq 0.05$ ) and women (OR 1.76,  $P \leq 0.05$ ). The

impact of a diagnosis of a number of common health conditions were investigated and it was observed that depression was significantly associated with a lack of sexual interest in both men (OR 3.19,  $P \leq 0.01$ ) and women (OR 2.08,  $P \leq 0.01$ ), erectile difficulties in men (OR 2.61,  $P \leq 0.01$ ) and lubrication difficulties (OR 2.42,  $P \leq 0.01$ ) and an inability to reach orgasm (OR 2.73,  $P \leq 0.01$ ) in women. Among men, a diagnosis of prostate disease was associated with a lack of sexual interest (OR 2.56,  $P \leq 0.05$ ).

#### Help-seeking behavior

The prevalence of selected help-seeking behaviors for sexual problems in the United States is summarized in Table 4. Of the respondents who were sexually active and reported experiencing at least

**Table 4** Prevalence of selected help-seeking behaviors for sexual problems by gender, USA, 2001–2002

|  | % (95% confidence interval) |
|--|-----------------------------|
| <i>Men</i>   |                             |
| Talked to partner  | 43.3 (38.3, 48.3)           |
| Talked to medical doctor   | 21.9 (18.0, 26.3)           |
| Taken drugs/used devices or talked to pharmacist   | 17.7 (14.1, 21.8)           |
| Looked for information anonymously (in books/magazines or by telephone help-line/internet) | 15.3 (11.9, 19.2)           |
| Talked to family member/friend   | 8.9 (6.3, 12.1)             |
| Talked to psychiatrist, psychologist or marriage counselor                                 | 3.7 (2.1, 6.0)              |
| Talked to a clergy person or religious adviser   | 2.5 (1.2, 4.5)              |
| Sought no help from a health professional  | 75.7 (71.3, 79.8)           |
| No action taken  | 45.2 (40.3, 50.3)           |
| <i>Women</i>   |                             |
| Talked to partner  | 43.4 (38.7, 48.1)           |
| Talked to medical doctor   | 16.1 (12.8, 19.7)           |
| Talked to family member/friend   | 15.6 (12.4, 19.2)           |
| Looked for information anonymously (in books/magazines or by telephone help-line/internet) | 15.0 (11.9, 18.6)           |
| Taken drugs/used devices or talked to pharmacist   | 13.7 (10.7, 17.2)           |
| Talked to psychiatrist, psychologist or marriage counselor                                 | 6.3 (4.3, 8.9)              |
| Talked to a clergy person or religious adviser   | 2.6 (1.3, 4.5)              |
| Sought no help from a health professional  | 79.7 (75.7, 83.3)           |
| No action taken  | 43.9 (39.3, 48.6)           |

Note: based on reports from respondents complaining of at least one sexual problem.

one sexual problem, 45.2% of men and 43.9% of women did not take any action (that is, they had not sought any help or advice). A slightly greater proportion of men (21.9%) than women (16.1%) reported talking to a medical doctor about their sexual problem(s), but overall the majority of men (75.7%) and women (79.7%) had sought no help from a health professional. Patterns of help-seeking behaviors were generally similar for men and women in the United States and talking to their partner was the most usual action taken by both men and women (43.3 and 43.4%, respectively).

#### *Factors associated with seeking medical help for sexual problems*

Some factors that might be associated with seeking medical help for sexual problems were investigated using logistic regression and the findings for both men and women in the United States are summarized in Table 5. A significant effect of age was seen only in men at age 60–69 years (OR 5.2,  $P \leq 0.01$ ), compared with the referent group aged 40–49 years. Certain sexual problems were associated with a greater likelihood of seeking medical help. Erectile

difficulties in men (OR 5.29,  $P \leq 0.001$ ) and lubrication difficulties in women (OR 2.09,  $P \leq 0.05$ ) were significant correlates of seeking medical help for sexual problems. A number of sexual beliefs and attitudes were significant correlates of seeking medical help for sexual problems. In men, these were ‘being very or somewhat dissatisfied with their own sexual functioning’ (OR 2.94,  $P \leq 0.01$ ), ‘a belief that decreased sexual ability would significantly affect their own self-esteem’ (OR 2.69,  $P \leq 0.05$ ) and ‘thinking that it is acceptable to use medical treatment for sexual problems’ (OR 13.52,  $P \leq 0.01$ ). In women, having been asked by a doctor about possible sexual difficulties during a routine visit in the past 3 years (OR 2.23,  $P \leq 0.05$ ), and thinking that a doctor should routinely ask patients about sexual function (OR 3.55,  $P \leq 0.05$ ) were significantly correlated with seeking medical help for sexual problems. Thinking that older people no longer want/have sex had quite a different effect for men and women. Although among women, this belief was associated with an increasing likelihood of seeking medical help for sexual problems (OR 3.14,  $P \leq 0.01$ ), men with this belief were less likely to seek medical help (OR 0.36,  $P \leq 0.05$ ).

#### *Attitudes and beliefs about diagnosis and treatment of sexual problems*

The most common reasons cited among respondents in the United States for not consulting a doctor about a sexual problem were thinking it was not very serious or waiting for the problem to go away (36.3% of men and 38.1% of women) and a belief that it is a normal part of aging or being comfortable as he/she is (25.4% of men and 28.2% of women) (Table 6). Lack of access to or affordability of medical care and embarrassment about discussing sexual problems with their medical doctor were cited as a reason by less than 5% of men and women, whereas a lack of belief that a sexual problem is a treatable medical condition was cited by about 15% of men and women. Few respondents in the United States had been asked by a doctor about possible sexual difficulties during a routine visit in the past 3 years (11.5% of men and 15.0% of women) but more than one-half of men (59.2%) and women (54.0%) thought that a doctor should routinely ask patients about their sexual function.

## Discussion

In this study, we report population-level data from middle-aged and older men and women in the United States concerning sexual behavior, the prevalence of sexual problems and associated help-seeking behaviors. The large cross-national population sample and the use of a common method of data collection represent two major strengths of

**Table 5** Factors associated with seeking medical help for sexual problems by gender, USA, 2001–2002

|   | Men                                     | Women                                |
|---|---|--------------------------------------|
| <i>Age (years)</i>  |   |                                      |
| 40–49   | Reference                               | Reference                            |
| 50–59   | 1.72 (0.63, 4.73)                       | 1.50 (0.70, 3.20)                    |
| 60–69   | <b>5.20 (1.82, 14.81)<sup>†</sup></b>   | 1.18 (0.47, 2.99)                    |
| 70–80   | 1.58 (0.52, 4.81)                       | 2.07 (0.64, 6.68)                    |
| <i>Education</i>  |   |                                      |
| Primary school or less  | Reference                               | Reference                            |
| Secondary/high school   | 0.99 (0.08, 11.69)                      | 2.51 (0.21, 30.37)                   |
| At least some college   | 1.20 (0.10, 14.31)                      | 2.89 (0.22, 37.53)                   |
| High/medium household income (vs low)   | 1.05 (0.32, 3.42)                       | 1.01 (0.42, 246)                     |
| <i>Sexual problems</i>  |   |                                      |
| Erectile difficulties   | <b>5.29 (2.42, 11.57)<sup>‡</sup></b>   |                                      |
| Early ejaculation   | 0.49 (0.23, 1.04)                       |                                      |
| Lack of sexual interest   | 1.29 (0.63, 2.65)                       | 1.13 (0.59, 2.17)                    |
| Inability to reach orgasm   |   | 1.20 (0.62, 2.32)                    |
| Lubrication difficulties  |   | <b>2.09 (1.11, 3.94)*</b>            |
| <i>General sexual attitudes</i>   |   |                                      |
| Have been asked by a doctor about possible sexual difficulties in a routine visit in the past 3 years | 1.35 (0.54, 3.35)                       | <b>2.23 (1.11, 4.48)*</b>            |
| Think a doctor should routinely ask patients about sexual function                                    | 0.93 (0.43, 2.00)                       | <b>3.55 (1.55, 8.13)*</b>            |
| Very/somewhat dissatisfied with sexual function   | <b>2.94 (1.31, 6.60)<sup>†</sup></b>    | 1.42 (0.58, 3.49)                    |
| Belief that decreased ability to perform sexually would significantly affect self-esteem              | <b>2.69 (1.12, 6.49)*</b>               | 1.65 (0.70, 3.87)                    |
| Belief that sex is an extremely/very important part of overall life                                   | 1.70 (0.47, 6.11)                       | 0.39 (0.08, 1.95)                    |
| Think it is ok to use medical treatment for sexual problems   | <b>13.52 (1.57, 116.15)<sup>†</sup></b> | 1.34 (0.47, 3.81)                    |
| Think that older people no longer want/have sex   | <b>0.36 (0.14, 0.97)*</b>               | <b>3.14 (1.33, 7.44)<sup>†</sup></b> |
| Belief in religion guiding sex  | 1.06 (0.54, 2.10)                       | 0.89 (0.47, 1.66)                    |

Odds ratios from logistic regression and 95% confidence intervals. Based on reports from respondents complaining of at least one sexual problem.

\* $P \leq 0.05$ ; <sup>†</sup> $P \leq 0.01$ ; <sup>‡</sup> $P \leq 0.001$ .

the GSSAB. Face-to-face interviews were not used to avoid causing respondents undue embarrassment when talking about private and sensitive issues, and to minimize the likelihood of respondents feeling obliged to give ‘socially desirable’ answers.<sup>22</sup> Only a sexual problem that persisted with moderate to higher frequency was considered to be a ‘dysfunction’. This method is essentially equivalent to using two sequential screening tests, and minimizes the risk of false-positive responses. It is likely, therefore, that the prevalence of sexual dysfunction may be under-reported in the GSSAB in comparison with studies that used more sensitive, but less specific methods.

The overall response rate in the United States (9.0%) was low, but similar to other countries in our study, namely, Austria, Canada, Germany, Spain, Italy, United Kingdom and Israel (<12%); and somewhat lower than the response rate in Australia, New Zealand, Sweden, France and Belgium (14–17%). In the data available, we did not find any of the characteristics to be correlated to participation in the survey. Although it is true that low completion rates can serve as a flag for the possibility of systematic biases in sample coverage,

it by no means guarantees or necessitates it. The prevalence of self-reported health conditions such as hypertension, diabetes and smoking in the GSSAB (these data are not shown here) was comparable with published values.<sup>23–25</sup> This suggests that contacts refused to participate in the study due to a general unwillingness to undergo a telephone interview, regardless of the subject matter and is therefore unlikely to have introduced a bias in the estimates of the prevalence of sexual behaviors and problems. It also appears to indicate that the study population was broadly representative of the US population. This assumption is further supported by the observation that the prevalence of erectile dysfunction among men in the US cohort of the GSSAB is comparable to values reported in published studies that were conducted in the United States among men aged 50 years or older in rural Central New York State,<sup>5</sup> and among US health professionals.<sup>6</sup> The prevalence of other sexual problems in our survey is also comparable to those yielded by the analysis of data from the National Health and Social Life Survey, a probability sample study of sexual behavior in a demographically

**Table 6** Attitudes, behaviors and beliefs about diagnosis of and treatment for sexual problems by gender, USA, 2001–2002

|  | % (95% confidence interval) |
|--|-----------------------------|
| <i>Men</i>   |                             |
| <i>Reasons for not consulting a doctor about the experienced sexual problem<sup>a</sup></i>                              |                             |
| Did not think it was very serious/<br>waiting if problem goes away   | 36.3 (30.8, 42.0)           |
| Normal with aging/I am comfortable<br>the way I am   | 25.4 (20.6, 30.8)           |
| Doctor cannot do much/do not think it<br>is a medical problem  | 14.9 (11.1, 19.5)           |
| Not comfortable talking to an MD/MD<br>is a close friend/MD is the wrong<br>gender                                       | 3.1 (1.4, 5.7)              |
| Do not have a regular physician/<br>doctor is expensive  | 2.4 (1.0, 4.8)              |
| Doctor uneasy to talk about sex  | 1.0 (0.2, 2.9)              |
| Have been asked by a doctor about<br>possible sexual difficulties in a routine<br>visit in the past 3 years <sup>b</sup> | 11.5 (8.0, 15.7)            |
| Think a doctor should routinely ask<br>patients about their sexual function <sup>b</sup>                                 | 59.2 (53.0, 65.1)           |
| <i>Women</i>   |                             |
| <i>Reasons for not consulting a doctor about the experienced sexual problem<sup>a</sup></i>                              |                             |
| Did not think it was very serious/<br>waiting if problem goes away   | 38.1 (32.9, 43.5)           |
| Normal with aging/I am comfortable<br>the way I am   | 28.2 (23.4, 33.2)           |
| Doctor cannot do much/do not think it<br>is a medical problem  | 13.5 (10.0, 17.6)           |
| Not comfortable talking to an MD/MD<br>is a close friend/MD is the wrong<br>gender                                       | 5.0 (2.9, 7.9)              |
| Do not have a regular physician/<br>doctor is expensive  | 2.3 (1.0, 4.6)              |
| Doctor uneasy to talk about sex  | 0.9 (0.2, 2.5)              |
| Have been asked by a doctor about<br>possible sexual difficulties in a routine<br>visit in the past 3 years <sup>b</sup> | 15.0 (11.4, 19.3)           |
| Think a doctor should routinely ask<br>patients about their sexual function <sup>b</sup>                                 | 54.0 (48.3, 59.6)           |

Abbreviations: MD, medical doctor.

<sup>a</sup>Based on reports from respondents complaining of at least one sexual problem who have not consulted a doctor.<sup>b</sup>Based on all respondents.

representative cohort of US adults.<sup>4</sup> In this study, an inability to reach orgasm, a lack of sexual interest and sex not pleasurable was reported by 22, 29 and 11% of the women and by 9, 16 and 7% of the men, respectively.

Some well-known risk factors (for example, smoking, diabetes, and so on) did not seem to affect sexual function/problems in our survey. This may be due to the limited power of our study, and to possible misclassification of the diagnosis of medical conditions as we had to rely on self-report. Despite that, we found that depression was a significant correlate of a number of sexual problems

in both men and women in the United States. Comorbidity between male erectile dysfunction and depression is known to be high but the nature of the relationship between the two conditions is unclear.<sup>26,27</sup> Although the distress of erectile dysfunction may contribute to the development of depression, it is also possible that depressive illness may lead to erectile difficulties. There have been data from a prospective study suggesting that the incidence of erectile dysfunction is increased among men with depression.<sup>28</sup> When considering the co-presence of depression and sexual dysfunction, the possible role of antidepressant treatments should not be overlooked. Sexual dysfunction is a common side effect of antidepressant therapy; however, the reported rates of dysfunction may vary between different agents.<sup>29,30</sup> The selective serotonin reuptake inhibitors appear to be associated with especially high rates of sexual dysfunction and whereas men taking selective serotonin reuptake inhibitors report higher rates of sexual side effects than women, the dysfunction experienced by women seems to be more severe.<sup>29,31</sup> Finally, just as cardiovascular problems are good markers for sexual problems (and vice versa), our data suggest that depression can be investigated in a similar way, that is, physicians who note that their patients are depressed may want to inquire about their sexual function and patients who are having problems with sex should be queried about their mood state.

Our findings indicate that feeling that the problem was not severe, or not being bothered by the problem may be deterring men and women in the United States from raising the subject of their sexual difficulties with their doctor. Furthermore, they show that doctors in the United States rarely ask patients about their sexual health during a routine consultation, even though this would be welcomed by the majority of men and women and would appear to encourage medical help-seeking for sexual problems, at least in women. If left untreated, sexual problems can greatly diminish a patient's quality of life and it is important that physicians—especially primary care physicians—incorporate questions about sexual functioning into their routine patient work-ups.<sup>32</sup> If handled sensitively, this should result not only in improved functioning for the patient but also an enhanced physician–patient relationship and greater professional satisfaction.

In conclusion, these findings indicate that middle-aged and elderly men and women in the United States continue to show sexual interest and activity, in spite of the high prevalence of several sexual dysfunctions. Few of the men and women who experience sexual difficulties seek medical help; this may be due in part to believing that the problem was not serious and/or not being bothered by the problem. Appropriate educational initiatives, aimed at both patients and healthcare professionals, may help to increase awareness and understanding

of sexual health issues and help them to identify and overcome potential barriers that their patients' might have in discussing and seeking help for sexual problems, thereby enabling more older adults to continue to enjoy a fulfilling sexual life.

## Acknowledgments

We acknowledge the contribution of our colleagues on the study's International Advisory Board: Alfredo Nicolosi (Italy), Bernard Levinson (South Africa), Clive Gingell (UK), Ferruh Simsek (Turkey), Gerald Brock (Canada), Jacques Buvat (France), Ken Marumo (Japan), Rosie King (Australia), Sae-Chul Kim (Korea) and Uwe Hartmann (Germany). The Global Study of Sexual Attitudes and Behaviors was funded by Pfizer Inc.

## Conflict of interest

Edward O Laumann and Edson D Moreira Jr are consultants for Pfizer Inc. Dale B Glasser is an employee and stockowner of Pfizer Inc. Raimundo C S Neves—none.

## References

- Helgason AR, Adolfsson J, Dickman P, Arver S, Fredrikson M, Gothberg M *et al*. Sexual desire, erection, orgasm and ejaculatory functions and their importance to elderly Swedish men: a population-based study. *Age Ageing* 1996; **25**: 285–291.
- Blanker MH, Bosch JL, Groeneveld FP, Bohnen AM, Prins A, Thomas S *et al*. Erectile and ejaculatory dysfunction in a community-based sample of men 50–78 years old: prevalence, concern, and relation to sexual activity. *Urology* 2001; **57**: 763–768.
- Giuliano F, Chevret-Measson M, Tsatsaris A, Reitz C, Murino M, Thonneau P. Prevalence of erectile dysfunction in France: results of an epidemiological survey of a representative sample of 1004 men. *Eur Urol* 2002; **42**: 382–389.
- Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999; **281**: 537–544.
- Ansong KS, Lewis C, Jenkins P, Bell J. Epidemiology of erectile dysfunction: a community-based study in rural New York State. *Ann Epidemiol* 2000; **10**: 293–296.
- Bacon CG, Mittleman MA, Kawachi I, Giovannucci E, Glasser DB, Rimm EB. Sexual function in men older than 50 years of age: results from the health professionals follow-up study. *Ann Intern Med* 2003; **139**: 161–168.
- Brock G, Moreira Jr ED, Glasser DB, Gingell C. Sexual disorders and associated help-seeking behaviors in Canada. *Can J Urol* 2006; **13**: 2953–2961.
- Moreira Jr ED, Abdo CH, Torres EB, Lobo CF, Fittipaldi JA. Prevalence and correlates of erectile dysfunction: results of the Brazilian study of sexual behavior. *Urology* 2001; **58**: 583–588.
- Moreira Jr ED, Lisboa Lobo CF, Villa M, Nicolosi A, Glasser DB. Prevalence and correlates of erectile dysfunction in Salvador, northeastern Brazil: a population-based study. *Int J Impot Res* 2002; **14**(Suppl 2): 3–9.
- Moreira Jr ED, Bestane WJ, Bartolo EB, Fittipaldi JA. Prevalence and determinants of erectile dysfunction in Santos, southeastern Brazil. *Sao Paulo Med J* 2002; **120**: 49–54.
- Hernandez Moreno PN, Mendoza Martinez R, Hernandez Marin I, Tovar Rodriguez JM, Ayala AR. Epidemiologic assessment of erectile function in a selected Mexican population. *Ginecol Obstet Mex* 2003; **71**: 332–342.
- Nolazco C, Bellora O, Lopez M, Surur D, Vazquez J, Rosenfeld C *et al*. Prevalence of sexual dysfunctions in Argentina. *Int J Impot Res* 2004; **16**: 69–72.
- Barlow DH, Cardozo LD, Francis RM, Griffin M, Hart DM, Stephens E *et al*. Urogenital ageing and its effect on sexual health in older British women. *Br J Obstet Gynaecol* 1997; **104**: 87–91.
- Abdo CH, Oliveira Jr WM, Moreira Jr ED, Fittipaldi JA. Prevalence of sexual dysfunctions and correlated conditions in a sample of Brazilian women—results of the Brazilian study on sexual behavior (BSSB). *Int J Impot Res* 2004; **16**: 160–166.
- Gott M, Hinchliff S. How important is sex in later life? The views of older people. *Soc Sci Med* 2003; **56**: 1617–1628.
- Matthias RE, Lubben JE, Atchison KA, Schweitzer SO. Sexual activity and satisfaction among very old adults: results from a community-dwelling Medicare population survey. *Gerontologist* 1997; **37**: 6–14.
- Nicolosi A, Moreira Jr ED, Shirai M, Bin Mohd Tambi MI, Glasser DB. Epidemiology of erectile dysfunction in four countries: cross-national study of the prevalence and correlates of erectile dysfunction. *Urology* 2003; **61**: 201–206.
- Nicolosi A, Laumann EO, Glasser DB, Moreira Jr ED, Paik A, Gingell C. Sexual behavior and sexual dysfunctions after age 40: the global study of sexual attitudes and behaviors. *Urology* 2004; **64**: 991–997.
- Laumann EO, Nicolosi A, Glasser DB, Paik A, Gingell C, Moreira E *et al*. Sexual problems among women and men aged 40–80 years: prevalence and correlates identified in the Global Study of Sexual Attitudes and Behaviors. *Int J Impot Res* 2005; **17**: 39–57.
- Moreira Jr ED, Brock G, Glasser DB, Nicolosi A, Laumann EO, Paik A *et al*. Help-seeking behaviour for sexual problems: the global study of sexual attitudes and behaviors. *Int J Clin Pract* 2005; **59**: 6–16.
- Moreira Jr ED, Glasser DB, King R, Duarte FG, Gingell C. Sexual difficulties and help-seeking among mature adults in Australia: results from the Global Study of Sexual Attitudes and Behaviours. *Sex Health* 2008; **5**: 227–234.
- (ASCF) ApiataAosbiF. A comparison between two models of investigation: telephone survey and face-to-face survey. *AIDS* 1992; **6**: 315–323.
- Balluz L, Ahluwalia IB, Murphy W, Mokdad A, Giles W, Harris VB. Surveillance for certain health behaviors among selected local areas—United States, Behavioral Risk Factor Surveillance System, 2002. *MMWR Surveill Summ* 2004; **53**: 1–100.
- Natarajan S, Nietert PJ. National trends in screening, prevalence, and treatment of cardiovascular risk factors. *Prev Med* 2003; **36**: 389–397.
- Whelton PK, He J, Muntner P. Prevalence, awareness, treatment and control of hypertension in North America, North Africa and Asia. *J Hum Hypertens* 2004; **18**: 545–551.
- Seidman SN. Exploring the relationship between depression and erectile dysfunction in aging men. *J Clin Psychiatry* 2002; **63**(Suppl 5): 5–12; discussion 23–25.
- Nicolosi A, Moreira Jr ED, Villa M, Glasser DB. A population study of the association between sexual function, sexual satisfaction and depressive symptoms in men. *J Affect Disord* 2004; **82**: 235–243.
- Moreira Jr ED, Lobo CF, Diament A, Nicolosi A, Glasser DB. Incidence of erectile dysfunction in men 40–69 years old: results from a population-based cohort study in Brazil. *Urology* 2003; **61**: 431–436.
- Clayton AH, Pradko JF, Croft HA, Montano CB, Leadbetter RA, Bolden-Watson C *et al*. Prevalence of sexual dysfunction among newer antidepressants. *J Clin Psychiatry* 2002; **63**: 357–366.
- Montgomery SA, Baldwin DS, Riley A. Antidepressant medications: a review of the evidence for drug-induced sexual dysfunction. *J Affect Disord* 2002; **69**: 119–140.
- Hensley PL, Nurnberg HG. SSRI sexual dysfunction: a female perspective. *J Sex Marital Ther* 2002; **28**(Suppl 1): 143–153.
- Sadovsky R. Integrating erectile dysfunction treatment into primary care practice. *Am J Med* 2000; **109**(Suppl 9A): 22–28; discussion 29–30.