

# ***A Possible Explanation of Low Energy gamma-ray Excess from Galactic Centre and Fermi Bubble by a Dark Matter Model with Two Real Scalars***


*Debasish Majumdar*

**Astroparticle Physics and Cosmology Division**  
**Saha Institute of Nuclear Physics**  
**Kolkata**

# Energy Budget of Universe


## PLANCK 2013 RESULTS !!! (March 21, 2013)

- Baryonic Matter are ~ 4.8%
- Dark Matter ~ 26.5%
- Dark Energy ~ 68.4%



# General Properties of Dark Matter

- Should be neutral
- Gravitationally interacting
- Stable
- Very weak interaction with other particles



Fermilab 95-759

- Major constituent is perhaps heavy (massive) particles (non-relativistic while decoupling)
- Mainly non-baryonic in nature

# *Dark Matter Hunt*

## Through Direct Detection

CDMS II

CoGeNT

CRESST II



XENON 100

LUX

## Through Indirect Detection

Final products of Dark Matter Annihilation, e.g. Gamma Ray, neutrinos etc.


# Direct Detection Results




# Indirect Detection (Signatures?)

- ➔ **Excess Gamma Emission from Galactic Centre Region**  
(*Fermi Gamma Ray Space Telescope (FGST) searches Gamma ray around 5 degrees surrounding GC*)
  - ➔ **Gamma Emission from *Fermi Bubble***
- ➔ **Excess positron fraction (AMS 02 @ ISS, PAMELA etc.)**
  - ➔ **Antiproton excess (BESS)**
  - ➔ **Neutrinos at ICECUBE, ANTARES ?**


# Hint of Dark Matter around 10 GeV



***Observed gamma flux bump by FGST***




# Fits to DM proposals







# Fermi Bubble



From end to end, the newly discovered gamma-ray bubbles extend 50,000 light-years, or roughly half of the Milky Way's diameter. Hints of the bubbles' edges were first observed in X-rays (blue) by ROSAT, a Germany-led mission operating in the 1990s. The gamma rays mapped by Fermi (magenta) extend much farther from the galaxy's plane.  
**Credit:** NASA's Goddard Space Flight Center



# **A Model for Dark Matter that may simultaneously explain**

 **Direct detection results**

 **Planck results for DM abundance**

 **10 GeV Gamma ray excess from  
Galactic Centre region**

 **Low energy gamma emission from  
Fermi Bubble**

# Two Scalar Singlet Model for Dark Matter

We Propose SM with additional 2 SM gauge singlets  $S_1, S_2$

Stability ensured by  $\mathbb{Z}_2 \times \mathbb{Z}_2$  or  $\mathbb{Z}_2 \times \mathbb{Z}'_2$

$$\begin{pmatrix} S \\ S' \end{pmatrix} \xrightarrow{\mathbb{Z}_2 \times \mathbb{Z}_2} \begin{pmatrix} -S \\ -S' \end{pmatrix} \quad S \xrightarrow{\mathbb{Z}_2} -S \quad \text{and} \quad S' \xrightarrow{\mathbb{Z}'_2} -S'$$

$$\mathbb{Z}_2 \times \mathbb{Z}_2$$

$$\begin{aligned}
 V(H, S, S') &= \frac{m^2}{2} H^\dagger H + \frac{\lambda}{4} (H^\dagger H)^2 \\
 &+ \frac{\delta_2}{2} H^\dagger H S^2 + \frac{k_2}{2} S^2 + \frac{k_4}{4} S^4 \\
 &+ \frac{\delta'_2}{2} H^\dagger H S'^2 + \frac{k'_2}{2} S'^2 + \frac{k'_4}{4} S'^4 \\
 &+ \frac{\delta''_2}{2} H^\dagger H S' S + \frac{k''_2}{2} S S' \\
 &+ \frac{1}{4} (k_4^a S S S' S' + k_4^b S S S S' + k_4^c S S' S' S')
 \end{aligned}$$

**After SSB**

$$M_{SS'} = \begin{pmatrix} k_2 + \delta_2 v^2/2 & \delta''_2 v^2/4 + k''_2/2 \\ \delta''_2 v^2/4 + k''_2/2 & k'_2 + \delta'_2 v^2/2 \end{pmatrix} \equiv \begin{pmatrix} M_{11} & M_{12} \\ M_{12} & M_{22} \end{pmatrix}$$

$$M_{S_1}^2 = \cos^2 \theta M_{11} + \sin^2 \theta M_{22} + 2 \cos \theta \sin \theta M_{12}$$

$$M_{S_2}^2 = \cos^2 \theta M_{22} + \sin^2 \theta M_{11} - 2 \cos \theta \sin \theta M_{12}$$

$$\tan 2\theta = \frac{2M_{12}}{M_{11} - M_{22}}$$

$$\mathbb{Z}_2 \times \mathbb{Z}'_2$$

$$\begin{aligned}
 V(H, S, S') &= \frac{m^2}{2} H^\dagger H + \frac{\lambda}{4} (H^\dagger H)^2 \\
 &+ \frac{\delta_2}{2} H^\dagger H S^2 + \frac{k_2}{2} S^2 + \frac{k_4}{4} S^4 \\
 &+ \frac{\delta'_2}{2} H^\dagger H S'^2 + \frac{k'_2}{2} S'^2 + \frac{k'_4}{4} S'^4 \\
 &+ \frac{1}{4} (k_4^a S S S' S')
 \end{aligned}$$

$$\begin{aligned}
 M_S^2 &= k_2 + \frac{\delta_2 v^2}{2} \\
 M_{S'}^2 &= k'_2 + \frac{\delta'_2 v^2}{2}
 \end{aligned}$$

## Vacuum Stability Bounds

$$\lambda \geq 0, k_4 \geq 0, k_4^p \geq 0, \quad \delta_2 + \sqrt{\lambda k_4} \geq 0,$$

$$\delta_2^p + \sqrt{\lambda k_4^p} \geq 0,$$

$$k_4^a + \sqrt{k_4 k_4^p} \geq 0,$$

$$\sqrt{\lambda k_4 k_4^p} + \delta_2 \sqrt{k_4^p} + \delta_2^p \sqrt{k_4} + 2k_4^p \sqrt{\lambda} + \sqrt{(\delta_2 + \sqrt{\lambda k_4})(\delta_2^p + \sqrt{\lambda k_4^p})(k_4^a + \sqrt{k_4 k_4^p})} \geq 0.$$

## Perturbative Unitarity Bounds

$$|\delta_2| \leq 8\pi \quad |k_4^a| \leq 8\pi$$

$$k_4 \leq \frac{8}{6}\pi \quad k_4' \leq \frac{8}{6}\pi$$

# The Potential

$$\begin{aligned}
 V(H, S, S') &= \frac{m^2}{2} H^\dagger H + \frac{\lambda}{4} (H^\dagger H)^2 \\
 &+ \frac{\delta_1}{2} H^\dagger H S + \frac{\delta_2}{2} H^\dagger H S^2 + \frac{\delta_1 m}{2\lambda} S + \frac{k_2}{2} S^2 + \frac{k_3}{3} S^3 + \frac{k_4}{4} S^4 \\
 &+ \frac{\delta'_1}{2} H^\dagger H S' + \frac{\delta'_2}{2} H^\dagger H S'^2 + \frac{\delta'_1 m}{2\lambda} S' + \frac{k'_2}{2} S'^2 + \frac{k'_3}{3} S'^3 + \frac{k'_4}{4} S'^4 \\
 &+ \frac{\delta''_2}{2} H^\dagger H S' S + \frac{k''_2}{2} S S' + \frac{1}{3} (k_3^a S S S' + k_3^b S S' S') \\
 &+ \frac{1}{4} (k_4^a S S S' S' + k_4^b S S S S' + k_4^c S S' S' S')
 \end{aligned}$$

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \longrightarrow \delta_1 = k_3 = \delta'_1 = k'_3 = k_3^a = k_3^b = 0$$

$$\mathbb{Z}_2 \times \mathbb{Z}'_2 \longrightarrow \delta''_2 = k''_2 = k_4^b = k_4^c = 0 \quad (\text{in addition})$$



# Relic Density Calculation for comparing with Planck result

$$0.1165 < \Omega_{\text{DM}} h^2 < 0.1227$$


## Boltzmann Eqns.

$$\begin{aligned} \frac{dn_1}{dt} + 3Hn_1 &= -\langle\sigma v\rangle_{11\rightarrow XX}(n_1^2 - n_{1\text{eq}}^2) - \langle\sigma v\rangle_{11\rightarrow 22} \left( n_1^2 - \frac{n_{1\text{eq}}^2}{n_{2\text{eq}}^2} n_2^2 \right) \\ \frac{dn_2}{dt} + 3Hn_2 &= -\langle\sigma v\rangle_{22\rightarrow XX}(n_2^2 - n_{2\text{eq}}^2) - \langle\sigma v\rangle_{22\rightarrow 11} \left( n_2^2 - \frac{n_{2\text{eq}}^2}{n_{1\text{eq}}^2} n_1^2 \right) \end{aligned}$$


$$\text{Total relic density } \Omega = \Omega_S + \Omega_{S'}$$

$$\langle\sigma v\rangle_{11\rightarrow XX}, \langle\sigma v\rangle_{22\rightarrow XX} \gg \langle\sigma v\rangle_{11\rightarrow 22} \Rightarrow \text{decoupled}$$

# Annihilation Channels




# Direct Detection




$$\sigma_{\text{nucleon}}^{\text{SI}} = (\delta_2)^2 \left( \frac{100 \text{ GeV}}{M_h} \right)^4 \left( \frac{50 \text{ GeV}}{M_S} \right)^2 (5 \times 10^{-42} \text{ cm}^2)$$

# Direct Detection Bounds



CDMS collaboration,  
[arXiv:1304.4279 \[hep-ex\]](https://arxiv.org/abs/1304.4279)

# CDMS II + PLANCK




*The hatched olive region is the parameter space allowed in the  $M_{S'} - \delta'_2$  (or  $M_S - \delta_2$ ) plane by CDMS II 90%CL data. Choosing  $(M_S = 8.6 \text{ GeV}, \delta_2 = 0.45)$  in the  $M_S - \delta_2$  plane, the only parameter space allowed in the  $M_{S'} - \delta'_2$  by Planck data is shown as the blue shaded region.*

# CDMS II + PLANCK


$M_S$ (or $M_{S'}$ ) (GeV)	$\delta_2$ (or $\delta'_2$ )	$\sigma^{\text{SI}}$ ( $\times 10^{-41}$ cm <sup>2</sup> )	$\langle\sigma v\rangle$ ( $\times 10^{-26}$ cm <sup>3</sup> /s)	Contribution in $\Omega_S$ (or $\Omega_{S'}$ )
8.6	0.45	1.9	3.2 { 2.6 ( $b\bar{b}$ ) 0.4 ( $c\bar{c}$ ) 0.2 ( $l\bar{l}$ )	81% 12% 7%
6.7	0.82	9.9	8.3 { 6.3 ( $b\bar{b}$ ) 1.2 ( $c\bar{c}$ ) 7.2 ( $l\bar{l}$ )	77% 15% 8%

# CoGeNT + PLANCK




$M_S$ (or $M_{S'}$ ) (GeV)	$\delta_2$ (or $\delta_2'$ )	$\sigma^{\text{SI}}$ ( $\times 10^{-41}$ cm $^2$ )	$\langle\sigma v\rangle$ ( $\times 10^{-26}$ cm $^3$ /s)	Contribution in $\Omega_S$ (or $\Omega_{S'}$ )
7.8	0.56	9.0	4.6 $\left\{ \begin{array}{l} 3.7 (b\bar{b}) \\ 0.6 (c\bar{c}) \\ 0.4 (l\bar{l}) \end{array} \right.$	80% 12.5% 7.5%
8.2	0.61	9.8	5.7 $\left\{ \begin{array}{l} 4.6 (b\bar{b}) \\ 0.7 (c\bar{c}) \\ 0.4 (l\bar{l}) \end{array} \right.$	81% 12% 7%

# CRESST II + PLANCK




$M_S$ (or $M_{S'}$ ) (GeV)	$\delta_2$ (or $\delta'_2$ )	$\sigma^{\text{SI}}$ ( $\times 10^{-41}$ cm $^2$ )	$\langle \sigma v \rangle$ ( $\times 10^{-26}$ cm $^3$ /s)	Contribution in $\Omega_S$ (or $\Omega_{S'}$ )	
25.3	0.36	1.6	3.0	$\left\{ \begin{array}{l} 2.4 (b\bar{b}) \\ 0.4 (c\bar{c}) \\ 0.2 (l\bar{l}) \end{array} \right.$	$\left\{ \begin{array}{l} 81\% \\ 12\% \\ 7\% \end{array} \right.$
23.3	0.47	3.2	4.8	$\left\{ \begin{array}{l} 3.9 (b\bar{b}) \\ 0.6 (c\bar{c}) \\ 0.3 (l\bar{l}) \end{array} \right.$	$\left\{ \begin{array}{l} 81\% \\ 12\% \\ 7\% \end{array} \right.$






# XENON 100 + PLANCK




# XENON 100 + PLANCK

$M_S$ (or $M_{S'}$ ) (GeV)	$\delta_2$ (or $\delta'_2$ )	$\sigma^{\text{SI}}$ ( $\times 10^{-45}$ cm <sup>2</sup> )	$\langle \sigma v \rangle$ ( $\times 10^{-26}$ cm <sup>3</sup> /s)	Contribution in $\Omega_S$ (or $\Omega_{S'}$ )
54.0	0.022	1.4	9.94	$\left\{ \begin{array}{l} 7.83 (b\bar{b}) \\ 1.29 (c\bar{c}) \\ 0.82 (l\bar{l}) \end{array} \right.$ 78.8% 13.2% 8.2%
56.0	0.011	0.8	3.95	$\left\{ \begin{array}{l} 3.11 (b\bar{b}) \\ 0.52 (c\bar{c}) \\ 0.32 (l\bar{l}) \end{array} \right.$ 78.7% 13.2% 8.1%
90.0	0.050	2.6	4.32	$\left\{ \begin{array}{l} 4.29 (W^+W^-) \\ 0.024 (b\bar{b}) \\ 0.004 (c\bar{c}) \\ 0.002 (l\bar{l}) \end{array} \right.$ 99.3% 0.55% 0.10% 0.05%
92.0	0.045	2.0	3.78	$\left\{ \begin{array}{l} 3.28 (W^+W^-) \\ 0.48 (ZZ) \\ 0.016 (b\bar{b}) \\ 0.003 (c\bar{c}) \\ 0.002 (l\bar{l}) \end{array} \right.$ 86.6% 12.8% 0.43% 0.08% 0.05%



# LUX + PLANCK



# Dark Matter Halo Profile

$$\rho(r) = \rho_0 F_{\text{halo}}(r) = \frac{\rho_0}{(r/r_c)^\gamma [1 + (r/r_c)^\gamma]^{(\beta-\gamma)/\alpha}}, \quad \rho_0 = 0.4 \text{ GeV/cm}^3$$


Halo Model	$\alpha$	$\beta$	$\gamma$	$r_c$ (Kpc)
Navarro, Frenk, White (NFW)	1	3	1	20
Moore	1.5	3	1.5	28
Isothermal	2	2	0	3.5

Einasto halo profile:


$$F_{\text{halo}}^{\text{Ein}}(r) = \exp \left[ \frac{-2}{\tilde{\alpha}} \left( \left( \frac{r}{r_\odot} \right)^{\tilde{\alpha}} - 1 \right) \right], \quad \tilde{\alpha} = 0.17$$

Isothermal  $\rightarrow$  flat profile, Moore, Einasto  $\rightarrow$  cuspy profile


# CDMS II + PLANCK for GC gamma excess




(a) Isothermal



(b) NFW







(c) Einasto







(d) Moore





# CoGeNT + PLANCK for GC gamma excess



# CRESST II+PLANCK for GC gamma excess




# XENON100+PLANCK for GC gamma excess







# XENON100+PLANCK for GC gamma excess




(a) Isothermal



(b) NFW




(c) Einasto




(d) Moore


# Comparing with Fermi Bubble




(a) Benchmark point 1




(b) Benchmark point 2



(c) Benchmark point 3



(d) Benchmark point 4A



(e) Benchmark point 4B

# Conclusions

- ➔ **Two real scalar SM gauge singlets odd under  $\mathbb{Z}_2 \times \mathbb{Z}'_2$**
- ➔ **(CDMS II, CoGeNT, CRESST II) + Planck + Galactic Centre + Fermi Bubble prefer flat DM halo profile like isothermal/Burkart**
- ➔ **(Xenon 100, LUX) + Planck + Galactic Centre + Fermi Bubble prefer cuspy nature of DM halo profiles such as Moore/Einasto**
- ➔ **This model provides an explanation for excess low energy gamma ray emission from Galactic Centre region as also from the low latitude Fermi Bubble region from the annihilation of Dark Matter**