

A primal-dual interior point method for nonlinear semidefinite programming ^{*†}

Hiroshi Yamashita[‡], Hiroshi Yabe[§] and Kouhei Harada[¶]

September 3, 2006 (revised August 31, 2007)

Abstract

In this paper, we consider a primal-dual interior point method for solving nonlinear semidefinite programming problems. By combining the primal barrier penalty function and the primal-dual barrier function, a new primal-dual merit function is proposed within the framework of the line search strategy. We show the global convergence property of our method. Finally some numerical experiments are given.

Key words. nonlinear semidefinite programming, primal-dual interior point method, barrier penalty function, primal-dual merit function, global convergence

1 Introduction

This paper is concerned with the following nonlinear semidefinite programming (SDP) problem:

$$(1) \quad \begin{array}{ll} \text{minimize} & f(x), \quad x \in \mathbb{R}^n, \\ \text{subject to} & g(x) = 0, \quad X(x) \succeq 0 \end{array}$$

where we assume that the functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $X : \mathbb{R}^n \rightarrow \mathbb{S}^p$ are sufficiently smooth, where \mathbb{S}^p denotes the set of p th order real symmetric matrices. By $X(x) \succeq 0$ and $X(x) \succ 0$, we mean that the matrix $X(x)$ is positive semidefinite and positive definite, respectively.

The problem (1) is an extension of the linear SDP problem. For the case of the linear SDP problems, the matrix $X(x)$ is defined by

$$X(x) = \sum_{i=1}^n x_i A_i - B$$

^{*}The second author was supported in part by the Grant-in-Aid for Scientific Research (C) 16510123 of Japan Society for the Promotion of Science.

[†]The original publication is available at www.springerlink.com.

[‡]Mathematical Systems Inc., 2-4-3, Shinjuku, Shinjuku-ku, Tokyo 160-0022, Japan. hy@msi.co.jp

[§]Department of Mathematical Information Science, Faculty of Science, Tokyo University of Science, 1-3, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan. yabe@rs.kagu.tus.ac.jp

[¶]Mathematical Systems Inc., 2-4-3, Shinjuku, Shinjuku-ku, Tokyo 160-0022, Japan. harada@msi.co.jp

with given matrices $A_i \in \mathbb{S}^p, i = 1, \dots, n$, and $B \in \mathbb{S}^p$. The linear SDP problems include linear programming problems, convex quadratic programming problems and second order cone programming problems, and they have many applications. Interior point methods for the linear SDP problems have been studied extensively by many researchers, see for example [1, 12, 13, 14, 16] and the references therein.

On the other hand, researches on numerical methods for nonlinear SDP are much more recent, and a few researchers have been studying these methods. For example, Kočvara and Stingl [9] developed a computer code PENNON for solving nonlinear SDP, in which the augmented Lagrangian function method was used. Correa and Ramirez [3] proposed an algorithm which used the sequentially linear SDP method. Related researches include Jarre [6], in which examples of nonlinear SDP problems were introduced, and Freund and Jarre [5]. Fares, Noll and Apkarian [4] applied the sequential linear SDP method to robust control problems. Recently Kanzow, Nagel, Kato and Fukushima [7] presented a successive linearization method with a trust region-type globalization strategy. However, no interior point type method for general nonlinear SDP problems has been proposed yet to our knowledge.

In this paper, we propose a globally convergent primal-dual interior point method for solving nonlinear SDP problems. The method is based on a line search algorithm in the primal-dual space. The present paper is organized as follows. In Section 2, the optimality conditions for problem (1) are described. In Sections 3 and 4, our primal-dual interior point method is proposed. Specifically, Section 3 presents the algorithm called SDPIP which constitutes the basic frame of primal-dual interior point methods. Section 4 gives the algorithm called SDPLS based on the line search strategy, which is an inner iteration of algorithm SDPIP given in Section 3. In Section 4.1, we describe the Newton method for solving nonlinear equations that are obtained by modifying the optimality conditions given in Section 2. In Section 4.2, we propose a new primal-dual merit function that consists of the primal barrier penalty function and the primal-dual barrier function. Then Section 4.3 presents algorithm SDPLS, and Section 5 shows its global convergence property. Furthermore, some numerical experiments are presented in Section 6. Finally, we give some concluding remarks in Section 7.

Throughout this paper, we define the inner product $\langle X, Z \rangle$ by $\langle X, Z \rangle = \text{tr}(XZ)$ for any matrices X and Z in \mathbb{S}^p , where $\text{tr}(M)$ denotes the trace of the matrix M . In this paper, $(v)_i$ denotes the i th element of the vector v if necessary.

2 Optimality conditions

Let the Lagrangian function of problem (1) be defined by

$$L(w) = f(x) - y^T g(x) - \langle X(x), Z \rangle,$$

where $w = (x, y, Z)$, and $y \in \mathbb{R}^m$ and $Z \in \mathbb{S}^p$ are the Lagrange multiplier vector and matrix which correspond to the equality and positive semidefiniteness constraints, respectively. We also define matrices

$$A_i(x) = \frac{\partial X}{\partial x_i}$$

for $i = 1, \dots, n$. Then Karush-Kuhn-Tucker (KKT) conditions for optimality of problem (1) are given by the following (see [2]):

$$(2) \quad r_0(w) \equiv \begin{pmatrix} \nabla_x L(w) \\ g(x) \\ X(x)Z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

and

$$(3) \quad X(x) \succeq 0, \quad Z \succeq 0.$$

Here $\nabla_x L(w)$ is given by

$$\begin{aligned} \nabla_x L(w) &= \nabla f(x) - A_0(x)^T y - \mathcal{A}^*(x)Z, \\ A_0(x) &= \begin{pmatrix} \nabla g_1(x)^T \\ \vdots \\ \nabla g_m(x)^T \end{pmatrix} \in \mathbb{R}^{m \times n}, \end{aligned}$$

where $\mathcal{A}^*(x)$ is an operator which yields

$$\mathcal{A}^*(x)Z = \begin{pmatrix} \langle A_1(x), Z \rangle \\ \vdots \\ \langle A_n(x), Z \rangle \end{pmatrix}.$$

In the following we will occasionally deal with the multiplication $X(x) \circ Z$ which is defined by

$$X(x) \circ Z = \frac{X(x)Z + ZX(x)}{2}$$

instead of $X(x)Z$. It is known that $X(x) \circ Z = 0$ is equivalent to the relation $X(x)Z = ZX(x) = 0$.

We call $w = (x, y, Z)$ satisfying $X(x) \succ 0$ and $Z \succ 0$ the interior point. The algorithm of this paper will generate such interior points. To construct an interior point algorithm, we introduce a positive parameter μ , and we replace the complementarity condition $X(x)Z = 0$ by $X(x)Z = \mu I$, where I denotes the identity matrix. Then we try to find a point that satisfies the barrier KKT (BKKT) conditions:

$$(4) \quad r(w, \mu) \equiv \begin{pmatrix} \nabla_x L(w) \\ g(x) \\ X(x)Z - \mu I \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

and

$$X(x) \succ 0, \quad Z \succ 0.$$

3 Algorithm for finding a KKT point

We first describe a procedure for finding a KKT point using the BKKT conditions. In this section, the subscript k denotes an iteration count of the outer iterations. We define

the norm $\|r(w, \mu)\|$ by

$$\|r(w, \mu)\| = \sqrt{\left\| \begin{pmatrix} \nabla_x L(w) \\ g(x) \end{pmatrix} \right\|^2 + \|X(x)Z - \mu I\|_F^2},$$

where $\|\cdot\|$ denotes the l_2 norm for vectors and $\|\cdot\|_F$ denotes the Frobenius norm for matrices. We also define $\|r_0(w)\|$ by $\|r_0(w)\| = \|r(w, 0)\|$.

Now we present the algorithm called SDPIP which calculates a KKT point.

Algorithm SDPIP

Step 0. (Initialize) Set $\varepsilon > 0$, $M_c > 0$ and $k = 0$. Let a positive sequence $\{\mu_k\}$, $\mu_k \downarrow 0$ be given.

Step 1. (Approximate BKKT point) Find an interior point w_{k+1} that satisfies

$$(5) \quad \|r(w_{k+1}, \mu_k)\| \leq M_c \mu_k.$$

Step 2. (Termination) If $\|r_0(w_{k+1})\| \leq \varepsilon$, then stop.

Step 3. (Update) Set $k := k + 1$ and go to Step 1. □

We note that the barrier parameter sequence $\{\mu_k\}$ in Algorithm SDPIP needs not be determined beforehand. The value of each μ_k may be set adaptively as the iteration proceeds. We call condition (5) the approximate BKKT condition, and call a point that satisfies this condition the approximate BKKT point.

If the matrix $A_0(x_*)$ is of full rank and there exists a nonzero vector $v \in \mathbb{R}^n$ such that

$$A_0(x_*)v = 0 \quad \text{and} \quad X(x_*) + \sum_{i=1}^n v_i A_i(x_*) \succ 0,$$

then we say that the Mangasarian-Fromovitz constraint qualification (MFCQ) condition is satisfied at a point x_* (see [3] for example).

The following theorem shows the convergence property of Algorithm SDPIP.

Theorem 1 *Assume that the functions f and g are continuously differentiable. Let $\{w_k\}$ be an infinite sequence generated by Algorithm SDPIP. Suppose that the sequence $\{x_k\}$ is bounded and that the MFCQ condition is satisfied at any accumulation point of the sequence $\{x_k\}$. Then the sequences $\{y_k\}$ and $\{Z_k\}$ are bounded, and any accumulation point of $\{w_k\}$ satisfies KKT conditions (2) and (3).*

Proof. To prove this theorem by contradiction, we suppose that either $\{y_k\}$ or $\{Z_k\}$ is not bounded, i.e.

$$(6) \quad \gamma_k \equiv \max \{|(y_k)_1|, \dots, |(y_k)_m|, \lambda_{\max}(Z_k)\} \rightarrow \infty,$$

where $\lambda_{\max}(Z_k)$ denotes the largest eigenvalue of the matrix Z_k . It follows from (5) that the boundedness of $\{x_k\}$ implies

$$\limsup_{k \rightarrow \infty} \|A_0(x_k)^T y_k + \mathcal{A}^*(x_k) Z_k\| < \infty.$$

Then we have $\|A_0(x_k)^T y_k / \gamma_k + \mathcal{A}^*(x_k) Z_k / \gamma_k\| \rightarrow 0$. Letting an arbitrary accumulation point of $\{x_k, y_k / \gamma_k, Z_k / \gamma_k\}$ be (x_*, y_*, Z_*) , we have

$$(7) \quad A_0(x_*)^T y_* + \mathcal{A}^*(x_*) Z_* = 0 \quad \text{and} \quad X_* Z_* = Z_* X_* = 0,$$

where $X_* = X(x_*)$. We will prove that $Z_* = 0$. For this purpose, we assume that $\lambda_{\max}(Z_*) > 0$ holds. Since the matrices X_* and Z_* commute, they share the same eigen-system. Thus the matrices X_* and Z_* can be transformed to the diagonal matrices by using the same orthogonal matrix P as follows:

$$\bar{X}_* \equiv P X_* P^T = \text{diag}(\lambda_1, \dots, \lambda_p) \quad \text{and} \quad \bar{Z}_* \equiv P Z_* P^T = \text{diag}(\tau_1, \dots, \tau_p),$$

where $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_p$ and $\tau_1 \leq \tau_2 \leq \dots \leq \tau_p$ are the nonnegative eigenvalues of X_* and Z_* , respectively. It follows from the assumption that there exists an integer p' such that $1 \leq p' < p$, $\lambda_{p'} = 0$ and $\lambda_{p'+1} > 0$ hold. Furthermore, the MFCQ condition implies that there exists a nonzero vector $v \in \mathbb{R}^n$ which satisfies

$$A_0(x_*) v = 0 \quad \text{and} \quad X_* + \sum_{i=1}^n v_i A_i(x_*) \succ 0.$$

Therefore, we get

$$(8) \quad (\bar{X}_*)_{jj} + \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} > 0$$

for $j = 1, \dots, p$, where $\bar{A}_i(x_*) = P A_i(x_*) P^T$. Since the following holds

$$0 = \lambda_j = (\bar{X}_*)_{jj} \quad \text{for} \quad j = 1, \dots, p',$$

equation (8) yields

$$(9) \quad \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} > 0 \quad \text{for} \quad j = 1, \dots, p'.$$

By premultiplying (7) by v^T , we have

$$\begin{aligned} 0 &= v^T A_0(x_*)^T y_* + v^T \mathcal{A}^*(x_*) Z_* = v^T \mathcal{A}^*(x_*) Z_* = \sum_{i=1}^n v_i \text{tr} \{A_i(x_*) Z_*\} \\ &= \sum_{i=1}^n v_i \text{tr} \{\bar{A}_i(x_*) \bar{Z}_*\} = \sum_{j=1}^p \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} \tau_j \\ &= \sum_{j=1}^{p'} \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} \tau_j + \sum_{j=p'+1}^p \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} \tau_j. \end{aligned}$$

Since the complementarity condition $\bar{X}_* \bar{Z}_* = 0$ implies $\tau_j = 0$ for $j = p' + 1, \dots, p$, the equation above yields

$$\sum_{j=1}^{p'} \sum_{i=1}^n v_i (\bar{A}_i(x_*))_{jj} \tau_j = 0.$$

By (9), we have $\tau_j = 0$ for $j = 1, \dots, p'$, which contradicts the assumption $\lambda_{\max}(Z_*) > 0$. Therefore we obtain $Z_* = 0$, which yields $A_0(x_*)^T y_* = 0$ from (7). Since the matrix $A_0(x_*)$ is of full rank, we have $y_* = 0$. This contradicts the fact that some element of y_* or Z_* is not zero by (6). Therefore, the sequences $\{y_k\}$ and $\{Z_k\}$ are bounded.

Let \hat{w} be any accumulation point of $\{w_k\}$. Since the sequences $\{w_k\}$ and $\{\mu_k\}$ satisfy (5) for each k and μ_k approaches zero, $r_0(\hat{w}) = 0$ follows from the definition of $r(w, \mu)$.

Therefore the proof is complete. \square

4 Algorithm for finding a barrier KKT point

As same as the case of linear SDP problems, we consider a scaling of the primal-dual pair $(X(x), Z)$ in applying the Newton method to the system of equations (4). In what follows, we denote $X(x)$ simply by X if it is not confused. We define a transformation $T \in \mathbb{R}^{p \times p}$, and we scale X and Z by

$$\tilde{X} = T X T^T \quad \text{and} \quad \tilde{Z} = T^{-T} Z T^{-1}$$

respectively. Using the transformation T , we replace the equation $XZ = \mu I$ by a form $\tilde{X} \circ \tilde{Z} = \mu I$, and deal with the scaled symmetrized residual:

$$(10) \quad \tilde{r}_S(w, \mu) \equiv \begin{pmatrix} \nabla_x L(w) \\ g(x) \\ \tilde{X} \circ \tilde{Z} - \mu I \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

instead of (4) to form Newton directions as described below.

4.1 Newton method

In this section, we consider a method for solving the BKKT conditions approximately for a given $\mu > 0$, which corresponds to the inner iterations of Step 1 of Algorithm SDPIP. Throughout this section, we assume that $X \succ 0$ and $Z \succ 0$ hold.

We apply a Newton-like method to the system of equations (10). Let the Newton directions for the primal and dual variables by $\Delta x \in \mathbb{R}^n$ and $\Delta Z \in \mathbb{S}^p$, respectively. We define $\Delta X = \sum_{i=1}^n \Delta x_i A_i(x)$ and we note that $\Delta X \in \mathbb{S}^p$. We also scale ΔX and ΔZ by

$$\Delta \tilde{X} = T \Delta X T^T \quad \text{and} \quad \Delta \tilde{Z} = T^{-T} \Delta Z T^{-1}.$$

Since $(\tilde{X} + \Delta \tilde{X}) \circ (\tilde{Z} + \Delta \tilde{Z}) = \mu I$ can be written as

$$(\tilde{X} + \Delta \tilde{X})(\tilde{Z} + \Delta \tilde{Z}) + (\tilde{Z} + \Delta \tilde{Z})(\tilde{X} + \Delta \tilde{X}) = 2\mu I,$$

neglecting the nonlinear parts $\Delta\tilde{X}\Delta\tilde{Z}$ and $\Delta\tilde{Z}\Delta\tilde{X}$ implies the Newton equation for (10)

$$(11) \quad G\Delta x - A_0(x)^T\Delta y - \mathcal{A}^*(x_*)\Delta Z = -\nabla_x L(x, y, Z)$$

$$(12) \quad A_0(x)\Delta x = -g(x)$$

$$(13) \quad \Delta\tilde{X}\tilde{Z} + \tilde{Z}\Delta\tilde{X} + \tilde{X}\Delta\tilde{Z} + \Delta\tilde{Z}\tilde{X} = 2\mu I - \tilde{X}\tilde{Z} - \tilde{Z}\tilde{X},$$

where G denotes the Hessian matrix of the Lagrangian function $L(w)$ or its approximation (see Remark 2 in Section 4.3).

Similarly to usual primal-dual interior point methods for linear SDP problems, we derive an explicit form of the direction $\Delta Z \in \mathbb{S}^p$ from equation (13) and substitute it into equation (11) in order to obtain the Newton direction $\Delta w = (\Delta x, \Delta y, \Delta Z) \in \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^p$. For this purpose, we make use of relations described in [1] and Appendix of [13]. For $U \in \mathbb{S}^p$, nonsingular $P \in \mathbb{R}^{p \times p}$ and $Q \in \mathbb{R}^{p \times p}$, we define the operator

$$(P \odot Q)U = \frac{1}{2}(PUQ^T + QUP^T)$$

and the symmetrized Kronecker product

$$(P \otimes_S Q)\text{svec}(U) = \text{svec}((P \odot Q)U),$$

where the operator svec is defined by

$$\text{svec}(U) = (U_{11}, \sqrt{2}U_{21}, \dots, \sqrt{2}U_{p1}, U_{22}, \sqrt{2}U_{32}, \dots, \sqrt{2}U_{p2}, U_{33}, \dots, U_{pp})^T \in \mathbb{R}^{p(p+1)/2}.$$

We note that, for any $U, V \in \mathbb{S}^p$,

$$(14) \quad \langle U, V \rangle = \text{tr}(UV) = \text{svec}(U)^T \text{svec}(V)$$

holds. By using the operator, the matrices \tilde{X} , \tilde{Z} , $\Delta\tilde{X}$ and $\Delta\tilde{Z}$ can be represented by

$$(15) \quad \tilde{X} = (T \odot T)X, \quad \tilde{Z} = (T^{-T} \odot T^{-T})Z,$$

$$(16) \quad \Delta\tilde{X} = (T \odot T)\Delta X \quad \text{and} \quad \Delta\tilde{Z} = (T^{-T} \odot T^{-T})\Delta Z.$$

Let $P' \in \mathbb{R}^{p \times p}$ and $Q' \in \mathbb{R}^{p \times p}$ be nonsingular, and $V \in \mathbb{S}^p$. By denoting the inverse operator of svec by smat , we have

$$(17) \quad (P \odot Q)U = \text{smat}((P \otimes_S Q)\text{svec}(U)).$$

We also define

$$(18) \quad (P \odot Q)^{-1}U = \text{smat}((P \otimes_S Q)^{-1}\text{svec}(U)).$$

The expressions above give

$$\begin{aligned} (P \odot Q)(P' \odot Q')U &= \text{smat}((P \otimes_S Q)\text{svec}((P' \odot Q')U)) \\ &= \text{smat}((P \otimes_S Q)(P' \otimes_S Q')\text{svec}(U)) \end{aligned}$$

and

$$\{(P \odot Q)(P' \odot Q')\}^{-1}U = (P' \odot Q')^{-1}(P \odot Q)^{-1}U.$$

Furthermore, we get

$$\begin{aligned}
\langle U, (P \odot Q)V \rangle &= \text{tr} \{U(P \odot Q)V\} \\
&= \frac{1}{2} \text{tr} \{U(PVQ^T + QVP^T)\} \\
&= \frac{1}{2} \text{tr} \{Q^T U P V + P^T U Q V\} \\
&= \text{tr} \{((P^T \odot Q^T)U)V\} \\
(19) \quad &= \langle (P^T \odot Q^T)U, V \rangle
\end{aligned}$$

and

$$\begin{aligned}
\langle U, (P \odot Q)^{-1}V \rangle &= \text{tr} \{U(P \odot Q)^{-1}V\} \\
&= \text{tr} \{((P^T \odot Q^T)(P^T \odot Q^T)^{-1}U)(P \odot Q)^{-1}V\} \\
&= \text{tr} \{((P^T \odot Q^T)^{-1}U)(P \odot Q)(P \odot Q)^{-1}V\} \\
&= \text{tr} \{((P^T \odot Q^T)^{-1}U)V\} \\
&= \langle (P^T \odot Q^T)^{-1}U, V \rangle.
\end{aligned}$$

Now we have the following theorem that gives the Newton directions.

Theorem 2 *Suppose that the operator $\tilde{X} \odot I$ is invertible. Then the direction $\Delta\tilde{Z} \in \mathbb{S}^p$ is given by the form*

$$(20) \quad \Delta\tilde{Z} = \mu\tilde{X}^{-1} - \tilde{Z} - (\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)\Delta\tilde{X},$$

or equivalently

$$(21) \quad \Delta Z = \mu X^{-1} - Z - (T^T \odot T^T)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)\Delta X,$$

and the directions $(\Delta x, \Delta y) \in \mathbb{R}^n \times \mathbb{R}^m$ satisfy

$$(22) \quad \begin{pmatrix} G + H & -A_0(x)^T \\ -A_0(x) & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta y \end{pmatrix} = - \begin{pmatrix} \nabla f(x) - A_0(x)^T y - \mu \mathcal{A}^*(x) X^{-1} \\ -g(x) \end{pmatrix},$$

where the elements of the matrix H are represented by the form

$$(23) \quad H_{ij} = \langle \tilde{A}_i(x), (\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)\tilde{A}_j(x) \rangle$$

with $\tilde{A}_i(x) = T A_i(x) T^T$.

Furthermore, if the matrix $G + H$ is positive definite and the matrix $A_0(x)$ is of full rank, then the Newton equations (11) – (13) give a unique search direction $\Delta w = (\Delta x, \Delta y, \Delta Z) \in \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^p$.

Proof. By equation (13), we have

$$2(\tilde{Z} \odot I)\Delta\tilde{X} + 2(\tilde{X} \odot I)\Delta\tilde{Z} = 2\mu(\tilde{X} \odot I)\tilde{X}^{-1} - 2(\tilde{X} \odot I)\tilde{Z},$$

which implies that

$$(\tilde{X} \odot I) \left(\tilde{Z} + \Delta\tilde{Z} - \mu\tilde{X}^{-1} \right) = -(\tilde{Z} \odot I)\Delta\tilde{X}.$$

Thus we obtain equation (20). Since $(T^{-T} \otimes_S T^{-T})^{-1} = (T^{-T})^{-1} \otimes_S (T^{-T})^{-1} = T^T \otimes_S T^T$ holds (see Appendix of [13]), it follows from (18) and (17) that for any $U \in \mathbb{S}^p$,

$$\begin{aligned} (T^{-T} \odot T^{-T})^{-1}U &= \text{smat} \left((T^{-T} \otimes_S T^{-T})^{-1} \text{svec}(U) \right) \\ &= \text{smat} \left((T^T \otimes_S T^T) \text{svec}(U) \right) \\ &= (T^T \odot T^T)U. \end{aligned}$$

By (15) and (16), equation (20) implies that

$$\Delta Z = \mu X^{-1} - Z - (T^T \odot T^T)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)\Delta X,$$

which means equation (21). Then we have

$$\begin{aligned} \mathcal{A}^*(x)\Delta Z &= \mu\mathcal{A}^*(x)X^{-1} - \mathcal{A}^*(x)Z - \mathcal{A}^*(x)(T^T \odot T^T)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)\Delta X \\ &= \mu\mathcal{A}^*(x)X^{-1} - \mathcal{A}^*(x)Z \\ &\quad - \sum_{j=1}^n \mathcal{A}^*(x)(T^T \odot T^T)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)A_j(x)\Delta x_j \\ (24) \quad &= \mu\mathcal{A}^*(x)X^{-1} - \mathcal{A}^*(x)Z - H\Delta x, \end{aligned}$$

where the elements of the matrix H are defined by the form

$$\begin{aligned} H_{ij} &= \text{tr} \left\{ A_i(x)(T^T \odot T^T)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)A_j(x) \right\} \\ &= \text{tr} \left\{ ((T \odot T)A_i(x))(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)(T \odot T)A_j(x) \right\} \\ &= \text{tr} \left\{ \tilde{A}_i(x)(\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)\tilde{A}_j(x) \right\} \\ &= \left\langle \tilde{A}_i(x), (\tilde{X} \odot I)^{-1}(\tilde{Z} \odot I)\tilde{A}_j(x) \right\rangle \end{aligned}$$

with $\tilde{A}_i(x) = TA_i(x)T^T$. This implies (23). By substituting (24) into (11), the Newton equations reduce to equation (22).

Furthermore, it is well known that the coefficient matrix of the linear system of equations (22) becomes nonsingular if the matrix $G + H$ is positive definite and the matrix $A_0(x)$ is of full rank.

Therefore the proof is complete. \square

We note that if the matrix G is updated by a positive definite quasi-Newton formula (see Remark 2 in Section 4.3) and the matrix H is chosen as a positive definite matrix, then Theorem 2 guarantees that the Newton direction is uniquely determined.

The following theorem shows the positive definiteness of the matrix H . In what follows, we assume that the matrices $A_1(x), \dots, A_n(x)$ are linearly independent, which means that $\sum_{i=1}^n v_i A_i(x) = 0$ implies $v_i = 0, i = 1, \dots, n$.

Theorem 3 *Suppose that \tilde{X} and \tilde{Z} are symmetric positive definite, and that $\tilde{X}\tilde{Z} + \tilde{Z}\tilde{X}$ is symmetric positive semidefinite. Suppose that the matrices $A_i(x), i = 1, \dots, n$ are linearly independent. Then the matrix H is positive definite.*

Furthermore, if $\tilde{X}\tilde{Z} = \tilde{Z}\tilde{X}$ holds, then H becomes a symmetric matrix.

Proof. If \tilde{X} is symmetric positive definite, then the operator $\tilde{X} \odot I$ is invertible (see Appendix 9 of [13]). Let $\tilde{U} = \sum_{i=1}^n u_i \tilde{A}_i(x)$ for any $u (\neq 0) \in \mathbb{R}^n$. Since the linear independence of the matrices $A_i(x)$ for $i = 1, \dots, n$ is equivalent to the linear independence of the matrices $\tilde{A}_i(x)$ for $i = 1, \dots, n$, $u \neq 0$ guarantees that $\tilde{U} \neq 0$. By defining $V = (\tilde{X} \odot I)^{-1} \tilde{U} \neq 0$, the quadratic form of H is written as

$$\begin{aligned} u^T H u &= \sum_{i=1}^n \sum_{j=1}^n u_i \operatorname{tr} \left\{ \tilde{A}_i(x) (\tilde{X} \odot I)^{-1} (\tilde{Z} \odot I) \tilde{A}_j(x) \right\} u_j \\ &= \operatorname{tr} \left\{ \tilde{U} (\tilde{X} \odot I)^{-1} (\tilde{Z} \odot I) \tilde{U} \right\} \\ &= \operatorname{tr} \left\{ ((\tilde{X} \odot I)^{-1} \tilde{U}) (\tilde{Z} \odot I) (\tilde{X} \odot I) (\tilde{X} \odot I)^{-1} \tilde{U} \right\} \\ &= \operatorname{tr} \left\{ V (\tilde{Z} \odot I) (\tilde{X} \odot I) V \right\} \\ &= \frac{1}{2} \left\{ \operatorname{tr} \left\{ V (\tilde{Z} \odot I) (\tilde{X} \odot I) V \right\} + \operatorname{tr} \left\{ V (\tilde{X} \odot I) (\tilde{Z} \odot I) V \right\} \right\}. \end{aligned}$$

It follows from property 6 of Symmetrized Kronecker product in Appendix of [13] and relation (14) that

(25)

$$\begin{aligned} u^T H u &= \frac{1}{4} \left\{ \operatorname{tr} \left\{ V ((\tilde{Z}\tilde{X} \odot I) + (\tilde{Z} \odot \tilde{X})) V \right\} + \operatorname{tr} \left\{ V ((\tilde{X}\tilde{Z} \odot I) + (\tilde{X} \odot \tilde{Z})) V \right\} \right\} \\ &= \frac{1}{4} \operatorname{svec}(V)^T \left(((\tilde{X}\tilde{Z} + \tilde{Z}\tilde{X}) \otimes_S I) + (\tilde{X} \otimes_S \tilde{Z}) + (\tilde{Z} \otimes_S \tilde{X}) \right) \operatorname{svec}(V). \end{aligned}$$

It follows from Property 11 of Symmetrized Kronecker product in Appendix of [13] that if \tilde{X} and \tilde{Z} are symmetric positive definite, then $\tilde{X} \otimes_S \tilde{Z}$ and $\tilde{Z} \otimes_S \tilde{X}$ are symmetric positive definite. It also follows from Property 9 that if $\tilde{X}\tilde{Z} + \tilde{Z}\tilde{X}$ is symmetric positive semidefinite, then $(\tilde{X}\tilde{Z} + \tilde{Z}\tilde{X}) \otimes_S I$ is symmetric positive semidefinite. Thus the matrix H is positive definite.

Next, we assume that $\tilde{X}\tilde{Z} = \tilde{Z}\tilde{X}$ holds. Since the relation $(\tilde{X} \odot I)(\tilde{Z} \odot I) = (\tilde{Z} \odot I)(\tilde{X} \odot I)$ holds, we have

$$(26) \quad (\tilde{X} \odot I)^{-1} (\tilde{Z} \odot I) = (\tilde{Z} \odot I) (\tilde{X} \odot I)^{-1}.$$

For any vectors $u, v \in \mathbb{R}^n$, we define

$$\tilde{U} \equiv \sum_{i=1}^n u_i \tilde{A}_i(x), \quad \tilde{V} \equiv \sum_{i=1}^n v_i \tilde{A}_i(x), \quad \tilde{U}' = (\tilde{X} \odot I)^{-1} \tilde{U} \quad \text{and} \quad \tilde{V}' = (\tilde{X} \odot I)^{-1} \tilde{V}.$$

Then in a similar way to the above, we obtain

$$\begin{aligned} u^T H v &= \text{tr} \left\{ \tilde{U} (\tilde{X} \odot I)^{-1} (\tilde{Z} \odot I) \tilde{V} \right\} \\ &= \text{tr} \left\{ \tilde{U} (\tilde{Z} \odot I) (\tilde{X} \odot I)^{-1} \tilde{V} \right\} \quad (\text{from (26)}) \\ &= \text{tr} \left\{ (\tilde{Z} \odot I) (\tilde{X} \odot I)^{-1} \tilde{V} \tilde{U} \right\} \\ &= \text{tr} \left\{ \tilde{V} (\tilde{X} \odot I)^{-1} (\tilde{Z} \odot I) \tilde{U} \right\} \quad (\text{from (19)}) \\ &= v^T H u. \end{aligned}$$

Letting $u = e_i$ and $v = e_j$ yields $H_{ij} = H_{ji}$, which implies that the matrix H is symmetric.

Therefore the theorem is proved. \square

We note that Theorems 2 and 3 correspond to Theorem 3.1 in [13].

The following theorem claims that a BKKT point is obtained if the Newton direction satisfies $\Delta x = 0$.

Theorem 4 *Assume that Δw solves (11) - (13). If $\Delta x = 0$, then $(x, y + \Delta y, Z + \Delta Z)$ is a BKKT point.*

Proof. It follows from the Newton equations that

$$\begin{aligned} \nabla f(x) - A_0(x)^T (y + \Delta y) - \mathcal{A}^*(x)(Z + \Delta Z) &= 0, \\ g(x) &= 0. \end{aligned}$$

Since equation (21) implies

$$Z + \Delta Z = \mu X^{-1},$$

we have

$$X \circ (Z + \Delta Z) = \mu I \quad \text{and} \quad Z + \Delta Z \succ 0.$$

Therefore the point $(x, y + \Delta y, Z + \Delta Z)$ satisfies the BKKT conditions. \square

In the subsequent discussions, we assume that the nonsingular matrix T is chosen so that \tilde{X} and \tilde{Z} commute, i.e., $\tilde{X}\tilde{Z} = \tilde{Z}\tilde{X}$. In this case, the matrices \tilde{X} and \tilde{Z} share the same eigensystem. To end this section, we give the two concrete choices of the transformation T that satisfy such a condition (see [13]).

(i) If we set $T = X^{-1/2}$, then we have $\tilde{X} = I$ and $\tilde{Z} = X^{1/2} Z X^{1/2}$. In this case, the matrices H and ΔZ can be represented by the form:

$$\begin{aligned} H_{ij} &= \text{tr} (A_i(x) X^{-1} A_j(x) Z), \\ \Delta Z &= \mu X^{-1} - Z - \frac{1}{2} (X^{-1} \Delta X Z + Z \Delta X X^{-1}). \end{aligned}$$

(ii) If we set $T = W^{-1/2}$ with $W = X^{1/2}(X^{1/2}ZX^{1/2})^{-1/2}X^{1/2}$, then we have $\tilde{X} = W^{-1/2}XW^{-1/2} = W^{1/2}ZW^{1/2} = \tilde{Z}$. Note that this choice is proposed by Nesterov and Todd. In this case, the matrices H and ΔZ can be represented by the form:

$$\begin{aligned} H_{ij} &= \text{tr} \{A_i(x)W^{-1}A_j(x)W^{-1}\}, \\ \Delta Z &= \mu X^{-1} - Z - W^{-1}\Delta XW^{-1}. \end{aligned}$$

4.2 Primal-dual merit function

In what follows, we assume that the matrix T is so chosen that $\tilde{X}\tilde{Z} = \tilde{Z}\tilde{X}$ is satisfied. To force the global convergence of the algorithm described in Section 4, we use a merit function in the primal-dual space. For this purpose, we propose the following merit function:

$$(27) \quad F(x, Z) = F_{BP}(x) + \nu F_{PD}(x, Z),$$

where $F_{BP}(x)$ and $F_{PD}(x, Z)$ are the primal barrier penalty function and the primal-dual barrier function, respectively, and they are given by

$$(28) \quad F_{BP}(x) = f(x) - \mu \log(\det X) + \rho \|g(x)\|_1,$$

$$(29) \quad F_{PD}(x, Z) = \langle X, Z \rangle - \mu \log(\det X \det Z),$$

where ν and ρ are positive parameters. It follows from the fact $\tilde{X}\tilde{Z} = TXZT^{-1}$ that $\langle \tilde{X}, \tilde{Z} \rangle = \langle X, Z \rangle$ and $F_{PD}(\tilde{x}, \tilde{Z}) = F_{PD}(x, Z)$ hold.

The following lemma gives a lower bound on the value of the primal-dual barrier function (29) and the asymptotic behavior of the function.

Lemma 1 *The primal-dual barrier function satisfies*

$$(30) \quad F_{PD}(x, Z) \geq p\mu(1 - \log \mu)$$

for any $X \succ 0$ and $Z \succ 0$. The equality holds in (30) if and only if $XZ = \mu I$ is satisfied. Furthermore, the following hold

$$(31) \quad \lim_{\langle X, Z \rangle \downarrow 0} F_{PD}(x, Z) = \infty \quad \text{and} \quad \lim_{\langle X, Z \rangle \uparrow \infty} F_{PD}(x, Z) = \infty.$$

Proof. Let λ_i and τ_i for $i = 1, \dots, p$ denote the eigenvalues of the matrices \tilde{X} and \tilde{Z} , respectively. We note that the matrices \tilde{X} and \tilde{Z} share the same eigensystem. Then the matrix $\tilde{X}\tilde{Z}$ has eigenvalues $\lambda_i\tau_i$, $i = 1, \dots, p$, and we have

$$\begin{aligned} (32) \quad F_{PD}(x, Z) &= \langle \tilde{X}, \tilde{Z} \rangle - \mu \log(\det \tilde{X} \det \tilde{Z}) \\ &= \sum_{i=1}^p \lambda_i \tau_i - \mu \log \left(\prod_{i=1}^p \lambda_i \tau_i \right) \\ &= \sum_{i=1}^p (\lambda_i \tau_i - \mu \log \lambda_i \tau_i). \end{aligned}$$

It is easily shown that the function $\phi(\xi) = \xi - \mu \log \xi$ ($\xi > 0$) is convex and achieves a minimum value at $\xi = \mu$. Thus we obtain

$$(33) \quad \begin{aligned} F_{PD}(x, Z) &\geq \sum_{i=1}^p (\mu - \mu \log \mu) \\ &= p(\mu - \mu \log \mu). \end{aligned}$$

It is clear that the equality holds in inequality (33) if and only if $\lambda_i \tau_i = \mu$, $i = 1, \dots, p$ are satisfied. Since \tilde{X} and \tilde{Z} commute, they can be represented by the forms $\tilde{X} = PD_X P^T$ and $\tilde{Z} = PD_Z P^T$ for an orthogonal matrix P , where D_X and D_Z are diagonal matrices whose diagonal elements are λ_i and τ_i , $i = 1, \dots, p$, respectively. Thus, by noting the relations $\tilde{X}\tilde{Z} = PD_X D_Z P^T$, we can show that $\tilde{X}\tilde{Z} = \mu I$ is equivalent to the equations $\lambda_i \tau_i = \mu$, $i = 1, \dots, p$. Furthermore, $\tilde{X}\tilde{Z} = \mu I$ is equivalent to $XZ = \mu I$. Therefore, the first part of this lemma is proved.

It follows from the algebraic and geometric mean $\frac{1}{p} \sum_{i=1}^p \lambda_i \tau_i \geq \left(\prod_{i=1}^p \lambda_i \tau_i \right)^{1/p}$ that

$$\begin{aligned} -\log \left(\prod_{i=1}^p \lambda_i \tau_i \right) &\geq -p \log \left(\sum_{i=1}^p \lambda_i \tau_i \right) + p \log p \\ &= -p \log \langle X, Z \rangle + p \log p. \end{aligned}$$

We use the inequality above and equation (32) to obtain

$$F_{PD}(x, Z) \geq \langle X, Z \rangle - \mu p \log \langle X, Z \rangle + \mu p \log p.$$

Therefore, the expressions (31) hold. This completes the proof. \square

Now we introduce the first order approximation F_l of the merit function by

$$F_l(x, Z; \Delta x, \Delta Z) = F(x, Z) + \Delta F_l(x, Z; \Delta x, \Delta Z),$$

which is used in the line search procedure. Here $\Delta F_l(x, Z; \Delta x, \Delta Z)$ corresponds to the directional derivative and it is defined by the form

$$\Delta F_l(x, Z; \Delta x, \Delta Z) = \Delta F_{BPl}(x; \Delta x) + \nu \Delta F_{PDl}(x, Z; \Delta x, \Delta Z),$$

where

$$(34) \quad \begin{aligned} \Delta F_{BPl}(x; \Delta x) &= \nabla f(x)^T \Delta x - \mu \text{tr}(X^{-1} \Delta X) \\ &\quad + \rho (\|g(x) + A_0(x) \Delta x\|_1 - \|g(x)\|_1), \\ \Delta F_{PDl}(x, Z; \Delta x, \Delta Z) &= \text{tr}(\Delta X Z + X \Delta Z - \mu X^{-1} \Delta X - \mu Z^{-1} \Delta Z). \end{aligned}$$

We show that the search direction is a descent direction for both the primal barrier penalty function (28) and the primal-dual barrier function (29). We first give an estimate of $\Delta F_{BPl}(x; \Delta x)$ for the primal barrier penalty function.

Lemma 2 Assume that Δw solves (11) – (13). Then the following holds

$$\Delta F_{BPI}(x; \Delta x) \leq -\Delta x^T (G + H) \Delta x - (\rho - \|y + \Delta y\|_\infty) \|g(x)\|_1.$$

Proof. It is clear from (12) and (34) that

$$(35) \quad \Delta F_{BPI}(x; \Delta x) = \nabla f(x)^T \Delta x - \mu \text{tr}(X^{-1} \Delta X) - \rho \|g(x)\|_1.$$

It follows from (11) that

$$\nabla f(x)^T \Delta x = -\Delta x^T G \Delta x + \Delta x^T A_0(x)^T (y + \Delta y) + \Delta x^T \mathcal{A}^*(x) (Z + \Delta Z).$$

Since $\mathcal{A}^*(x) (Z + \Delta Z) = \mu \mathcal{A}^*(x) X^{-1} - H \Delta x$ holds by (24), the preceding expression implies that

$$\nabla f(x)^T \Delta x = -\Delta x^T (G + H) \Delta x - g(x)^T (y + \Delta y) + \mu \Delta x^T \mathcal{A}^*(x) X^{-1}.$$

By using the relations

$$\Delta x^T \mathcal{A}^*(x) X^{-1} = \sum_{i=1}^n \Delta x_i \text{tr}(A_i(x) X^{-1}) = \text{tr}\left(\left(\sum_{i=1}^n \Delta x_i A_i(x)\right) X^{-1}\right) = \text{tr}(X^{-1} \Delta X),$$

equation (35) yields

$$\begin{aligned} \Delta F_{BPI}(x; \Delta x) &= -\Delta x^T (G + H) \Delta x - g(x)^T (y + \Delta y) \\ &\quad + \mu \text{tr}(X^{-1} \Delta X) - \mu \text{tr}(X^{-1} \Delta X) - \rho \|g(x)\|_1 \\ &\leq -\Delta x^T (G + H) \Delta x - (\rho - \|y + \Delta y\|_\infty) \|g(x)\|_1. \end{aligned}$$

The proof is complete. \square

Next we estimate the difference $\Delta F_{PDI}(x, Z; \Delta x, \Delta Z)$ for the primal-dual barrier function (29).

Lemma 3 Assume that Δw solves (11) – (13). Then the following holds

$$(36) \quad \Delta F_{PDI}(x, Z; \Delta x, \Delta Z) \leq 0.$$

The equality holds in (36) if and only if the matrices X and Z satisfy the relation $XZ = \mu I$.

Proof. It follows from the Newton equation (13) that

$$\begin{aligned} \Delta F_{PDI}(x, Z; \Delta x, \Delta Z) &= \text{tr} \left\{ (I - \mu \tilde{X}^{-1} \tilde{Z}^{-1}) (\tilde{Z} \Delta \tilde{X} + \tilde{X} \Delta \tilde{Z}) \right\} \\ &= \frac{1}{2} \text{tr} \left\{ (I - \mu \tilde{X}^{-1} \tilde{Z}^{-1}) (\tilde{Z} \Delta \tilde{X} + \tilde{X} \Delta \tilde{Z} + \Delta \tilde{X} \tilde{Z} + \Delta \tilde{Z} \tilde{X}) \right\} \\ &= \text{tr} \left\{ (I - \mu \tilde{X}^{-1} \tilde{Z}^{-1}) (\mu I - \tilde{X} \tilde{Z}) \right\} \\ &= -\text{tr} \left\{ \tilde{X}^{-1} \tilde{Z}^{-1} (\mu I - \tilde{X} \tilde{Z})^2 \right\} \\ &= -\text{tr} \left\{ (\tilde{X} \tilde{Z})^{-1/2} (\mu I - \tilde{X} \tilde{Z})^2 (\tilde{X} \tilde{Z})^{-1/2} \right\}. \end{aligned}$$

Since the matrix $(\tilde{X}\tilde{Z})^{-1/2}(\mu I - \tilde{X}\tilde{Z})^2(\tilde{X}\tilde{Z})^{-1/2}$ is symmetric positive semidefinite, we have

$$\Delta F_{PDI}(x, Z; \Delta x, \Delta Z) \leq 0.$$

It is clear that the equality holds in the above if and only if the matrix $\mu I - \tilde{X}\tilde{Z}$ becomes the zero matrix. Therefore the proof is complete. \square

Now we obtain the following theorem by using the two lemmas given above. This theorem shows that the Newton direction Δw becomes a descent search direction for the proposed primal-dual merit function (27).

Theorem 5 *Assume that Δw solves (11) – (13) and that the matrix $G + H$ is positive definite. Suppose that the penalty parameter ρ satisfies $\rho > \|y + \Delta y\|_\infty$. Then the following hold:*

- (i) *The direction Δw becomes a descent search direction for the primal-dual merit function $F(x, Z)$, i.e. $\Delta F_l(x, Z; \Delta x, \Delta Z) \leq 0$.*
- (ii) *If $\Delta x \neq 0$, then $\Delta F_l(x, Z; \Delta x, \Delta Z) < 0$.*
- (iii) *$\Delta F_l(x, Z; \Delta x, \Delta Z) = 0$ holds if and only if $(x, y + \Delta y, Z)$ is a BKKT point.*

Proof. (i) and (ii) : It follows directly from Lemmas 2 and 3 that

$$(37) \quad \begin{aligned} \Delta F_l(x, Z; \Delta x, \Delta Z) &\leq -\Delta x^T(G + H)\Delta x \\ &\quad -(\rho - \|y + \Delta y\|_\infty)\|g(x)\|_1 \\ &\leq 0. \end{aligned}$$

The last inequality becomes a strict inequality if $\Delta x \neq 0$. Therefore the results hold.

(iii) If $\Delta F_l(x, Z; \Delta x, \Delta Z) = 0$ holds, then $\Delta F_{BPI}(x; \Delta x) = 0$ and $\Delta F_{PDI}(x, Z; \Delta x, \Delta Z) = 0$ are satisfied, and equation (37) yields

$$\Delta x = 0 \quad \text{and} \quad g(x) = 0.$$

It follows from Lemma 3 that $\Delta F_{PDI}(x, Z; \Delta x, \Delta Z) = 0$ implies $X \circ Z = \mu I$, i.e. $XZ = \mu I$. Thus equation (21) yields $\Delta Z = 0$. Then equation (11) implies that $\nabla f(x) - A_0(x)^T(y + \Delta y) - \mathcal{A}^*(x)Z = 0$. Hence $(x, y + \Delta y, Z)$ is a BKKT point.

Conversely, suppose that $(x, y + \Delta y, Z)$ is a BKKT point. Equations (11) and (24) imply that

$$G\Delta x - \mathcal{A}^*(x)\Delta Z = 0 \quad \text{and} \quad \mathcal{A}^*(x)\Delta Z = -H\Delta x.$$

It follows that $(G + H)\Delta x = 0$ holds, which yields $\Delta x = 0$. Using equation (35) and Lemma 3, we have

$$\Delta F_{BPI}(x; \Delta x) = 0 \quad \text{and} \quad \Delta F_{PDI}(x, Z; \Delta x, \Delta Z) = 0,$$

which implies $\Delta F_l(x, Z; \Delta x, \Delta Z) = 0$. Therefore, the theorem is proved. \square

4.3 Algorithm SDPLS that uses the line search procedure

To obtain a globally convergent algorithm to a BKKT point for a fixed $\mu > 0$, we modify the basic Newton iteration. Our iterations take the form

$$x_{k+1} = x_k + \alpha_k \Delta x_k, \quad Z_{k+1} = Z_k + \alpha_k \Delta Z_k \quad \text{and} \quad y_{k+1} = y_k + \Delta y_k$$

where α_k is a step size determined by the line search procedure described below. Throughout this section, the index k denotes the inner iteration count for a given $\mu > 0$. We note that $X_k \succ 0$ and $Z_k \succ 0$ are maintained for all k in the following. We also denote $X(x_k)$ by X_k for simplicity.

The main iteration is to decrease the value of the merit function (27) for fixed μ . Thus the step size is determined by the sufficient decrease rule of the merit function. Specifically, we adopt Armijo's rule. At the current point (x_k, Z_k) , we calculate the initial step size by

$$(38) \quad \bar{\alpha}_{xk} = \begin{cases} -\frac{\gamma}{\lambda_{\min}(X_k^{-1} \Delta X_k)} & \text{if } X(x) \text{ is linear} \\ 1 & \text{otherwise} \end{cases}$$

and

$$(39) \quad \bar{\alpha}_{zk} = -\frac{\gamma}{\lambda_{\min}(Z_k^{-1} \Delta Z_k)},$$

where $\lambda_{\min}(M)$ denotes the minimum eigenvalue of the matrix M , and $\gamma \in (0, 1)$ is a fixed constant. If the minimum eigenvalue in either expression is positive, we ignore the corresponding term.

A step to the next iterate is given by

$$\alpha_k = \bar{\alpha}_k \beta^{l_k}, \quad \bar{\alpha}_k = \min \{ \bar{\alpha}_{xk}, \bar{\alpha}_{zk}, 1 \},$$

where $\beta \in (0, 1)$ is a fixed constant, and l_k is the smallest nonnegative integer such that the sufficient decrease condition

$$(40) \quad F(x_k + \bar{\alpha}_k \beta^{l_k} \Delta x_k, Z_k + \bar{\alpha}_k \beta^{l_k} \Delta Z_k) \leq F(x_k, Z_k) + \varepsilon_0 \bar{\alpha}_k \beta^{l_k} \Delta F_l(x_k, Z_k; \Delta x_k, \Delta Z_k)$$

and the positive definiteness condition

$$(41) \quad X(x_k + \bar{\alpha}_k \beta^{l_k} \Delta x_k) \succ 0,$$

where $\varepsilon_0 \in (0, 1)$. Lemma 4 (ii) given below guarantees that an integer l_k exists.

Now we give a line search algorithm called Algorithm SDPLS. This algorithm should be regarded as the inner iteration of Algorithm SDPIP (see Step 1 of Algorithm SDPIP). We also note that ε' given below corresponds to $M_c \mu$ in Algorithm SDPIP.

Algorithm SDPLS

Step 0. (Initialize) Let $w_0 \in \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{S}^p$ ($X_0 \succ 0, Z_0 \succ 0$), and $\mu > 0, \rho > 0, \nu > 0$. Set $\varepsilon' > 0, \gamma \in (0, 1), \beta \in (0, 1)$ and $\varepsilon_0 \in (0, 1)$. Let $k = 0$.

Step 1. (Termination) If $\|r(w_k, \mu)\|_* \leq \varepsilon'$, then stop.

Step 2. (Compute direction) Calculate the matrix G_k and the transformation T_k . Determine the direction Δw_k by solving (11) – (13).

Step 3. (Step size) Find the smallest nonnegative integer l_k that satisfies the criteria (40) and (41), and calculate

$$\alpha_k = \bar{\alpha}_k \beta^{l_k}.$$

Step 4. (Update variables) Set

$$x_{k+1} = x_k + \alpha_k \Delta x_k, \quad Z_{k+1} = Z_k + \alpha_k \Delta Z_k \quad \text{and} \quad y_{k+1} = y_k + \Delta y_k.$$

Step 5. Set $k := k + 1$ and go to Step 1. □

Remark 1. Theorem 2 can be used to calculate the direction Δw_k in Step 2. Specifically, we compute the directions $(\Delta x_k, \Delta y_k)$ by solving linear system of equations (22), and we obtain ΔZ_k from equation (21). It follows from Theorem 4 that if $\Delta x_k = 0$ is obtained, then we can get the BKKT point $(x_k, y_k + \Delta y_k, Z_k + \Delta Z_k)$ and stop the procedure of the algorithm.

Remark 2. When the matrix G_k approximates the Hessian matrix $\nabla_x^2 L(w_k)$ of the Lagrangian function by using the quasi-Newton updating formula in Step 2, we have the following secant condition

$$G_{k+1} s_k = q_k,$$

where $s_k = x_{k+1} - x_k$ and

$$\begin{aligned} q_k &= \nabla_x L(x_{k+1}, y_{k+1}, Z_{k+1}) - \nabla_x L(x_k, y_{k+1}, Z_{k+1}) \\ &= (\nabla f(x_{k+1}) - A_0(x_{k+1})^T y_{k+1} - \mathcal{A}^*(x_{k+1}) Z_{k+1}) - (\nabla f(x_k) - A_0(x_k)^T y_{k+1} - \mathcal{A}^*(x_k) Z_{k+1}) \\ &= \nabla f(x_{k+1}) - \nabla f(x_k) - (A_0(x_{k+1}) - A_0(x_k))^T y_{k+1} - (\mathcal{A}^*(x_{k+1}) - \mathcal{A}^*(x_k)) Z_{k+1}. \end{aligned}$$

We note that it is easy to calculate the vector q_k . In order to preserve the positive definiteness of the matrix G_k , we can use the modified BFGS update proposed by Powell, which is given by the form

$$G_{k+1} = G_k - \frac{G_k s_k s_k^T G_k}{s_k^T G_k s_k} + \frac{\hat{q}_k \hat{q}_k^T}{s_k^T \hat{q}_k},$$

where

$$\begin{aligned} \hat{q}_k &= \psi_k q_k + (1 - \psi_k) G_k s_k, \\ \psi_k &= \begin{cases} 1 & \text{if } s_k^T q_k \geq 0.2 s_k^T G_k s_k \\ \frac{0.8 s_k^T G_k s_k}{s_k^T (G_k s_k - q_k)} & \text{otherwise.} \end{cases} \end{aligned}$$

Remark 3. If we want to use the Hessian matrix $\nabla_x^2 L(w_k)$ for the matrix G_k , we adopt Levenberg-Marquardt type modification of $\nabla_x^2 L(w_k)$ to obtain a positive semi-definite G_k for global convergence property shown in the next section. Namely, we compute a parameter $\beta \geq 0$ which gives a positive semidefinite $\nabla_x^2 L(w_k) + \beta I$. The procedure used in the numerical experiments below is as follows:

Step 0. Calculate the Cholesky decomposition of $\nabla_x^2 L(w_k)$. If it is successful, set $\beta = 0$, and stop. If not, set $\beta = 1.0$, and go to Step 1.

Step 1. Calculate the Cholesky decomposition of $\nabla_x^2 L(w_k) + \beta I$. If it is successful, go to Step 2. Otherwise go to Step 3.

Step 2. Repeat $\beta := \beta/2$ until the Cholesky decomposition fails. Set $\beta := 2\beta$, and stop.

Step 3. Repeat $\beta := 2\beta$ until the Cholesky decomposition succeeds. Stop. \square

This method is used to solve large scale nonconvex problems in the following.

Remark 4. If the matrix $X(x)$ is linear with respect to x , the step size rules (38) and (39) are usual ones used for the linear SDP problems. From the viewpoint of the numerical accuracy and computational cost, we calculate the minimum eigenvalue of the symmetric matrix $L^{-1}\Delta ZL^{-T}$ in computing the minimum eigenvalue in (39) based on the fact that the spectrum of the nonsymmetric matrix $Z^{-1}\Delta Z$ is same as that of the symmetric matrix $L^{-1}\Delta ZL^{-T}$, where $Z = LL^T$ is the Cholesky factorization of Z . We can also calculate the minimum eigenvalue of the matrix $X^{-1}\Delta X$ in equation (38) in the same way if $X(x)$ is linear.

5 Global convergence to a barrier KKT point

In this section, we prove global convergence of Algorithm SDPLS. For this purpose, we make the following assumptions.

Assumptions

- (A1) The functions $f, g_i, i = 1, \dots, m$, and X are twice continuously differentiable.
- (A2) The sequence $\{x_k\}$ generated by Algorithm SDPLS remains in a compact set Ω of \mathbb{R}^n .
- (A3) For all k on Ω , the matrix $A_0(x_k)$ is of full rank and the matrices $A_1(x_k), \dots, A_n(x_k)$ are linearly independent.
- (A4) The matrix G_k is uniformly bounded and positive semidefinite.
- (A5) The transformation T_k is chosen such that \tilde{X}_k and \tilde{Z}_k commute, and both of the sequences $\{T_k\}$ and $\{T_k^{-1}\}$ are bounded.
- (A6) The penalty parameter ρ is sufficiently large so that $\rho > \|y_k + \Delta y_k\|_\infty$ holds for all k .

\square

Assumption (A2) assures the existence of an accumulation point of the generated sequence $\{x_k\}$. The boundedness of the generated sequence $\{x_k\}$ is derived if there exist upper and lower bounds on the variable x , which is a reasonable assumption in practice. We should note that if a quasi-Newton approximation is used for computing the matrix G_k , then we only need the continuity of the first order derivatives of functions in assumption (A1).

In order to show the global convergence property, we first present the following lemma that gives a base for Armijo's line search rule. The merit function is differentiable except for the part $\|g(x)\|_1$, so we can prove this lemma in the same way as Lemmas 2 and 3 in [17].

Lemma 4 *Let $d_x \in \mathbb{R}^n$ and $D_z \in \mathbb{R}^{p \times p}$ be given. Define $F'(x, Z; d_x, D_z)$ by*

$$F'(x, Z; d_x, D_z) = \lim_{t \downarrow 0} \frac{F(x + td_x, Z + tD_z) - F(x, Z)}{t}.$$

Then the following hold:

(i) *There exists a $\theta \in (0, 1)$ such that*

$$F(x + d_x, Z + D_z) \leq F(x, Z) + F'(x + \theta d_x, Z + \theta D_z; d_x, D_z),$$

whenever $X(x + d_x) \succ 0$ and $Z + D_z \succ 0$.

(ii) *Let $\varepsilon_0 \in (0, 1)$ be given. If $\Delta F_l(x, Z; d_x, D_z) < 0$, then*

$$F(x + \alpha d_x, Z + \alpha D_z) - F(x, Z) \leq \varepsilon_0 \alpha \Delta F_l(x, Z; d_x, D_z),$$

for sufficiently small $\alpha > 0$. □

The following lemma shows the boundedness of the sequence $\{w_k\}$ and the uniformly positive definiteness of the matrix H_k .

Lemma 5 *Suppose that assumptions (A1), (A2) and (A6) are satisfied. Let the sequence $\{w_k\}$ be generated by Algorithm SDPLS. Then the following hold.*

(i) *$\liminf_{k \rightarrow \infty} \det(X_k) > 0$ and $\liminf_{k \rightarrow \infty} \det(Z_k) > 0$.*

(ii) *The sequence $\{w_k\}$ is bounded.*

In addition, if assumptions (A3), (A4) and (A5) are satisfied, the following hold.

(iii) *There exists a positive constant M such that*

$$\frac{1}{M} \|v\|^2 \leq v^T (G_k + H_k) v \leq M \|v\|^2 \quad \text{for any } v \in \mathbb{R}^n$$

for all $k \geq 0$.

(iv) *The sequence $\{\Delta w_k\}$ is bounded.*

Proof. (i) Since the sequence $\{F_{PD}(x_k, Z_k)\}$ is bounded below from Lemma 1, the sequence $\{F_{BP}(x_k)\}$ is bounded above, because the function value of $F(x_k, Z_k)$ decreases monotonically. Therefore it follows from the log barrier term in $F_{BP}(x)$ that $\det X_k$ is bounded away from zero, and we have $\liminf_{k \rightarrow \infty} \det X_k > 0$. This implies that $\liminf_{k \rightarrow \infty} \det Z_k > 0$ also holds, because $\{F_{PD}(x_k, Z_k)\}$ is bounded above and below and $\langle X_k, Z_k \rangle \geq 0$ is satisfied.

(ii) The boundedness of the sequences $\{Z_k\}$ and $\{y_k\}$ follows from assumptions (A2), (A6) and the monotone decreasing of $F(x_k, Z_k)$. Therefore the sequence $\{w_k\}$ is bounded.

(iii) From Appendix 9 of [13] that the operator $\tilde{X} \odot I$ is invertible. For the vector V defined in the proof of Theorem 3, $\text{svec}(V)$ can be represented by the form

$$\begin{aligned} \text{svec}(V) &= \text{svec} \left(\text{smat}((\tilde{X} \otimes_S I)^{-1} \tilde{U}) \right) \\ &= (\tilde{X} \otimes_S I)^{-1} \sum_{i=1}^n u_i \text{svec}(\tilde{A}_i(x)), \end{aligned}$$

where $\tilde{U} \equiv \sum_{i=1}^n u_i \tilde{A}_i(x) \neq 0$. Letting

$$\tilde{A}(x) = \left(\text{svec}(\tilde{A}_1(x)), \dots, \text{svec}(\tilde{A}_n(x)) \right) \in \mathbb{R}^{p(p+1)/2 \times n}$$

and

$$u = (u_1, \dots, u_n)^T,$$

we have

$$\text{svec}(V) = (\tilde{X} \otimes_S I)^{-1} \tilde{A}(x)u.$$

Therefore it follows from (25) that

$$u^T H_k u = u^T \tilde{A}(x_k)^T ((\tilde{X}_k \otimes_S I)^{-1})^T \hat{H}_k (\tilde{X}_k \otimes_S I)^{-1} \tilde{A}(x_k) u,$$

where

$$\hat{H}_k = ((\tilde{X}_k \tilde{Z}_k + \tilde{Z}_k \tilde{X}_k) \otimes_S I) + (\tilde{X}_k \otimes_S \tilde{Z}_k) + (\tilde{Z}_k \otimes_S \tilde{X}_k).$$

The boundedness of the sequence $\{w_k\}$ guarantees the uniformly positive definiteness and boundedness of the matrix $((\tilde{X}_k \otimes_S I)^{-1})^T \hat{H}_k (\tilde{X}_k \otimes_S I)^{-1}$. Since the linear independence of the matrices $A_i(x_k)$ for $i = 1, \dots, n$ is equivalent to the linear independence of the vectors $\text{svec}(\tilde{A}_i(x_k))$ for $i = 1, \dots, n$, the matrix $\tilde{A}(x_k)$ is of column full rank. This implies that there exist positive constants λ_{min} and λ_{max} , which are independent of k , such that

$$\lambda_{min} \|u\|^2 \leq u^T H_k u \leq \lambda_{max} \|u\|^2$$

holds. Thus by assumption (A4), we obtain the result.

(iv) Since, by results (ii) and (iii) shown above, the sequence $\{w_k\}$ is bounded and $\{G_k + H_k\}$ is uniformly bounded and positive definite, Theorem 2 guarantees the desired result. \square

By Theorem 4, $\Delta x_k = 0$ guarantees that $(x_k, y_k + \Delta y_k, Z_k + \Delta Z_k)$ is a BKKT point. Thus in what follows, we assume that $\Delta x_k \neq 0$ for any $k \geq 0$. The following theorem gives the global convergence of an infinite sequence generated by Algorithm SDPLS.

Theorem 6 *Suppose that assumptions (A1) – (A6) hold. Let an infinite sequence $\{w_k\}$ be generated by Algorithm SDPLS. Then there exists at least one accumulation point of $\{w_k\}$, and any accumulation point of the sequence $\{w_k\}$ is an BKKT point.*

Proof. In the proof, we define the following notations

$$u_k = \begin{pmatrix} x_k \\ Z_k \end{pmatrix} \quad \text{and} \quad \Delta u_k = \begin{pmatrix} \Delta x_k \\ \Delta Z_k \end{pmatrix}$$

for simplicity. By Lemma 5 (ii), the sequence $\{w_k\}$ has at least one accumulation point. The boundedness of the sequence $\{w_k\}$ implies that all eigenvalues of X_k and Z_k are bounded above. It follows from Lemma 5 (i) that each smallest eigenvalue of X_k and Z_k is bounded away from zero. By Lemma 5 (iv), $\|\Delta w_k\|_*$ is uniformly bounded above. Hence, we have $\liminf_{k \rightarrow \infty} \bar{\alpha}_k > 0$. Furthermore, the sequence $\{l_k\}$ that satisfies $X(x_k + \bar{\alpha}_k \beta^{l_k} \Delta x_k) \succ 0$ is uniformly bounded above.

It follows from Lemma 5 (iii) that there exists a positive constant M such that

$$\frac{1}{M} \|v\|^2 \leq v^T (G_k + H_k) v \leq M \|v\|^2$$

for any $v \in \mathbb{R}^n$ and all $k \geq 0$. Thus by (37), we have

$$\Delta F_l(u_k; \Delta u_k) \leq -\frac{\|\Delta x_k\|^2}{M} < 0,$$

and inequality (40) yields

$$\begin{aligned} (42) \quad F(u_{k+1}) - F(u_k) &\leq \varepsilon_0 \bar{\alpha}_k \beta^{l_k} \Delta F_l(u_k; \Delta u_k) \\ &\leq -\varepsilon_0 \bar{\alpha}_k \beta^{l_k} \frac{\|\Delta x_k\|^2}{M} \\ &< 0. \end{aligned}$$

Because the sequence $\{F(u_k)\}$ is monotonically decreasing and bounded below, the left-hand side of (42) converges to 0, which implies that

$$\lim_{k \rightarrow \infty} \beta^{l_k} \Delta F_l(u_k; \Delta u_k) = 0.$$

If there exists a finite number N such that $l_k < N$ for all k , then we have $\lim_{k \rightarrow \infty} \Delta F_l(u_k; \Delta u_k) = 0$.

Now we suppose that there exists a subsequence $K \subset \{0, 1, \dots\}$ such that $l_k \rightarrow \infty, k \in K$. Then we can assume $l_k > 0$ for sufficiently large $k \in K$ without loss of generality, which means that the point $u_k + \theta'_k \alpha_k \Delta u_k / \beta$ does not satisfy condition (40) for some $\theta'_k \in (0, 1)$. Thus, we get

$$(43) \quad F(u_k + \theta'_k \alpha_k \Delta u_k / \beta) - F(u_k) > \varepsilon_0 \theta'_k \alpha_k \Delta F_l(u_k; \Delta u_k) / \beta.$$

By Lemma 4, there exists a $\theta_k \in (0, 1)$ such that for $k \in K$,

$$\begin{aligned} (44) \quad F(u_k + \theta'_k \alpha_k \Delta u_k / \beta) - F(u_k) &\leq \theta'_k \alpha_k F'(u_k + \theta_k \theta'_k \alpha_k \Delta u_k / \beta; \Delta u_k) / \beta \\ &\leq \theta'_k \alpha_k \Delta F_l(u_k + \theta_k \theta'_k \alpha_k \Delta u_k / \beta; \Delta u_k) / \beta. \end{aligned}$$

Then, from (43) and (44), we see that

$$\varepsilon_0 \Delta F_l(u_k; \Delta u_k) < \Delta F_l(u_k + \theta_k \theta'_k \alpha_k \Delta u_k / \beta; \Delta u_k).$$

This inequality yields

$$(45) \quad \Delta F_l(u_k + \theta_k \theta'_k \alpha_k \Delta u_k / \beta; \Delta u_k) - \Delta F_l(u_k; \Delta u_k) > (\varepsilon_0 - 1) \Delta F_l(u_k; \Delta u_k) > 0.$$

Thus by the fact $l_k \rightarrow \infty$, $k \in K$, we have $\alpha_k \rightarrow 0$ and thus $\|\theta_k \theta'_k \alpha_k \Delta u_k / \beta\|_* \rightarrow 0$, $k \in K$, because $\|\Delta u_k\|_*$ is uniformly bounded. Here $\|\Delta u_k\|_*$ is defined by $\|\Delta u_k\|_* = \|\Delta x_k\| + \|\Delta Z_k\|_F$. This implies that the left-hand side of (45) and therefore $\Delta F_l(u_k; \Delta u_k)$ converges to zero when $k \rightarrow \infty$, $k \in K$.

By the discussions above, we have proved that

$$(46) \quad \lim_{k \rightarrow \infty} \Delta F_l(u_k; \Delta u_k) = 0.$$

Since equation (46) implies that

$$\Delta F_{BPl}(x_k; \Delta x_k) \rightarrow 0 \quad \text{and} \quad \Delta F_{PDl}(x_k, z_k; \Delta x_k, \Delta z_k) \rightarrow 0,$$

it follows from equations (37), (12) and Lemma 3 that

$$\Delta x_k \rightarrow 0, \quad g(x_k) \rightarrow 0, \quad X_k Z_k \rightarrow \mu I \quad (\tilde{X}_k \tilde{Z}_k \rightarrow \mu I).$$

Therefore, equation (21) yields

$$\Delta Z_k \rightarrow 0.$$

By equation (11), we have

$$\nabla_x L(x_k, y_k + \Delta y_k, Z_k) \rightarrow 0,$$

which implies that

$$r(x_k, y_k + \Delta y_k, Z_k, \mu) \rightarrow 0.$$

Since $x_{k+1} = x_k + \alpha_k \Delta x_k$, $Z_{k+1} = Z_k + \alpha_k \Delta Z_k$, $\Delta x_k \rightarrow 0$, $\Delta Z_k \rightarrow 0$ and $y_{k+1} = y_k + \Delta y_k$, the result follows. Therefore, the theorem is proved. \square

The preceding theorem guarantees that any accumulation point of the sequence $\{(x_k, y_k, Z_k)\}$ satisfies the BKKT conditions. If we adopt a common step size α_k as $w_{k+1} = w_k + \alpha_k \Delta w_k$ in Step 4 of Algorithm SDPLS, where α_k is determined in Step 3, then the result of the theorem is replaced by the statement that any accumulation point of the sequence $\{(x_k, y_k + \Delta y_k, Z_k)\}$ satisfies the BKKT conditions.

6 Numerical Experiments

The algorithm of this paper is implemented and brief numerical experiments are done in order to verify the theoretical results of the proposed algorithm. The program is written in C++, and is run on 3.2GHz Pentium IV PC with LINUX OS.

In the following experiments, initial values of various quantities are set as follows: $\mu_0 = 1.0, X_0 = I, Z_0 = I$. The barrier parameter is updated by the rule $\mu_{k+1} = \mu_k/10.0$ after an approximate barrier KKT point is obtained in Step 1 of Algorithm SDPIP where we set $M_c = 0.1$ and $\gamma = 0.9$, and the transformation is set to be $T = X^{-1/2}$ at each iteration of Algorithm SDPLS.

The first problem is Gaussian channel capacity problem which is described in [15]:

$$\begin{aligned} & \text{minimize} && \frac{1}{2}(\log \det(X + R) - \log \det R), \\ & \text{subject to} && \text{tr}(X) \leq nP, X \succeq 0, \end{aligned}$$

where noise covariance $R \in \mathbb{S}^n$ is known and given, and input covariance $X \in \mathbb{S}^n$ is the variable to be determined. The parameter $P \in \mathbb{R}$ gives a limit on the average total power in the input. If all channels are independent, i.e., all covariances are diagonal, and the noise covariance depends on X as $R_{ii} = r_i + a_i X_{ii}$, $a_i > 0$ (case of near-end cross-talk), the above problem can be written as

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \sum_{i=1}^n \log\left(1 + \frac{X_{ii}}{r_i + a_i X_{ii}}\right), \\ & \text{subject to} && \sum_{i=1}^n X_{ii} \leq nP, X_{ii} \geq 0. \end{aligned}$$

This problem can be transformed to SDP type one:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \sum_{i=1}^n \log(1 + t_i), \\ & \text{subject to} && \sum_{i=1}^n X_{ii} \leq nP, X_{ii} \geq 0, t_i \geq 0, \\ & && \begin{pmatrix} 1 - a_i t_i & \sqrt{r_i} \\ \sqrt{r_i} & a_i X_{ii} + r_i \end{pmatrix} \succeq 0, i = 1, \dots, n. \end{aligned}$$

In our experiment, r_i and a_i are set to uniform random numbers between 0 and 1. P is set to 1. We solved problems with $n = 10, 20, \dots, 10240$ using exact Hessian of the Lagrangian as the matrix G , and the results are shown below.

Table 1. Gaussian channel capacity problem

n	iteration	CPU (sec)
10	28	0.03
20	26	0.17
40	31	0.11
80	39	0.32
160	48	1.07
320	52	3.8
640	40	10.2
1280	44	41.3
2560	38	137
5120	43	607
10240	45	2559

The second problem is minimization of the minimal eigenvalue problem defined as:

$$\begin{aligned} & \text{minimize} && \lambda_{\min}(M(q)), \\ & \text{subject to} && q \in Q, \end{aligned}$$

where $q \in \mathbb{R}^n$, $Q \subset \mathbb{R}^n$, and $M \in \mathbb{S}^m$ is a function of q . We formulate this problem as follows:

$$\begin{aligned} & \text{minimize} && \text{tr}(\Pi M(q)), \\ & && \text{tr}(\Pi) = 1, \\ & \text{subject to} && \Pi \succeq 0, \\ & && q \in Q, \end{aligned}$$

where $\Pi \in \mathbb{S}^m$ is an additional Matrix variable (Leibfritz and Maruhn 2005). In our experiment, we set $q = (x, y)^T$, and $M = xyA + xB + yC$ with given $A, B, C \in \mathbb{S}^m$. The elements of matrices A, B and C are set from uniform random numbers in $[-5, 5]$. The constraint region Q for the variable q is set to $[-1, 1] \times [-1, 1]$. We solved problems with the sizes of M, Π, A, B, C equal to 10, 20, 40, 80 respectively, with BFGS quasi-Newton update for the matrix G .

Table 2. Minimization of the minimal eigenvalue problem

m	iteration	CPU (sec)
10	30	0.12
20	32	0.88
40	69	46.9
80	56	1176

The third problem is a real financial one and taken from [8]. The model is to discriminate failure and non-failure companies by a Logit model using a positive semidefinite quadratic discriminant function. The problem for learning is defined by

$$\begin{aligned} & \text{maximize} && \sum_{i=1}^M (y_i z(x_i) - \log(1 + e^{z(x_i)})), a \in \mathbb{R}, b \in \mathbb{R}^q, Q \in \mathbb{S}^q, \\ & \text{subject to} && Q \succeq 0, \end{aligned}$$

where $z(x) = a + b^T x + \frac{1}{2} x^T Q x$, and $x_i = (x_1, \dots, x_q)_i$ gives financial data of each company $i = 1, \dots, M$. The value of y_i gives failure or non-failure as follows:

$$\begin{aligned} y_i &= 0 \Leftrightarrow (x_1, \dots, x_q)_i \in M_0(\text{non - failure}), \\ y_i &= 1 \Leftrightarrow (x_1, \dots, x_q)_i \in M_1(\text{failure}). \end{aligned}$$

In [8], Konno et.al. proposed a method that used a cutting plane approximation of positive semidefinite condition and solved resulting linearly constrained problems using an interior point NLP algorithm in NUOPT. In Tables 1 and 2, we list two examples. Tables 1 and 2 show the results with both BFGS update (bfgs) and exact Hessian (hesse) for the matrix G . In each table, the algorithms used, the final objective function value, the minimum eigenvalue of the obtained matrix Q , the total inner iteration counts and the run time (sec) are given. The learning experiments were done by Japan Credit Rating Agency, Ltd. with their own financial data including the data provided by Tokyo Shoko Research, Ltd. These tables show that our methods solve the problems efficiently and that our method (hesse) performs better than our method (bfgs).

Table 3.1. Logit model/Example 1: number of variables = 28,
 $q = 6, M = 6084, M_0 = 6053$

algorithm	final objective	final $\lambda_{\min}(Q)$	iteration	time (sec)
cutting plane	-153.0808	-9.59e-05	—	7.77
ours (bfgs)	-153.0828	1.76e-09	117	1.65
ours (hesse)	-153.0828	1.77e-09	27	0.80

Table 3.2. Logit model/Example 2: number of variables = 45,
 $q = 8, M = 6084, M_0 = 6053$

algorithm	final objective	final $\lambda_{\min}(Q)$	iteration	time (sec)
cutting plane	-143.7445	-9.17e-05	—	30.3
ours (bfgs)	-143.7468	3.88e-09	233	4.2
ours (hesse)	-143.7468	4.01e-09	30	1.5

Tables 3.1 and 3.2 show the required iteration counts for each value of μ . It is clear that majority of iterations are required at the first few values of μ .

Table 3.3. Logit model: iteration counts for each μ in Example 1

μ	bfgs	hesse
1.0e0	75	17
1.0e-1	25	2
1.0e-2	14	2
1.0e-3	4	2
1.0e-4	3	1
1.0e-5	3	2
1.0e-6	1	1
1.0e-7	1	1

Table 3.4. Logit model: iteration counts for each μ in Example 2

1.0e0	150	19
1.0e-1	35	3
1.0e-2	23	2
1.0e-3	9	1
1.0e-4	11	2
1.0e-5	3	2
1.0e-6	2	1
1.0e-7	1	1

The forth problem in our experiment is from the nearest correlation matrix problem:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|X - A\|_F, \\ & \text{subject to} && X \succeq \epsilon I, \\ & && X_{ii} = 1, i = 1, \dots, n, \end{aligned}$$

where $A \in \mathbb{S}^n$ is given, and we want to obtain $X \in \mathbb{S}^n$ which is nearest to A , and satisfies the given constraints. In the above problem, eigenvalues of X should not be less than

$\epsilon > 0$, and the diagonals of X is equal to 1. There exists special purpose algorithms for solving this type of problem (e.g., [10]). Therefore we add additional constraints which gives an upper bound on the condition number of the matrix X :

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|X - A\|_F, \\ & && zI \preceq X \preceq yI, \\ & \text{subject to} && y \leq \kappa z, \quad z \geq \epsilon \\ & && X \succeq \epsilon I, \\ & && X_{ii} = 1, i = 1, \dots, n, \end{aligned}$$

where y and z denote the maximal and minimal eigenvalue of X respectively, and the upper bound of their ratio (condition number) $\kappa > 0$ is given. Elements of the matrix A are uniform random numbers in $[-1, 1]$ with $A_{ii} = 1, i = 1, \dots, n$. We set $\epsilon = 10^{-3}, \kappa = 10.0$. Results of various values of n are given below, where the exact Hessian is used for the matrix G .

Table 4. Nearest correlation matrix problem

n	iteration	CPU (sec)
10	22	0.05
20	19	0.80
40	18	24.88
80	19	594.08

The fifth problem area is the so called static output feedback (SOF) problems from *COMPL_eib* library. The following is the SOF- \mathcal{H}_2 type problem:

$$\begin{aligned} & \text{minimize} && \text{tr}(X), \\ & \text{subject to} && Q \succeq 0, \\ & && A(F)Q + QA(F)^T + B_1B_1^T \preceq 0, \\ & && \begin{pmatrix} X & C(F)Q \\ QC(F)^T & Q \end{pmatrix} \succeq 0, \end{aligned}$$

where $X \in \mathbb{S}^{n_z \times n_z}, F \in \mathbb{R}^{n_u \times n_y}$ and $Q \in \mathbb{S}^{n_x \times n_x}$ are variable matrices to be determined. The matrices $A \in \mathbb{R}^{n_x \times n_x}, B \in \mathbb{R}^{n_x \times n_u}, B_1 \in \mathbb{R}^{n_x \times n_w}, C \in \mathbb{R}^{n_y \times n_x}, C_1 \in \mathbb{R}^{n_z \times n_x}, D_{11} \in \mathbb{R}^{n_z \times n_w}, D_{12} \in \mathbb{R}^{n_z \times n_u}$ and $D_{21} \in \mathbb{R}^{n_y \times n_w}$ are give constant matrices, and form the matrices $A(F), B(F), C(F), D(F)$ which appear in the problem definition as follows:

$$\begin{aligned} A(F) &= A + BFC, \\ B(F) &= B_1 + BFD_{21}, \\ C(F) &= C_1 + D_{12}FC, \\ D(F) &= D_{11} + D_{12}FD_{21}. \end{aligned}$$

The initial interior points are not known for this type of problem, and it turns out that it is not easy to find them. So we try various starting points, and solve the problems for which we can find initial interior points. We list the results for these problems below. Iterations are stopped when the norm of KKT conditions is less than 10^{-6} . In [11],

numerical results for these problems performed by PENBMI, a specialized BMI-version of PENNON is reported. We list CPU data of PENBMI multiplied by a factor 2.5/3.2 which is a ratio of CPU speeds used in two experiments. We note that the various conditions of these experiments are not equal, so the PENBMI's CPU data is listed to crudely observe how our algorithm performs compared with PENBMI.

Table 5.1. SOF- \mathcal{H}_2 problem

problem	n	n_x	n_y	n_u	n_w	n_z	iteration	CPU(sec)	CPU(PENBMI)
AC1	27	5	3	3	3	2	38	0.11	0.62
AC2	39	5	3	3	3	5	138	0.64	1.25
AC3	38	5	4	2	5	5	41	0.19	0.56
AC6	64	7	4	2	7	7	68	0.69	2.53
AC17	22	4	2	1	4	4	117	0.26	0.27
HE1	15	4	1	2	2	2	174	0.31	0.17
HE2	24	4	2	2	4	4	33	0.09	0.59
HE3	115	8	6	4	1	10	269	7.94	1.53
REA1	26	4	3	2	4	4	76	0.21	0.74
DIS1	88	8	4	4	1	8	47	0.93	5.04
DIS2	16	3	2	2	3	3	43	0.08	0.18
DIS3	58	6	4	4	6	6	252	2.33	1.93
DIS4	66	6	6	4	6	6	30	0.38	2.91
AGS	160	12	2	2	12	12	43	2.28	130
BDT1	96	11	3	3	1	6	46	1.07*	2.78
MFP	26	4	2	3	4	4	112	0.33	0.46
EB1	59	10	1	1	2	2	55	0.68	16.2
EB2	59	10	1	1	2	2	50	0.61	21.0
PSM	49	7	3	2	2	5	46	0.29	2.01
NN2	7	2	1	1	2	2	27	0.03	0.22
NN4	26	4	3	2	4	4	32	0.09	0.30
NN8	16	3	2	2	3	3	63	0.12	0.27
NN11	157	16	5	3	3	3	188	12.19*	223
NN15	20	3	2	2	1	4	64	0.13	0.27
NN16	62	8	4	4	8	4	124	1.51	36.4

The CPU data with * means norm tolerance is set to 10^{-5} .

We next describe the results for SOF- \mathcal{H}_∞ problem which is defined by the following:

$$\begin{aligned}
& \text{minimize } \gamma, \\
& \text{subject to } Q \succeq 0, \\
& \gamma \geq 0, \\
& \begin{pmatrix} A(F)^T Q + Q A(F) & Q B(F) & C(F)^T \\ B(F)^T Q & -\gamma I & D(F)^T \\ C(F) & D(F) & -\gamma I \end{pmatrix} \succeq 0,
\end{aligned}$$

where $Q \in \mathbb{S}^{n_x \times n_x}$ and $F \in \mathbb{R}^{n_u \times n_y}$ are variable matrices to be determined. As in the SOF- \mathcal{H}_2 type problems, we report the results for problems with feasible initial point obtained.

Table 5.2. SOF- \mathcal{H}_∞ problems

problem	n	n_x	n_y	n_u	n_w	n_z	iteration	CPU(sec)	CPU(PENBMI)
AC4	13	4	2	1	2	2	188	0.34	0.64
HE2	15	4	2	2	4	4	64	0.15	0.13
DIS2	11	3	2	2	3	3	156	0.24	8.00
AGS	83	12	2	2	12	12	116	6.84	3.27
MFP	17	4	2	3	4	4	102	0.27	0.42
EB1	57	10	1	1	2	2	277	4.63	1.43
EB2	57	10	1	1	2	2	74	1.21	1.79
PSM	35	7	3	2	2	5	78	0.39	0.58
NN2	5	2	1	1	2	2	27	0.03	0.06

The last set of problems is obtained from SDPLIB to check our algorithms for large scale problems. SDPLIB is a library for linear SDP problems. We add the quadratic term $x^T Q x$ to the original linear objective function $c^T x$ to form nonlinear objective function $x^T Q x / 2 + c^T x$ where the matrix Q is sparse and symmetric positive definite. The values of the diagonal elements of Q are set to 1, and those of the off-diagonal elements are uniform random numbers from $[0, 1]$, and if generated random number is greater than 0.03, the value is set to 0. Therefore the density of nonzero elements of the matrix Q is approximately 3%.

Table 6. SDPLIB with nonlinear objective

problem	n	p	iteration	CPU(sec)
arch8	174	335	51	14.10
control7	136	45	33	95.38
maxG11	800	800	27	252.44
mcp500-1	500	500	39	84.17
qap10	1021	101	35	65.85
ss30	132	426	47	44.71
theta6	4375	300	68	3695.86
truss8	496	628	31	14.89

From the above set of experiments, we think the proposed method works as described in this paper, and hope the method is similarly efficient as existing primal-dual interior point methods for ordinary nonlinear programming [17].

7 Concluding Remarks

In this paper, we have proposed a primal-dual interior point method for solving nonlinear semidefinite programming problems. Within the line search strategy, we have proposed the primal-dual merit function that consists of the primal barrier penalty function and the primal-dual barrier function, and we have proved the global convergence property of our method. Reported numerical experiments show the practical efficiency of our method.

Analysis of the rate of convergence and more extensive numerical experiments of our method are under further research. In addition, we plan to construct a method within the framework of the trust region globalization strategy.

Acknowledgement. The authors would like to thank Japan Credit Rating Agency, Ltd. for providing us the results of their numerical experiments.

References

- [1] Alizadeh, F., Haeberly, J. A., Overton, M. L.: Primal-dual interior-point methods for semidefinite programming: Convergence rates, stability and numerical results. *SIAM Journal on Optimization* **8**, 746-768 (1998)
- [2] Boyd, S., Vandenberghe, L.: *Convex Optimization*. Cambridge University Press, 2004.
- [3] Correa, R., Ramirez, C. H.: A global algorithm for nonlinear semidefinite programming. *SIAM Journal on Optimization* **15**, 303-318 (2004)
- [4] Fares, B., Noll, D., Apkarian, P.: Robust control via sequential semidefinite programming. *SIAM Journal on Control and Optimization* **40**, 1791-1820 (2002)
- [5] Freund, R. W., Jarre, F. : A sensitivity analysis and a convergence result for a sequential semidefinite programming method. Numerical Analysis Manuscript No. 03-4-08, Bell Laboratories, April 2003.
- [6] Jarre, F.: A QQP-minimization method for semidefinite and smooth nonconvex programs. Technical report, Institute für Angewandte Mathematik und Statistik, Universität Würzburg, August 1998.
- [7] Kanzow, C., Nagel, C., Kato, H., Fukushima, M.: Successive linearization methods for nonlinear semidefinite programs. *Computational Optimization and Applications* **31**, 251-273 (2005)
- [8] Konno, H., Kawadaï, N., Wu, D.: Estimation of failure probability using semi-definite Logit model. *Computational Management Science* **1**, 59-73 (2004)
- [9] Kočvara, M., Stingl, M.: PENNON : A code for convex nonlinear and semidefinite programming. *Optimization Methods and Software* **18**, 317-333 (2003)
- [10] Qi, H., Sun, D.: A quadratically convergent newton method for computing the nearest correlation matrix. *SIAM journal on matrix analysis and applications* **28**, 360-385 (2007)
- [11] Stingl, M.: On the Solution of Nonlinear Semidefinite Programs by Augmented Lagrangian Methods. Dissertation, Shaker Verlag, Aachen, (2006)
- [12] Todd, M. J.: Semidefinite optimization. *Acta Numerica* **10**, 515-560 (2001)
- [13] Todd, M. J., Toh, K. C., Tütüncü, R. H.: On the Nesterov-Todd direction in semidefinite programming. *SIAM Journal on Optimization* **8**, 769-796 (1998)

- [14] Vandenberghe, L., Boyd, S.: Semidefinite programming. *SIAM Review* **38**, 49-95 (1996)
- [15] Vandenberghe, L., Boyd, S., and Wu, S.-P.: Determinant maximization with linear matrix inequality constraints. *SIAM Journal on Matrix Analysis and Applications* **19**, 499-533 (1998)
- [16] Wolkowitz, H., Saigal, R., Vandenberghe, L. (eds.) *Handbook of Semidefinite Programming: Theory, Algorithms and Applications*, Kluwer International Series in Operations Research and Management Science, Kluwer, Boston, 2000.
- [17] Yamashita, H.: A globally convergent primal-dual interior point method for constrained optimization. *Optimization Methods and Software* **10**, 443-469 (1998)