

A QUASI-POLYNOMIAL BOUND FOR THE DIAMETER OF GRAPHS OF POLYHEDRA

GIL KALAI AND DANIEL J. KLEITMAN

ABSTRACT. The diameter of the graph of a d -dimensional polyhedron with n facets is at most $n^{\log d+2}$

Let P be a convex polyhedron. The *graph* of P denoted by $G(P)$ is an abstract graph whose vertices are the extreme points of P and two vertices u and v are adjacent if the interval $[v, u]$ is an extreme edge (= 1-dimensional face) of P . The diameter of the graph of P is denoted by $\delta(P)$.

Let $\Delta(d, n)$ be the maximal diameter of the graphs of d -dimensional polyhedra P with n facets. (A facet is a $(d-1)$ -dimensional face.) Thus, P is the set of solutions of n linear inequalities in d variables. It is an old standing problem to determine the behavior of the function $\Delta(d, n)$. The value of $\Delta(d, n)$ is a lower bound for the number of iterations needed for Dantzig's simplex algorithm for linear programming with any pivot rule.

In 1957 Hirsch conjectured [2] that $\Delta(d, n) \leq n - d$. Klee and Walkup [6] showed that the Hirsch conjecture is false for unbounded polyhedra. They proved that for $n \geq 2d$, $\Delta(d, n) \geq n - d + [d/5]$. This is the best known lower bound for $\Delta(d, n)$. The statement of the Hirsch conjecture for bounded polyhedra is still open. For a recent survey on the Hirsch conjecture and its relatives, see [5].

In 1967 Barnette proved [1, 3] that $\Delta(d, n) \leq n^{3^{d-3}}$. An improved upper bound, $\Delta(d, n) \leq n^{2^{d-3}}$, was proved in 1970 by Larman [7]. Barnette's and Larman's bounds are linear in n but exponential in the dimension d . In 1990 the first author [4] proved a subexponential bound $\Delta(d, n) \leq 2^{\sqrt{(n-d)\log(n-d)}}$.

The purpose of this paper is to announce and to give a complete proof of a quasi-polynomial upper bound for $\Delta(d, n)$. Such a bound was proved by the first author in March 1991. The proof presented here is a substantial simplification that was subsequently found by the second author. See [4] for the original proof and related results. The existence of a polynomial (or even linear) upper bound for $\Delta(d, n)$ is still open. Recently, the first author found a randomized pivot rule for linear programming which requires an expected $n^{4\sqrt{d}}$ (or less) arithmetic operations for every linear programming problem with d variables and n constraints.

Theorem 1.

$$(1) \quad \Delta(d, n) \leq n^{\log d+2}.$$

Received by the editors July 1, 1991.

1991 *Mathematics Subject Classification.* Primary 52A25, 90C05.

The first author was supported in part by a BSF grant by a GIF grant. The second author was supported by an AFOSR grant.

Proof. Let P be a d -dimensional polyhedron with n facets, and let v and u be two vertices of P . Let k_v [k_u] be the maximal positive number such that the union of all vertices in all paths in $G(P)$ starting from v [u] of length at most k_v [k_u] are incident to at most $n/2$ facets. Clearly, there is a facet F of P so that we can reach F by a path of length $k_v + 1$ from v and a path of length $k_u + 1$ from u . We claim now that $k_v \leq \Delta(d, [n/2])$. Indeed, let Q be the polyhedron obtained from P by ignoring all the inequalities that correspond to facets that cannot be reached from v by a path of length at most k_v . Let ω be a vertex in $G(P)$ whose distance from v is k_v . We claim that the distance of ω from v in $G(Q)$ is also k_v . To see this consider the shortest path between v and ω in $G(Q)$. If the length of this path is smaller than k_v there must be an edge in the path that is not an edge of P . Consider the first such edge E . Since E is not an edge of P , it intersects a hyperplane H that corresponds to one of the inequalities that was ignored. This gives a path in P of length smaller than k_v from v to the facet of P determined by H , which is a contradiction.

We obtained that the distance from v to u is at most $\Delta(d - 1, n - 1) + 2\Delta(d, [n/2]) + 2$. This gives the inequality $\Delta(d, n) \leq \Delta(d - 1, n - 1) + 2\Delta(d, [n/2]) + 2$, which implies the statement of the theorem.

REFERENCES

1. D. W. Barnette, *W_v paths on 3-polytopes*, J. Combin. Theory 7 (1969), 62–70.
2. G. B. Dantzig, *Linear programming and extensions*, Princeton Univ. Press, Princeton, NJ 1963.
3. B. Grünbaum, *Convex polytopes*, Wiley Interscience, London, 1967.
4. G. Kalai, *Upper bounds for the diameter and height of polytopes*, Discrete Comput. Geom. 7 (1992) (to appear).
5. V. Klee and P. Kleinschmidt, *The d -steps conjecture and its relatives*, Math. Operation Research 12 (1987), 718–755.
6. V. Klee and D. Walkup, *The d -step conjecture for polyhedra of dimension $d < 6$* , Acta Math. 133 (1967), 53–78.
7. D. G. Larman, *Paths on polytopes*, Proc. London Math. Soc. (3) 20 (1970), 161–178.

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, ISRAEL AND
IBM ALMADEN RESEARCH CENTER, SAN JOSE, CALIFORNIA 95120

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE,
MASSACHUSETTS 02139