

Stephen L. Bloom

A REPRESENTATION THEOREM FOR THE LATTICE OF STANDARD CONSEQUENCE OPERATIONS

This paper will be published in *Studia Logica*.

§1. Introduction

A *sentential language* S is an algebra of finite type (i.e. a set equipped with a finite number of finitary functions) which is absolutely free in the class of all similar algebras, freely generated by a set of “(sentential) variables”. A consequence operation on S is a function C from the power set of S , PS , into PS satisfying the following three conditions:

- i) $X \subseteq C(X)$, all $X \subseteq S$;
- ii) $X \subseteq Y \Rightarrow C(X) \subseteq C(Y)$, all $X, Y \subseteq S$;
- iii) $C(C(X)) \subseteq C(X)$, all $X \subseteq S$.

A consequence operation C is algebraic (or finite) if

- iv) $C(X) = U\{C(Y) \mid Y \subseteq X, Y \text{ finite}\}$

and is *structural* if for all endomorphisms h of S ,

$$hC(X) \subseteq C(h(X)), \text{ all } X \subseteq S.$$

If C is both algebraic and structural, C is called *standard* (see [W] and [B]).

It is known that the collection ST (for “standard”) of all standard consequence operations on S forms a complete lattice when the ordering is defined by:

$$C \leq C' \text{ if } C(X) \subseteq C'(X), \text{ all } X \subseteq S.$$

Indeed, if $C_i, i \in I$ is a subset of ST , then $\bigvee(C_i, i \in I)$ is the consequence operation C satisfying:

$$\alpha \in C(X) \text{ iff, for some } i_1, \dots, i_n \in I,$$

$$\alpha \in C_{i_1}(C_{i_2}(\dots(C_{i_n}(X))\dots))$$

In this note, a representation of the lattice ST will be given.

For any sentential language S , let $L(S)$ be the first-order language (without equality) having one unary predicate symbol T , whose *terms* form an algebra isomorphic to S . (Thus we identify the formulas of S with the terms of $L(S)$). Let H denote the collection of all sentences of $L(S)$ which are the *universal closures* of formulas of the form

$$(*) \quad T(\tau_1) \wedge T(\tau_2) \wedge \dots \wedge T(\tau_k) \cdot \rightarrow T(\tau)$$

where $k \geq 0$, τ_i, τ are terms. When $k = 0$ (*) becomes just $T(\tau)$. H is thus the collection of (strict) basic Horn sentences (see [CK]).

For a subset Γ of H , let $\bar{\Gamma} = \{\sigma \in H \mid \Gamma \models \sigma\}$, where \models is the classical logical consequence operation. It is easily seen that the collection HN (for ‘‘Horn’’) of subsets of H of the form $\bar{\Gamma}, \Gamma \subseteq H$, is a complete lattice when ordered by set inclusion. Indeed, if $\bar{\Gamma}_i, i \in I$ are in HN , then

$$\Lambda(\Gamma_i, i \in I) = \bigcap(\bar{\Gamma}_i : i \in I),$$

i.e. the meet operation in HN is just intersection.

In the next section, we will prove the following

THEOREM. *The lattices ST and HN are isomorphic.*

COROLLARY. *ST is a complete, compactly generated (‘‘algebraic’’) lattice (i.e. every element is a join of compact elements).*

Indeed, the lattice HN is algebraic, since $\bar{\Gamma} = \bigvee(\bar{\Gamma}_f : \Gamma_f \subseteq \Gamma, \Gamma_f \text{ finite})$, and the sets $\bar{\Gamma}_f$, with Γ_f finite are compact in HN .

§2. Proof of the theorem

A matrix M is a pair (A, T) consisting of an algebra A similar to S and a subset T of A (we use the same letter for an algebra and its underlying set.)

Equivalently, a matrix is just a $L(S)$ -structure. Any matrix $M = (A, T)$ determines a structural consequence operation C_M on S by: for $X \subseteq S$, $\tau \in S$,

$\tau \in C_M(X)$ if, for any homomorphism $h : S \rightarrow A$, $h(\tau) \in T$ whenever $h(X) \subseteq T$.

For any consequence operation C on S , let $K(C)$ be the class of all matrices M such that $C \leq C_M$.

LEMMA 1. *A class K of matrices is $K(C)$ for some standard C iff $K = \text{Mod}\bar{\Gamma}$, for some $\Gamma \subseteq H$.*

PROOF. If $K = K(C)$, C standard, then by ([B], Theorem 2.6) it follows that $K = \text{Mod}\bar{\Gamma}$, where Γ consists of all sentences

$$(**) \quad \forall \vec{x} [T(\tau_1) \dots T(\tau_k) \rightarrow T(\tau)]$$

such that $\tau \in C(\tau_1, \dots, \tau_k)$.

Conversely, if $K = \text{Mod}\bar{\Gamma}$, define C by:

$\tau \in C(X)$ iff $\tau \in C_M(X)$, all $M \in K$. By ([B], Theorem 2.9) C is standard, and it is easily seen that $K = K(C)$.

REMARK. It is well-known [G] that an axiomatizable class K of matrices is $\text{Mod}\bar{\Gamma}$, some $\Gamma \subseteq H$ iff K is closed under arbitrary products and substructures.

LEMMA 2. *Suppose C_i are standard consequence operations on S , Γ_i are subsets of H such that $K(C_i) = \text{Mod}\bar{\Gamma}_i$, $i = 1, 2$. Then $C_1 \leq C_2$ iff $\bar{\Gamma}_1 \subseteq \bar{\Gamma}_2$.*

PROOF. By [W], Theorem 3.1) it follows that $C_1 \leq C_2$ iff $K(C_2) \subseteq K(C_1)$. The lemma thus follows easily.

From Lemmas 1, 2 it follows that the function $C \mapsto \Gamma_C$ taking the standard consequence C to the subset $\Gamma_C = \bar{\Gamma}_C$ of H with $K(C) = \text{Mod}\bar{\Gamma}_C$ is a lattice isomorphism $ST \rightarrow HN$. Notice that the sets $\bar{\Gamma}$, with Γ finite, correspond to the ‘‘finitely based’’ [B] consequence operations on S , i.e. those definable from a finite number of structural rules [W]. These consequence operations are precisely the compact elements of the lattice ST .

We close with several problems. It is known that if M is a finite matrix, C_M is standard, but we do not know whether C_M is compact on ST . Further, we don't know if the meet of two compact elements is compact.

References

- [B] S. L. Bloom, *Some theorems on Structural Consequence Operations*, IBM Research Report RC 4620, to appear in **Studia Logica**.
- [CK] C. C. Chang, H. J. Keisler, **Model Theory**, North Holland, Amsterdam 1973.
- [G] G. Grätzer, **Universal Algebra**, Van Nostrand, Princeton 1968.
- [W] R. Wójcicki, *Matrix approach in Methodology of Sentential Calculi*, **Studia Logica** 32 (1973), pp. 7–37.

Stevens Institute of Technology
Hoboken, N.Y., USA.