

A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators

Article

Published Version

Creative Commons: Attribution 3.0 (CC-BY)

Open Access

Godfray, H.C. J., Blacquiere, T., Field, L. M., Hails, R. S., Petrokofsky, G., Potts, S. G., Raine, N. E., Vanbergen, A. J. and McLean, A. R. (2014) A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proceedings of the Royal Society B-Biological Sciences*, 281 (1786). 20140558. ISSN 1471-2954 doi: <https://doi.org/10.1098/rspb.2014.0558> Available at <https://centaur.reading.ac.uk/37205/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://rspb.royalsocietypublishing.org/content/281/1786/20140558.abstract>

To link to this article DOI: <http://dx.doi.org/10.1098/rspb.2014.0558>

Publisher: The Royal Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators

H. Charles J. Godfray, Tjeerd Blacquière, Linda M. Field, Rosemary S. Hails, Gillian Petrokofsky, Simon G. Potts, Nigel E. Raine, Adam J. Vanbergen and Angela R. McLean

Proc. R. Soc. B 2014 **281**, 20140558, published 21 May 2014

Oxford Martin School Restatement Project No. 2.

Restatements are reviews of the natural science evidence base underlying areas of current policy concern and controversy. They are written in as policy neutral terms as possible and are designed to be read by an informed but not technically specialist audience. The evidence base is summarised in a series of paragraphs each of which includes an assessment of the nature of the evidence. Restatements are produced by a writing team reflecting the breadth of opinion on the topic in the science community and involve wide consultation with interested stakeholders. The final version is submitted for peer review in a scientific journal.

This pdf contains:

- | | |
|-------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Pages 2-4 | A short paper describing the restatement project |
| Pages 4-10 | The restatement itself which is the formal appendix of the paper |
| Pages 11-24 | An annotated bibliography of the evidence underlying the restatement (formally the Electronic Supplementary Material accompanying the paper) |

The paper is open access and can be freely distributed in its original version.



rspb.royalsocietypublishing.org

Review



Cite this article: Godfray HJ, Blacquière T, Field LM, Hails RS, Petrokofsky G, Potts SG, Raine NE, Vanbergen AJ, McLean AR. 2014 A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proc. R. Soc. B* **281**: 20140558.

<http://dx.doi.org/10.1098/rspb.2014.0558>

Received: 7 March 2014

Accepted: 7 May 2014

Subject Areas:

environmental science

Keywords:

insecticides, neonicotinoids, pollination, honeybee, bumblebee, pollinator

Authors for correspondence:

H. Charles J. Godfray and Angela R. McLean
e-mail: neonics@oxfordmartin.ox.ac.uk

[†]Present address: School of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada N1G 2W1.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2014.0558> or via <http://rspb.royalsocietypublishing.org>.

A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators

H. Charles J. Godfray¹, Tjeerd Blacquière², Linda M. Field³, Rosemary S. Hails⁴, Gillian Petrokofsky¹, Simon G. Potts⁵, Nigel E. Raine^{6,†}, Adam J. Vanbergen⁷ and Angela R. McLean¹

¹Oxford Martin School, c/o Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK

²Plant Research International, Wageningen University and Research, PO Box 16, 6700 AA Wageningen, The Netherlands

³Rothamsted Research, Harpenden, Herts AL5 2JQ, UK

⁴NERC Centre for Ecology and Hydrology, Crowmarsh Gifford, Wallingford OX10 8BB, UK

⁵School of Agriculture, Policy and Development, University of Reading, Reading, UK

⁶School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK

⁷NERC Centre for Ecology and Hydrology, Bush Estate, Penicuik, Edinburgh EH26 0QB, UK

There is evidence that in Europe and North America many species of pollinators are in decline, both in abundance and distribution. Although there is a long list of potential causes of this decline, there is concern that neonicotinoid insecticides, in particular through their use as seed treatments are, at least in part, responsible. This paper describes a project that set out to summarize the natural science evidence base relevant to neonicotinoid insecticides and insect pollinators in as policy-neutral terms as possible. A series of evidence statements are listed and categorized according to the nature of the underlying information. The evidence summary forms the appendix to this paper and an annotated bibliography is provided in the electronic supplementary material.

1. Introduction

Neonicotinoid insecticides are a highly effective tool to reduce crop yield losses owing to insect pests. Since their introduction in the 1990s, their use has expanded so that today they comprise about 30% by value of the global insecticide market [1]. They are commonly applied to crops as seed treatments, with the insecticide taken up systemically by the growing plant, so that it can be present in all plant parts, including nectar and pollen that bees and other pollinating insects collect and consume. Pollinators can potentially be exposed to neonicotinoids in other ways, for example through plant exudates, dust from planting machines and contamination of soil and water.

There is evidence that in Europe and North America many species of pollinators are in decline; both in abundance and distribution. There is a long list of potential causes for these declines, including parasites, disease, adverse weather and loss of habitat [2,3]. However, there has been particular concern about the impact on pollinators of the relatively recently introduced neonicotinoids and the European Union (EU) imposed a partial restriction on their use in December 2013. This decision has been criticized on the grounds that the benefits of neonicotinoid use outweigh any detriment they might cause.

The tension between the agricultural and environmental consequences of neonicotinoid use, and the recent EU restriction, has made this topic one of the most controversial involving science and policy. Here, we describe a project that aimed to provide a ‘restatement’ of the relevant natural science evidence base expressed in a succinct way that is comprehensible to non-expert readers. We have tried to be policy-neutral though are aware that complete neutrality is

impossible. The evidence restatement forms appendix A to this paper and is accompanied in the electronic supplementary material by a detailed annotated bibliography that provides an entry into the technical literature. The restatement is divided into six sections: after a description of the methodology and the importance of pollinators and insecticides, successive sections consider evidence for exposure paths, laboratory evidence for lethal and sublethal effects, the occurrence of residues in pollinators and their products in the environment, experiments conducted in the field, and consequences for pollinators at colony and population levels.

Experiments to establish the effect of defined doses of insecticides upon individual pollinators are required by regulatory authorities and can be carried out under laboratory conditions. These laboratory studies have the strength of allowing carefully controlled experiments to be performed on individual insects subjected to well-defined exposure. However, because they are conducted under artificial conditions, it is hard to assess a number of processes that may be relevant in the field. For example, neonicotinoids may affect the sensitivity of insects to other stressors; pollinators may actively avoid food contaminated by insecticide and responses at the colony or population level may mitigate or exacerbate the loss or impairment of individual insects. Nevertheless, such experiments provide important information about the range of concentrations where death or sublethal effects are to be expected.

Purely observational surveys in the field are used to establish the levels of exposure that occur under normal use. A number of large surveys in different countries have measured neonicotinoid residues in wild-foraging honeybees and unmanaged pollinators, as well as in nectar, pollen, honey and wax within bee colonies. These data are heavily weighted towards honeybees, and long time series are seldom available.

Experiments in the field are used to establish the impact of different doses of insecticide on pollinator behaviour, mortality and colony performance. They may be conducted as part of the registration process or for general research. One class of experiment involves bees artificially exposed to neonicotinoids and then observed to forage in the field. These are designed to discover whether neonicotinoids affect the performance of individual pollinators (and where appropriate their colonies) under field conditions. The critical issue here is whether the experimental exposure to insecticides is representative of what pollinators are actually likely to experience. The second class of experiment involves placing bee colonies in the environment in situations where they are exposed to crops treated with neonicotinoids, with suitable controls. These are large, difficult experiments where the unit of replication is typically the field site and where there are potentially many confounding factors to be taken into consideration. So far only one such study has been concluded successfully. The statistical power of this type of experiment is likely to be constrained by the expense and logistics of high levels of replication.

To understand the consequences of changing neonicotinoid use, it is important to consider pollinator colony- and population-level processes, the likely effect on pollination ecosystem services, as well as how farmers might change their agronomic practices in response to restrictions on neonicotinoid use. While all these areas are currently being researched, there is at present a relatively limited evidence base to guide policy-makers.

2. Material and methods

The literature on pollinators and neonicotinoids was reviewed and a first draft evidence summary produced by a subset of the authors. At a workshop, all authors met to discuss the different evidence components and to assign to each a description of the nature of the evidence using a restricted set of terms. We considered several options to describe the nature of the evidence we summarize including the GRADE [4] system widely used in the medical sciences, or the restricted vocabulary used by the International Panel on Climate Change [5]. However, none precisely matched our needs and instead we used a scoring system based on one previously developed for another 'restatement' project concerning bovine tuberculosis [6]. The categories we used are:

- [D_{ata}] a strong evidence base involving experimental studies or field *data* collection, with appropriate detailed statistical or other quantitative analysis;
- [E_{xp_op}] a consensus of *expert opinion* extrapolating results from related ecological systems and well-established ecological principles;
- [S_{upp_ev}] some *supporting evidence* but further work would improve the evidence base substantially; and
- [P_{rojns}] *projections* based on the available evidence for which substantial uncertainty often exists that could affect outcomes.

These categories are explicitly not in rank order.

A revised evidence summary was produced and further debated electronically to produce a consensus draft. This was sent out to 34 stakeholders or stakeholder groups including scientists involved in pollinator research, representatives of the farming and agrochemical industries, non-governmental organizations concerned with the environment and conservation, and UK government departments and statutory bodies responsible for pollinator policy. The document was revised in the light of much helpful feedback. Though many groups were consulted, the project was conducted completely independently of any stakeholder and was funded by the Oxford Martin School (part of the University of Oxford).

3. Results

The summary of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators is given in appendix A, with an annotated bibliography provided as the electronic supplementary material.

4. Discussion

The purpose of this project is not to conclude whether neonicotinoids are 'safe' or 'dangerous' but to try to help set out the existing evidence base. When neonicotinoids are used as seed dressing on crops visited by pollinators there is no doubt that these systemic insecticides are typically present in pollen and nectar and so bees and other pollinators can be exposed to them [7,8]. The concentrations in pollen and nectar are nearly always some way below those that would cause immediate death. The great problem is to understand whether the sublethal doses received by pollinators in the field lead to significant impairment in individual performance, and whether the cumulative effect on colonies and populations affects

pollination in farmed and non-farmed landscapes and the viability of pollinator populations [3].

For this topic, the published literature is a small fraction of the evidence that has been collected. The process of registering a new insecticide requires the production of detailed environmental risk assessments (see <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:093:0001:0084:EN:PDF> and <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:093:0085:0152:EN:PDF>). These include substantial evidence on toxicity to non-target organisms (including honeybees) and a range of further studies that will, in some cases, escalate to full-scale field trials of toxicity. The data generated in such studies are not typically in the public domain, or only in a form summarized by the regulatory agencies, and hence we have not been able to include reference to them. There are understandable commercial reasons for the withholding of this information, though the chief reason is not that it contains proprietary intellectual property but that the information would be commercially advantageous to a competitor in registering the compound when it is out of licence. We wonder if registration rules might be amended to allow this type of data to be published, a clear public good, without disadvantaging companies that had invested in its collection.

If neonicotinoids are not available, then farmers will have to choose alternative pest-management strategies, alternative crops or accept greater losses. The impact upon pollinators of withdrawing neonicotinoids will be greatly influenced by such choices. Farmers' likely strategies when faced with restrictions on the use of neonicotinoids are being researched, but there is currently only limited evidence to guide policy-makers in what changes to expect. This is just one aspect of human behaviour, economics and other social science that may be relevant to questions about threats to pollinators. However, it was not the purpose of this review to summarize the social science literature in this area (the annotated bibliography provides an entry into this literature).

There is clear evidence of the great value of neonicotinoids in agriculture [1] as well as the importance of the ecosystem services provided to agriculture by managed and wild pollinators [9]. Pollinators also have intrinsic importance as components of natural biodiversity that cannot, or can only inexactly, be accorded economic value. In some cases, intelligent regulation of insecticide use can provide 'win-wins' that improve both agricultural and biodiversity outcomes but in other cases there will be trade-offs, both within and between different agricultural and environmental objectives. Different stakeholders will quite naturally differ in the weightings they attach to the variety of objectives affected by insecticide use, and there is no unique answer to the question of how best to regulate neonicotinoids, an issue that inevitably has both economic and political dimensions. But economic and political arguments need to be consistent with the natural science evidence base, even though the latter will always be less complete than desirable. We hope that our attempt to set out this evidence base in as policy-neutral a manner as possible will stimulate discussion within the science community about whether our assessments are fair and where investment most needs to be made to strengthen them. We hope it will also make the evidence base less contested and so help stakeholders from all perspectives develop coherent policy and policy recommendations.

Acknowledgements. We are very grateful for extremely valuable comment and criticism from David Aston, Giles Budge, Peter Campbell, Norman

Carreck, Claire Carvell, Mark Clook, Christopher Connolly, Nicolas Desneux, Lynn Dicks, Adrian Dixon, Falko Driifhout, Dave Goulson, Matt Heard, Gemma Harper, Chris Hartfield, Emma Hockridge, Julie Howarth, Reed Johnson, Ged Kerins, Rebecca Lawrence, Paul Leonard, Marco Lodesani, Stephen Martin, Christian Maus, Piotr Medrzycki, Jane Memmott, Chris Mullin, John Mumford, Andy Musgrove, Ralf Nauen, Jeff Ollerton, Juliet Osborne, Robert Paxton, Deborah Procter, Stuart Roberts, Lucy Rothstein, Helen Roy, Cynthia Scott-Dupree, Fabio Sgolastra, Matt Shardlow, Richard Shore, Lisa Smith, Dave Spurgeon, Steve Sunderland, David Williams and Paul de Zylva. Their insights have strongly shaped the final document, but not all their helpful suggestions were or could be included and the final version is the responsibility of the authors alone.

Appendix A. A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators

For an annotated bibliography of the evidence supporting each statement, see the electronic supplementary material.

(a) Introduction and aims

- (1) Wild and managed insect pollinators play a critical role in the production of a variety of different foods (and in the case of honeybees also produce various 'hive products' of which the most important is honey) and are an important functional and cultural component of biodiversity. Insecticides are applied to crops to control insect pests and make a very important contribution to achieving high yields. Insecticides kill insects and thus clearly have both positive and negative effects on different aspects of food security and the environment. Concern has been expressed by a number of bodies that neonicotinoid insecticides may be harming pollinators and a partial restriction on their use in the EU came into force across all 28 member states in December 2013 (to be reviewed after 2 years). Other bodies have criticized this decision, arguing that the benefits of neonicotinoid use outweigh their costs.
- (2) The aim here is to provide a succinct summary of the evidence base relevant to policy-making in this area as of April 2014. It also provides a consensus judgement by the authors on the nature of the different evidence components; a consensus arrived at using the studies listed in the annotated bibliography. We use the following descriptions, which explicitly are not a ranking, indicated by abbreviated codes. Statements are considered to be supported by:
 - [Data] a strong evidence base involving experimental studies or field *data* collection, with appropriate detailed statistical or other quantitative analysis;
 - [Exp_op] a consensus of *expert opinion* extrapolating results from related ecological systems and well-established ecological principles;
 - [Supp_ev] some *supporting evidence* but further work would improve the evidence base substantially; and
 - [Projns] *projections* based on the available evidence for which substantial uncertainty often exists that could affect outcomes.
- (3) The review focuses on the natural science evidence relevant to pollinator policy in the EU but includes relevant data from other regions; its scope does not include

evidence from social sciences and economics. The statements are based on the evidence in the peer-reviewed scientific literature, though the annotated bibliography also notes the existence of information in non-reviewed reports and industry studies.

(b) Pollinators and neonicotinoid insecticides

- (4) Insect pollinators are required to achieve optimum yield and quality for a number of important food crops. The most economically significant crops in the UK include oilseed rape (canola), soft fruits (strawberry, raspberry, etc.), top fruits (apple, pear, plum, etc.) and vegetables (courgettes, runner beans, tomato, etc.), whereas in continental Europe sunflower, peaches, melon and other crops are also important. Insect pollinators are important for both field crops and those grown under glass, though in their absence some crops can, to differing extents, be wind- or self-pollinated without the involvement of insects. Many plant species in pastureland and non-agricultural habitats require insect pollinators for successful reproduction [D_{ata}].
- (5) A lack of pollinators can reduce crop yields and quality [D_{ata}], and there is some evidence that pollinator diversity can reduce the variance in pollination and hence improve crop yield stability [S_{upp_ev}]. Where insect-pollinated crops are grown in glasshouses or 'polytunnels' the introduction of pollinators can be particularly important for both quality and quantity of yield [D_{ata}]. There is emerging evidence for the potential of economically significant pollination deficits in some UK field crops in some years [S_{upp_ev}], but data do not currently exist to determine whether observed changes in pollinator abundance and diversity (see para. 7) have affected the economic value of crop yields [E_{xp_op}].
- (6) Pollination may be carried out by wild or managed insects. The most important pollinators for crops include honeybees, which are native to Europe (their status in the British Isles is unclear [E_{xp_op}]) but are now almost entirely managed, bumblebees, solitary bees and true flies (including hoverflies).¹ Other pollinators such as butterflies and moths are not as important for crop pollination, particularly in northern temperate regions, but do pollinate wild plant species. Wild pollinators can be viewed as an element of natural capital² that provides (with managed species) pollination, an ecosystem service of economic importance to society. Pollinators are also an important component of a nation's biodiversity [D_{ata}].
- (7) Data from volunteer recording schemes that record species presence (but not abundance or absence) have revealed changes in the diversity and distribution of pollinators. In Great Britain, The Netherlands and Belgium (where the best data exist) the average numbers of species of bumblebees, butterfly and moths, and solitary bees in different areas have declined since the 1950s [D_{ata}]. There is some evidence of a recent slowdown in the rate of decline in species richness (for bumblebees in all three European countries) and also some increases (solitary bees in Great Britain and The Netherlands but not in Belgium where the decline continues) [D_{ata}]. The data for hoverflies are more complex with species richness reported to have increased, decreased or remained unchanged depending on location and the geographical scale of the analysis.
- (8) Honeybees throughout Europe (and elsewhere) have been severely affected by the introduction of the *Varroa destructor* mite which both parasitizes bees and acts as a vector for a number of debilitating and paralytic honeybee viruses [D_{ata}]. In addition, honeybee colony losses have increased in frequency across Europe and the USA because of overwintering mortality [D_{ata}] which is thought to arise from multiple factors, including adverse weather, poor nutrition as well as parasites and disease [S_{upp_ev}]. Some of these losses in the USA have been ascribed to a particular syndrome, colony collapse disorder, though its precise nature is debated [E_{xp_op}]. Not all parts of the world have experienced recent increases in overwintering colony mortality [D_{ata}].
- (9) Neonicotinoids are a relatively new class of insecticide, introduced in the early 1990s. They target the nicotinic acetylcholine receptor (nAChR) with high affinity for insect receptors and low affinity for mammalian receptors and have relatively low (but not zero) mammalian and bird toxicity. They can be used as sprays, applied to soils as drenches or in granular form, introduced into irrigation water or injected into trees. However, they are most frequently (approx. 90% by volume in the UK) applied as seed treatments with the insecticide being taken up systemically by the growing plant. The convenience and cost-effectiveness of seed treatments, the development of resistance to other classes of insecticide by many insect pests, and restrictions on the use of other compounds, have resulted in neonicotinoids capturing 28.5% of the global insecticides market (2011; worth US\$3.6B) and their wide use in Europe [D_{ata}].
- (10) Five neonicotinoids are approved for use in the EU: three from the *N*-nitroguanidine group—clothianidin, imidacloprid and thiamethoxam (metabolized to clothianidin in the plant, insect and environment); and two from the *N*-cyanoamidine group: thiacloprid and acetamiprid. Concern over their possible effects on pollinators has focused on the first three because they are the most used compounds, they have greater honeybee toxicity and they are used as seed treatments so can be present in the pollen and nectar of treated crops [D_{ata}].
- (11) In Europe (and elsewhere), environmental risk assessments of pesticides including all neonicotinoids are required before a product can come to market. A tiered approach has been adopted to ensure cost-effectiveness and proportionality. The tiers start with laboratory tests to determine hazard to a standard set of seven non-target organisms (including honeybees) and, if potential hazards are identified, may progress through more complex semi-field experiments and modelling to simulate exposure under different more realistic conditions, culminating with full-scale toxicity assessments to identify potential risks in the field. Field trials were conducted during the original environmental risk assessment process

for neonicotinoids. Extensive data are often generated during the registration process but typically is not placed in the public domain, except in summary form [D_{ata}].

(c) Exposure of pollinators to neonicotinoid insecticides

- (12) Neonicotinoids have been widely used in Europe as a seed treatment for oilseed rape, sunflowers, maize, potato, soya bean (and other crops such as cereals and beets not visited by pollinators).
 - (a) A single treated oilseed rape seed is typically treated with approximately 35 µg neonicotinoids and a maize seed with 1.2 mg (see Endnote 3) [D_{ata}].
 - (b) Pollinators may be exposed to neonicotinoids applied as sprays. The use of *N*-nitroguanidine neonicotinoids at flowering time is restricted in most countries though acetamiprid and thiacloprid (from the less toxic *N*-cyanoamidine group) are sprayed on raspberries, fruit trees and oilseed rape at flowering time [D_{ata}].
- (13) The plant absorbs some of the insecticide from the seed treatment and as it grows the insecticide spreads to all plant parts including the nectar and pollen that bees and other pollinators collect and consume [D_{ata}].
 - (a) Estimates of the concentration of neonicotinoids in the pollen and nectar of seed-treated crops vary considerably with average maximum levels (from 20 published studies) of 1.9 (nectar) and 6.1 (pollen) ng g⁻¹. Concentrations vary across crops and can be appreciably higher if neonicotinoids are applied as foliar sprays, soil drenches or through drip irrigation [D_{ata}].
- (14) Some plants secrete droplets of liquid (xylem sap) called guttation fluid at leaf tips or margins. High concentrations of neonicotinoids have been measured in the guttation fluid of seed-treated plants (up to 10⁴–10⁵ times that in nectar), especially when plants are young [D_{ata}]. There has been concern that were pollinators to use guttation fluid as a source of water they would ingest highly toxic levels of insecticides. The available evidence does not suggest that pollinators collect guttation fluid containing neonicotinoids to any great extent, in part because it chiefly is present at times of the year when crops are unattractive to pollinators and other sources of water are present [E_{xp_op}].
- (15) Dust emitted from seed drilling machines can contain high concentrations of neonicotinoids; as well as being deposited on the soil, the dust can drift to contaminate neighbouring flowering crops and natural vegetation as well as surface waters. Sporadic incidents of mass honeybee mortality in several EU countries, the USA and Canada have been caused by dust from seed drilling machines [D_{ata}].
 - (a) Issues concerning dust chiefly involve the formulation of the insecticide, in particular, how it is made to 'stick' to the seed. EU and national regulations on formulation and seed drilling have been introduced to reduce the risks of these problems [D_{ata}].
- (16) Neonicotinoids introduced into the environment as seed treatments can affect soil insects and other invertebrates, effects considered in insecticide evaluation and registration. They persist in the environment with

typical half-lives estimated to be of the order 15–300 days (with some longer estimates from laboratory studies and in the field under drought and freezing conditions). There is evidence that neonicotinoids can accumulate in soils when treated crops are grown repeatedly in the same field. Neonicotinoids can sometimes, but not always, be detected in weeds or in subsequent crops grown in the same soil, though when present the concentrations are considerably lower than in the target crop. Neonicotinoids have been detected in surface or groundwater around fields where they have been used as seed treatments [S_{upp_ev}].

- (17) Bees bring pollen and nectar (which in social bees is often extensively modified post-ingestion) to their hives or nests to feed their developing larvae [D_{ata}] which thus may have different patterns of exposure and susceptibility compared with adults (see also para. 24) [S_{upp_ev}].
- (18) The risk of exposure to neonicotinoids for different pollinator species will be influenced by many aspects of their biology and ecology including body size, flower preference, whether they are a social species, and whether the time of year at which they are active (or in the case of social species experiencing rapid colony growth) coincides with the flowering of neonicotinoid-treated crops. There may also be differences in the physiological susceptibility of different pollinator species to neonicotinoids [E_{xp_op}].
- (19) The exposure of pollinators to neonicotinoids will be affected by the distribution of flowering crops in the landscape, the fraction that are treated with neonicotinoids, the length of time the treated crops are in flower, and the availability of alternative, suitable floral resources (including weeds and managed resources in floral strips, wildflower headlands, untreated crops, etc.) and whether they are contaminated with insecticide. Over multiple years the frequency of treated crops in agricultural rotations will affect long-term population exposure [E_{xp_op}].
- (20) The distance between treated fields and nest sites or honeybee hives will affect insect exposure to neonicotinoids [E_{xp_op}].
 - (a) Pollinators can forage over a large area: the maximum foraging distance for bumblebees is 2–3 km from the colony (though with considerable variation) and for honeybees 10–15 km (median distances are 1–6 km); some solitary bees may only forage a few hundred metres or less. Observed foraging distances are strongly influenced by the distribution of flowering plants [D_{ata}].
- (21) *Summary*. There are several proven pathways through which pollinators may be exposed to neonicotinoid insecticides applied as seed treatments (or in other ways). Quantitative information about the extent and significance of these different routes in the published literature is poor [E_{xp_op}].

(d) Laboratory studies of lethal and sublethal effects of neonicotinoids

- (22) Estimates of LD₅₀s (see Endnote 4) for different neonicotinoid-pollinator combinations are available,

although a majority of the studies have considered only the honeybee [D_{ata}].

- (a) The acute oral LD₅₀s for the major neonicotinoids have been estimated (by EFSA⁵) to be 3.7 ng per honeybee for imidacloprid, 3.8 ng per honeybee for clothianidin and 5.0 ng per honeybee for thiamethoxam (these estimates are used in the calculations below). A meta-analysis of 14 studies of imidacloprid estimated an LD₅₀ of 4.5 ng per honeybee (95% confidence limits 3.9–5.2 ng) [D_{ata}].
 - (b) Equivalent acute contact LD₅₀s have been estimated (by EFSA) to be 81 ng per honeybee for imidacloprid, 44 ng per honeybee for clothianidin and 24 ng per honeybee for thiamethoxam [D_{ata}].
 - (c) There is considerable variation among LD₅₀s measured across different bee species, and this is influenced by type of neonicotinoid and mode of application [D_{ata}]. This complicates simple comparison with honeybee data [E_{xp_op}].
 - (d) A honeybee, returning to the hive after foraging, typically carries 25–40 mg nectar or 10–30 mg pollen. If nectar or pollen is contaminated with insecticide at the concentrations described in Para. 13a, then these loads will contain approximately 0.06 ng (nectar) or 0.12 ng (pollen) of insecticide. Depending on the type of neonicotinoid this is 1–3% of the LD₅₀ acute oral dose (though note that none of the pollen and hardly any of the nectar is metabolized by the forager). A colony of 10 000 workers was observed to store 750 g of pollen in four days. If all the pollen was similarly contaminated this equates to 8–11% of the acute oral LD₅₀ [P_{rojns}].
 - (e) Maximum pollen consumption is found among nursing honeybees that can consume 7.2 mg d⁻¹. If the pollen contains 6.1 ng g⁻¹ neonicotinoid the daily intake is 0.044 ng or, depending on the compound, 0.8–1.1% of the acute oral toxicity LD₅₀. Maximum nectar consumption is found among nectar-foraging honeybees and can be 32–128 mg d⁻¹. If nectar contains 1.9 ng g⁻¹ neonicotinoid the daily intake is 0.061–0.243 ng, or 1.2–6.7% of the LD₅₀ acute oral [P_{rojns}].
 - (f) Honeybee colonies collect pollen and nectar from multiple sources, which dilutes the effects of foraging on neonicotinoid-treated crops [D_{ata}]. For this reason and because they are based on the average maximum neonicotinoid concentrations in Para. 13a, the calculations in subparagraphs d and e above should be viewed as a worst-case scenario [E_{xp_op}].
- (23) Prolonged exposure of pollinators in the laboratory to doses of neonicotinoids that do not cause immediate death can reduce longevity (chronic toxicity). Because chronic effects can be estimated in many different ways, comparisons are harder than for acute toxicity [D_{ata}].
 - (a) For honeybees and bumblebees, chronic lethal effects have typically been reported when bees are fed diets containing 10–20 ng g⁻¹ neonicotinoid over 10–20 days, although some studies with higher doses have not observed such effects [D_{ata}].
 - (b) These neonicotinoid concentrations are higher than the worst-case assumptions of maximum insecticide consumption in para. 22e [P_{rojns}].
 - (24) Effects of neonicotinoids on adult pollinators have been detected in the laboratory at doses substantially below those that cause death. At the lowest doses responses involve metabolic changes (for example, in acetylcholinesterase activity) and subtle neurological and behavioural responses. As doses increase (including concentrations in food similar to that observed in the nectar and pollen of treated crops) olfactory learning, memory and feeding behaviour can be affected, though there is considerable variability in the results reported in different studies. When doses approach lethal concentrations substantial neurological and locomotory impairment can occur [D_{ata}].
 - (a) The majority of studies have involved honeybees; where comparisons of honeybees with bumblebees and solitary bees have been made differences are frequently observed, although these depend on species, assay and type of neonicotinoid and general patterns are difficult to discern [S_{upp_ev}].
 - (b) There has been debate in the literature as to the extent that neonicotinoids accumulate in pollinators; recent studies have suggested that bees have a substantial capacity to extrude neonicotinoids from cells and tissue (honeybees were estimated to clear 2 ng d⁻¹ imidacloprid from their body—approximately 50% of oral LD₅₀—and larger bumblebees 7 ng d⁻¹) [D_{ata}].
 - (25) Sublethal effects on larval development and colony productivity have been identified in the laboratory.
 - (a) Delayed larval and pupal developments have been observed in honeybees though at neonicotinoid concentrations higher than those expected to occur in the field [D_{ata}].
 - (b) Increases in development time, and reductions in worker egg laying, worker production, worker longevity and male and new queen (gyne) production have been observed in bumblebee colonies when food is provided containing concentrations of neonicotinoids towards the high end of those observed in nectar and pollen in treated crops in the field. Similar results have been found for larval development and reproductive output in solitary bees [S_{upp_ev}].
 - (26) Stressed pollinators tend to be more susceptible to neonicotinoids (and vice versa), although data are largely restricted to honeybees [S_{upp_ev}].
 - (a) Honeybees stressed by disease are more susceptible (lethal and sublethal effects occur at lower doses) to neonicotinoids, whereas in bumblebees synergistic effects of neonicotinoids and parasites on queen longevity, but not other colony parameters, have been observed. Neonicotinoids can modulate insect innate immunity negatively affecting anti-viral and other defences [D_{ata}].
 - (b) Laboratory molecular biological studies show a potential for the presence of other pesticides (targeted at fungi and *Varroa*) to exacerbate the effects of neonicotinoids though there is limited evidence for such effects from studies with live insects [S_{upp_ev}].
 - (c) It is likely that pollinators exposed to poorer diets are more susceptible to neonicotinoids (and other stressors) [E_{xp_op}].
 - (27) In interpreting these laboratory results, the following issues need to be considered:
 - (a) There is extensive information on the acute lethality of major neonicotinoids in honeybees, but data on other effects, on other pollinators and with the full range of neonicotinoids, are more limited [E_{xp_op}].

- (b) Stress affects insect responses to neonicotinoids and laboratory conditions may be more or less stressful than in the field, an effect that is probably pollinator-species specific and rarely directly assessed in experiments [E_{xp_op}].
 - (c) Laboratory experiments normally involve feeding pollinators with sugar solution or mixed pollen which may affect insects differently to naturally collected food [E_{xp_op}].
 - (d) Chronic and sublethal effects will depend on the pattern of dietary consumption and the rate at which ingested neonicotinoids are cleared from the body [E_{xp_op}]. In addition, neonicotinoids can act as anti-feedants and hence may affect pollinators through reduced food intake, though typically at concentrations higher than expected in the field. How insecticide treated food is presented to pollinators in laboratory experiments, and whether the insects have access to alternative foods, will thus influence the observed responses [S_{upp_ev}].
 - (e) It is challenging to study the impacts of neonicotinoids on entire colonies in the laboratory (particularly for honeybees). As a result, the majority of laboratory studies examine effects on individual bees or queenless groups (often referred to as micro-colonies in bumblebee studies). These results need careful interpretation when assessing how they might translate to whole colony impacts for social bees in the field [E_{xp_op}].
- (28) *Summary.* The strengths of laboratory studies are that they allow carefully controlled experiments to be performed on individual insects subjected to well-defined exposure. The weaknesses are that they are conducted under very artificial conditions (which may affect tolerance to external stress), any avoidance response by the insect is limited and hence the exposure dose and form is determined solely by the experimenter, and responses at the colony or population level are both difficult to study and to extrapolate to the field. Nevertheless, they provide important information about the range of concentrations where death or sublethal effects may be expected to occur [E_{xp_op}].

(e) Neonicotinoid residues observed in pollinators in the field

- (29) Nectar and pollen collected from bees constrained to feed on treated crops have similar insecticide concentrations to those found in samples taken from the plant [D_{ata}].
- (30) There have been few surveys of pesticide and metabolite levels in honeybees in the field. Two studies in Belgium (sample size, $n = 48$ and 99) and one in the USA ($n = 140$) found no honeybees with residues, while a survey in France conducted in 2002–2005 ($n = 187$) detected imidacloprid in 11% of honeybees (at concentrations of 0.03–1.0 ng per bee) [D_{ata}]. We are aware of no data on other pollinators [E_{xp_op}].
- (31) Insecticide residues are more likely to be found in nectar and pollen collected by honeybees and in honey than in the insects themselves. Thus, the French study that found imidacloprid residues in 11% of the bees sampled

also found residues in 22% of honey samples and 40% of pollen samples (mean and range: 0.9, 0.2–5.7 ng g⁻¹). Some large surveys (e.g. a Spanish study with $n = 1021$) found no contaminated pollen; a German study that surveyed hives ($n = 215$) after oilseed rape flowering found low incidence of those neonicotinoids used in seed treatments (though higher incidence of thiacloprid); an American study found imidacloprid in 3% of pollen ($n = 350$) and 1% of wax samples ($n = 208$) [D_{ata}].

- (32) *Summary.* Neonicotinoids can be detected in wild pollinators as well as honeybee and bumblebee colonies but data are relatively few and restricted to a limited number of species. Studies to date have found low levels of residues in surveys of honeybees and honeybee products. Observed residues in bees and the products they collect will depend critically on details of spatial and temporal sampling relative to crop treatment and flowering [E_{xp_op}].

(f) Experiments conducted in the field

- (33) This section discusses recent studies that have explored the consequences of providing bee colonies placed in the field with food containing insecticide, as well as experiments where the performance of colonies placed adjacent to fields treated or not treated with neonicotinoids are compared. Some earlier studies with limited statistical power are listed in the annotated bibliography [E_{xp_op}].
- (34) Schneider *et al.* 2012 [10]. Individual honeybees were given single sublethal doses of imidacloprid or clothianidin and their foraging behaviour was monitored. Reductions in foraging activity and longer time foraging flights were not observed at field-relevant doses although negative effects were seen at doses greater or equal to 0.5 ng per bee (clothianidin) or 1.5 ng per bee (imidacloprid) [D_{ata}].
 - (a) These doses are higher than those likely to be encountered by honeybees foraging on nectar from treated plants (see calculations in para. 22e) [E_{xp_op}].
- (35) Henry *et al.* 2012 [11]. Honeybees fed a single high dose of thiamethoxam (1.34 ng, equivalent to 27% of the LD₅₀) and then released away from the hive were significantly less likely to return successfully than controls. The return rate depended on the local landscape structure and the extent of the honeybees' experience of the landscape. The failure to return per trip was estimated to be up to twice the expected background daily mortality [D_{ata}].
 - (a) The rate of forager loss per trip (15%) was analysed as if it were excess daily mortality but as foraging honeybees make 10–30 trips per day real loss rates would be very much higher, reflecting the high dose of insecticide used in the experiment (see para. 22e for calculation of likely field doses) [E_{xp_op}].
 - (b) Assuming honeybees were exposed every day to this dose rate (much higher than expected from observed residues in pollen and nectar), mathematical modelling of colony development predicted severe decline within a season though this conclusion depends critically on poorly understood aspects of honeybee colony dynamics [P_{rojns}].
- (36) Whitehorn *et al.* 2012 [12]. Bumblebee (*Bombus terrestris*) colonies fed exclusively on imidacloprid-treated sugar water (at two concentrations: 0.7 or 1.4 ng g⁻¹) and

pollen (either 6 or 12 ng g⁻¹) for two weeks in the laboratory before being placed in the field (for six weeks) showed reductions in growth rate and queen production. A subsequent study [13] using the same concentrations of imidacloprid found the bumblebees' capacity to forage for pollen (but not nectar) was impaired [D_{ata}].

- (a) The concentrations of insecticide are at the high end of those observed in the nectar and pollen of treated plants (Para. 13a) and are likely to be greater than most bees will receive in the field because alternative food sources were not available [E_{xp_op}].
- (37) Gill *et al.* 2012 [14]. Bumblebee (*B. terrestris*) colonies given access to sugar water containing imidacloprid (10 ng g⁻¹) and allowed to forage for pollen and nectar in the field grew more slowly than controls; individual foragers from imidacloprid-treated colonies were less successful at collecting pollen, and treated colonies sent out more workers to forage and lost more foragers, compared to controls. Combined exposure to imidacloprid and a second pesticide of a different class (a pyrethroid) tended to reduce further colony performance and increase the chances of colony failure [D_{ata}].
- (a) The concentration of insecticide in the sugar water is within the range observed in nectar in the field but considerably higher than the average (1.9 ng g⁻¹; Para. 13a). The actual amount of imidacloprid consumed by individual bumblebees was not measured but will be diluted through foraging from other sources (no pollen was provided). Although it is difficult to make precise comparisons, the pyrethroid concentrations used were towards the upper end of recommended application rates for field or fruit crops [E_{xp_op}].
- (38) Thompson *et al.* 2013 [15]. Bumblebee (*B. terrestris*) colonies were placed adjacent to single oilseed rape fields grown from seeds that were treated with clothianidin, imidacloprid or had no insecticidal seed treatment. No relationship between the oilseed rape treatment and insecticide residues was observed, presumably because the bees were foraging over spatial scales larger than a field. Insecticide residues varied among colonies and the authors reported no evidence of a correlation with colony performance [D_{ata}].
- (a) The experimental design, in particular the lack of replication at field level and absence of a clear effect of treatment, allows only limited inference about the effects of neonicotinoids in the field [E_{xp_op}].
- (39) Pilling *et al.* 2013 [16]. Over a 4 year period, honeybee colonies (six per 2 ha field) were placed beside thiamethoxam-treated or control fields of maize (three replicates) or oilseed rape (two replicates) for between 5 and 8 days (first 3 years) or 19 and 23 days (fourth year) to coincide with the crop flowering period (at other times the colonies were kept in woodland presumed to have no local exposure to insecticides). Honeybees from treatment hives had higher concentrations of insecticide residues, but no differences in multiple measures of colony performance or overwintering survival were observed [D_{ata}].
- (a) Levels of replication precluded formal statistical analysis though the lack of any differences between treatment and control was reasonably consistent across field sites [E_{xp_op}].

- (40) *Summary.* The experiments described in Paras. 33–37 involve bees artificially exposed to neonicotinoids and observed to forage in the field. They show the potential for neonicotinoids to affect the performance of individual pollinators and pollinator colonies in the field. The main issue for their interpretation is the extent to which the doses received by the bees are representative of what they will receive under normal use of neonicotinoids in the field. It appears that most studies have used concentrations at the high end of those expected in the field. The experiments described in Paras. 38 and 39 are true field experiments in the sense that the treatments involve the normal use of neonicotinoids, though only the Pilling *et al.* [16] study was successfully concluded and found no effects of neonicotinoids, but with limited statistical power to detect differences [E_{xp_op}].

(g) Consequences of neonicotinoid use

- (41) At the colony or population level, there may be processes that can compensate for the deaths of individual insects which would mitigate the potential effects of mortality caused by neonicotinoid insecticides. Thus, the deaths of individual pollinators may not lead to a simple proportionate decrease in the overall numbers of that pollinator species. In the case of rare species, extra mortality caused by insecticides could lead to a threshold population density being crossed below which the species declines to extinction, hence magnifying their effects. However, there is a weak evidence base to help understand the presence and magnitude of these effects in the field. Models of honeybee and bumblebee colony dynamics, as well as population-level models of all pollinators, are important tools to explore these effects [E_{xp_op}].
- (42) There is evidence that some crops do not always receive sufficient pollination [D_{ata}], and further limited evidence that this has increased in recent decades [S_{upp_ev}]; but the information available does not allow us to determine whether or not this has been influenced by the increased use of neonicotinoids [E_{xp_op}]. Whether pollination deficits in wild plants have increased is not known [E_{xp_op}].
- (43) Declines in the populations of many insect species in general and pollinators in particular have been observed (para. 7) although the decline in bees predate by some decades the introduction of neonicotinoid insecticides, and there is some evidence of a recent abatement in the rate of decline for some groups [D_{ata}]. Habitat alteration (especially in farmland) is widely considered to be the most important factor responsible. The evidence available does not allow us to say whether neonicotinoid use has had an effect on these trends since their introduction [E_{xp_op}].
- (44) There have been marked increases in overwintering mortality of managed honeybee populations in recent decades (para. 8) [D_{ata}]. It has been suggested that insecticides (particularly neonicotinoids) may be wholly or partly responsible. The weak evidence base cannot at present resolve this question although honeybee declines began before the wide use of neonicotinoids and there is poor geographical correlation between neonicotinoid use and honeybee decline [E_{xp_op}]. Two studies using different structured methodologies have explored this question.

- (a) Cresswell *et al.* 2012 [17]. Used ‘Hill’s epidemiological “causality criteria”’ and concluded that the evidence base did not currently support a role for dietary neonicotinoids in honeybee decline but that this conclusion should be seen as provisional [E_{xp_op}].
- (b) Staveley *et al.* 2014 [18]. Used ‘causal analysis’ methodology and concluded that neonicotinoids were ‘unlikely’ to be the sole cause of honeybee decline but could be a contributing factor [E_{xp_op}].
- (45) Neonicotinoids are efficient plant protection compounds and if their use is restricted farmers may switch to other pest-management strategies (for example, different insecticides applied in different ways or non-chemical control measures) that may have effects on pollinator populations that could overall be more or less damaging than neonicotinoids. Alternatively, they may choose not to grow the crops concerned, which will reduce exposure of pollinators to neonicotinoids but also reduce the total flowers available to pollinators [E_{xp_op}].
- (46) *Summary.* To understand the consequences of changing neonicotinoid use, it is important to consider pollinator colony-level and population processes, the likely effect on pollination ecosystem services, as well as how

farmers might change their agronomic practices in response to restrictions on neonicotinoid use. While all these areas are currently being researched there is at present a limited evidence base to guide policy-makers [E_{xp_op}].

Endnotes

¹The honeybee is *Apis mellifera* (Apidae); bumblebees are *Bombus* species (Apidae), while solitary bees belong to a number of different, related families (Apiformes). Bees belong to the order Hymenoptera, while true flies are in the order Diptera (hoverflies are in the family Syrphidae) and butterflies and moths in the order Lepidoptera.

²Natural capital describes the components of the natural environment that produce value (directly and indirectly) for people; the actual benefits are called ecosystem services (which can be thought of as the flows that arise from natural capital stocks).

³A milligram (mg) is one thousandth (10^{-3}) of a gram (g); a microgram (μg) is one millionth (10^{-6}) of a gram and a nanogram (ng) is one billionth (10^{-9}) of a gram. We express concentrations as nanograms insecticide in 1 g of substance and hence in units of ng g^{-1} (the equivalent metrics ‘one part per billion’ or $1 \mu\text{g kg}^{-1}$ are frequently used in the literature). Concentrations are also sometimes expressed per volume ($\mu\text{g l}^{-1}$); for neonicotinoids 1 ng g^{-1} is approximately $1.3 \mu\text{g l}^{-1}$ in a 50% weight for weight sugar solution.

⁴The LD₅₀ (lethal dose 50%) is the amount of a substance that kills 50% of exposed organisms.

⁵European Food Safety Authority.

References

- Jeschke P, Nauen R, Schindler M, Elbert A. 2011 Overview of the status and global strategy for neonicotinoids. *J. Agric. Food Chem.* **59**, 2897–2908. (doi:10.1021/jf101303g)
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353. (doi:10.1016/j.tree.2010.01.007)
- Vanbergen AJ, Insect Pollinators Initiative. 2013 Threats to an ecosystem service: pressures on pollinators. *Front. Ecol. Environ.* **11**, 251–259. (doi:10.1890/120126)
- Guyatt GH, Oxman AD, Vist G, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ. 2008 GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *Br. Med. J.* **336**, 924–926. (doi:10.1136/bmj.39489.470347.AD)
- IPCC. 2007 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change (Contribution of Working Group II)*. Cambridge, UK: Cambridge University Press.
- Godfray HJ *et al.* 2013 A restatement of the natural science evidence base relevant to the control of bovine tuberculosis in Great Britain. *Proc. R. Soc. B* **280**, 20131634. (doi:10.1098/rspb.2013.1634).
- Blacquière T, Smaghe G, Van Gestel CAM, Mommaerts V. 2012 Neonicotinoids in bees: A review on concentrations, side-effects and risk assessment. *Ecotoxicology* **21**, 973–992. (doi:10.1007/s10646-012-0863-x)
- Goulson D. 2013 An overview of the environmental risks posed by neonicotinoid insecticides. *J. Appl. Ecol.* **50**, 977–987. (doi:10.1111/1365-2664.12111)
- Klein AM, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T. 2007 Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. B* **274**, 303–313. (doi:10.1098/rspb.2006.3721)
- Schneider CW, Tautz J, Grünwald B, Fuchs S. 2012 RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. *PLoS ONE* **7**, e30023. (doi:10.1371/journal.pone.0030023)
- Henry M, Béguin M, Requier F, Rollin O, Odoux JF, Aupinel P, Aptel J, Tchamitchian S, Decourtye A. 2012 A common pesticide decreases foraging success and survival in honey bees. *Science* **336**, 348–350. (doi:10.1126/science.1215039)
- Whitehorn PR, O’Connor S, Wackers FL, Goulson D. 2012 Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* **336**, 351–352. (doi:10.1126/science.1215025)
- Feltham H, Park K, Goulson D. In press. Field realistic doses of pesticide imidacloprid reduce bumblebee pollen foraging efficiency. *Ecotoxicology* **23**, 317–323. (doi:10.1007/s10646-014-1189-7)
- Gill RJ, Ramos-Rodriguez O, Raine NE. 2012 Combined pesticide exposure severely affects individual and colony-level traits in bees. *Nature* **491**, 105–108. (doi:10.1038/nature11585)
- Thompson H, Harrington P, Wilkins W, Pietravalle S, Sweet D, Jones A. 2013 Effects of neonicotinoid seed treatments on bumble bee colonies under field conditions. See <http://www.fera.co.uk/ccss/documents/defraBumbleBeeReportPS2371V4a.pdf>.
- Pilling E, Campbell P, Coulson M, Ruddle N, Tornier I. 2013 A four-year field program investigating long-term effects of repeated exposure of honey bee colonies to flowering crops treated with thiamethoxam. *PLoS ONE* **8**, e77193. (doi:10.1371/journal.pone.0077193)
- Cresswell JE, Desneux N, van Engelsdorp D. 2012 Dietary traces of neonicotinoid pesticides as a cause of population declines in honey bees: an evaluation by Hill’s epidemiological criteria. *Pest Manag. Sci.* **68**, 819–827. (doi:10.1002/ps.3290)
- Staveley JP, Law SA, Fairbrother A, Menzie CA. 2014 A causal analysis of observed declines in managed honey bees (*Apis mellifera*). *Hum. Ecol. Risk Assess.* **20**, 566–591. (doi:10.1080/10807039.2013.831263)

Annotated bibliography to accompany:

A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators

H. Charles J. Godfray, Tjeerd Blacquière, Linda M. Field, Rosemary S. Hails, Gillian Petrokofsky, Simon G. Potts, Nigel E. Raine, Adam J. Vanbergen & Angela R. McLean

Paragraph numbering corresponds to those in the main document; full references at end. Website URLs were accessed 7 March 2014. Any corrections and clarification will be posted at <http://www.futureoffood.ox.ac.uk/news/neonics>.

(a) Introduction and aims

- (1) References below where topics discussed in more detail. EU partial restriction is Regulation (EU) No. 485/2013 European Commission (2013). For concerns about the restriction of neonicotinoid use see Campbell (2013), Walters (2013).
- (2) Categories developed by authors, influenced by scheme in Godfray *et al.* (2013).
- (3) For an introduction to the social science and economic literature on pollinators see Kevan and Phillips (2001), Losey and Vaughan (2006), Kremen *et al.* (2007), Zhang *et al.* (2007), Gallai *et al.* (2009), Keitt (2009), Kuldna *et al.* (2009), Osgathorpe *et al.* (2011), UK National Ecosystem Assessment (2011), Noleppa and Hahn (2013), Vanbergen and Insect Pollinators Initiative (2013). Goulson (2013) reviews studies comparing yields on crops protected by neonicotinoids or through other means.

(b) Pollinators and neonicotinoid insecticides

- (4) UK data from UK National Ecosystem Assessment (2011) (updated in Vanbergen *et al.*, 2014) which also estimates economic importance of pollinator services. For summary of European crops dependent on pollinators see Table 2 of the SOM in Leonhardt *et al.* (2013) and for US see Calderone (2012). Ollerton *et al.* (2011) estimate that 87% of all plant species are animal pollinated; Klein *et al.* (2007) calculate that 87 of the 115 most important food plant species for man require animal pollination, and though these make up only 35-40% of human food by volume; Eilers *et al.* (2011) show that some essential human nutrients come largely from pollinated crop plants; Lautenbach *et al.* (2012) map global distribution of crops needing pollination and hence likely benefits of pollinators; Aizen *et al.* (2008) note proportion of pollinating crops increasing.
- (5) Benefits of insect pollination for crops shown by Klein *et al.* (2007), Hoehn *et al.* (2008), Garibaldi *et al.* (2011b), Bommarco *et al.* (2012b), Brittain *et al.* (2013a), Brittain *et al.* (2013b), Garratt *et al.* (2014a, 2014b); and in enclosures by Dag (2008). Aizen and Harder (2009) show that globally crop demand for pollination is growing faster than the supply of honeybees.
- (6) For overview see Free (1993). Importance of wild pollinators to crops discussed by Greenleaf and Kremen (2006a), Greenleaf and Kremen (2006b), Winfree *et al.* (2007), Jauker *et al.* (2012), Klein *et al.* (2012), Brittain *et al.* (2013b), Garibaldi *et al.* (2013). Pollination as an ecosystem service is explored by Losey and Vaughan (2006), Boyd and Banzhaf (2007), Kremen *et al.* (2007), Zhang *et al.* (2007), Aizen *et al.* (2008), Brosi *et al.* (2008), Aizen *et al.* (2009), Lonsdorf *et al.*

- (7) Biesmeijer *et al.* (2006), Committee on the Status of Pollinators in North America (2007), Goulson *et al.* (2008), Potts *et al.* (2010a), Potts *et al.* (2010b), Gonzalez-Varo *et al.* (2013), Lebuhn *et al.* (2013), Vanbergen and Insect Pollinators Initiative (2013) discuss the evidence for pollinator declines, and the multiple factors that may be responsible. See also Grixti *et al.* (2009), Cameron *et al.* (2011) (US bumblebees) and Bommarco *et al.* (2012a) (changes in bumblebee community composition in Sweden). Evidence of historic declines and recent slowdowns and reversals is in Carvalheiro *et al.* (2013). For butterfly and moth data see The State of Britain's Larger Moths 2013 (<http://butterfly-conservation.org/files/state-of-britains-larger-moths-2013-report.pdf>), Conrad *et al.* (2004) and Asher *et al.* (2001).
- (8) Rosenkranz *et al.* (2010) review the consequences of the introduction of *Varroa* to Europe; see also Carreck *et al.* (2010). Elevated overwintering mortality in Belgium: Nguyen *et al.* (2010); in the US: VanEngelsdorp *et al.* (2011), VanEngelsdorp *et al.* (2012). Martin *et al.* (2012) showed massive increase of a single strain of a bee virus after *Varroa* introduction to Hawaii, indicating that *Varroa* is both vectoring viruses but also driving selection for higher viral virulence. US beekeepers have compensated for overwintering losses by initiating more colonies; VanEngelsdorp *et al.* (2011). For colony collapse disorder in the US and for discussion about its contribution to colony losses as well as its presence or absence in Europe see vanEngelsdorp *et al.* (2009), Ratnieks and Carreck (2010), vanEngelsdorp *et al.* (2010), Williams *et al.* (2010), Frazier *et al.* (2011), Smith *et al.* (2014). For global patterns in honeybee numbers see Aizen and Harder (2009).
- (9) Kollmeyer (1999), Maienfisch *et al.* (2001b) & Jeschke *et al.* (2013) describe neonicotinoid discovery; mode of action discussed by Maienfisch *et al.* (2001a), Matsuda *et al.* (2001), Elbert *et al.* (2008) and uptake by plant by Sur and Stork (2003); invertebrate/vertebrate toxicity by Tomizawa *et al.* (2000), Tomizawa and Casida (2003), Tomizawa and Casida (2005), Goulson (2013) Table S1. Rise and current use of neonicotinoids described by Jeschke and Nauen (2008), Jeschke *et al.* (2011), Jeschke *et al.* (2013). For details of pesticide usage on UK crops in 2012 see <http://pusstats.fera.defra.gov.uk/>. Nauen and Denholm (2005) discuss the evolution of resistance.
- (10) See the EU pesticides database at http://ec.europa.eu/sanco_pesticides/public/?event=activesubstance.selection, thiamethoxam metabolism: Nauen *et al.* (2003).
- (11) For Europe see European Food Safety Authority (2013e) and the US Environmental Protection Agency. Kohler and Triebkorn (2013) review general issues of wildlife

ecotoxicology. The data required for pesticide active substances and associated products under Regulation 1107/2009 are outlined in <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:093:0001:0084:EN:P> DF and <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:093:0085:0152:EN:PDF>.

(c) Exposure of pollinators to neonicotinoid insecticides

- (12) Klein *et al.* (2007), Leonhardt *et al.* (2013), Vanbergen and Insect Pollinators Initiative (2013).
- (a) Velasco *et al.* (1999) estimate an average rapeseed weighs 5.9 mg and Environment Protection Agency (2003) give figures of 600g active ingredient per 100kg seeds giving 35µg per seed. Typical figures quoted for much larger maize seed (~200 mg) are 1.2mg (Environment Protection Agency (2003)).
- (b) Larson *et al.* (2013) show that if clothianidin is sprayed on turf with clover in bloom (against label recommendation) to control lawn pests, the growth of colonies of bumblebees that forage on the clover is impaired, see also Gels *et al.* (2002). Zimmer and Nauen (2011) on use of thiacloprid on oilseed rape. We are aware of on-going studies of the consequences to bees of spraying *N*-cyanoamidine neonicotinoids on flowering raspberries and oilseed rape and when published we shall note at <http://www.futureoffood.ox.ac.uk/news/neonics>.
- (13) For action of neonicotinoids see Maienfisch *et al.* (2001a), Matsuda *et al.* (2001), Elbert *et al.* (2008). Sur and Stork (2003) estimate 1-20% neonicotinoids absorbed by plant.
- (a) Blacqui re *et al.* (2012) review studies on sunflower and maize; see also Schmuck *et al.* (2001), Bonmatin *et al.* (2005), Rortais *et al.* (2005), Halm *et al.* (2006), Cutler and Scott-Dupree (2007), Cresswell (2011), Krupke *et al.* (2012), Pohorecka *et al.* (2012). Concentrations may differ on other crops and be influenced by application method, for example see work by Stoner and Eitzer (2012) on squash and Dively and Kamel (2012) on pumpkin who found relatively higher concentrations compared to seed treatments.
- (14) For studies of guttation fluid see Girolami *et al.* (2009), Reetz *et al.* (2011), Schenke *et al.* (2011), Tapparo *et al.* (2011), Hoffmann and Castle (2012), Pistorius *et al.* (2012) and EFSA (2012).
- (15) Bortolotti *et al.* (2003), Alix *et al.* (2009), Girolami *et al.* (2009), Pistorius *et al.* (2009), Thompson (2010), Tremolada *et al.* (2010), Marzaro *et al.* (2011), Girolami *et al.* (2012), Krupke *et al.* (2012), Pochi *et al.* (2012), Tapparo *et al.* (2012), Girolami *et al.* (2013).
- (a) Nuyttens *et al.* (2013) review relevant issues of formulation and sowing equipment configuration; see EFSA (2013) for regulations; air deflectors have become mandatory for certain products in the Netherlands, France, Belgium and Germany.
- (16) Goulson (2013), Table 1, lists 35 estimates from 12 published studies of soil half-lives (dissipation times) for different neonicotinoids (of which 10 were conducted in the field, the rest in the laboratory). More information has been collected by industry for regulatory purposes but is not in the public domain though summarised in regulatory documents (EFSA). For accumulation in soil see Anon (2006) discussed by Goulson (2013) in Figure 2. For leaching and contamination of groundwater see Gupta *et al.* (2008), Selim *et al.* (2010), Starner and Goh (2012) and Goulson (2013). Krupke *et al.* (2012) found weed flowers (dandelion, *Taraxacum*) visited by pollinators near treated feeds contained neonicotinoids (though whether through dust or soil is not known). Schmuck *et al.* (2001); Charvet *et al.* (2004) found untreated sunflower and maize grown on soils containing imidacloprid from earlier treatments do not have detectable residues in pollen and nectar.
- (17) Winston (1987), Seeley (1995), Brodschneider and Crailsheim (2010); studies of larval mortality discussed in para. (24).
- (18) Thompson and Hunt (1999), Hoyle *et al.* (2007), Cresswell *et al.* (2012b) for seasonal susceptibility of honeybees.
- (19) Authors' summary.
- (20) Jha and Kremen (2013) discuss bumblebee foraging distances and behaviour, see also Osborne *et al.* (1999), Darvill *et al.* (2004), Knight *et al.* (2005), Osborne *et al.* (2008), Hagen *et al.* (2011), Carvell *et al.* (2012). For honey bees see Visscher and Seeley (1982); Beekman and Ratnieks (2000) (which includes maximum 15km estimate); Steffan-Dewenter and Kuhn (2003). For solitary bees Gathmann and Tschardtke (2002), Greenleaf *et al.* (2007), Zurbuchen *et al.* (2010a), Zurbuchen *et al.* (2010b).
- (21) Authors' summary.

(d) Laboratory studies of lethal and sublethal effects of neonicotinoids

- (22) Recent review in Blacqui re *et al.* (2012), see also Decourtye and Devillers (2010) and meta-analysis by Arena and Sgolastra (2014).
- (a) The oral sensitivities reported here come from European Food Safety Authority (2013c), European Food Safety Authority (2013b), European Food Safety Authority (2013a) though we note that not all the studies upon which these assessments are based are in the public domain. The meta-analysis is Cresswell (2011) which included 13 peer-reviewed and one non-peer-reviewed studies.
- (b) The contact sensitivities reported here also come from European Food Safety Authority (2013c), European Food Safety Authority (2013b), European Food Safety Authority (2013a). A study by Iwasa *et al.* (2004) compared honeybee acute contact LD₅₀s for imidacloprid (18 ng bee⁻¹), clothianidin (22 ng bee⁻¹), thiamethoxam (30 ng bee⁻¹) and, for neonicotinoids from the cyano-substituted group, acetamiprid (7 µg bee⁻¹), thiacloprid (15 µg bee⁻¹). These figures differ somewhat from those in EFSA though are in the same general range.
- (c) A meta-analysis by Arena and Sgolastra (2014) compares the differential sensitivity of bee species to different classes of insecticides and includes data on neonicotinoid exposure to nine bee species including the honeybee; the latter comes near the middle of the range of sensitivities with studies on bumblebees consistently reporting higher sensitivities (a similar pattern was found with other insecticide classes). In a direct comparison of acute contact toxicity across non-*Apis* bee species, Scott-Dupree *et al.* (2009) showed variation of up to 19 times for imidacloprid (up to 4 times for clothianidin); see also Stark *et al.* (1995), Biddinger *et al.* (2013).
- (d) For general honeybee economics see Winston (1987), Seeley (1995), Brodschneider and Crailsheim (2010); Rortais

- et al.* (2005) observed the (relatively small) colony of 10,000. We have calculated these figures and those in the next two subparagraphs to provide a general guide to the relative concentrations involved. Since most of the nectar and pollen in the foraging honeybee stomach is not metabolized (Fournier *et al.* 2014) the insect is only exposed to a small fraction of any insecticide it contains, almost all of which will be carried to the hive.
- (e) Winston (1987), Seeley (1995), Brodschneider and Crailsheim (2010) and authors' calculations.
- (f) Winston (1987), Seeley (1995), Brodschneider and Crailsheim (2010), Decourtye *et al.* (2010) and authors' conclusions.
- (23) Recent review in Blacquière *et al.* (2012).
- (a) See Decourtye *et al.* (2001), Suchail *et al.* (2001), Decourtye *et al.* (2003), and European Food Safety Authority (2013c), European Food Safety Authority (2013b), European Food Safety Authority (2013a) for studies of chronic lethality in the laboratory.
- (b) Authors' conclusions.
- (24) Recent review in Blacquière *et al.* (2012) and see Decourtye and Devillers (2010) for general issues of sublethal effects on non-target insects Desneux *et al.* (2007); see also European Food Safety Authority (2012), European Food Safety Authority (2013c), European Food Safety Authority (2013b), European Food Safety Authority (2013a). See Belzunces *et al.* (2012) and Boily *et al.* (2013) for imidacloprid stimulation of acetylcholinesterase activity; for neuronal inactivation Palmer *et al.* (2013), on learning performance and behaviour, Decourtye *et al.* (2004a), Decourtye *et al.* (2005a), Decourtye *et al.* (2004b), El Hassani *et al.* (2008), Yang *et al.* (2008), Aliouane *et al.* (2009), Matsumoto (2013a), Matsumoto (2013b). The proboscis extension response is discussed by Bitterman *et al.* (1983) and Giurfa and Sandoz (2012).
- (a) For sublethal effects on bumble bees see Tasei *et al.* (2000), Tasei *et al.* (2001), Morandin and Winston (2003), Franklin *et al.* (2004), Mommaerts *et al.* (2010), Cresswell *et al.* (2012b), Laycock *et al.* (2012), Bryden *et al.* (2013), Elston *et al.* (2013), Fauser-Misslin *et al.* (2013) (the last two of which found no effects at estimated field concentrations), Laycock *et al.* (2014); and on solitary bees Abbott *et al.* (2008), Thompson and Hunt (1999), Scott-Dupree *et al.* (2009), Brittain and Potts (2011), Biddinger *et al.* (2013), Sandrock *et al.* (2014) compare risk factors for different bee species.
- (b) For debate see Suchail *et al.* (2004a), Suchail *et al.* (2004b), Tennekes (2010), Maus and Nauen (2011), Tennekes and Sánchez-Bayo (2012); figures from Cresswell *et al.* (2013). Hawthorne and Dively (2011) describe how multi-drug resistance transporters actively remove neonicotinoids from the cytoplasm so reducing their toxicity.
- (25) Recent review in Blacquière *et al.* (2012).
- (a) Studies on honeybees: Decourtye *et al.* (2005b) (imidacloprid added at 5 ng g^{-1} ; at the high end of what observed in field prior to any processing of the food by the adult), Yang *et al.* (2012) (observed effects when the doses given to larvae were near the LD_{50} reported from larger adults); for effects on cellular physiology see Gregorc and Ellis (2011). Wu *et al.* (2011) studied larvae reared in old contaminated combs; the combs contained many different chemical residues, the most abundant of which were acaricides (used by the beekeepers against *Varroa*); three types of neonicotinoid were present at low levels, but in only one out of the 13 replicates. Retarded larval development and reduced longevity of emerging bees was observed. This study has been cited as evidence of the effects of neonicotinoids on larval honeybees, in the authors' view incorrectly.
- (b) Bumblebees: Elston *et al.* (2013) found thiamethoxam reduced worker consumption of sugar water solution and their production of wax cells at both 1 and 10 ng g^{-1} , and at the higher dosage workers laid fewer eggs and no (male) larvae were produced in micro-colonies; see also Laycock *et al.* (2012), Laycock and Cresswell (2013). Fauser-Misslin *et al.* (2013) studied chronic dietary exposure of whole bumblebee colonies. Solitary bees: Abbott *et al.* (2008), Sandrock *et al.* (2014).
- (26) European Food Safety Authority (2012), Thompson (2012).
- (a) Abrol (2007), Alaux *et al.* (2010), Vidau *et al.* (2011) showed honeybees infected by the microsporidian *Nosema ceranae* have lower neonicotinoid LD_{50} , and Pettis *et al.* (2012) that prior exposure to neonicotinoids in the hive led to a higher *N. ceranae* spore load in artificially infected bees in the lab (though no effect in the hive). Pettis *et al.* (2013) found that honeybees fed pollen contaminated by neonicotinoids were significantly less likely to become infected by *Nosema* (fungicides used in hives had the opposite effect). Fauser-Misslin *et al.* (2013) found chronic exposure of bumblebees to thiamethoxam and clothianidin increased the negative effects of infection by the trypanosome *Crithidia bombi*. Di Prisco *et al.* (2013) showed neonicotinoids reduced antiviral defences via effects on $\text{NF-}\kappa\text{B}$ signalling pathways in honeybees. See also European Food Safety Authority (2012, p. 108). Doublet *et al.* (2014) find that in the laboratory sublethal doses of the *N*-cyanoamidine group neonicotinoid thiacloprid can increase the detrimental effects on honeybees of microsporidian and viral pathogens.
- (b) Review in Blacquière *et al.* (2012); see also Iwasa *et al.* (2004), European Food Safety Authority (2012, p. 113). Hawthorne and Dively (2011) show that honeybees from colonies treated with the in-hive antibiotic oxytetracycline (not allowed in the EU) may be sensitised to acaricides and insecticides (including neonicotinoids); oxytetracycline inhibits multi-drug resistance transporters that clear cells of toxins. Williamson *et al.* (2013) found little additive acute effects of imidacloprid and the organophosphate acaricide coumaphos (a modest improvement in memory) while chronic multiple exposure caused significant impairment Williamson and Wright (2013).
- (c) For increased sensitivity to pesticides (lower LD_{50}) when honeybees are stressed, for example when raised on protein deficient diets, see Von der Ohe and Janke (2009), Wehling *et al.* (2009).
- (27) This paragraph chiefly authors' judgment.
- (a) Walters (2013) comments on lack of data on full range of neonicotinoids.
- (b) See Visser and Blacquière (2010) for discussion and preliminary experiments.
- (c) Authors' comment.
- (d) For issue of insecticide clearance see Para. (24)(b); for anti-feedant behaviour see Suchail *et al.* (2004a), Department for Environment (2007), Cresswell *et al.* (2013), and in bumblebees Laycock *et al.* (2012), Elston *et al.* (2013).
- (e) Authors' comment.
- (28) Authors' summary.

(e) Neonicotinoid residues observed in pollinators in the field

- (29) Pilling *et al.* (2013).
- (30) Comprehensive review in Blacquière *et al.* (2012); the Belgium studies are by Pirard *et al.* (2007) and Nguyen *et al.* (2009); the US study by Mullin *et al.* (2010); and the French study by Chauzat *et al.* (2006), Chauzat *et al.* (2009), Chauzat *et al.* (2011); see Wiest *et al.* (2011) for recent technical advances.
- (31) Comprehensive review in Blacquière *et al.* (2012); French studies see para. (30); Spanish study is by Bernal *et al.* (2010), the German by Genersch *et al.* (2010). Škerl *et al.* (2009) found pollen collected from hives near recently sprayed apple trees had high levels of thiacloprid (a neonicotinoid less harmful to bees) after apple tree spraying, a compound found relatively frequently in the surveys of Škerl *et al.* (2009), Genersch *et al.* (2010). As part of an experiment Thompson *et al.* (2013) placed bumblebee colonies near treated fields and recorded residues in nests (though often not of the substance applied to the adjacent field). Mullin *et al.* (2010) found neonicotinoid residues in less than 2% of wax samples ($n = 208$) and less than 5% of pollen samples in North America ($n = 350$) though relatively high levels of agricultural chemicals were observed. See also Bonmatin *et al.* (2003), Brittain and Potts (2011), Krupke *et al.* (2012).
- (32) Authors' summary.

(f) Experiments conducted in the field

- (33) For earlier studies see Schmuck *et al.* (2001), Faucon *et al.* (2005), Cutler and Scott-Dupree (2007); see also Cresswell (2011) for a meta-analysis of 13 laboratory and semi-field studies of imidacloprid that concluded field-realistic doses did not cause honey bee death but could reduce colony performance by 6-20%.
- (34) Schneider *et al.* (2012).
- (a) Authors' comment.
- (35) Henry *et al.* (2012a) gave bees an acute dose of thiamethoxam and released them 1km away from the colony carrying Radio Frequency Identification (RFID) tags that were read at the hive entrance (the use of RFID tag technology to monitor bee foraging activity is described by Streit *et al.* (2003) and Molet *et al.* (2008)).
- (a) As foragers typically make multiple trips from the hive to collect food each day, reported homing failure rates of 10.2-31.6% per trip would rapidly reduce the forager workforce. Henry *et al.* (2012a) measured the dose given to bees but Food and Environment Research Agency (2013) questioned the realism of exposing the bees to the equivalent of a daily dose in one meal.
- (b) Assumptions in original model in Henry *et al.* (2012a) were debated by Cresswell and Thompson (2012) and Henry *et al.* (2012b).
- (36) Whitehorn *et al.* (2012) allowed bumblebees to feed on pollen and sugar water containing 6 and 0.7 ng g⁻¹ imidacloprid respectively (low dose) and double this (high dose) and measured colony mass and queen production in the field. The insecticide affected both measures of performance with little difference between low and high doses. Feltham *et al.* (2014) used RFID tag technology to assess foraging success of dosed and non-dosed bees. Whitehorn *et al.* (2012) and Gill *et al.* (2012) provided food containing neonicotinoids at rates informed by those observed in pollen and nectar though Food

and Environment Research Agency (2013) has argued that in practice these are higher than would occur in the field.

- (a) Authors' comment. There was considerable variability in queen production among colonies, with many colonies producing no queens; the replication was sufficient to detect differences despite this variability that might reflect that the colonies were stressed. Imidacloprid can cause anti-feedant behaviour affecting energy intake but typically at higher concentrations than used here.
- (37) Gill *et al.* (2012) placed bumblebees in nests with an antechamber where they had access to sugar water and monitored colony size and foraging (by RFID tagging workers and measuring pollen loads) over a 28-day period. The experiment had four treatments: sugar water containing imidacloprid, antechamber with surface sprayed with pyrethroid (λ -cyhalothrin); the two combined, and a control.
- (a) Authors' comment. The bumblebees walked over filter paper sprayed with pyrethroid at a concentration that if extrapolated to the whole environment would be equivalent to an application rate of about 40g ha⁻¹. This is considerably higher than the 7.5 g ha⁻¹ EU maximum for oilseed rape though higher concentrations (up to 30 g ha⁻¹) are allowed on some field crops and up to 125g ha⁻¹ in grapes, cane fruit and hops (http://ec.europa.eu/food/fs/sc/scp/out01_ppp_en.html). Exposure by walking over filter paper (through tarsal contact) may be less than an animal encounters in the field, for example through spray drift.
- (38) Thompson *et al.* (2013) is a non-peer reviewed report published on the web in March 2013 with the version currently (March 2014) available containing a supplementary explanatory note added in June 2013. The study involved placing ten bumblebee nests adjacent to three different fields (hence there was no replication of treatments) though these could not be carried out at the same time (introducing the possibility of systematic error).
- (a) Authors' comment. The design of the study was criticised by several commentators and in a detailed review by European Food Safety Authority (2013d). The addendum added on 14th June 2013 states "This study was not designed as a definitive statistically robust study" but to look quickly for "major effects". Reanalysis of some of the data reported to the Advisory Committee on Pesticides has questioned the lack of a colony-level relationship between neonicotinoid residues and colony performance (<http://www.pesticides.gov.uk/guidance/industries/pesticides/advisory-groups/acp/ACP-News/ACP-359-29-January-2013-Detailed-Record-of-Discussion>) but this too has not been peer-reviewed.
- (39) Pilling *et al.* (2013). Neonicotinoid treatments were typical for France where the experiments were carried out. The short exposure time in the first three years coincided with oilseed rape flowering. It is possible that bees from a single hive exploit asynchronously flowering fields and so experience longer exposure.
- (40) Authors' summary.
- ### (g) Consequences of neonicotinoid use
- (41) The observed abundance of a species is influenced by factors that have the same effect on death rates (or birth rates) irrespective of population size (density-independent factors, a

- typical example would be the effects of bad weather) and those whose magnitudes depend on population size (density-dependent factors; a typical example would be starvation due to competition for food). The reduction in pollinator population size caused by insecticides (a density-independent factor) may be less if they result in reduced density-dependent mortality through factors acting subsequently. Such effects are likely to be common in pollinator population dynamics though very difficult to predict in the absence of detailed study. "Allee effects" are density-dependent factors that increase in severity at low densities (a typical example would be reduced fecundity due to failure to mate at low population densities). If insecticide application reduced population density to a level where an Allee effect occurred then population extinction could occur. We do not know of examples of Allee effects from pollinator populations. Kohler and Triebkorn (2013) review population consequences of pesticide use. For population dynamic models of honeybee colonies see Khoury *et al.* (2011), Becher *et al.* (2013), Khoury *et al.* (2013) and bumblebee colonies see Bryden *et al.* (2013).
- (42) Garratt *et al.* (2014a), Garratt *et al.* (2014b) report deficits in UK apple pollination and Breeze *et al.* (2014) suggest a lack of availability of honeybees may affect pollination of oilseed rape in Europe. Bommarco *et al.* (2012b) provide evidence of a decrease in clover seed yield through time linked to pollinator loss. Garibaldi *et al.* (2011b) show decreases in crop yield and stability with distance from native vegetation. Aizen *et al.* (2008) for global comparison of yield growth of insect- and wind-pollinated plants (no difference suggesting no pollination deficit). Dicks *et al.* (2013) survey main evidence needs for pollinator conservation. Ashman *et al.* (2004) review pollination deficits in wild flowers and Holzschuh *et al.* (2011) shows mass-flowering crops can attract pollinators away from wild species. Our assessment of available evidence.
- (43) See references in Para. (7); importance of habitat loss: Carvell *et al.* (2006), Ricketts *et al.* (2008), Brown and Paxton (2009), Goulson *et al.* (2010), Carvalheiro *et al.* (2011), Kennedy *et al.* (2013), though see Gonzalez-Varo *et al.* (2013) and Vanbergen (2013) for importance of considering multiple effects. Authors' assessment of available evidence.
- (44) See references in Para. (8) . Authors' assessment of available evidence.
- (a) Cresswell *et al.* (2012a); see also Maxim and van der Sluijs (2010).
- (b) Staveley *et al.* (2014).
- (45) Authors' assessment of available evidence.
- (46) Authors' summary.

References

- Abbott V.A., Nadeau J.L., Higo H.A., Winston M.L. 2008 Lethal and sublethal effects of imidacloprid on *Osmia lignaria* and clothianidin on *Megachile rotundata* (Hymenoptera: Megachilidae). *Journal of Economic Entomology* **101**, 784-796. (doi:10.1603/0022-0493(2008)101[784:laseoi]2.0.co;2).
- Abrol D.P. 2007 Honeybees and rapeseed: a pollinator-plant interaction *Advances in Botanical Research* **45**, 337-367 (doi: 10.1016/S0065-2296(07)45012-1).
- Aizen M.A., Garibaldi L.A., Cunningham S.A., Klein A.M. 2009 How much does agriculture depend on pollinators? Lessons from long-term trends in crop production. *Annals of Botany* **103**, 1579-1588. (doi: 10.1093/Aob/Mcp076).
- Aizen M.A., Garibaldi L.A., Cunningham S.A., Klein A.M. 2008 Long-term global trends in crop yield and production reveal no current pollination shortage but increasing pollinator dependency. *Current Biology* **18**, 1572-1575. (doi:10.1016/j.cub.2008.08.066).
- Aizen M.A., Harder L.D. 2009 The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology* **19**, 915-918. (doi:10.1016/J.Cub.2009.03.071).
- Alaux C., Brunet J.L., Dussaubat C., Mondet F., Tchamitchan S., Cousin M., Brillard J., Baldy A., Belzunces L.P., Le Conte Y. 2010 Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environmental Microbiology* **12**, 774-782. (doi:10.1111/j.1462-2920.2009.02123.x10.1051/apido/2009039).
- Aliouane Y., el Hassani A.K., Gary V., Armengaud C., Lambin M., Gauthier M. 2009 Subchronic exposure of honeybees to sublethal doses of pesticides: effects on behavior. *Environmental Toxicology and Chemistry* **28**, 113-122. (doi:10.1897/08-110.1).
- Alix A., Vergnet C., Mercier T. 2009 Risks to bees from dusts emitted at sowing of coated seeds: concerns, risk assessment and risk management. *Julius Kuhn Archiv* **423**, 131-132.
- Anon. 2006 Draft assessment Report (DAR) – public version – Initial risk assessment provided by the rapporteur Member State Germany for the existing active substance imidacloprid. Volume 3, Annex B, B. 8, February 2006, European Community.
- Arena M., Sgolastra F. 2014 A meta-analysis comparing the sensitivity of bees to pesticides. *Ecotoxicology*. (January 2014) (doi:10.1007/s10646-10014-11190-10641).
- Asher J., Warren M.S., Fox R., Harding P., Jeffcoate G., Jeffcoate S. 2001 *The Millennium Atlas of Butterflies in Britain & Ireland*. Oxford, Oxford University Press.
- Ashman T.L., Knight T.M., Steets J.A., Amarasekare P., Burd M., Campbell D.R., Dudash M.R., Johnston M.O., Mazer S.J., Mitchell R.J., Morgan, M.T., Wilson, W.G. 2004 Pollen limitation of plant reproduction: ecological and evolutionary causes and consequences. *Ecology* **85**, 2408-2421. (doi: 10.1890/03-8024).
- Becher M.A., Osborne J.L., Thorbek P., Kennedy P.J., Grimm V. 2013 Towards a systems approach for understanding honeybee decline: a stocktaking and synthesis of existing models. *Journal of Applied Ecology* **50**, 868-880. (doi:10.1111/1365-2664.12112).
- Beekman M., Ratnieks F.L.W. 2000 Long-range foraging by the honey-bee, *Apis mellifera* L. *Functional Ecology* **14**, 490-496. (doi: 10.1046/j.1365-2435.2000.00443.x).

15. Belzunces L.P., Tchamitchian S., Brunet J.L. 2012 Neural effects of insecticides in the honey bee. *Apidologie* **43**, 348-370. (doi:10.1007/s13592-012-0134-0).
16. Bernal J., Garrido-Bailón E., Del Nozal M.J., González-Porto A.V., Martín-Hernández R., Diego J.C., Jiménez J.J., Bernal J.L., Higes M. 2010 Overview of pesticide residues in stored pollen and their potential effect on bee colony (*Apis mellifera*) losses in Spain. *Journal of Economic Entomology* **103**, 1964-1971. (doi:10.1603/EC10235).
17. Biddinger D.J., Robertson J.L., Mullin C., Frazier J., Ashcraft S.A., Rajotte E.G., Joshi N.K., Vaughn M. 2013 Comparative toxicities and synergism of apple orchard pesticides to *Apis mellifera* (L.) and *Osmia cornifrons* (Radoszkowski). *PLoS ONE* **8**(9): e72587. (doi:10.1371/journal.pone.0072587).
18. Biesmeijer J.C., Roberts S.P.M., Reemer M., Ohlemüller R., Edwards M., Peeters T., Schaffers A.P., Potts S.G., Kleukers R., Thomas C.D., Settele J., Kunin, W. E. 2006 Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* **313**, 351-354. (doi:10.1126/science.1127863).
19. Bitterman M.E., Menzel R., Fietz A., Schafer S. 1983 Classical conditioning of proboscis extension in honeybees (*Apis mellifera*). *Journal of Comparative Psychology* **97**, 107-119. (doi: 10.1037/0735-7036.97.2.107).
20. Blacquièrre T., Smaghe G., Van Gestel C.A.M., Mommaerts V. 2012 Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology* **21**, 973-992. (doi:10.1007/s10646-012-0863-x).
21. Boily M., Sarrasin B., Deblois C., Aras P., Chagnon M. 2013 Acetylcholinesterase in honey bees (*Apis mellifera*) exposed to neonicotinoids, atrazine and glyphosate: laboratory and field experiments. *Environmental Science and Pollution Research International* **20**, 5603-5614. (doi:10.1007/s11356-013-1568-2).
22. Bommarco R., Lundin O., Smith H.G., Rundlöf M. 2012a Drastic historic shifts in bumble-bee community composition in Sweden. *Proceedings of the Royal Society B: Biological Sciences* **279**, 309-315. (doi:10.1098/rspb.2011.0647).
23. Bommarco R., Marini L., Vaissière B.E. 2012b Insect pollination enhances seed yield, quality, and market value in oilseed rape. *Oecologia* **169**, 1025-1032. (doi:10.1007/s00442-012-2271-6).
24. Bonmatin J.M., Marchand P.A., Charvet R., Moineau I., Bengsch E.R., Colin M.E. 2005 Quantification of imidacloprid uptake in maize crops. *Journal of Agricultural and Food Chemistry* **53**, 5336-5341. (doi:10.1021/jf0479362).
25. Bonmatin J.M., Moineau I., Charvet R., Fleche C., Colin M.E., Bengsch E.R. 2003 A LC/APCI-MS/MS method for analysis of imidacloprid in soils, in plants, and in pollens. *Analytical Chemistry* **75**, 2027-2033. (doi:10.1021/ac020600b).
26. Bortolotti L., Montanari R., Marcelino J., Medrzycki P., Maini S., Porrini C. 2003 Effects of sub-lethal imidacloprid doses on the homing rate and foraging activity of honey bees. *Bulletin of Insectology* **56**, 63-67.
27. Boyd J., Banzhaf S. 2007 What are ecosystem services? The need for standardized environmental accounting units. *Ecological Economics* **63**, 616-626. (doi:10.1016/j.ecolecon.2007.01.002).
28. Breeze TD, Vaissière B.E, Bommarco R, Petanidou T, Seraphides N, Kozák, L., Scheper, J., Biesmeijer, J.C., Kleijn, D., Gyldenkerne, S., Moretti, M., Holzschuh, A., Steffan-Dewenter, I., Stout, J.C., Pärtel, M., M., Zobel, M., Potts, S.G. 2014 Agricultural policies exacerbate honeybee pollination service supply- demand mismatches across europe. *PLoS ONE* **9**(1): e82996. (doi:10.1371/journal.pone.0082996).
29. Brittain C., Kremen C., Klein A.M. 2013a Biodiversity buffers pollination from changes in environmental conditions. *Global Change Biology* **19**, 540-547. (doi:10.1111/gcb.12043).
30. Brittain C., Potts S.G. 2011 The potential impacts of insecticides on the life-history traits of bees and the consequences for pollination. *Basic and Applied Ecology* **12**, 321-331. (doi:10.1016/j.baae.2010.12.004).
31. Brittain C., Williams N., Kremen C., Klein A.M. 2013b Synergistic effects of non-*Apis* bees and honey bees for pollination services. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122767. (doi:10.1098/rspb.2012.2767).
32. Brodschneider R., Crailsheim K. 2010 Nutrition and health in honey bees. *Apidologie* **41**, 278-294. (doi:10.1051/apido/2010012).
33. Brosi B.J., Armsworth P.R., Daily G.C. 2008 Optimal design of agricultural landscapes for pollination services. *Conservation Letters* **1**, 27-36.
34. Brown M.J.F., Paxton R.J. 2009 The conservation of bees: a global perspective. *Apidologie* **40**, 410-416. (doi:10.1051/apido/2009019).
35. Bryden J., Gill R.J., Mitton R.A.A., Raine N.E., Jansen V.A.A. 2013 Chronic sublethal stress causes bee colony failure. *Ecology Letters* **16**, 1463-1469. (doi:10.1111/ele.12188).
36. Calderone N.W. 2012 Insect pollinated crops, insect pollinators and US agriculture: trend analysis of aggregate data for the period 1992-2009. *PLoS ONE* **7**(5): e37235. (doi:10.1371/journal.pone.0037235).
37. Cameron S.A., Lozier J.D., Strange J.P., Koch J.B., Cordes N., Solter L.F., Griswold T.L. 2011 Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 662-667. (doi:10.1073/pnas.1014743108).
38. Campbell P.J. 2013 Declining European bee health: banning the neonicotinoids is not the answer. *Outlooks on Pest Management* **24**, 52-57. (doi:10.1564/v24_apr_02).
39. Carreck N.L. 2008 Are honey bees (*Apis mellifera* L.) native to the British Isles? *Journal of Apicultural Research* **47**, 318-322. (doi:10.3896/ibra.1.47.4.15).
40. Carreck N.L., Bell B.V., Martin S.J. 2010 Honey bee colony collapse and changes in viral prevalence associated with *Varroa destructor*. *Journal of Apicultural Research* **49**, 93-94. (doi:10.3896/ibra.1.49.1.13).
41. Carvalheiro L.G., Kunin W.E., Keil P., Aguirre-Gutierrez J., Ellis W.N., Fox R., Groom Q., Hennekens S., Van Landuyt W., Maes D., Van de Meutter, F., Michez, D., Rasmont, P., Ode, B., Potts, S.G., Reemer, M., Roberts, S.P., Schaminee, J., WallisDeVries, M.F., Biesmeijer, J.C. 2013 Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology Letters* **16**, 870-878. (doi:10.1111/ele.12121).
42. Carvalheiro L.G., Veldtman R., Shenkute A.G., Tesfay G.B., Pirk C.W., Donaldson J.S., Nicolson S.W. 2011 Natural and within-farmland biodiversity enhances crop productivity. *Ecology Letters* **14**, 251-259. (doi:10.1111/j.1461-0248.2010.01579.x).
43. Carvell C., Jordan W.C., Bourke A.F.G., Pickles R., Redhead J.W., Heard M.S. 2012 Molecular and spatial analyses reveal links between colony-specific foraging distance and landscape-level resource availability in two bumblebee species. *Oikos* **121**, 734-742. (doi:10.1111/j.1600-0706.2011.19832.x).

44. Carvell C., Roy D.B., Smart S.M., Pywell R.F., Preston C.D., Goulson D. 2006 Declines in forage availability for bumblebees at a national scale. *Biological Conservation* **132**, 481-489. (doi:10.1016/j.biocon.2006.05.008).
45. Charvet R., Katouzian-Safadi M., Colin M.E., Marchand P.A., Bonmatin J.M. 2004 Systemic insecticides: new risk for pollinator insects. *Insecticides systémiques: de nouveaux risques pour les insectes pollinisateurs* **62**, 29-35.
46. Chauzat M., Martel A., Cougoule N., Porta P., Lachaize J., Zeggane S., Aubert M., Carpentier P., Faucon J. 2011 An assessment of honeybee colony matrices, *Apis mellifera* (Hymenoptera: Apidae) to monitor pesticide presence in continental France. *Environmental Toxicology and Chemistry* **30**, 103-111. (doi:10.1002/etc.361).
47. Chauzat M.P., Carpentier P., Martel A.C., Bougeard S., Cougoule N., Porta P., Lachaize J., Madec F., Aubert M., Faucon J.P. 2009 Influence of pesticide residues on honey bee (Hymenoptera: Apidae) colony health in France. *Environmental Entomology* **38**, 514-523. (doi:10.1603/022.038.0302).
48. Chauzat M.P., Faucon J.P., Martel A.C., Lachaize J., Cougoule N., Aubert M. 2006 A survey of pesticide residues in pollen loads collected by honey bees in France. *Journal of Economic Entomology* **99**, 253-262.
49. Conrad K.F., Woiwod I.P., Parsons M., Fox R., Warren M.S. 2004 Long-term population trends in widespread British moths. *Journal of Insect Conservation* **8**, 119-136.
50. Cresswell J.E. 2011 A meta-analysis of experiments testing the effects of a neonicotinoid insecticide (imidacloprid) on honey bees. *Ecotoxicology* **20**, 149-157. (doi:10.1007/s10646-010-0566-0).
51. Cresswell J.E., Desneux N., vanEngelsdorp D. 2012a Dietary traces of neonicotinoid pesticides as a cause of population declines in honey bees: an evaluation by Hill's epidemiological criteria. *Pest Management Science* **68**, 819-827. (doi:10.1002/ps.3290).
52. Cresswell J.E., Page C.J., Uygun M.B., Holmbergh M., Li Y., Wheeler J.G., Laycock I., Pook C.J., de Ibarra N.H., Smirnov N., Tyler, C.R. 2012b Differential sensitivity of honey bees and bumble bees to a dietary insecticide (imidacloprid). *Zoology* **115**, 365-371. (doi:10.1016/j.zool.2012.05.003).
53. Cresswell J.E., Robert F.X., Florance H., Smirnov N. 2013 Clearance of ingested neonicotinoid pesticide (imidacloprid) in honey bees (*Apis mellifera*) and bumblebees (*Bombus terrestris*). *Pest Management Science* (doi:10.1002/ps.3569).
54. Cresswell J.E., Thompson H.M. 2012 Comment on "a common pesticide decreases foraging success and survival in honey bees". *Science* **337**, 1453-b. (doi:10.1126/science.1224618).
55. Cutler G.C., Scott-Dupree C.D. 2007 Exposure to clothianidin seed-treated canola has no long-term impact on honey bees. *Journal of Economic Entomology* **100**, 765-772. (doi:10.1603/0022-0493(2007)100[765:ETCSCH]2.0.CO;2).
56. Dag A. 2008 Bee pollination of crop plants under environmental conditions unique to enclosures. *Journal of Apicultural Research* **47**, 162-165.
57. Darvill B., Knight M.E., Goulson D. 2004 Use of genetic markers to quantify bumblebee foraging range and nest density. *Oikos* **107**, 471-478.
58. Decourtye A., Armengaud C., Renou M., Devillers J., Cluzeau S., Gauthier M., Pham-Delegue M.H. 2004a Imidacloprid impairs memory and brain metabolism in the honeybee (*Apis mellifera* L.). *Pesticide Biochemistry and Physiology* **78**, 83-92. (doi:10.1016/J.Pestbp.2003.10.001).
59. Decourtye A., Devillers J. 2010 Ecotoxicity of neonicotinoid insecticides to bees. In *Advances in Experimental Medicine and Biology*, Vol 683 (ed. Thany S.H.), pp. 85-95. ISBN 978-1-4419-6445-8.
60. Decourtye A., Devillers J., Cluzeau S., Charreton M., Pham-Delegue M.H. 2004b Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. *Ecotoxicology and Environmental Safety* **57**, 410-419. (doi:10.1016/J.Ecoenv.2003.08.001).
61. Decourtye A., Devillers J., Genecque E., Le Menach K., Budzinski H., Cluzeau S., Pham-Delègue M.H. 2005a Comparative sublethal toxicity of nine pesticides on olfactory learning performances of the honeybee *Apis mellifera*. *Archives of Environmental Contamination and Toxicology* **48**, 242-250. (doi:10.1007/s00244-003-0262-7).
62. Decourtye A., Lacassie E., Pham-Delègue M.H. 2003 Learning performances of honeybees (*Apis mellifera* L) are differentially affected by imidacloprid according to the season. *Pest Management Science* **59**, 269-278. (doi:10.1002/ps.631).
63. Decourtye A., Le Metayer M., Pottiau H., Tisseur M., Odoux J.F., Pham-Delegue M.H. 2001 Impairment of olfactory learning performances in the honey bee after long term ingestion of imidacloprid. *Hazards of Pesticides to Bees; INRA Colloquium*, 113-117.
64. Decourtye A., Mader E., Desneux N. 2010 Landscape enhancement of floral resources for honey bees in agro-ecosystems. *Apidologie* **41**, 264-277. (doi:10.1051/Apido/2010024).
65. Decourtye A., Tisseur M., Taséi J.N., Pham-Delègue, M.H. 2005b Toxicité et risques liés à l'emploi de pesticides chez les pollinisateurs: cas de l'abeille domestique. In *Enjeux phytosanitaires pour l'agriculture et l'environnement au XXI^e siècle* (ed. Regnault-Roger C.), pp. 283-299. Paris, Tec et Doc Lavoisier.
66. Department for Environment Food and Rural Affairs 2007 Assessment of the Risk Posed to Honeybees by Systemic Pesticides (Project no. PS2322). London, Defra.
67. Derecka, K., Blythe, M.J., Malla, S., Genereux, D.P., Guffanti, A., Pavan, P., Moles, A., Snart, C., Ryder, T., Ortori, C.A., Barrett, D.A., Schuster, E., Stöger, R. 2013 Transient exposure to low levels of insecticide affects metabolic networks of honeybee larvae. *PLoS ONE* **8**(7): e68191. (doi:10.1371/journal.pone.0068191).
68. Desneux N., Decourtye A., Delpuech J.M. 2007 The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology* **52**, 81-106. (doi:10.1146/annurev.ento.52.110405.091440).
69. Di Prisco G., Cavaliere V., Annoscia D., Varricchio P., Caprio E., Nazzi F., Gargiulo G., Pennacchio F. 2013 Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. *Proceedings of the National Academy of Sciences of the United States of America* (doi:10.1073/pnas.1314923110).
70. Dicks L.V., Abrahams A., Atkinson J., Biesmeijer J., Bourn N., Brown C., Brown M.J.F., Carvell C., Connolly C., Cresswell J.E., Croft, P., Darvill, B., De Zylva, P., Effingham, P., Fountain, M., Goggin, A., Harding, D., Harding, T., Hartfield, C., Heard, M.S., Heathcote, R., Heaver, D., Holland, J., Howe, M., Hughes, B., Huxley, T., Kunin, W. E., Little, J., Mason, C., Memmott, J., Osborne, J., Pankhurst, T., Paxton, R.J., Pocock, M.J.O., Potts,

- S.G., Power, E.F., Raine, N.E., Ranelagh, E., Roberts, S., Saunders, R., Smith, K., Smith, R.M., Sutton, P., Tilley, L.A.N., Tinsley, A., Tonhasca, A., Vanbergen, A.J., Webster, S., Wilson, A., Sutherland, W. J. 2013 Identifying key knowledge needs for evidence-based conservation of wild insect pollinators: a collaborative cross-sectoral exercise. *Insect Conservation and Diversity* **6**, 435-446. (doi:10.1111/j.1752-4598.2012.00221.x).
71. Dively G.P., Kamel A. 2012 Insecticide residues in pollen and nectar of a cucurbit crop and their potential exposure to pollinators. *Journal of Agricultural and Food Chemistry* **60**, 4449-4456. (doi:10.1021/jf205393x).
72. Doublet, V., Labruassias, M., de Miranda, J.R., Moritz, R.F.A., Paxton, R.J. 2014. Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. *Environmental Microbiology* doi:10.1111/1462-2920.12426.
73. Eilers E.J., Kremen C., Greenleaf S.S., Garber A.K., Klein A.M. 2011 Contribution of pollinator-mediated crops to nutrients in the human food supply. *PLoS ONE* **6**(6): e21363. (doi:10.1371/journal.pone.0021363).
74. El Hassani A.K., Dacher M., Gary V., Lambin M., Gauthier M., Armengaud C. 2008 Effects of sublethal doses of acetamiprid and thiamethoxam on the behavior of the honeybee (*Apis mellifera*). *Archives of Environmental Contamination and Toxicology* **54**, 653-661. (doi:10.1007/s00244-007-9071-8).
75. Elbert A., Haas M., Springer B., Thielert W., Nauen R. 2008 Applied aspects of neonicotinoid uses in crop protection. *Pest Management Science* **64**, 1099-1105. (doi:10.1002/ps.1616).
76. Elston C., Thompson H.M., Walters K.F.A. 2013 Sub-lethal effects of thiamethoxam, a neonicotinoid pesticide, and propiconazole, a DMI fungicide, on colony initiation in bumblebee (*Bombus terrestris*) micro-colonies. *Apidologie* **44**, 563-574. (doi:10.1007/s13592-013-0206-9).
77. Environment Protection Agency. 2003 *Pesticide Fact Sheet: Clothianidin*. Environment Protection Agency, Washington, USA.
78. European Commission. 2013 Commission implementing regulation (EU) No 485/2013 of 24 May 2013 amending Implementing Regulation (EU) No 540/2011, as regards the conditions of approval of the active substances clothianidin, thiamethoxam and imidacloprid, and prohibiting the use and sale of seeds treated with plant protection products containing those active substances. *Official Journal of the European Union* **139**, 12-26.
79. European Food Safety Authority. 2013a Conclusion on the peer review of the pesticide risk assessment for bees for the active substance clothianidin. *EFSA Journal* **11**, 58. (doi:10.2903/j.efsa.2013.3066).
80. European Food Safety Authority. 2013b Conclusion on the peer review of the pesticide risk assessment for bees for the active substance imidacloprid. *EFSA Journal* **11**, 55. (doi:10.2903/j.efsa.2013.3068).
81. European Food Safety Authority. 2013c Conclusion on the peer review of the pesticide risk assessment for bees for the active substance thiamethoxam. *EFSA Journal* **11**, 68. (doi:10.2903/j.efsa.2013.3067).
82. European Food Safety Authority. 2013d Evaluation of the FERA study on bumble bees and consideration of its potential impact on the EFSA conclusions on neonicotinoids. *EFSA Journal* **11**, 20. (doi:10.2903/j.efsa.2013.3242).
83. European Food Safety Authority. 2013e Guidance on the risk assessment of plant protection products on bees (*Apis mellifera*, spp. and solitary bees). *EFSA Journal* **11**, 266. (doi:10.2903/j.efsa.2013.3295).
84. European Food Safety Authority. 2012 Statement on the findings in recent studies investigating sub-lethal effects in bees of some neonicotinoids in consideration of the uses currently authorised in Europe. *EFSA Journal* **10**, 27. (doi:10.2903/j.efsa.2012.2752).
85. Faucon J.P., Aurières C., Drajnudel P., Mathieu L., Ribière M., Martel A.C., Zeggane S., Chauzat M.P., Aubert M.F.A. 2005 Experimental study on the toxicity of imidacloprid given in syrup to honey bee (*Apis mellifera*) colonies. *Pest Management Science* **61**, 111-125. (doi:10.1002/ps.957).
86. Fauser-Misslin A., Sadd B.M., Neumann P., Sandrock C. 2013 Influence of combined pesticide and parasite exposure on bumblebee colony traits in the laboratory. *Journal of Applied Ecology*, n/a-n/a. (doi:10.1111/1365-2664.12188).
87. Feltham H., Park K., Goulson D. 2014 Field realistic doses of pesticide imidacloprid reduce bumblebee pollen foraging efficiency. *Ecotoxicology*. (doi:10.1007/s10646-014-1189-7).
88. Food and Environment Research Agency. 2013 *An Assessment of Key Evidence about Neonicotinoids and Bees*. Food and Environment Research Agency, Sand Hutton, Yorks, UK.
89. Fournier, A., Rollin, O., Le Féon, V., Decourtye, A., Henry, M. 2014 Crop-emptying rate and the design of pesticide risk assessment schemes in the honey bee and wild bees (Hymenoptera: Apidae). *Journal of Economic Entomology*, **107**, 38-46. (doi:10.1603/EC13087).
90. Franklin M.T., Winston M.L., Morandin L.A. 2004 Effects of clothianidin on *Bombus impatiens* (Hymenoptera: Apidae) colony health and foraging ability. *Journal of Economic Entomology* **97**, 369-373.
91. Frazier J., Mullin C., Frazier M., Ashcraft S. 2011 Managed pollinator CAP Coordinated Agricultural Project - Pesticides and their involvement in colony collapse disorder. *American Bee Journal* **151**, 779-784.
92. Free J.B. 1993 *Insect Pollination of Crops*. London, UK, Academic Press, Ltd.
93. Gallai N., Salles J.M., Settele J., Vaissière B.E. 2009 Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* **68**, 810-821 (doi: 10.1016/j.ecolecon.2008.06.014).
94. Garibaldi L.A., Aizen M.A., Klein A.M., Cunningham S.A., Harder L.D. 2011a Global growth and stability of agricultural yield decrease with pollinator dependence. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 5909-5914. (doi:10.1073/pnas.1012431108).
95. Garibaldi L.A., Steffan-Dewenter I., Kremen C., Morales J.M., Bommarco R., Cunningham S.A., Carvalheiro L.G., Chacoff N.P., Dudenhoffer J.H., Greenleaf S.S., Holzschuh, A., Isaacs, R., Krewenka, K., Mandelik, Y., Mayfield, M.M., Morandin, L.A., Potts, S.G., Ricketts, T.H., Szentgyorgyi, H., Viana, B.F., Westphal, C., Winfree, R., Klein, A.M. 2011b Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters* **14**, 1062-1072. (doi:10.1111/j.1461-0248.2011.01669.x).
96. Garibaldi L.A., Steffan-Dewenter I., Winfree R., Aizen M.A., Bommarco R., Cunningham S.A., Kremen C., Carvalheiro L.G., Harder L.D., Afik O., Bartomeus, I., Benjamin, F., Boreux, V., Cariveau, D., Chacoff, N.P., Dudenhoffer, J.H., Freitas, B.M., Ghazoul, J., Greenleaf, S., Hipólito, J., Holzschuh, A., Howlett, B., Isaacs, R., Javorek, S.K., Kennedy, C.M., Krewenka, K.M., Krishnan, S., Mandelik, Y., Mayfield, M.M., Motzke, I., Munyuli,

- T., Nault, B.A., Otieno, M., Petersen, J., Pisanty, G., Potts, S.G., Rader, R., Ricketts, T.H., Rundlöf, M., Seymour, C.L., Schüepp, C., Szentgyörgyi, H., Taki, H., Tschamntke, T., Vergara, C.H., Viana, B.F., Wanger, T.C., Westphal, C., Williams, N., Klein, A.M. 2013 Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* **340**, 1608-1611. (doi:10.1126/science.1230200).
97. Garratt M.P.D., Breeze T., Jenner N., Polce C., Biesmeijer J.C., Potts S.G. 2014 Avoiding a bad apple: insect pollination enhances fruit quality and economic value. *Agriculture Ecosystems & Environment* **184**, 34-40. (doi: 10.1016/j.agee.2013.10.032)
98. Garratt M.P.D., Truslove C.L., Coston D.J., Evans R.L., Moss E.D., Dodson C., Jenner N., J.C. B., Potts S.G. 2014 Pollination deficits in UK apple orchards. *Journal of Pollination Ecology* **12**, 9-14.
99. Gathmann A., Tschamntke T. 2002 Foraging ranges of solitary bees. *Journal of Animal Ecology* **71**, 757-764. (doi:10.1046/J.1365-2656.2002.00641.X).
100. Gels J.A., Held D.W., Potter D.A. 2002 Hazards of insecticides to the bumble bees *Bombus impatiens* (Hymenoptera : Apidae) foraging on flowering white clover in turf. *Journal of Economic Entomology* **95**, 722-728. (doi:10.1603/0022-0493-95.4.722).
101. Genersch E., Von Der Ohe W., Kaatz H., Schroeder A., Otten C., Büchler R., Berg S., Ritter W., Mühlen W., Gisder S., Meixner, M., Liebig, G., Rosenkranz, P. 2010 The German bee monitoring project: a long term study to understand periodically high winter losses of honey bee colonies. *Apidologie* **41**, 332-352. (doi: 10.1051/apido/2010014)
102. Gill R.J., Ramos-Rodriguez O., Raine N.E. 2012 Combined pesticide exposure severely affects individual-and colony-level traits in bees. *Nature* **491**, 105-108. (doi:10.1038/nature11585).
103. Girolami V., Marzaro M., Vivan L., Mazzon L., Giorio C., Marton D., Tapparo A. 2013 Aerial powdering of bees inside mobile cages and the extent of neonicotinoid cloud surrounding corn drillers. *Journal of Applied Entomology* **137**, 35-44. (doi:10.1111/J.1439-0418.2012.01718.X).
104. Girolami V., Marzaro M., Vivan L., Mazzon L., Greatti M., Giorio C., Marton D., Tapparo A. 2012 Fatal powdering of bees in flight with particulates of neonicotinoids seed coating and humidity implication. *Journal of Applied Entomology* **136**, 17-26. (doi:10.1111/j.1439-0418.2011.01648.x).
105. Girolami V., Mazzon L., Squartini A., Mori N., Marzaro M., Bernardo A.D., Greatti M., Giorio C., Tapparo A. 2009 Translocation of neonicotinoid insecticides from coated seeds to seedling guttation drops: A novel way of intoxication for bees. *Journal of Economic Entomology* **102**, 1808-1815. (doi:10.1603/029.102.0511).
106. Giurfa M., Sandoz J.C. 2012 Invertebrate learning and memory: fifty years of olfactory conditioning of the proboscis extension response in honeybees. *Learning & Memory* **19**, 54-66. (doi:10.1101/Lm.024711.111).
107. Godfray H.C.J., Donnelly C.A., Kao R.R., Macdonald D.W., McDonald R.A., Petrokofsky G., Wood J.L.N., Woodroffe R., Young D.B., McLean A.R. 2013 A restatement of the natural science evidence base relevant to the control of bovine tuberculosis in Great Britain. *Proceedings of the Royal Society B: Biological Sciences* **280**. (doi:10.1098/rspb.2013.1634).
108. Gonzalez-Varo J.P., Biesmeijer J.C., Bommarco R., Potts S.G., Schweiger O., Smith H.G., Steffan-Dewenter I., Szentgyörgyi H., Woyciechowski M., Vila M. 2013 Combined effects of global change pressures on animal-mediated pollination. *Trends in Ecology and Evolution* **28**, 524-530. (doi:10.1016/j.tree.2013.05.008).
109. Goulson D. 2013 An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology* **50**, 977-987. (doi:10.1111/1365-2664.12111).
110. Goulson D., Lepais O., O'Connor S., Osborne J.L., Sanderson R.A., Cussans J., Goffe L., Darvill B. 2010 Effects of land use at a landscape scale on bumblebee nest density and survival. *Journal of Applied Ecology* **47**, 1207-1215. (doi:10.1111/j.1365-2664.2010.01872.x).
111. Goulson D., Lye G.C., Darvill B. 2008 Decline and conservation of bumble bees. *Annual Review of Entomology* **53**, 191-208. (doi:10.1146/annurev.ento.53.103106.093454).
112. Greenleaf S.S., Kremen C. 2006a Wild bee species increase tomato production and respond differently to surrounding land use in Northern California. *Biological Conservation* **133**, 81-87. (doi:10.1016/J.Biocon.2006.05.025).
113. Greenleaf S.S., Kremen C. 2006b Wild bees enhance honey bees' pollination of hybrid sunflower. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 13890-13895. (doi:10.1073/Pnas.0600929103).
114. Greenleaf S.S., Williams N.M., Winfree R., Kremen C. 2007 Bee foraging ranges and their relationship to body size. *Oecologia* **153**, 589-596. (doi:10.1007/S00442-007-0752-9).
115. Gregorc A., Ellis J.D. 2011 Cell death localization in situ in laboratory reared honey bee (*Apis mellifera* L.) larvae treated with pesticides. *Pesticide Biochemistry and Physiology* **99**, 200-207. (doi:10.1016/J.Pestbp.2010.12.005).
116. Grixti J.C., Wong L.T., Cameron S.A., Favret C. 2009 Decline of bumble bees (*Bombus*) in the North American Midwest. *Biological Conservation* **142**, 75-84. (doi:10.1016/j.biocon.2008.09.027).
117. Gupta S., Gajbhiye V.T., Gupta R.K. 2008 Soil dissipation and leaching behavior of a neonicotinoid insecticide thiamethoxam. *Bulletin of Environmental Contamination and Toxicology* **80**, 431-437. (doi:10.1007/s00128-008-9420-y).
118. Hagen M., Wikelski M., Kissling W.D. 2011 Space use of bumblebees (*Bombus* spp.) revealed by radio-tracking. *PLoS ONE* **6**(5): e19997. (doi:10.1371/journal.pone.0019997).
119. Halm M.P., Rortais A., Arnold G., Tasei J.N., Rault S. 2006 New risk assessment approach for systemic insecticides: the case of honey bees and imidacloprid (Gaucho). *Environmental Science & Technology* **40**, 2448-2454. (doi:10.1021/Es051392i).
120. Hawthorne D.J., Dively G.P. 2011 Killing them with kindness? In-hive medications may inhibit xenobiotic efflux transporters and endanger honey bees. *PLoS ONE* **6**(11): e26796. (doi:10.1371/journal.pone.0026796).
121. Henry M., Béguin M., Requier F., Rollin O., Odoux J.F., Aupinel P., Aptel J., Tchamitchian S., Decourtye A. 2012a A common pesticide decreases foraging success and survival in honey bees. *Science* **336**, 348-350. (doi:10.1126/science.1215039).
122. Henry M., Béguin M., Requier F., Rollin O., Odoux J.F., Aupinel P., Aptel J., Tchamitchian S., Decourtye A. 2012b Response to comment on "a common pesticide decreases foraging success and survival in honey bees". *Science* **337**, 1453. (doi:10.1126/science.1224930).
123. Hoehn P., Tschamntke T., Tyljanakis J.M., Steffan-Dewenter I. 2008 Functional group diversity of bee pollinators increases crop yield. *Proceedings of the Royal Society B: Biological Sciences* **275**, 2283-2291. (doi:10.1098/rspb.2008.0405).

124. Hoffmann E.J., Castle S.J. 2012 Imidacloprid in melon guttation fluid: a potential mode of exposure for pest and beneficial organisms. *Journal of Economic Entomology* **105**, 67-71. (doi:10.1603/Ec11251).
125. Holzschuh A., Dormann C.F., Tscharntke T., Steffan-Dewenter I. 2011 Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the Royal Society B: Biological Sciences* **278**, 3444-3451. (doi:10.1098/rspb.2011.0268).
126. Hoyle M., Hayter K., Cresswell J.E. 2007 Effect of pollinator abundance on self-fertilization and gene flow: application to GM canola. *Ecological Applications* **17**, 2123-2135. (doi:10.1890/06-1972.1).
127. Iwasa T., Motoyama N., Ambrose J.T., Roe R.M. 2004 Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Protection* **23**, 371-378. (doi:10.1016/j.cropro.2003.08.018).
128. Jauker F., Bondarenko B., Becker H.C., Steffan-Dewenter I. 2012 Pollination efficiency of wild bees and hoverflies provided to oilseed rape. *Agricultural and Forest Entomology* **14**, 81-87. (doi:10.1111/j.1461-9563.2011.00541.x).
129. Jeschke P., Nauen R. 2008 Neonicotinoids - from zero to hero in insecticide chemistry. *Pest Management Science* **64**, 1084-1098. (doi:10.1002/ps.1631).
130. Jeschke P., Nauen R., Beck M.E. 2013 Nicotinic acetylcholine receptor agonists: a milestone for modern crop protection. *Angewandte Chemie International Edition* **52**, 9464-9485. (doi:10.1002/anie.201302550).
131. Jeschke P., Nauen R., Schindler M., Elbert A. 2011 Overview of the status and global strategy for neonicotinoids. *Journal of Agricultural and Food Chemistry* **59**, 2897-2908. (doi:10.1021/jf101303g).
132. Jha S., Kremen C. 2013 Resource diversity and landscape-level homogeneity drive native bee foraging. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 555-558. (doi:10.1073/pnas.1208682110).
133. Keitt T.H. 2009 Habitat conversion, extinction thresholds, and pollination services in agroecosystems. *Ecological Applications* **19**, 1561-1573. (doi:10.1890/08-0117.1).
134. Kennedy C.M., Lonsdorf E., Neel M.C., Williams N.M., Ricketts T.H., Winfree R., Bommarco R., Brittain C., Burley A.L., Cariveau D., Carvalheiro, L.G., Chacoff, N.P., Cunningham, S.A., Danforth, B.N., Dudenhoffer, J.H., Elle, E., Gaines, H.R., Garibaldi, L.A., Gratton, C., Holzschuh, A., Isaacs, R., Javorek, S.K., Jha, S., Klein, A.M., Krewenka, K., Mandelik, Y., Mayfield, M.M., Morandin, L., Neame, L.A., Otieno, M., Park, M., Potts, S.G., Rundlof, M., Saez, A., Steffan-Dewenter, I., Taki, H., Viana, B.F., Westphal, C., Wilson, J.K., Greenleaf, S.S., Kremen, C. 2013 A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters* **16**, 584-599. (doi:10.1111/Ele.12082).
135. Kevan P.G., Phillips T.P. 2001 The economic impacts of pollinator declines: an approach to assessing the consequences. *Conservation Ecology* **5**, art. no.-8.
136. Khoury D.S., Barron A.B., Myerscough M.R. 2013 Modelling food and population dynamics in honey bee colonies. *PLoS ONE* **8**(5): e59084. (doi:10.1371/journal.pone.0059084).
137. Khoury D.S., Myerscough M.R., Barron A.B. 2011 A quantitative model of honey bee colony population dynamics. *PLoS ONE* **6**(4): e18491. (doi:10.1371/journal.pone.0018491).
138. Klein A.M., Brittain C., Hendrix S.D., Thorp R., Williams N., Kremen C. 2012 Wild pollination services to California almond rely on semi-natural habitat. *Journal of Applied Ecology*, **49**, 723-732. (doi:10.1111/j.1365-2664.2012.02144.x).
139. Klein A.M., Vaissière B.E., Cane J.H., Steffan-Dewenter I., Cunningham S.A., Kremen C., Tscharntke T. 2007 Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303-313. (doi:10.1098/rspb.2006.3721).
140. Knight M.E., Martin A.P., Bishop S., Osborne J.L., Hale R.J., Sanderson A., Goulson D. 2005 An interspecific comparison of foraging range and nest density of four bumblebee (*Bombus*) species. *Molecular Ecology* **14**, 1811-1820. (doi:10.1111/J.1365-294x.2005.02540.X).
141. Kohler H.R., Triebkorn R. 2013 Wildlife ecotoxicology of pesticides: can we track effects to the population level and beyond? *Science* **341**, 759-765. (doi:10.1126/science.1237591).
142. Kollmeyer W.D., Flattum, R.F., Foster, J.P., Powell, J.E., Schroeder, M.E., Soloway, S.B. 1999 Discovery of the nitromethylene heterocycle insecticides. In *Nicotinoid insecticides and the nicotinic acetylcholine receptor* (eds. Yamamoto I., Casida J.), pp. 71-89. Tokyo, Springer-Verlag.
143. Kremen C., Williams N.M., Aizen M.A., Gemmill-Herren B., LeBuhn G., Minckley R., Packer L., Potts S.G., Roulston T., Steffan-Dewenter I., Vazquez, D.P., Winfree, R., Adams, L., Crone, E.E., Greenleaf, S.S., Keitt, T.H., Klein, A.M., Regetz, J., Ricketts, T.H. 2007 Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology Letters* **10**, 299-314. (doi:10.1111/J.1461-0248.2007.01018.X).
144. Krupke C.H., Hunt G.J., Eitzer B.D., Andino G., Given K. 2012 Multiple routes of pesticide exposure for honey bees living near agricultural fields. *PLoS ONE* **7**(1): e29268. (doi:10.1371/journal.pone.0029268).
145. Kuldna P., Peterson K., Poltinae H., Luig J. 2009 An application of DPSIR framework to identify issues of pollinator loss. *Ecological Economics* **69**, 32-42. (doi:10.1016/J.Ecolecon.2009.01.005).
146. Larson J.L., Redmond C.T., Potter D.A. 2013 Assessing insecticide hazard to bumble bees foraging on flowering weeds in treated lawns. *PLoS ONE* **8**(6): e66375. (doi:10.1371/journal.pone.0066375).
147. Lautenbach S., Seppelt R., Liebscher J., Dormann C.F. 2012 Spatial and temporal trends of global pollination benefit. *PLoS ONE* **7**(4): e35954. (doi:10.1371/journal.pone.0035954).
148. Laycock, I., Cotterell, K.C., O'Shea-Wheller, T.A., Cresswell, J.E. 2014 Effects of the neonicotinoid pesticide thiamethoxam at field-realistic levels on microcolonies of *Bombus terrestris* worker bumble bees. *Ecotoxicology and Environmental Safety* **100**, 153-158. (doi:10.1016/j.ecoenv.2013.10.027)
149. Laycock I., Cresswell J.E. 2013 Repression and recuperation of brood production in *Bombus terrestris* bumble bees exposed to a pulse of the neonicotinoid pesticide imidacloprid. *PLoS ONE* **8**(11): e79872. (doi:10.1371/journal.pone.0079872).
150. Laycock I., Lenthall K.M., Barratt A.T., Cresswell J.E. 2012 Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *Ecotoxicology* **21**, 1937-1945. (doi:10.1007/s10646-012-0927-y).
151. LeBuhn G., Droege S., Connor E.F., Gemmill-Herren B., Potts S.G., Minckley R.L., Griswold T., Jean R., Kula E., Roubik D.W., Cane, J., Wright, K.W., Frankie, G., Parker, F. 2013 Detecting insect pollinator declines on regional and global scales.

- Conservation Biology* **27**, 113-120. (doi:10.1111/j.1523-1739.2012.01962.x).
152. Leonhardt S.D., Gallai N., Garibaldi L.A., Kuhlmann M., Klein A.M. 2013 Economic gain, stability of pollination and bee diversity decrease from southern to northern Europe. *Basic and Applied Ecology* **14**, 461-471. (doi:10.1016/j.Baae.2013.06.003).
 153. Lonsdorf E., Kremen C., Ricketts T., Winfree R., Williams N., Greenleaf S. 2009 Modelling pollination services across agricultural landscapes. *Annals of Botany* **103**, 1589-1600. (doi: 10.1093/aob/mcp069).
 154. Losey J.E., Vaughan M. 2006 The economic value of ecological services provided by insects. *BioScience* **56**, 311-323. (doi:10.1641/0006-3568(2006)56[311:Tevoes]2.0.Co;2).
 155. Maienfisch P., Angst M., Brandl F., Fischer W., Hofer D., Kayser H., Kobel W., Rindlisbacher A., Senn R., Steinemann A., Widmer, H. 2001a Chemistry and biology of thiamethoxam: a second generation neonicotinoid. *Pest Management Science* **57**, 906-913. (doi:10.1002/Ps.365).
 156. Maienfisch P., Huerlimann H., Rindlisbacher A., Gsell L., Dettwiler H., Haettenschwiler J., Sieger E., Walti M. 2001b The discovery of thiamethoxam: a second-generation neonicotinoid. *Pest Management Science* **57**, 165-176. (doi:10.1002/1526-4998(200102)57:2<165::Aid-Ps289>3.0.Co;2-G).
 157. Martin S.J., Highfield A.C., Brettell L., Villalobos E.M., Budge G.E., Powell M., Nikaido S., Schroeder D.C. 2012 Global honey bee viral landscape altered by a parasitic mite. *Science* **336**, 1304-1306. (doi:10.1126/science.1220941).
 158. Marzaro M., Vivan L., Targa A., Mazzon L., Mori N., Greatti M., Toffolo E.P., di Bernardo A., Giorio C., Marton D., Tapparo, A., Girolami, V. 2011 Lethal aerial powdering of honey bees with neonicotinoids from fragments of maize seed coat. *Bulletin of Insectology* **64**, 119-126. (doi:10.1039/c1em10085h).
 159. Matsuda K., Buckingham S.D., Kleier D., Rau J.J., Grauso M., Sattelle D.B. 2001 Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends in Pharmacological Sciences* **22**, 573-580. (doi:10.1016/s0165-6147(00)01820-4).
 160. Matsumoto T. 2013a Reduction in homing flights in the honey bee *Apis mellifera* after a sublethal dose of neonicotinoid insecticides. *Bulletin of Insectology* **66**, 1-9.
 161. Matsumoto T. 2013b Short- and long-term effects of neonicotinoid application in rice fields, on the mortality and colony collapse of honeybees (*Apis mellifera*). *Journal of Apicultural Science* **57**, 21-35. (doi:10.2478/Jas-2013-0014).
 162. Maus C., Nauen R. 2011 Response to the publication: Tennekes, H.A. (2010): The significance of the Druckrey-Küpfmüller equation for risk assessment-The toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology* **280**, 176-177. (doi:10.1016/j.tox.2010.11.014).
 163. Maxim L., van der Sluijs J.P. 2010 Expert explanations of honeybee losses in areas of extensive agriculture in France: Gaucho (R) compared with other supposed causal factors. *Environmental Research Letters* **5**, 014006. (doi:10.1088/1748-9326/5/1/014006).
 164. Molet, M., Chittka, L., Stelzer, R.J., Streit, S.D., Raine, N.E. 2008 Colony nutritional status modulates worker responses to foraging recruitment pheromone in the bumblebee *Bombus terrestris*. *Behavioral Ecology and Sociobiology* **62**, 1919-1926. (doi:10.1007/s00265-008-0623-3).
 165. Mommaerts V., Reynders S., Boulet J., Besard L., Sterk G., Smagge G. 2010 Risk assessment for side-effects of neonicotinoids against bumblebees with and without impairing foraging behavior. *Ecotoxicology* **19**, 207-215. (doi:10.1007/s10646-009-0406-2).
 166. Morandin L.A., Winston M.L. 2003 Effects of novel pesticides on bumble bee (Hymenoptera: Apidae) colony health and foraging ability. *Environmental Entomology* **32**, 555-563. (doi:10.1603/0046-225X-32.3.555)
 167. Mullin C.A., Frazier M., Frazier J.L., Ashcraft S., Simonds R., vanEngelsdorp D., Pettis J.S. 2010 High Levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *PLoS ONE* **5**(3): e9754 (doi:10.1371/journal.pone.0009754).
 168. National Research Council Committee on the Status of Pollinators in North America. 2007. *Status of Pollinators in North America*. USA, Washington, DC, National Academies Press.
 169. Nauen R., Denholm I. 2005 Resistance of insect pests to neonicotinoid insecticides: current status and future prospects. *Archives of Insect Biochemistry and Physiology* **58**, 200-215. (doi:10.1002/Arch.20043).
 170. Nauen R., Ebbinghaus-Kintscher U., Salgado V.L., Kausmann M. 2003 Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pesticide Biochemistry and Physiology* **76**, 55-69. (doi:10.1016/S0048-3575(03)00065-8).
 171. Nguyen B.K., Mignon J., Laget D., De Graaf D.C., Jacobs F.J., VanEngelsdorp D., Brostaux Y., Saegerman C., Haubruge E. 2010 Honey bee colony losses in Belgium during the 2008-9 winter. *Journal of Apicultural Research* **49**, 337-339. (doi:10.3896/IBRA.1.49.4.07).
 172. Nguyen B.K., Saegerman C., Pirard C., Mignon J., Widart J., Thirionet B., Verheggen F.J., Berkvens D., De Pauw E., Haubruge E. 2009 Does imidacloprid seed-treated maize have an impact on honey bee mortality? *Journal of Economic Entomology* **102**, 616-623. (doi:10.1603/029.102.0220).
 173. Noleppa S., Hahn S. 2013 The value of neonicotinoid seed treatment in the European Union. A socio-economic, technological and environmental review. In *HFFA Working Paper 01/2013* (Germany, Berlin, Humboldt Forum for Food and Agriculture e.V. (HFFA)).
 174. Nuyttens D., Devarrewaere W., Verboven P., Foque D. 2013 Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review. *Pest Management Science* **69**, 564-575. (doi:10.1002/ps.3485).
 175. Ollerton J., Winfree R., Tarrant S. 2011 How many flowering plants are pollinated by animals? *Oikos* **120**, 321-326. (doi:10.1111/j.1600-0706.2010.18644.x).
 176. Osborne J.L., Clark S.J., Morris R.J., Williams I.H., Riley J.R., Smith A.D., Reynolds D.R., Edwards A.S. 1999 A landscape-scale study of bumble bee foraging range and constancy, using harmonic radar. *Journal of Applied Ecology* **36**, 519-533. (doi:10.1046/J.1365-2664.1999.00428.X).
 177. Osborne J.L., Martin A.P., Carreck N.L., Swain J.L., Knight M.E., Goulson D., Hale R.J., Sanderson R.A. 2008 Bumblebee flight distances in relation to the forage landscape. *Journal of Animal Ecology* **77**, 406-415. (doi:10.1111/J.1365-2656.2007.01333.X).
 178. Osgathorpe L.M., Park K., Goulson D., Acs S., Hanley N. 2011 The trade-off between agriculture and biodiversity in marginal areas: can crofting and bumblebee conservation be

- reconciled? *Ecological Economics* **70**, 1162-1169. (doi:10.1016/J.Econ.2011.01.010).
179. Palmer M.J., Moffat C., Saranzewa N., Harvey J., Wright G.A., Connolly C.N. 2013 Cholinergic pesticides cause mushroom body neuronal inactivation in honeybees. *Nature Communications* **4**, 1634. (doi:10.1038/ncomms2648).
180. Pettis J.S., Lichtenberg E.M., Andree M., Stitzinger J., Rose R., vanEngelsdorp D. 2013 Crop pollination exposes honey bees to pesticides which alters their susceptibility to the gut pathogen *Nosema ceranae*. *PLoS ONE* **8**(7): e70182. (doi:10.1371/journal.pone.0070182).
181. Pettis J.S., vanEngelsdorp D., Johnson J., Dively G. 2012 Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften* **99**, 153-158. (doi:10.1007/S00114-011-0881-1).
182. Pilling E., Campbell P., Coulson M., Ruddle N., Tornier I. 2013 A four-year field program investigating long-term effects of repeated exposure of honey bee colonies to flowering crops treated with thiamethoxam. *PLoS ONE* **8**(10): e77193. (doi:10.1371/journal.pone.0077193).
183. Pirard C., Widart J., Nguyen B.K., Deleuze C., Heudt L., Haubruge E., De Pauw E., Focant J.F. 2007 Development and validation of a multi-residue method for pesticide determination in honey using on-column liquid-liquid extraction and liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A* **1152**, 116-123. (doi:10.1016/j.chroma.2007.03.035).
184. Pistorius J., Bischoff G., Heimbach U., Stahler M. 2009 Bee poisoning incidents in Germany in spring 2008 caused by abrasion of active substance from treated seeds during sowing of maize. In "Hazards of pesticides to bees" - *Proceedings of the 10th International Symposium of the ICP-Bee Protection Group, October 8-10, 2008, Bucharest, Romania*. (eds. Oomen, P.A., Thompson, H.M.) *Julius Kuhn Archiv* **423**, 118-126.
185. Pistorius J., Brobyn T., Campbell P., Forster R., Lortsch J.A., Marolleau F., Maus C., Lückmann J., Suzuki H., Wallner K., Becker, R. 2012 Assessment of risks to honey bees posed by guttation. In "Hazards of pesticides to bees" - *Proceedings of the 11th International Symposium of the ICP-BR Bee Protection Group, Wageningen, The Netherlands, November 2-4, 2011*. (eds. Oomen, P.A., Thompson, H.M.) *Julius Kuhn Archiv* **437**, 199 -208 (doi:10.5073/jka.2012.437.056).
186. Pochi D., Biocca M., Fanigliulo R., Pulcini P., Conte E. 2012 Potential exposure of bees, *Apis mellifera* L., to particulate matter and pesticides derived from seed dressing during maize sowing. *Bulletin of Environmental Contamination and Toxicology* **89**, 354-361. (doi:10.1007/S00128-012-0664-1).
187. Pohorecka K., Skubida P., Miszczak A., Semkiw P., Sikorski P., Zagibajło K., Teper D., Kołtowski Z., Skubida M., Zdrańska D., Bober, A. 2012 Residues of neonicotinoid insecticides in bee collected plant materials from oilseed rape crops and their effect on bee colonies. *Journal of Apicultural Science* **56**, 115-134. (doi:10.2478/v10289-012-0029-3).
188. Potts S.G., Biesmeijer J.C., Kremen C., Neumann P., Schweiger O., Kunin W.E. 2010a Global pollinator declines: trends, impacts and drivers. *Trends in Ecology and Evolution* **25**, 345-353. (doi:10.1016/j.tree.2010.01.007).
189. Potts S.G., Roberts S.P.M., Dean R., Marris G., Brown M.A., Jones R., Neumann P., Settele J. 2010b Declines of managed honey bees and beekeepers in Europe. *Journal of Apicultural Research* **49**, 15-22. (doi:10.1016/j.tree.2010.01.007).
190. Rader R., Howlett B.G., Cunningham S.A., Westcott D.A., Edwards W. 2012 Spatial and temporal variation in pollinator effectiveness: do unmanaged insects provide consistent pollination services to mass flowering crops? *Journal of Applied Ecology* **49**, 126-134. (doi:10.1111/j.1365-2664.2011.02066.x).
191. Rader R., Howlett B.G., Cunningham S.A., Westcott D.A., Newstrom-Lloyd L.E., Walker M.K., Teulon D.A.J., Edwards W. 2009 Alternative pollinator taxa are equally efficient but not as effective as the honeybee in a mass flowering crop. *Journal of Applied Ecology* **46**, 1080-1087. (doi:10.1111/j.1365-2664.2009.01700.x).
192. Ratnieks F.L.W., Carreck N.L. 2010 Clarity on honey bee collapse? *Science* **327**, 152-153. (doi:10.1126/Science.1185563).
193. Reetz J.E., Zuhlke S., Spitteller M., Wallner K. 2011 Neonicotinoid insecticides translocated in guttated droplets of seed-treated maize and wheat: a threat to honeybees? *Apidologie* **42**, 596-606. (doi:10.1007/S13592-011-0049-1).
194. Ricketts T.H., Regezt J., Steffan-Dewenter I., Cunningham S.A., Kremen C., Bogdanski A., Gemmill-Herren B., Greenleaf S.S., Klein A.M., Mayfield M.M., Morandin, L.A., Ochieng, A., Viana, B.F. 2008 Landscape effects on crop pollination services: are there general patterns? *Ecology Letters* **11**, 499-515. (doi:10.1111/J.1461-0248.2008.01157.X).
195. Rortais A., Arnold G., Halm M.-P., Touffet-Briens F. 2005 Modes of honeybees exposure to systemic insecticides: estimated amounts of contaminated pollen and nectar consumed by different categories of bees. *Apidologie* **36**, 71-83. (doi:10.1051/apido:2004071).
196. Rosenkranz P., Aumeier P., Ziegelmann B. 2010 Biology and control of *Varroa destructor*. *Journal of Invertebrate Pathology* **103**, S96-S119. (doi:10.1016/j.jip.2009.07.016).
197. Sandrock C., Tanadini L., Pettis J., Biesmeijer J.C., Potts S.G., Neumann P. 2014 Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success. *Agricultural and Forest Entomology*. (doi:10.1111/afe.12041).
198. Schenke D., Joachimsmeier I., Pistorius J., Heimbach U. 2011 Pesticides in guttation droplets following seed treatment - field studies. In *Navigating environmental challenges: historical lessons guiding future directions*, Society of Environmental Toxicology and Chemistry North America (SETAC) 32nd annual meeting, Boston, USA (pp. 124-125, Society of Environmental Toxicology and Chemistry North America
199. Schmuck R., Schöning R., Stork A., Schramel O. 2001 Risk posed to honeybees (*Apis mellifera* L., Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest Management Science* **57**, 225-238. (doi:10.1002/ps.270).
200. Schneider C.W., Tautz J., Grünwald B., Fuchs S. 2012 RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. *PLoS ONE* **7**(1): e30023. (doi:10.1371/journal.pone.0030023).
201. Scott-Dupree C.D., Conroy L., Harris C.R. 2009 Impact of currently used or potentially useful insecticides for canola agroecosystems on *Bombus impatiens* (Hymenoptera: Apidae), *Megachile rotundata* (Hymenoptera: Megachilidae), and *Osmia lignaria* (Hymenoptera: Megachilidae). *Journal of Economic Entomology* **102**, 177-182. (doi:10.1603/029.102.0125).
202. Seeley T.D. 1995 *The Wisdom of the Hive*. Cambridge, Mass, USA, Harvard University Press.

203. Selim H.M., Jeong C.Y., Elbana T.A. 2010 Transport of imidacloprid in soils: miscible displacement experiments. *Soil Science* **175**, 375-381. (doi:10.1097/Ss.0b013e3181ebc9a2).
204. Škerl M.I.S., Velikonja Bolta S., Baša Česnik H., Gregorc A. 2009 Residues of pesticides in honeybee (*Apis mellifera carnica*) bee bread and in pollen loads from treated apple orchards. *Bulletin of Environmental Contamination and Toxicology* **83**, 374-377. (doi:10.1007/s00128-009-9762-0).
205. Smith, K.M., Loh, E.H., Rostal, M.K., Zambrana-Torrelío, C.M., Mendiola, L., Daszak, P. 2014. Pathogens, pests, and economics: drivers of honey bee colony declines and losses. *EcoHealth*, 1-12. (doi: 10.1007/s10393-013-0870-2).
206. Stark J.D., Jepson P.C., Mayer D.F. 1995 Limitations to use of topical toxicity data for predictions of pesticide side-effects in the field. *Journal of Economic Entomology* **88**, 1081-1088.
207. Starner K., Goh K.S. 2012 Detections of the neonicotinoid insecticide imidacloprid in surface waters of three agricultural regions of California, USA, 2010-2011. *Bulletin of Environmental Contamination and Toxicology* **88**, 316-321. (doi:10.1007/s00128-011-0515-5).
208. Staveley J.P., Law S.A., Fairbrother A., Menzie C.A. 2014 A causal analysis of observed declines in managed honey bees (*Apis mellifera*). *Human and Ecological Risk Assessment* **20**, 566-591. (doi:10.1080/10807039.2013.831263).
209. Steffan-Dewenter I., Kuhn A. 2003 Honeybee foraging in differentially structured landscapes. *Proceedings of the Royal Society of London Series B, Biological Sciences* **270**, 569-575. (doi:10.1098/rspb.2002.2292).
210. Stoner K.A., Eitzer B.D. 2012 Movement of soil-applied imidacloprid and thiamethoxam into nectar and pollen of squash (*Cucurbita pepo*). *PLoS ONE* **7**(6): e39114. (doi:10.1371/journal.pone.0039114).
211. Streit, S., Bock, F., Pirk, C.W.W., Tautz, J. 2003 Automatic life-long monitoring of individual insect behaviour now possible. *Zoology* **106**, 169-171. (doi: 10.1078/0944-2006-00113).
212. Suchail S., Debrauwer L., Belzunces L.P. 2004a Metabolism of imidacloprid in *Apis mellifera*. *Pest Management Science* **60**, 291-296. (doi:10.1002/ps.772).
213. Suchail S., Guez D., Belzunces L.P. 2001 Discrepancy between acute and chronic toxicity induced by imidacloprid and its metabolites in *Apis mellifera*. *Environmental Toxicology and Chemistry* **20**, 2482-2486.
214. Suchail S., Sousa G.d., Rahmani R., Belzunces L.P. 2004b *In vivo* distribution and metabolisation of ¹⁴C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Management Science* **60**, 1056-1062. (doi:10.1002/ps.895).
215. Sur R., Stork A. 2003 Uptake, translocation and metabolism of imidacloprid in plants. *Bulletin of Insectology* **56**, 35-40.
216. Tapparo A., Giorio C., Marzaro M., Marton D., Solda L., Girolami V. 2011 Rapid analysis of neonicotinoid insecticides in guttation drops of corn seedlings obtained from coated seeds. *Journal of Environmental Monitoring* **13**, 1564-1568. (doi:10.1039/C1em10085h).
217. Tapparo A., Marton D., Giorio C., Zanella A., Soldà L., Marzaro M., Vivan L., Girolami V. 2012 Assessment of the environmental exposure of honeybees to particulate matter containing neonicotinoid insecticides coming from corn coated seeds. *Environmental Science and Technology* **46**, 2592-2599. (doi:10.1021/es2035152).
218. Tasei J.N., Lerin J., Ripault G. 2000 Sub-lethal effects of imidacloprid on bumblebees, *Bombus terrestris* (Hymenoptera: Apidae), during a laboratory feeding test. *Pest Management Science* **56**, 784-788.
219. Tasei J.N., Ripault G., Rivault E. 2001 Hazards of imidacloprid seed coating to *Bombus terrestris* (Hymenoptera: Apidae) when applied to sunflower. *Journal of Economic Entomology* **94**, 623-627.
220. Tennekes H.A. 2010 The significance of the Druckrey-Küpfmüller equation for risk assessment - the toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology* **276**, 1-4. (doi:10.1016/j.tox.2010.07.005).
221. Tennekes H.A., Sánchez-Bayo F. 2012 Time-dependent toxicity of neonicotinoids and other toxicants: implications for a new approach to risk assessment. *Journal of Environmental & Analytical Toxicology* **54:001**. (doi:10.4172/2161-0525.S4-001).
222. Thompson H., Harrington P., Wilkins W., Pietravalle S., Sweet D., Jones A. 2013 Effects of neonicotinoid seed treatments on bumble bee colonies under field conditions. (<http://fera.co.uk/ccss/documents/defraBumbleBeeReportPS2371V4a.pdf>.)
223. Thompson H.M. 2012 Interaction between pesticides and other factors in effects on bees. Supporting Publications 2012:EN-340. UK, York, Food and Environment Research Agency; 204 pp. p.
224. Thompson H.M. 2010 Risk assessment for honey bees and pesticides-recent developments and 'new issues'. *Pest Management Science* **66**, 1157-1162. (doi:10.1002/ps.1994).
225. Thompson H.M., Hunt L. 1999 Extrapolating from honeybees to bumblebees in pesticide risk assessment. *Ecotoxicology* **8**, 147-166. (doi:10.1023/a:1026444029579).
226. Tomizawa M., Casida J.E. 2005 Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annual Review of Pharmacology and Toxicology* **45**, 247-268. (doi:10.1146/Annurev.Pharmtox.45.120403.095930).
227. Tomizawa M., Casida J.E. 2003 Selective toxicity of neonicotinoids attributable to specificity of insect and mammalian nicotinic receptors. *Annual Review of Entomology* **48**, 339-364. (doi:10.1146/Annurev.Ento.48.091801.112731).
228. Tomizawa M., Lee D.L., Casida J.E. 2000 Neonicotinoid insecticides: molecular features conferring selectivity for insect versus mammalian nicotinic receptors. *Journal of Agricultural and Food Chemistry* **48**, 6016-6024. (doi:10.1021/Jf000873c).
229. Tremolada P., Mazzoleni M., Saliu F., Colombo M., Vighi M. 2010 Field trial for evaluating the effects on honeybees of corn sown using cruiser ® and celest xl ® treated seeds. *Bulletin of Environmental Contamination and Toxicology* **85**, 229-234. (doi:10.1007/s00128-010-0066-1).
230. UK National Ecosystem Assessment. 2011 *The UK National Ecosystem Assessment: synthesis of the key findings*. (UK, Cambridge, UNEP-WCMC).
231. Vanbergen A.J., Insect Pollinators Initiative. 2013 Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment* **11**, 251-259. (doi:10.1890/120126).
232. Vanbergen, A.J., Heard, M.S., Breeze, T., Potts, S.G., Hanley, N. 2014 Status and value of pollination services – a report for the Department for Environment, Food and Rural Affairs. Defra Contract number: PH0514.
233. VanEngelsdorp D., Caron D., Hayes J., Underwood R., Henson M., Rennich K., Spleen A., Andree M., Snyder R., Lee K,

- Roccascecca, K., Wilson, M., Wilkes, J., Lengerich, E., Pettis, J. 2012 A national survey of managed honey bee 2010-11 winter colony losses in the USA: results from the Bee Informed Partnership. *Journal of Apicultural Research* **51**, 115-124. (doi:10.3896/IBRA.1.51.1.14).
234. VanEngelsdorp D., Evans J.D., Saegerman C., Mullin C., Haubruge E., Nguyen B.K., Frazier M., Frazier J., Cox-Foster D., Chen Y.P., Underwood, R., Tarpay, D.R., Pettis, J. S. 2009 Colony collapse disorder: a descriptive study. *PLoS ONE* **4**(8): e6481. (doi:10.1371/journal.pone.0006481).
235. VanEngelsdorp D., Hayes Jr J., Underwood R.M., Caron D., Pettis J. 2011 A survey of managed honey bee colony losses in the USA, fall 2009 to winter 2010. *Journal of Apicultural Research* **50**, 1-10. (doi:10.3896/IBRA.1.50.1.01).
236. VanEngelsdorp D., Speybroeck N., Evans J.D., Nguyen B.K., Mullin C., Frazier M., Frazier J., Cox-Foster D., Chen Y.P., Tarpay D.R., Haubruge, E., Pettis, J.S., Saegerman, C. 2010 Weighing risk factors associated with bee colony collapse disorder by classification and regression tree analysis. *Journal of Economic Entomology* **103**, 1517-1523. (doi:10.1603/Ec09429).
237. Velasco L., Mollers C., Becker H.C. 1999 Estimation of seed weight, oil content and fatty acid composition in intact single seeds of rapeseed (*Brassica napus* L.) by near-infrared reflectance spectroscopy. *Euphytica* **106**, 79-85. (doi:10.1023/A:1003592115110).
238. Vidau C., Diogon M., Aufauvre J., Fontbonne R., Viguès B., Brunet J.L., Texier C., Biron D.G., Blot N., Alaoui H., Belzunces, L.P., Delbac, F. 2011 Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by nosema ceranae. *PLoS ONE* **6**(6): e21550. (doi:10.1371/journal.pone.0021550).
239. Visscher P.K., Seeley T.D. 1982 Foraging strategy of honeybee colonies in a temperate deciduous forest. *Ecology* **63**, 1790-1801. (doi:10.2307/1940121).
240. Visser A., Blacquière T. 2010 Survival rate of honeybee (*Apis mellifera*) workers after exposure to sublethal concentrations of imidacloprid. *Proceedings Netherland Entomological Society Meetings* **21**, 29-34.
241. Von der Ohe W., Janke M. 2009 Bienen im Stress. Schäden entstehen wenn verschiedene Faktoren zusammen kommen. *Allgemeine Deutsche Imkerzeitung* **2009/4**, 10-11.
242. Walters K. 2013 Data, data everywhere but we don't know what to think? Neonicotinoid Insecticides and Pollinators. *Outlooks on Pest Management* **24**, 151-155. (doi:10.1564/v24_aug_02).
243. Wehling M., von der Ohe W., Brasse D., Forster R. 2009 Colony losses - interactions of plant protection products and other factors. In "Hazards of pesticides to bees" - *Proceedings of the 10th International Symposium of the ICP-Bee Protection Group, October 8-10, 2008, Bucharest, Romania*. (eds. Oomen, P.A., Thompson, H.M.) *royJulius Kuhn Archiv* **423**, 153-154.
244. Whitehorn P.R., O'Connor S., Wackers F.L., Goulson D. 2012 Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* **336**, 351-352. (doi:10.1126/science.1215025).
245. Whitfield C.W., Behura S.K., Berlocher S.H., Clark A.G., Johnston J.S., Sheppard W.S., Smith D.R., Suarez A.V., Weaver D., Tsutsui N.D. 2006 Thrice out of Africa: ancient and recent expansions of the honey bee, *Apis mellifera*. *Science* **314**, 642-645. (doi:10.1126/Science.1132772).
246. Wiest L., Buleté A., Giroud B., Fratta C., Amic S., Lambert O., Pouliquen H., Arnaudguilhem C. 2011 Multi-residue analysis of 80 environmental contaminants in honeys, honeybees and pollens by one extraction procedure followed by liquid and gas chromatography coupled with mass spectrometric detection. *Journal of Chromatography A* **1218**, 5743-5756. (doi:10.1016/j.chroma.2011.06.079).
247. Williams G.R., Tarpay D.R., VanEngelsdorp D., Chauzat M.P., Cox-Foster D.L., Delaplane K.S., Neumann P., Pettis J.S., Rogers R.E.L., Shutler D. 2010 Colony Collapse Disorder in context. *Bioessays* **32**, 845-846. (doi:10.1002/Bies.201000075).
248. Williamson S.M., Baker D.D., Wright G.A. 2013 Acute exposure to a sublethal dose of imidacloprid and coumaphos enhances olfactory learning and memory in the honeybee *Apis mellifera*. *Invert Neurosci* **13**, 63-70. (doi:10.1007/s10158-012-0144-7).
249. Williamson S.M., Wright G.A. 2013 Exposure to multiple cholinergic pesticides impairs olfactory learning and memory in honeybees. *Journal of Experimental Biology* **216**, 1799-1807. (doi:10.1242/jeb.083931).
250. Winfree R., Kremen C. 2009 Are ecosystem services stabilized by differences among species? A test using crop pollination. *Proc Biol Sci* **276**, 229-237. (doi:10.1098/rspb.2008.0709).
251. Winfree R., Williams N.M., Dushoff J., Kremen C. 2007 Native bees provide insurance against ongoing honey bee losses. *Ecology Letters* **10**, 1105-1113. (doi:10.1111/J.1461-0248.2007.01110.X).
252. Winston M.L. 1987 *The biology of the honey bee*. Cambridge, Mass ; London : Harvard University Press.
253. Wu J.Y., Anelli C.M., Sheppard W.S. 2011 Sub-lethal effects of pesticide residues in brood comb on worker honey bee (*Apis mellifera*) development and longevity. *PLoS ONE* **6**(2): e14720. (doi:10.1371/journal.pone.0014720).
254. Yang E.C., Chang H.C., Wu W.Y., Chen Y.W. 2012 Impaired olfactory associative behavior of honeybee workers due to contamination of imidacloprid in the larval stage. *PLoS ONE* **7**(11): e49472. (doi:10.1371/journal.pone.0049472).
255. Yang E.C., Chuang Y.C., Chen Y.L., Chang L.H. 2008 Abnormal foraging behavior induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *Journal of Economic Entomology* **101**, 1743-1748. (doi:10.1603/0022-0493-101.6.1743).
256. Zhang W., Ricketts T.H., Kremen C., Carney K., Swinton S.M. 2007 Ecosystem services and dis-services to agriculture. *Ecological Economics* **64**, 253-260. (doi:10.1016/J.Ecoecon.2007.02.024).
257. Zimmer C.T., Nauen R. 2011 Pyrethroid resistance and thiacloprid baseline susceptibility of European populations of *Meligethes aeneus* (Coleoptera: Nitidulidae) collected in winter oilseed rape. *Pest Management Science* **67**, 599-608. (doi:10.1002/Ps.2137).
258. Zurbuchen A., Cheesman S., Klaiber J., Muller A., Hein S., Dorn S. 2010a Long foraging distances impose high costs on offspring production in solitary bees. *Journal of Animal Ecology* **79**, 674-681. (doi:10.1111/J.1365-2656.2010.01675.X).
259. Zurbuchen A., Landert L., Klaiber J., Muller A., Hein S., Dorn S. 2010b Maximum foraging ranges in solitary bees: only few individuals have the capability to cover long foraging distances. *Biological Conservation* **143**, 669-676. (doi:10.1016/J.Biocon.2009.12.003).