

 Open access • Journal Article • DOI:10.1016/J.GLOPLACHA.2016.11.012

A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects

— [Source link](#) 

Enrique Doblas-Miranda, Rocío Alonso, Xavier Arnan, V. Bermejo ...+18 more authors

Institutions: Complutense University of Madrid, University of Castilla–La Mancha, Spanish National Research Council, University of Granada ...+3 more institutions

Published on: 01 Jan 2017 - Grid and Pervasive Computing

Topics: Land degradation, Land use, Global change, Climate change and Population

Related papers:

- [Reassessing global change research priorities in mediterranean terrestrial ecosystems: how far have we come and where do we go from here?](#)
- [R: A language and environment for statistical computing.](#)
- [Landscape – wildfire interactions in southern Europe: Implications for landscape management](#)
- [Biodiversity hotspots for conservation priorities](#)
- [Resilience of Mediterranean terrestrial ecosystems and fire severity in semiarid areas: Responses of Aleppo pine forests in the short, mid and long term.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/a-review-of-the-combination-among-global-change-factors-in-590ea3g89j>

1 **A review of the combination among global change factors in forests,**
2 **shrublands and pastures of the Mediterranean Region: beyond drought**
3 **effects**

4
5 *E Doblas-Miranda*^{1*}, *R Alonso*², *X Arnan*¹, *V Bermejo*², *L Brotons*^{1,3}, *J de las Heras*⁴, *M*
6 *Estiarte*^{1,5}, *JA Hódar*⁶, *P Llorens*⁷, *F Lloret*^{1,8}, *FR López-Serrano*⁴, *J Martínez-Vilalta*^{1,8}, *D*
7 *Moya*⁴, *J Peñuelas*^{1,5}, *J Pino*^{1,8}, *A Rodrigo*^{1,8}, *N Roura-Pascual*^{3,9}, *F Valladares*^{9,11}, *M Vilà*¹²,
8 *R Zamora*⁶, *J Retana*^{1,8}

9

10 1 CREAM, Cerdanyola del Vallès 08193, Spain.

11 2 Ecotoxicology of Air Pollution, CIEMAT, Avda. Complutense 22, 28040 Madrid, Spain.

12 3 Forestry Technology Centre of Catalonia (CTFC), St. Llorenç de Morunys km 2, 25280 Solsona,
13 Spain.

14 4 Technical School of Agricultural and Forestry Engineering, University of Castilla la Mancha,
15 Campus Universitario s/n, 02071 Albacete, Spain.

16 5 CSIC, Cerdanyola del Vallès 08193, Spain.

17 6 Terrestrial Ecology Group, Animal Biology and Ecology Department, University of Granada, E-
18 18071 Granada, Spain.

19 7 Institute of Environmental Assessment and Water Research (IDAEA), CSIC, 08034 Barcelona,
20 Spain.

21 8 Universitat Autònoma de Barcelona, Cerdanyola del Vallès 08193, Spain

22 9 Animal Biology Area, Environmental Sciences Department, University of Girona, Campus
23 Montilivi, 17071 Girona, Spain.

24 10 National Museum of Natural Sciences (MNCN), CSIC, Serrano 115 dpdo. E-28006 Madrid,
25 Spain.

26 11 Departamento de Biología y Geología, ESCET, Universidad Rey Juan Carlos, c) Tulipán s/n,
27 28933 Móstoles, Madrid, Spain.

28 12 Doñana Biological Station (EBD-CSIC), Américo Vespucio s/n, Isla de la Cartuja, 41092 Sevilla,
29 Spain.
30 * Corresponding author: +34 935814664, e.doblas@creaf.uab.es
31

This is the author's version of a work that was accepted for publication in Global and planetary change (Ed. Elsevier). Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Doblas-Miranda, E. et al. "A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: beyond drought effects" in Global and planetary change, vol. 148 (Jan. 2017), p. 42-54. DOI 10.1016/j.gloplacha.2016.11.012

32 **Abstract**

33 Climate change, alteration of atmospheric composition, land abandonment in some areas and
34 land use intensification in others, wildfires and biological invasions threaten forests,
35 shrublands and pastures all over the world. However, the impacts of the combinations
36 between global change factors are not well understood despite its pressing importance. Here
37 we posit that reviewing global change factors combination in an exemplary region can
38 highlight the necessary aspects in order to better understand the challenges we face, warning
39 about the consequences, and showing the challenges ahead of us. The forests, shrublands and
40 pastures of the Mediterranean Basin are an ideal scenario for the study of these combinations
41 due to its spatial and temporal heterogeneity, increasing and diverse human population and
42 the historical legacy of land use transformations. The combination of multiple global change
43 factors in the Basin shows different ecological effects. Some interactions alter the effects of
44 a single factor, as drought enhances or decreases the effects of atmospheric components on
45 plant ecophysiology. Several interactions generate new impacts: drought and land use
46 changes, among others, alter water resources and lead to land degradation, vegetation
47 regeneration decline, and expansion of forest diseases. Finally, different factors can occur
48 alone or simultaneously leading to further increases in the risk of fires and biological
49 invasions. The transitional nature of the Basin between temperate and arid climates involves
50 a risk of irreversible ecosystem change towards more arid states. However, combinations
51 between factors lead to unpredictable ecosystem alteration that goes beyond the particular
52 consequences of drought. Complex global change scenarios should be studied in the
53 Mediterranean and other regions of the world, including interregional studies. Here we show
54 the inherent uncertainty of this complexity, which should be included in any management
55 strategy.

56

57 **Keywords:** Atmospheric composition alteration, biological invasions, climate change, global
58 change factors interaction, land use intensification, land abandonment, natural resilience,
59 novel ecosystems, wildfires

60 **1 Introduction**

61 The Earth system is subject to a wide range of new planetary-forces that are originated in
62 human activities, ranging from the emission of greenhouse gases to the transformation of
63 landscapes and the loss of biota. The magnitude and rates of human-induced changes to the
64 global environment –a phenomenon known as global change- has accelerated since the
65 second half of the last century (Steffen et al., 2004; Vitousek, 1994). There is general
66 agreement about the factors of global environmental change and their ecological
67 consequences on terrestrial ecosystems. They imply extreme climatic events, atmospheric
68 chemical pollution, land use modifications, frequent fires and biological invasions, among
69 others (Lindner et al., 2010; Sala et al., 2000). However, uncertainty prevails in our capacity
70 to understand and predict the impact of their combination (Langley and Hungate 2014;
71 Scherber 2015). Therefore, there is a growing interest in understanding not only the factors
72 of global change and derived disturbances, but also the combinations among them (Moreira
73 et al., 2011; Rosenblatt and Schmitz, 2014).

74 Having a good knowledge of the factors of global environmental change and their
75 interactions is crucial to understand local to global implications, anticipate effects, prepare
76 for changes and reduce the risks of decision-making in a changing environment (Sternberg
77 and Yakir, 2015). This is especially certain in areas where many factors are involved and
78 intermingled, as in the Mediterranean Basin (Mooney et al., 2001; Sala et al., 2000). The
79 heterogeneity and transitional nature of the Mediterranean biogeography and the long history
80 of human alterations result in a spatially-structured landscape mosaic (Blondel et al., 2010;
81 Scarascia-Mugnozza et al., 2000; Woodward, 2009). All these aspects combined have
82 contributed to sustain a rich biota, which make the Mediterranean Basin a global biodiversity
83 hotspot (Myers et al., 2000), and to provide a scenario where historical legacies may have a
84 greater effect on present ecological processes than current factors (Dambrine et al., 2007).
85 However, future scenarios indicate that global change in the Mediterranean Basin will likely

86 involve a great risk of biodiversity loss (Malcolm et al., 2006; Sala et al., 2000) and a decline
87 of other ecosystem services, such as water and food resources, and carbon uptake (MEA,
88 2005; Schröter et al., 2005).

89 Numerous studies have examined the factors of global change on terrestrial ecosystems of
90 the highly diverse Mediterranean Basin (as it could be appreciated in the following review),
91 but a systematic revision of the effects of all factors of global change and their combination
92 is lacking. Here we first review the current and future impacts of the main global change
93 factors (drought and other climatic events, alteration of atmospheric composition, land use
94 intensification and abandonment, wildfires and biological invasions) on forests, shrublands
95 and pastures of the Mediterranean Basin (although the present work is focussed in terrestrial
96 ecosystems for practical reasons, we highly recommend Coll et al., 2010, as start point to a
97 similar review in the Mediterranean Sea) to then provide an assessment of the main types of
98 combinations among these factors. Our principal objectives are to show the impending
99 challenges of global change in the Mediterranean Basin and to warn about the potential
100 consequences of different combinations of global change factors.

101

102 **2 Main global change factors in the Mediterranean Basin**

103 *2.1 Drought and other climatic events*

104 Current aridity levels in the Mediterranean Basin appear to be unprecedented in the last 500
105 years (Nicault et al., 2008). Most climate models forecast substantial increases in
106 temperature and declines in precipitation, which will increase heat stress and largely reduce
107 water availability in the Basin (Gao and Giorgi, 2008; Hoerling et al., 2011). Models also
108 predict increases in climatic variability, with more extreme temperature and precipitation
109 events (Gao et al., 2006; Solomon et al., 2007).

110 Recent changes in precipitation have already been related to field data on tree growth
111 decreases (Sarris et al., 2007), increased growth variability (Vieira et al., 2010) and crown
112 defoliation on Mediterranean forests, in contrast to northern Europe (Carnicer et al., 2011).
113 Modelling exercises also project important changes in forest growth, although they also
114 highlight the complexity of the interactions involved (Fyllas et al., 2010; Sabaté et al., 2002).
115 Several drought simulation experiments have shown that water (Limousin et al., 2009) and
116 carbon fluxes (Matteucci et al., 2010; Misson et al., 2010) are highly sensitive to reductions
117 in precipitation. At the same time, phenology (Klein et al., 2013; Morin et al., 2010), nutrient
118 allocation and accumulation (Simoes et al., 2008) and key soil processes (e.g., Curiel-Yuste
119 et al., 2011; Sherman et al., 2012) have been shown to be affected by rainfall and
120 temperature manipulations. Described effects on plant communities should affect faunal
121 communities, as in the case of seed feeders (e.g., Sánchez-Humanes and Espelta, 2011) and
122 fauna affected by habitat loss (e.g., Scalercio, 2009). The effects of other climate extremes,
123 such as cold temperatures, have been less studied, although they may also be important
124 (Valladares et al., 2008).

125 Although evidence from both observational (e.g., Kazakis et al., 2007; Vennetier and Ripert,
126 2009) and experimental studies (e.g., De Dato et al., 2008; Matías et al., 2012) suggests that
127 changes in species composition can occur, studying these changes is difficult because they
128 require long-term monitoring. At the same time, some reports highlight the importance of
129 intraspecific variability, phenotypic plasticity and local adaptation (Poirier et al., 2012;
130 Ramírez-Valiente et al., 2010), among a plethora of stabilizing processes that may prevent
131 vegetation shifts from eventually occurring (cf. Lloret et al. 2012). Drought has also been
132 shown to affect the composition of soil fauna (e.g., Legakis and Adamopoulou, 2005;
133 Tsiafouli et al., 2005) and butterfly communities (Parmesan et al., 1999).

134

135 *2.2 Alteration of atmospheric composition*

136 The orography of the Mediterranean Basin provokes that in summer a stagnant layer of air
137 acts as a reservoir where most pollutants are transformed. Moreover, emissions in the Basin
138 could be drive directly into the mid and upper troposphere, being transported toward the
139 region (Moreno and Fellous, 1997). The impact of atmospheric composition changes in
140 Mediterranean Basin forests has scarcely been studied, despite the fact that these forests are
141 considered a significant carbon sink (Valentini et al., 2000).

142 Although short-term carbon dioxide (CO₂)-enrichment experiments in temperate forests
143 show an increase in net primary production (Norby et al., 2005), several tree-ring studies
144 have reported a general decrease in tree growth in the Mediterranean Basin (Nicault et al.,
145 2008). The controversy may be due to the constraints imposed by water or nutrient scarcity
146 on plant growth, affecting the overall impact of increased CO₂ effects (Leonardi et al., 2012;
147 Zhao and Running, 2010). In addition, photosynthetic acclimation to high CO₂ cannot be
148 ruled out (Peñuelas et al., 2011).

149 In the Western Mediterranean Basin, herbaria analysis shows a decrease in nitrogen (N)
150 concentration in leaf tissues throughout the 20th century (Peñuelas and Estiarte, 1997). The
151 increase in N deposition during recent decades in Europe (Galloway et al., 2008), can, at
152 least partially, offset N limitation and sustain the growth promoted by the CO₂ fertilization
153 (Milne and van Oijen, 2005). Nevertheless, other nutrients, such as phosphorus (P), will
154 remain unaltered and immobilized in biomass and soils, limiting further plant growth and
155 generating a significant imbalance in the N:P ratio (Peñuelas et al., 2012). Furthermore, N
156 deposition causes changes in soil quality, plant physiology and community composition, and
157 has been recognized as an important driver in biodiversity loss (Dias et al., 2011; Ochoa-
158 Hueso et al., 2011). Total annual estimates of N deposition in the Mediterranean Basin are
159 higher than those promoting adverse effects (Im et al., 2013).

160 Climatic conditions in the Mediterranean Basin favour Tropospheric ozone (O₃) formation
161 and persistence (Cristofanelli and Bonasoni, 2009; Hodnebrog et al., 2012). Mediterranean
162 woody vegetation seems to be in general tolerant to O₃ adverse effects due to its
163 sclerophyllous leaf structure, low gas exchange rates, BVOCs emissions and active
164 antioxidant defences (Paoletti, 2006). However, leaf senescence, increases in leaf mass per
165 area and spongy parenchyma thickness, decreases in photochemical maximal efficiency and
166 in the chlorophyll content, and biomass reduction caused by O₃ have been described in some
167 Mediterranean forest species (Paoletti, 2006; Ribas et al., 2005). Interactive effects between
168 CO₂ and O₃ are very variable as they depend on pollutant concentrations, species sensitivity
169 and interactions with other stresses such as plant competition, drought and nutrient
170 availability (Karnosky et al., 2007; Wittig et al., 2009).

171 The Mediterranean Basin is one of the hotspots of biogenic volatile organic compounds
172 (BVOC) emissions in Europe (Steinbrecher et al., 2009). BVOCs can act as a chemical sink
173 for O₃ at the leaf level, protecting vegetation from its negative effects (Fares et al., 2008;
174 Loreto et al., 2004), or enhancing O₃ production in the atmosphere through photochemical
175 reactions in the presence of N oxides (Peñuelas and Staudt, 2010). Increasing emissions of
176 BVOCs have, in any case, ecological impacts on Mediterranean life, given their key role in
177 plant defence and communication with other organisms (Peñuelas and Staudt, 2010). Rising
178 temperatures increase BVOC emission rates by enhancing their synthesis and by facilitating
179 vaporization (Peñuelas and Llusà, 2001), which likely results in an increasing feedback to
180 warming. BVOC emission rates present a broad range among plant species and therefore
181 will be largely affected by changes in vegetation biomass, vegetation types and land uses.

182

183 *2.3 Land use intensification and abandonment*

184 In the Mediterranean Basin region, contrasting patterns of recent land use changes appear
185 (Petit et al., 2001) with both abandonment and intensification co-occurring in the northern
186 areas, while deforestation and intense use of forest resources is still dominant in the southern
187 rim (Grove and Rackham, 2001) (Figure 1).

188 In the southern part of the Mediterranean Basin, the increasing rates of deforestation threaten
189 the scarce forest resources and ecological services of the region (Grove and Rackham, 2001).
190 Even if the amount of deforestation in the southern Mediterranean in the 1990s was low
191 compared to Latin America or Tropical Asia, the rate of increase compared to the '80s was
192 four times higher (Hansen and DeFries, 2004). Consequences of deforestation in this region
193 go beyond ecological effects, implying whole ecosystem change (Zaimeche, 1994).

194 In the northern Mediterranean Basin, metropolitan coastal landscapes are one of the most
195 altered in the world (Hepcan et al., 2012; Myers et al., 2000). Simultaneously, forests around
196 northern Mediterranean cities are suffering increasing ecological impact due to intense use
197 for leisure and progressive forest fragmentation resulting from urban sprawl (Jomaa et al.,
198 2008; Salvati et al., 2014). However, land use intensification of lowland regions is
199 encompassed with afforestation of low productive uplands (Falcucci et al., 2007; Roura-
200 Pascual et al., 2005) due to crop and pasture abandonment (Debussche et al., 1999; Tomaz et
201 al., 2013), and also to deliberate reforestation (Hansen and DeFries, 2004). These changes
202 are linked to profound socioeconomic shifts that led to a rural exodus and a decrease in
203 many of the traditional uses of forests (Grove and Rackham, 2001; Hill et al., 2008). As a
204 result, the northern Mediterranean forest landscapes have undergone large-scale changes, not
205 only in their general extent, but also in terms of vegetation structure, composition and
206 dynamics (Roura-Pascual et al., 2005). Novel forests composed of pioneer and introduced
207 species, and with relatively unknown structural and functional attributes, have proliferated
208 (Eldridge et al., 2011; Hobbs et al., 2006). These forests are becoming essential for the
209 restoration of landscape corridors between what remains of the historical forests and for the

210 recovery of forest species (Sirami et al., 2008). However, forest recovery could be heavily
211 influenced by the long-term effects of past land uses, which might determine soil fertility, or
212 by landscape impacts of current fire disturbance regimes (Puerta-Piñero et al., 2012). In fact,
213 past land uses could be a key factor altering the effects of current global changes and thus
214 differentiating the Basin from other Mediterranean regions of the world.

215

216 *2.4 Wild fires*

217 Wild fires of the Mediterranean Basin represent a dramatic hazard due to the dense human
218 population of the region (Dwyer et al., 2000). Moreover, historical alteration of fire patterns
219 in the Basin has modified vegetation resilience, differentiating it from the flora of other
220 Mediterranean regions (Pausas, 1999). Although in recent decades there has been a steady
221 increase in the resources invested in fire prevention and suppression, the number and extent
222 of wildfires have increased over the same period (Carmo et al., 2011; Piñol et al., 1998).
223 Climate has been the main driver of global biomass burning for the past two millennia
224 (Marlon et al., 2009). In the Mediterranean region, predictions indicate a general rise in fire
225 risk due to current warming (Moriondo et al., 2006).

226 Changes in the fire regime modify Mediterranean communities and their resilience to fire
227 (Paula et al., 2009; Tessler et al., 2014) in two ways. First, non-resilient tree species
228 dominant in sub-Mediterranean regions (Lloret et al., 2005) show very low regeneration after
229 large wildfires and are replaced by oak forests, shrublands or grasslands (Bendel et al., 2006;
230 Retana et al., 2002). Second, the higher fire frequency and intensity in fire-prone areas might
231 result in: (i) a decrease in the resprouting ability of plants and reduced resilience at the
232 landscape level of forests dominated by resprouters (Díaz-Delgado et al., 2002; Marzano et
233 al., 2012); (ii) a failure of obligate seeders regeneration when time intervals between fires are

234 shorter than the time required for a sufficient seed bank to build up ('immaturity risk', *sensu*
235 Zedler, 1995).

236 Additionally, wildfire events have major influences on the release of N and other air
237 pollutants and on the water quality of burned catchments (Johnson et al., 2007). Moreover,
238 increases in fire recurrence can affect ecosystem processes including long-term reductions in
239 primary production (Delitti et al., 2005; Dury et al., 2011) and increases in erosion (Thornes,
240 2009) as a consequence of a slow recovery of the soil organic layers (Shakesby, 2011) and
241 changes in microbial properties (Guénon et al., 2011). These changes frequently lead to
242 changes in plant and animal communities favoured by open areas (e.g., Broza and Izhaki,
243 1997; Fattorini, 2010; Kiss et al., 2004).

244

245 *2.5 Biological invasions*

246 Patterns of recent invasions (i.e. neophytes) among habitat types seem to be quite consistent
247 across Europe (Chytrý et al., 2008) and therefore across the Mediterranean Basin. The invasion
248 patterns differ considerably amongst taxonomy groups, although they tend to mostly occupy
249 anthropogenic habitats, while natural and semi-natural woody habitats are relatively resistant
250 to invasions (Arianoutsou et al., 2010; DAISIE, 2009). As in other regions worldwide, the
251 increase in the establishment of non-native species in the Mediterranean Basin will continue
252 due to the expanding transport of goods and people. Currently, the information available on
253 non-native species in the Basin is not complete and the number of non-native species across
254 taxonomic groups is underestimated (DAISIE, 2009). Detailed information about their
255 distribution and ecological impacts is necessary to determine exactly the current status of
256 biological invasions in the Mediterranean region.

257 We are starting to identify the ecological and economic consequences of invasions in
258 terrestrial ecosystems of the Mediterranean Basin. Non-native plants compete with native

259 species, decreasing local diversity and changing community composition (Vilà et al., 2006).
260 Changes in ecosystem functioning have been less explored, but they include alterations in
261 decomposition rates (De Marco et al., 2013) and changes in soil C and N pools (Vilà et al.,
262 2006). Even though the number of successful invaders seems to be higher in plants, the
263 consequences caused by animal invasions are not of a lower magnitude. The presence of
264 non-native vertebrates poses severe threats to native biodiversity through competition for
265 resources, predation and hybridization with native species, as well as economic impacts
266 (DAISIE, 2009). Most non-native terrestrial invertebrate species established in Europe are
267 known to be potential pests for agriculture and forestry products, while around 7 % affect
268 human and animal health (DAISIE, 2009). The ecological consequences of non-native
269 invertebrates have received less attention. Certain ants, such as *Linepithema humile* or
270 *Wasmannia auropunctata*, are known to have a dramatic effect on native invertebrate
271 communities (Blight et al., 2014; Vonshak et al., 2010).

272

273 **3 The combinations among factors alter the impacts of global change in the** 274 **Mediterranean Basin**

275 By addressing the principal global change factors affecting the Mediterranean Basin
276 separately, we have already covered how different pollutants can interact and how their
277 fluxes depend on forest cover, while current increases in fire frequency imply further
278 atmospheric alterations. In order to disentangle the possible effects of global change
279 combinations, we have crossed the different factors among them (Table1), and different
280 kinds of combinations have emerged (Figure 2). In the following sections we review the
281 potential combined effects of the various processes identified in the Region (following the
282 numbering in Table 1), boosted in many cases by the effects of drought. First, one factor can
283 alter the effect of another factor: for instance, the effects of atmospheric chemical

284 compounds on plant ecophysiology can be enhanced or decreased by drought (Figure 2a;
285 Section 3.1). Second, several interactions among factors trigger new impacts, such as the
286 alteration of water resources, land degradation, regeneration decline, and expansion of forest
287 diseases (Figure 2b; Sections 3.2, 3.3, 3.4, 3.5). Finally, different factors, alone or
288 simultaneously, can enhance the risk of other factors, as in the case of wildfire or invasion
289 risk (Figure 2c; Sections 3.6, 3.7).

290

291 *3.1 Modification of plant ecophysiology by interactions between atmospheric alteration and* 292 *drought*

293 Water availability is the main factor limiting biological activity in Mediterranean ecosystems
294 and, thus, modulating the response to changes in atmospheric chemistry. The direct effects
295 of higher atmospheric CO₂ include stomatal closure and enhancement of plant water-use
296 efficiency (WUE). WUE can alleviate the effects of drought on plant physiology and slow
297 down the depletion of soil water during drought progression (Morgan et al., 2004) (Figure
298 2a). Observations of naturally grown Mediterranean forests show a clear increase in WUE
299 during the 20th century, suggesting that the unobserved CO₂-fertilization benefits in growth
300 have likely been counteracted by drought (Peñuelas et al., 2011) (Figure 2a).

301 The reduction in plant growth caused by drought might be due to less N absorption. In this
302 sense, foliar N concentration has been found to have a positive correlation with precipitation
303 (Nahm et al., 2006). Also, drought affects soil microbial activity, leading to a reduction in N
304 mineralization and thus in absorption of deposited N (Rutigliano et al., 2009). All these
305 factors can increase soil N accumulation in oxidized forms and result in greater N losses
306 through leaching after torrential storms (Avila et al., 2010; MacDonald et al., 2002).

307 Depending on the level of stress, drought results in both decreases and increases in BVOC
308 emission rates (Peñuelas and Staudt, 2010). Mild heat stress may increase BVOC emissions

309 by making the isoprenoid synthesis pathway more competitive than carbon fixation
310 (Niinemets, 2010). On the contrary, severe drought may greatly decrease emissions because
311 of detrimental effects on protein levels and substrate supplies (Fortunati et al., 2008).
312 Drought stress protects plants against O₃ by inducing stomatal closure and pollutant uptake.
313 Indeed, high summer O₃ levels in the Mediterranean Basin occur when the seasonal drought
314 is more intense and plants are less physiologically active (Gerosa et al., 2009; Safieddine et
315 al., 2014). However, the additive effects of drought and O₃ have been described mainly
316 through an O₃-induced lose of stomatal regulation favouring drought stress (McLaughlin et
317 al., 2007). Ambient O₃ concentrations can thus increase water use by forest trees,
318 contributing to reduce water availability and thus amplifying the effects of climate change
319 (Alonso et al., 2014).

320

321 *3.2 Alteration of water resources by interactions between land use change and climate* 322 *change*

323 Water resources are very important in the densely populated and water-limited
324 Mediterranean Basin. The future of water resources in catchments must be assessed not only
325 in view of climate-forcing predictions, but also considering land-cover changes (Bates et al.,
326 2008), especially woody plant encroachment in mountain areas. A large set of catchment
327 experiments demonstrates that changes in land cover from grassed to forested areas involve a
328 reduction in runoff (i.e. Bosch and Hewlett, 1982; Brown et al., 2005). However, some
329 debate exists concerning larger catchments, where the role of forest cover is not always
330 clearly identifiable in the flow records (Andréassian, 2004; Oudin et al., 2008).

331 Historical records of large catchments studied in southern Europe show decreasing annual
332 trends and changes in flow regimes (e.g. Dahmani and Meddi 2009; Lespinas et al., 2010).

333 These trends are attributed to climatic shifts, increasing water consumption and

334 encroachment of forest cover due to land abandonment (García-Ruiz et al., 2011; Otero et
335 al., 2011). There seems to be a forest expansion threshold over which the effect of forest
336 cover on river discharges can be detected. In catchments with large and rapid forest
337 expansion, the effects of forest encroachment in the reduction of river discharges are well
338 documented (e.g., Gallart et al., 2011; Niedda et al. 2014). However, for other catchments,
339 the effects of forest advance on runoff are not so clear, as for example in some mountain
340 catchments in southern France or in catchments distributed from South to Central Italy (e.g.
341 Lespinas et al., 2010; Preti et al. 2011).

342 Considering only climate predictions and water consumption scenarios, the frequency of
343 floods is not expected to increase in Mediterranean Europe, except due to extreme climatic
344 events (Lehner et al., 2006). However, the influence of land-cover changes on floods, even at
345 the small catchment scale, is particularly difficult to assess in Mediterranean catchments
346 (Wittenberg et al., 2007). Among other factors, less is known about the rainfall partitioning
347 process in typical open woodlands, savannah-type ecosystems, isolated trees and shrub
348 formations than in closed forests (Latron et al., 2009; Llorens and Domingo, 2007).

349

350 *3.3 Land degradation favoured by interactions between either land use change or fire and* 351 *climatic events*

352 The loss of ecological and economical soil productivity is directly controlled by vegetation
353 cover, but can be aggravated by dry and variable climates (Imeson and Emmer, 1995;
354 Kosmas et al., 2002). Mediterranean ecosystems couple extreme climatic events with
355 materials that are highly susceptible to erosion (Poesen and Hooke, 1997). Current
356 predictions are that climate change, in combination with farmland abandonment, unsuitable
357 plantations, deforestation, overgrazing and fire, can overload the resilience of natural
358 ecosystem to erosion (Thornes, 2009).

359 While erosion is the initial process leading to soil and productivity losses, desertification is
360 the irreversible positive feedback loop of overexploitation favoured in certain dryland
361 systems (Kéfi et al., 2007; Puigdefábregas, 1995). There is a threshold over which the effects
362 of erosion are irreversible and the ecosystem cannot recover original biomass levels
363 (Puigdefábregas and Mendizabal, 1998). Desertification can be intensified and extended by
364 prolonged droughts (Kosmas et al., 2002), but also by potential human demographic
365 explosions in south-eastern Mediterranean regions (Le Houérou, 1992; Naveh, 2007).

366 Among the aforementioned factors, farmland abandonment increases the risk of gully
367 development when artificial systems are no longer maintained (Koulouri and Giourga, 2007;
368 Lesschen et al., 2007). The reduction in forest cover by clear-felling or fire increases water
369 runoff and sediment yields, especially when the organic layer is extensively affected (Imeson
370 and Emmer, 1995; Thornes, 2009). Vegetation-cover loss caused by overgrazing also results
371 in soil compaction, gully development and ultimately erosion hotspots (Thornes, 2005).

372 Overgrazing can result in greater impacts as climate become drier, combining both
373 disturbances in a negative feedback cycle (Köchy et al., 2008).

374 Drought induces impacts on vegetation that may result in erosion intensification (Thornes
375 and Brandt, 1994). The most direct effect of climate change may be increased rainfall
376 erosivity in the Mediterranean Basin, where the total rainfall will decrease but rainfall
377 intensity during certain events will increase (Nunes and Nearing, 2011). Aridity can also
378 affect soil biota negatively and slow down soil decomposition processes, decreasing the
379 content of organic matter (Curiel-Yuste et al., 2011; Imeson and Emmer, 1995). Appropriate
380 vegetation recovery after abandonment, disturbance or management should prevent soil and
381 nutrient loss (Duran Zuazo and Rodriguez Pleguezuelo, 2008; Fox et al., 2006).

382

383 3.4 Regeneration decline promoted by interactions between either land intensification or fire
384 and drought

385 Forest resilience is based on both the forest capacity to recover the pre-disturbance state and
386 the rate of plant growth. In this context, an increase in drought events might cause adverse
387 impacts on plant regeneration. Recurrent droughts affect woody species performance
388 differently, depending on species or functional type-specific sensitivity, leading to changes
389 in species composition and structure (De Dato, 2008; Galiano et al., 2010).

390 Herbivory can inhibit or exacerbate plant responses to climate-change conditions (Post and
391 Pedersen, 2008; Speed et al., 2010). In recent decades, the populations of wild ungulates
392 have increased beyond carrying capacities in the Mediterranean Basin, particularly in
393 protected areas and mountain regions (Noy-Meir et al., 1989). Where animals are selective
394 consumers of saplings and resprouts (such as goats), overgrazing severely affects forest
395 regeneration. This effect is aggravated in Mediterranean areas, where species such as *Pinus*
396 *sylvestris* present low sapling growth rates in comparison with those of northern latitudes
397 due to water limitation (Danell et al., 2003; Edenius et al., 1995). Furthermore, browsing on
398 saplings and resprouts in the Mediterranean Basin is more severe in summer and dry years,
399 when other food resources for ungulates are less abundant, diminishing the time for recovery
400 from damage (Herrero et al., 2012; Hester et al., 2004).

401 Fragmentation can also lead to regeneration decline in combination with drought. Smaller
402 patches not necessarily affect plant growth, which seems to be related to water stress, but
403 definitely affect reproduction (Matesanz et al., 2009). Considering the functionality of the
404 plant-soil-microbial system, small patches could even ameliorate the negative impacts of
405 drought through increasing the capacity of the soil to retain water due to higher soil organic
406 matter content than large patches. However, expected climatic changes in the already water-
407 limited Mediterranean Basin will overcome these processes (Flores-Rentería et al., 2015).

408 Post-fire forest regeneration depends on the identity and the regeneration capabilities of
409 dominant species (Buhk et al., 2007; Seligman and Henkin, 2000), which drives the
410 regeneration pattern of the whole plant community (Montès et al., 2004). First, in forests
411 dominated by seeders (such as several serotinous pine species, including *P. halepensis*, *P.*
412 *pinaster* and *P. brutia*), post-fire regeneration can be affected by drought since seed
413 germination requires imbibition of the embryo after the first autumn rains (Tsitsoni, 1997).
414 Higher aridity may lead to a reduction in reproduction effort and diminished seed bank
415 viability (Espelta et al., 2011; Keeley et al., 2005). Second, post-fire recovery of non-
416 serotinous pines such as *P. sylvestris* and *P. nigra* depends mainly on seed dispersal from
417 adjacent unburned patches. Therefore, frequent and intense fires might favour species shifts
418 (Retana et al., 2002). Finally, the resprouting ability of broadleaved forests can also decrease
419 due to long drought periods and low soil moisture (Castellari and Artale, 2010).

420

421 *3.5 Disease expansions induced by interactions between land use change and climate* 422 *change*

423 There is common agreement that climate change will favour forest pest species, since
424 survival of many arthropods depends on low temperature thresholds (Williams and Liebhold,
425 1995), while fungi or pathogens are also benefited by dry conditions (Ayres and
426 Lombardero, 2000; Jactel et al., 2012). However, the role of forest structure and composition
427 in disease expansion is more controversial (Figure 2b).

428 A Mediterranean example of insect pest is the pine processionary moth (PPM)
429 (*Thaumetopoea pityocampa*/*T. wilkinsoni* complex, Notodontidae), a well-known case due to
430 its ecological, economic and medical importance (Erkan, 2011; Gatto et al., 2009; Vega et
431 al., 2000). European cold-temperate species like the oak moth (*T. processionea*) and the
432 summer pine processionary moth (*T. pinivora*) have increased the intensity of their outbreaks

433 during the last two or three decades (Aimi et al., 2008; Groenen and Meurisse, 2012).
434 Meanwhile, the PPM has expanded in altitude (Battisti et al., 2005; Hódar and Zamora,
435 2004) and latitude (Battisti et al., 2005; Kerdelhué et al., 2009). PPM is a paradigm case of
436 sensitivity to global change for three reasons. First, due to its particular life cycle, with the
437 larval development occurring during winter (instead of spring-summer as is usual in
438 Lepidoptera), PPM is strongly dependent on minimum winter temperatures (Seixas Arnaldo
439 et al., 2011). Second, PPM has also shown a high capacity for local adaptation, with some
440 populations shifting to a summer cycle in cool areas and tolerating high temperatures at its
441 southern limit of distribution (Pimentel et al., 2006; Santos et al., 2011). And third, extensive
442 substitutions of broadleaved woodlands to pine plantations all over the Mediterranean have
443 created a situation in which PPM can thrive (Jactel et al., 2009; Kerdelhué et al., 2009).
444 Many other insect pests are showing similar dynamics and their importance is expected to
445 increase in the coming years, although reliable estimates are still not available (Battisti,
446 2005).

447 The story is different for fungus pathogens, which will benefit from the physiological
448 responses to temperature increase in combination with drought effects on plants. Cases such
449 as charcoal disease (*Biscogniauxia mediterranea*; Desprez-Loustau et al., 2006), Dutch elm
450 disease (*Ophiostoma ulmi*; Resco de Dios et al., 2007), chestnut blight (*Cryphonectria*
451 *parasitica*; Waldboth and Oberhuber, 2009) or oak decline (*Phytophthora cinnamomi*;
452 Brasier and Scott, 1994) are illustrative of the threats facing a large part of the Mediterranean
453 woodlands. For example, the combination of longer drought periods and fire may extend the
454 distribution of several diseases (such as *P. cinnamomi*) that affect forest stands in southern
455 Europe (Bergot et al., 2004). However, the possible effects that host range expansion and
456 forest connectivity increase have on pathogen dispersal have yet to be probed (Pautasso et
457 al., 2010).

458

459 *3.6 Increase of fire risk by the combination with drought and/or land-use change*

460 There is increasing evidence to show that high temperatures and low air humidity conditions
461 have become more common in recent decades and have been correlated with an increase in
462 the total burned surface (Dimitrakopoulos et al., 2011). Models predict that these climatic
463 conditions are going to become more frequent (Moriondo et al., 2006), determining changes
464 in the fire regime (Mouillot et al., 2002). Wildfires are expected to be more frequent at
465 higher altitudes and northern regions of the Mediterranean Basin, where they occurred only
466 occasionally in the past (for the Southern Alps, Reinhard et al., 2005). This pattern will
467 result in important consequences as dominant species of these areas often lack efficient post-
468 fire regeneration mechanisms (Vacchiano et al., 2014; Vilà-Cabrera et al., 2012), but may
469 also lead to more heterogeneous landscapes that have greater resilience to further
470 disturbances.

471 The social and ecological impacts of wildfires are related to the implementation of large-
472 scale, organized fire suppression strategies at the national level. These strategies decrease the
473 area burned in the short term, but lead to contrasting results in the long term due to fuel
474 accumulation (Piñol et al., 2005). In addition to climate, fuel is in fact the other main
475 physical driver of fire. Extensive agricultural abandonment during the past century has led to
476 extensive successional shrublands and forests mostly dominated by pines. The low
477 investment in fuel reduction practices has favoured high fuel load and vertical continuity
478 promoting high-intensity crown fires (Lloret et al., 2009; Mitsopoulos and Dimitrakopoulos,
479 2007). Crown fires have also affected large areas of managed pine woodlands, probably as a
480 result of fuel continuity across the landscape and the mountainous nature of the territory.
481 Also, in some areas, land use transformation to extensive grazing and human leisure
482 activities can easily give rise to fires, while rural exodus prevents early fire extinction.

483 In summary, the conjunction of a trend towards a homogeneous landscape dominated by
484 fuel-loaded vegetation (Loepfe et al., 2010) and a very active fire suppression policy is
485 favouring fuel accumulation (Lloret et al., 2009). This state of affairs, together with the
486 increasing climatic fire risk, is likely changing the fire regime to a set of large, frequent and
487 intense wildfires, thus challenging the resilience of the Mediterranean vegetation (Moreira et
488 al., 2011; Tsitsoni, 1997). To some extent, we may be contemplating wildfires as the catalyst
489 for the adjustment of many Mediterranean Basin ecosystems to a new climate-driven status
490 closer to semi-arid.

491

492 *3.7 Increase of invasion risk by the combination with drought, land-use change, atmospheric* 493 *alteration or fire*

494 Climate change can enhance biological invasions through increasing survival, reproduction
495 and spread of non-native species from warm climates (Walther et al., 2009). In the
496 Mediterranean Basin terrestrial ecosystems, many non-native species from temperate and
497 cold climates might only be able to shift their ranges northward or to expand in altitude.
498 However, the empirical evidence that this is occurring is anecdotal. Non-native species
499 whose native ranges are drier and warmer than their introduced ranges can be at an
500 advantage due to physiological or reproductive adaptations (for insects, Bale and Hayward,
501 2010). Still, model simulations and experiments suggest that changes in temperature alone
502 do not determine non-native plant distribution and fitness (Gritti et al., 2006; Ross et al.,
503 2008). In fact, recent studies stress the important influence of land-cover change in
504 accelerating invasions (Boulant, et al., 2009; Polce et al., 2011).

505 Future projections of changes in land use highlight that the invasion levels of terrestrial
506 ecosystems will increase regardless of the socioeconomic scenario (Chytrý et al., 2012).

507 Open areas favoured by land-use changes frequently provide “windows of opportunity” for

508 invasion as they increase propagule pressure and favour non-native species adapted to take
509 advantage of resource release (Ross et al., 2008; Roura-Pascual et al., 2009). In the
510 Mediterranean Basin, past crop uses explain the distribution and abundance of invasive
511 species in recently recovered forests and shrublands after a process of land abandonment
512 (Pretto et al., 2012). Moreover, certain land-use changes increase the fragmentation and
513 isolation of forest landscapes, which are more invaded than large continuous forests
514 (Malavasi et al., 2014). This landscape configuration enhances levels of invasion at forest
515 edges with urbanized or agricultural areas (Carpintero et al., 2004).

516 The interaction of atmospheric N deposition and plant invasion has not yet been explored in
517 the Mediterranean Basin, but it has been in other Mediterranean ecosystems (Padgett and
518 Allen, 1999). Fertilization experiments in arid scrublands of California indicate that areas
519 with high N deposition are more susceptible to non-native grass invasions, particularly in
520 wet years (Rao and Allen, 2010).

521 Fire has been proven to increase the expansion of non-native perennial grasses in the
522 Mediterranean Basin (Vilà et al., 2001; although see Dimitrakopoulos et al., 2005 for
523 contrasting results) which could feed back to increase the burnt area (Grigulis et al., 2005).

524 Some non-native plants invade recently burnt forests but disappear later on as their
525 persistence is constrained by the recovery of the native vegetation (Pino et al., 2013). On the
526 other hand, little information is available on the increasing pool of plant species able to
527 invade deeply shaded undisturbed forests (Martin et al., 2009). There are no similar studies
528 for non-native fauna, but fires are expected to create new opportunities for the expansion of
529 non-native animals already inhabiting the surroundings of the burned areas.

530 Combinations between environmental change and biological invasions are still largely
531 unknown. However, as the interaction of different global change factors can alter historical
532 succession patterns of native species (Keeley et al., 2005), similar interactions might lead to

533 more frequent and resilient invasions, challenging the resistance of the Mediterranean
534 terrestrial ecosystems.

535

536 *3.8 Potential combinations between more than two factors of global change*

537 Apart of the suggested combinations, more than two factors can interact generating even
538 more complex effects. It has been already mentioned the complex feedbacks between
539 climate, fire and atmospheric CO₂, the first increasing fire risk, which contributes to higher
540 CO₂ concentration in the atmosphere, which can in turn increase global warming (Stavros et
541 al., 2014). More specific are the studies of Dury et al. (2011) and Hodnebrog et al. (2012),
542 where other interactions between changes in atmospheric composition, climate and fire are
543 shown. Modelling the interaction between increasing levels of CO₂, drought and fire
544 frequency shows dramatic effects on forest productivity and distribution (Dury et al., 2011).
545 Also, the combined effects of fires, climate warming and different biogenic emissions affect
546 atmospheric ozone levels (Hodnebrog et al., 2012). Gil-Tena et al. (2011) show how fire,
547 land use changes and climate change can affect the distribution of bird species, while these
548 effects that can not be predicted by studying only one of these factors (Clavero et al. 2011).
549 Similarly, Mariota et al. (2014) have modelled how the combined effects of climate change
550 and fire on vegetation could be modified by land use changes.

551 Unfortunately, the few studies including three factors interaction mentioned in the previous
552 paragraph are not selected examples but the only ones found after a meticulous search (lists
553 of keywords related with each factor were included together and in all the potential different
554 combinations of four and three factors by using different fields on the ISI Web of Science in
555 the search of published research articles related to global change factors interaction in the
556 Mediterranean region, from 1900 to 2015). Moreover, although interactions between more

557 than three factors are also likely, we were not able to find any study considering this
558 possibility in Mediterranean forests, shrublands or pastures.

559

560 **4 Concluding remarks: global change combination in the Mediterranean Basin**

561 Different global change factors combine and interact causing unprecedented ecological
562 effects, which can be hardly predicted by the analysis of each factor in isolation. These
563 combinations and interactions bring some inherent uncertainty, which should be considered
564 in future research guidelines and when applying forest management strategies (Doblas-
565 Miranda et al., 2015). Principal sources of uncertainty are the contrasting effects between
566 atmospheric pollutants and drought, the role of forest cover in water availability, floods and
567 pest expansion and the thresholds of irreversibility that lead the change from one ecosystem
568 to another. In addition, much more complex interactions arise when combinations occur
569 together. For example, through altering forest extension and density, reforestation can
570 decrease erosion but may also reduce water availability, while drought can enhance erosion
571 and decrease water reserves. Moreover, both reforestation and drought may also indirectly
572 contribute to erosion by increasing fire risk (Figure 3). Uncertainty should be faced by
573 developing balanced adaptive strategies that account for the most likely consequences of the
574 major expected impacts and the inclusion of such information in any decision making
575 process (McCarthy and Possingham, 2007).

576 Comparative studies across regions and ecosystems by multisite approaches are necessary to
577 understand the impacts of global change. Particularly in the Mediterranean, previous
578 evaluations of the effects of global change have been performed (Lavorel et al., 1998; MEA,
579 2005; Sala et al., 2000), but new considerations need to be addressed. Climate change, and
580 especially drought, emerges as a crucial factor in most of the reviewed interactions and
581 therefore it should be considered when it comes to designing and applying international

582 management policies. For example, drought effects must be present when assessing critical
583 levels of several pollutants or mitigation effects of carbon sequestration in forests. The
584 ecological transitional nature of the Mediterranean Basin between temperate and arid regions
585 supposes a delicate equilibrium for multiple ecosystems, where a combination of global
586 change factors can balance their development to new arid states. Novel communities
587 associated to new global change factors, such as land abandonment and new fire regimes,
588 will be more prevalent, while our information about them remains scarce (Hobbs et al.,
589 2006). The identification of transition states leading to novel systems and the understanding
590 of the driving forces behind them remains a key priority for further research.

591 The information compiled in the present review highlights the potential relevance and
592 impact of interactions among emerging global change factors in the Mediterranean Basin.
593 Although global change is unavoidable in many cases, change does not necessarily mean
594 catastrophe, but adaptation. The enormous challenge of conserving Mediterranean terrestrial
595 ecosystems and the services they provide can only be met by means of a collective effort
596 involving not only the scientific community, but also forest managers and owners, decision
597 makers and the civic responsibility of society at large.

598

599 **Acknowledgements**

600 The present review is an outcome of the research project MONTES-Consolider (CSD2008-
601 00040), funded by the Spanish Ministry of Economy and Competitiveness. We thank
602 Jacque Minnett for her professional review as a native English speaker. Three anonymous
603 reviewers provided useful insights that were included in the current version.

604

605 **Bibliography**

606 Aimi, A., Larsson, S., Ronnås, C., Frazão, J., Battisti, A., 2008. Growth and survival of
607 larvae of *Thaumetopoea pinivora* inside and outside a local outbreak area. *Agricultural*
608 *and Forest Entomology* 10, 225–232.

609 Alonso, R. Elvira, S. González-Fernández I., Calvete, H. García-Gómez H. & Bermejo V.,
610 2014. Drought stress does not protect *Quercus ilex* L. from ozone effects: results from a
611 comparative study of two subspecies differing in ozone sensitivity. *Plant Biology* 16,
612 375–384.

613 Andréassian, V., 2004. Waters and forests: from historical controversy to scientific debate.
614 *Journal of Hydrology* 291, 1–27.

615 Arianoutsou, M., Delipetrou, P., Celesti-Grapow, L., Basnou, C., Bazos, I., Kokkoris, Y.,
616 Blasi, C., Vilà, M., 2010. Comparing naturalized alien plants and recipient habitats across
617 an east–west gradient in the Mediterranean Basin. *Journal of Biogeography* 37, 1811–
618 1823.

619 Ayres, M.P., Lombardero, M.J., 2000. Assessing the consequences of global change for
620 forest disturbance from herbivores and pathogens. *Science of the Total Environment* 262,
621 263–286.

622 Bale, J.S., Hayward, S.A.L., 2010. Insect overwintering in a changing climate. *Journal of*
623 *Experimental Biology* 213, 980–994.

624 Bates, B.C., Kundzewicz, Z.W., Wu, S., Palutikof, J.P., 2008. Climate change and water.
625 Technical Paper of the Intergovernmental Panel on Climate Change. IPCC Secretariat,
626 Geneva, 210 pp.

627 Battisti, A., 2005. Overview of entomological research concerning the forest ecosystems of
628 the northern rim of the Mediterranean Sea. In: Lieutier, F., Ghaioule, D. (Eds.),
629 *Entomological Research in Mediterranean Forest Ecosystems*. INRA Editions, Versailles,
630 pp. 15–20.

631 Battisti, A., Stastny, M., Netherer, S., Robinet, C., Schopf, A., Roques, A., Larsson, S.,
632 2005. Expansion of geographic range in the pine processionary moth caused by increased
633 winter temperatures. *Ecological Applications* 15, 2084–2096.

634 Bendel, M., Tinner, W., Ammann, B., 2006. Forest dynamics in the Pfyn forest in recent
635 centuries (Valais, Switzerland, Central Alps): interaction of pine (*Pinus sylvestris*) and
636 oak (*Quercus* sp.) under changing land use and fire frequency. *Holocene* 16, 81–89.

637 Bergot, M., Cloppet, E., Péronaud, V., Déqué, M., Desprez-Loustau, M.L., 2004.
638 Simulation of potential range expansion of oak disease caused by *Phytophthora*
639 *cinnamomi* under climate change. *Global Change Biology* 10, 1539–1552.

640 Blight, O., Orgeas, J., Torre, F., Provost, E., 2014. Competitive dominance in the
641 organisation of Mediterranean ant communities. *Ecological Entomology* 39, 595–602.

642 Blondel, J., Aronson, J., Bodiou, J.-Y., Boeuf, G., 2010. *The Mediterranean Region:*
643 *Biological Diversity in Space and Time*. Oxford University Press, Oxford, 376 pp.

644 Bosch, J.M., Hewlett, J.D., 1982. A review of catchment experiments to determine the effect
645 of vegetation changes on water yield and evapotranspiration. *Journal of Hydrology* 55, 3–
646 23.

647 Boulant, N., Garnier, A., Curt, T., Lepart, J., 2009. Disentangling the effects of land use,
648 shrub cover and climate on the invasion speed of native and introduced pines in
649 grasslands. *Diversity & distributions* 15, 1047–1059.

650 Brasier, C.M., Scott, J.K., 1994. European oak declines and global warming: a theoretical
651 assessment with special reference to the activity of *Phytophthora cinnamomi*. *EPPO*
652 *Bulletin* 24, 221–232.

653 Brown, A.E., Zhang, L., McMahon, T.A., Western, A.W., Vertessy, R.A., 2005. A review of
654 paired catchment studies for determining changes in water yield resulting from alterations
655 in vegetation. *Journal of Hydrology* 310, 28–61.

656 Broza, M., Izhaki, I., 1997. Post-fire arthropod assemblages in Mediterranean forest soils in
657 Israel. *International Journal of Wildland Fire* 7, 317–325.

658 Buhk, C., Meyn, A., Jentsch, A., 2007. The challenge of plant regeneration after fire in the
659 Mediterranean Basin: scientific gaps in our knowledge on plant strategies and evolution
660 of traits. *Plant Ecology* 192, 1–19.

661 Carmo, M., Moreira, F., Casimiro, P., Vaz, P., 2011. Land use and topography influences on
662 wildfire occurrence in northern Portugal. *Landscape and Urban Planning* 100, 169–176.

663 Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., Peñuelas, J., 2011. Widespread
664 crown condition decline, food web disruption, and amplified tree mortality with increased
665 climate change-type drought. *Proceedings of the National Academy of Sciences USA*
666 108, 1474–1478.

667 Carpintero, S., Reyes-Lopez, J., de Reyna, L.A., 2004. Impact of human dwellings on the
668 distribution of the exotic Argentine ant: a case study in the Doñana National Park, Spain.
669 *Biological Conservation* 115, 279–289.

670 Castellari, S., Artale, V., 2010. *Climate change in Italy: evidence, impacts and vulnerability.*
671 *Euro-Mediterranean Centre for Climate Change – CMCC – Bononia University Press,*
672 *Rome.*

673 Chytrý, M., Maskell, L.C., Pino, J., Pyšek, P., Vilà, M., Font, X., Smart, S.M., 2008. Habitat
674 invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental
675 and oceanic regions of Europe. *Journal of Applied Ecology* 45, 448–458.

676 Chytrý, M., Wild, J., Pyšek, P., Jarošík, V., Dendoncker, N., Reginster, I., Pino, J., Maskell,
677 L.C., Vilà, M., Pergl, J., 2012. Projecting trends in plant invasions in Europe under
678 different scenarios of future land-use change. *Global Ecology and Biogeography* 21, 75–
679 87.

680 Clavero, M., Villero, D., Brotons, L., 2011. Climate change or land use dynamics: Do we
681 know what climate change indicators indicate? *PLOS ONE* 6, e18581.

682 Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Lasram, F.B., Aguzzi, J., Ballesteros, E.,
683 Bianchi, C.N., Corbera, J., Dailianis, T., Danovaro, R., Estrada, M., Froggia, C., Galil,
684 B.S., Gasol, J.M., Gertwagen, R., Gil, J., Guilhaumon, F., Kesner-Reyes, K., Kitsos,
685 M.S., Koukouras, A., Lampadariou, N., Laxamana, E., López-Fé de la Cuadra, C.M.,
686 Lotze, H.K., Martin, D., Mouillot, D., Oro, D., Raicevich, S., Rius-Barile, J., Saiz-
687 Salinas, J.I., San Vicente, C., Somot, S., Templado, J., Turon, X., Vafidis, D., Villanueva,
688 R., Voultsiadou, E., 2010. The Biodiversity of the Mediterranean Sea: Estimates, Patterns,
689 and Threats. PLOS ONE 5, e11842.

690 Cristofanelli, P., Bonasoni, P., 2009. Background ozone in the Southern Europe and
691 Mediterranean area: influence of the transport processes. Environmental Pollution 157,
692 1399–1406.

693 Curiel-Yuste, J., Peñuelas, J., Estiarte, M., Garcia-Mas, J., Mattana, S., Ogaya, R., Pujol, M.,
694 Sardans, J., 2011. Drought-resistant fungi control soil organic matter decomposition and
695 its response to temperature. Global Change Biology 17, 1475–1486.

696 Dahmani, A., Meddi, M., 2009. Climate Variability and its Impact on Water Resources in
697 the Catchment Area of Wadi Fekan Wilaya of Mascara (West Algeria). European Journal
698 of Scientific Research 36, 458–472.

699 DAISIE, 2009. Handbook of Alien Species in Europe. Springer, Berlin, 400 pp.

700 Dambrine, E., Dupouey, J.L., Laüt, L., Humbert, L., Thinon, M., Beaufils, T., Richard, H.,
701 2007. Present forest biodiversity patterns in France related to former Roman agriculture.
702 Ecology 88, 1430–1439.

703 Danell, K., Bergstrom, R., Edenius, L., Ericsson, G., 2003. Ungulates as drivers of tree
704 population dynamics at module and genet levels. Forest Ecology and Management 181,
705 67–76.

706 Debussche, M., Lepart, J., Dervieux, A., 1999. Mediterranean landscape changes: evidences
707 from old postcards. Global Ecology and Biogeography 8, 3–15.

708 De Dato, G., Pellizzaro, G., Cesaraccio, C., Sirca, C., De Angelis, P., Duce, P., Spano, D.,
709 Mugnozza, G.S., 2008. Effects of warmer and drier climate conditions on plant
710 composition and biomass production in a Mediterranean shrubland community. *iForest* 1,
711 39–48.

712 De Marco, A., Arena, C., Giordano, M., Virzo, A.D.S., 2013. Impact of the invasive tree
713 black locust on soil properties of Mediterranean stone pine-helm oak forests. *Plant & Soil*
714 372, 473–486.

715 Delitti, W., Ferran, A., Trabaud, L., Vallejo, V.R., 2005. Effects of fire recurrence in
716 *Quercus coccifera* L. shrublands of the Valencia Region (Spain): I. plant composition and
717 productivity. *Plant Ecology* 177, 57–70.

718 Desprez-Loustau, M.L., Marçais, B., Nageleisen, L.M., Piou, D., Vanini, A., 2006.
719 Interactive effects of drought and pathogens in forest trees. *Annals of Forest Science* 63,
720 595–610.

721 Dias, T., Malveiro, S., Martins-Loução, M.A., Sheppard, L.J., Cruz, C., 2011. Linking N-
722 driven biodiversity changes with soil N availability in a Mediterranean ecosystem. *Plant*
723 *and Soil* 341, 125–136.

724 Díaz-Delgado, R., Lloret, F., Pons, X., Terradas, J., 2002. Satellite evidence of decreasing
725 resilience in Mediterranean plant communities after recurrent wildfires. *Ecology* 83,
726 2293–2303.

727 Dimitrakopoulos, A.P., Vlahou, M., Anagnostopoulou, C.G., Mitsopoulos, I.D., 2011.
728 Impact of drought on wildland fires in Greece: implications of climatic change? *Climatic*
729 *Change* 109, 331–347.

730 Dimitrakopoulos, P.G., Galanidis, A., Siamantziouras, A.S.D., Troumbis, A.Y., 2005. Short-
731 term invasibility patterns in burnt and unburnt experimental Mediterranean grassland
732 communities of varying diversities. *Oecologia* 143, 428–437.

733 Doblas-Miranda, E., Martinez-Vilalta, J., Lloret, F., Alvarez, A., Avila, A., Bonet, F.J.,
734 Brotons, L., Castro, J., Curiel Yuste, J., Diaz, M., Ferrandis, P., Garcia-Hurtado, E.,
735 Iriondo, J.M., Keenan, T.F., Latron, J., Llusia, J., Loepfe, L., Mayol, M., More, G., Moya,
736 D., Penuelas, J., Pons, X., Poyatos, R., Sardans, J., Sus, O., Vallejo, V.R., Vayreda, J.,
737 Retana, J., 2015. Reassessing global change research priorities in Mediterranean
738 terrestrial ecosystems: How far have we come and where do we go from here? *Global*
739 *Ecology and Biogeogr* 24, 25–43.

740 Duran Zuazo, V.H., Rodriguez Pleguezuelo, C.R., 2008. Soil-erosion and runoff prevention
741 by plant covers. A review. *Agronomy for sustainable development* 28, 65–86.

742 Dury, M., Hambuckers, A., Warnant, P., Henrot, A., Favre, E., Ouberdous, M., Francois, L.,
743 2011. Responses of European forest ecosystems to 21st century climate: assessing changes
744 in interannual variability and fire intensity. *iForest* 4, 82–99.

745 Dwyer, E., Pinnok, S., Gregoire, J.-M., Pereira, J.M.C., 2000. Global spatial and temporal
746 distribution of vegetation fire as determined from satellite observations. *International*
747 *Journal of Remote Sensing* 21, 1289–1302.

748 Edenius, L., Danell, K., Nyquist, H., 1995. Effects of simulated moose browsing on growth,
749 mortality, and fecundity on Scots pine: relations to plant productivity. *Canadian Journal*
750 *of Forest Research* 25, 529–535.

751 Eldridge, D.J., Bowker, M.A., Maestre, F.T., Roger, E., Reynolds, J.F., Whitford, W.G.,
752 2011. Impacts of shrub encroachment on ecosystem structure and functioning: towards a
753 global synthesis. *Ecology Letters* 14, 709–722.

754 Erkan, N., 2011. Impact of pine processionary moth (*Thaumetopoea wilkinsoni* Tams) on
755 growth of Turkish red pine (*Pinus brutia* Ten.). *African journal of agricultural research* 6,
756 4983–4988.

757 Espelta, J.M., Arnan, X., Rodrigo, A., 2011. Non-fire induced seed release in a weakly
758 serotinous pine: climatic factors, maintenance costs or both? *Oikos* 120, 1752–1760.

759 Falcucci, A., Maiorano, L., Boitani, L., 2007. Changes in land-use/land-cover patterns in
760 Italy and their implications for biodiversity conservation. *Landscape Ecology* 22, 617–
761 631.

762 Fares, S., Loreto, F., Kleist, E., Wildt, J., 2008. Stomatal uptake and stomatal deposition of
763 ozone in isoprene and monoterpene emitting plants. *Plant Biology* 10, 44–54.

764 Fattorini, S., 2010. Effects of fire on tenebrionid communities of a *Pinus pinea* plantation: a
765 case study in a Mediterranean site. *Biodiversity and Conservation* 19, 1237–1250.

766 Flores-Rentería, D., Curiel Yuste, J., Rincón, A., Brearley, F.Q., García-Gil, J.C.,
767 Valladares, F., 2015. Habitat fragmentation can modulate drought effects on the plant-
768 soil-microbial system in Mediterranean holm oak (*Quercus ilex*) forests. *Microbial*
769 *Ecology* 69, 798–812.

770 Fyllas, N.M., Politi, P.I., Galanidis, A., Dimitrakopoulos, P.G. Arianoutsou, M., 2010.
771 Simulating regeneration and vegetation dynamics in Mediterranean coniferous forests.
772 *Ecological Modelling* 221, 1494–1504.

773 Fortunati, A., Barta, C., Brilli, F., Centritto, M., Zimmer, I., Schnitzler, J.P., Loreto, F.,
774 2008. Isoprene emission is not temperature-dependent during and after severe drought-
775 stress: a physiological and biochemical analysis. *Plant Journal* 55, 687–697.

776 Fox, D., Berolo, W., Carrega, P., Darboux, F., 2006. Mapping erosion risk and selecting sites
777 for simple erosion control measures after a forest fire in Mediterranean France. *Earth*
778 *Surface Processes and Landforms* 31, 606–621.

779 Gallart, F., Delgado, J., Beatson, S.W., Posner, H., Llorens, P., Marcé, R., 2011. Analysing
780 the effect of global change on the historical trends of water resources in the headwaters of
781 the Llobregat and Ter river basins (Catalonia, Spain). *Physics and Chemistry of the Earth*
782 36, 655–661.

783 Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R.,
784 Martinelli, L.A., Seitzinger, S.P., Sutton, M.A., 2008. Transformation of the nitrogen
785 cycle: recent trends, questions, and potential solutions. *Science* 320, 889–892.

786 Gao, X., Giorgi, F., 2008. Increased aridity in the Mediterranean region under greenhouse
787 gas forcing estimated from high resolution simulations with a regional climate model.
788 *Global and Planetary Change* 62, 195–209.

789 Gao, X., Pal, J.S., Giorgi, F., 2006. Projected changes in mean and extreme precipitation
790 over the Mediterranean region from high resolution double nested RCM simulations.
791 *Geophysical Research Letters* 33, L03706.

792 García-Ruiz, J.M., López-Moreno, J.I., Vicente-Serrano, S.M., Lasanta-Martínez, T.,
793 Beguería, S., 2011. Mediterranean water resources in a global change scenario. *Earth-*
794 *Science Reviews* 105, 121–139.

795 Gatto, P., Zocca, A., Battisti, A., Barrento, M.J., Branco, M., Paiva, M.R., 2009. Economic
796 assessment of managing processionary moth in pine forests: A case-study in Portugal.
797 *Journal of Environmental Management* 90, 683–691.

798 Gerard, F., Petit, S., Smith, G., Thomson, A., Brown, N., Manchester, S., Wadsworth, R.,
799 Bugar, G., Halada, L., Bezák, P., Boltiziar, M., De badts, E., Halabuk, A., Mojses, M.,
800 Petrovic, F., Gregor, M., Hazeu, G., Múcher, C.A., Wachowicz, M., Huitu, H., Tuominen,
801 S., Köhler, R., Olschofsky, K., Ziese, H., Kolar, J., Sustera, J., Luque, S., Pino, J., Pons,
802 X., Roda, F., Roscher, M., Feranec, J., 2010. Land cover change in Europe between 1950
803 and 2000 determined employing aerial photography. *Progress in Physical Geography* 34,
804 183–205.

805 Gerosa, G., Finco, A., Mereu, S., Vitale, M., Manes, F., Denti, A.B., 2009. Comparison of
806 seasonal variations of ozone exposure and fluxes in a Mediterranean Holm oak forest
807 between the exceptionally dry 2003 and the following year. *Environmental Pollution* 157,
808 1737–1744.

809 Gil-Tena, A., Fortin, M.J., Brotons, L., Saura, S., 2011. Forest Avian Species Richness
810 Distribution and Management Guidelines under Global Change in Mediterranean
811 Landscapes. In Li, C., Laforteza, R., Chen, J. (Eds.) Landscape Ecology in Forest
812 Management and Conservation: Challenges and Solutions for Global Change. Springer,
813 Berlin, 231–251.

814 Grigulis, K., Lavorel, S., Davies, I.D., Dossantos, A., Lloret, F., Vilà, M., 2005. Landscape-
815 scale positive feedbacks between fire and expansion of the large tussock grass,
816 *Ampelodesmos mauritanica* in Catalan shrublands. *Global Change Biology* 11, 1042–
817 1053.

818 Gritti, E.S., Smith B., Sykes, M.T., 2006. Vulnerability of Mediterranean Basin ecosystems
819 to climate change and invasion by exotic plant species. *Journal of Biogeography* 33, 145–
820 157.

821 Groenen, F., Meurisse, N., 2012. Historical distribution of the oak processionary moth
822 *Thaumetopoea processionea* in Europe suggests recolonization instead of expansion.
823 *Agricultural and Forest Entomology* 14, 147–155.

824 Grove, A.T., Rackman, O., 2001. The nature of Mediterranean Europe. Yale University
825 Press, China, 384 pp.

826 Guénon, R., Vennetier, M., Dupuy, N., Ziarelli, F., Gros, R., 2011. Soil organic matter
827 quality and microbial catabolic functions along a gradient of wildfire history in a
828 Mediterranean ecosystem. *Applied Soil Ecology* 48, 81–93.

829 Hansen, M.C., DeFries, R.S., 2004. Detecting long-term global forest change using
830 continuous fields of tree-cover maps from 8-km advanced very high resolution radiometer
831 (AVHRR) data for the years 1982–99. *Ecosystems* 7, 695–716.

832 Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A.,
833 Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A.,
834 Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st-

835 century forest cover change. *Science* 342, 850–853. Data available on-line from:
836 <http://earthenginepartners.appspot.com/science-2013-global-forest>.

837 Herrero, A., Zamora, R., Castro, J., Hódar, J.A., 2012. Limits of pine forest distribution at
838 the treeline: herbivory matters. *Plant Ecology* 213, 459–469.

839 Hepcan, S., Hepcan, C.C., Kilicaslan, C., Ozkan, M.B., Kocan, N., 2013 Analyzing
840 Landscape Change and Urban Sprawl in a Mediterranean Coastal Landscape: A Case
841 Study from Izmir, Turkey. *Journal of Coastal Research* 29, 301–310.

842 Hester, A.J., Millard, P., Baillie, G.J., Wendler, R., 2004. How does timing of browsing
843 affect above- and below-ground growth of *Betula pendula*, *Pinus sylvestris* and *Sorbus*
844 *aucuparia*? *Oikos* 105, 536–550.

845 Hill, J., Stellmes, M., Udelhoven, T., Röder, A., Sommer, S., 2008. Mediterranean
846 desertification and land degradation. Mapping related land use change syndromes based
847 on satellite observations. *Global and Planetary Change* 64, 146–157.

848 Hobbs, R.J., Arico, S., Aronson, J., Baron, J.S., Bridgewater, P., Cramer, V.A., Epstein,
849 P.R., Ewel, J.J., Klink, C.A., Lugo, A.E., Norton, D., Ojima, D., Richardson, D.M.,
850 Sanderson, E.W., Valladares, F., Vila, M., Zamora, R., Zobel, M., 2006. Novel
851 ecosystems: theoretical and management aspects of the new ecological world order.
852 *Global Ecology and Biogeography* 15, 1–7.

853 Hódar, J.A., Zamora, R., 2004. Herbivory and climatic warming: a Mediterranean
854 outbreaking caterpillar attacks a relict, boreal pine species. *Biodiversity and Conservation*
855 13, 493–500.

856 Hodnebrog, O, Solberg, S., Stordal, F., Svendby, T.M., Simpson, D., Gauss, M., Hilboll, A.,
857 Pfister, G.G., Turquety, S., Richter, A., Burrows, J.P., van der Gon, H.A.C.D., 2012.
858 Impact of forest fires, biogenic emissions and high temperatures on the elevated Eastern
859 Mediterranean ozone levels during the hot summer of 2007. *Atmospheric Chemistry and*
860 *Physics* 12, 8727–8750.

861 Hoerling, M., Eischeid, J., Perlwitz, J., Quan, X., Zhang, T., Pegion, P., 2011. On the
862 increased frequency of Mediterranean drought. *Journal of Climate* 25, 2146–2161.

863 Im, U., Christodoulaki, S., Violaki, K., Zarmas, P., Kocak, M., Daskalakis, N.,
864 Mihalopoulos, N., Kanakidou, M., 2013. Atmospheric deposition of nitrogen and sulfur
865 over southern Europe with focus on the Mediterranean and the Black Sea. *Atmospheric*
866 *Environment* 81, 660–670.

867 Imeson, A.C., Emmer, I.M., 1995. Implications of climate change on land degradation in the
868 Mediterranean. In: Jeftić, L. Milliman, J.D., Sestini, G. (Eds.), *Climate Change and the*
869 *Mediterranean*. UNEP, Arnold, Boston, pp. 95–128.

870 Jactel, H., Nicoll, B.C., Branco, M., Gonzalez-Olabarria, J.R., Grodzki, W., Långström, B.,
871 Moreira, F., Netherer, S., Orazio, C., Piou, D., Santos, H., Schelhaas, M.J., Tojic, K.,
872 Vodde, F., 2009. The influences of forest stand management on biotic and abiotic risks of
873 damage. *Annals of Forest Science* 66, 701.

874 Jactel, H., Petit, J., Desprez-Loustau, M.L., Delzon, S., Piou, D., Battisti, A., Koricheva, J.,
875 2012. Drought effects on damage by forest insects and pathogens: a meta-analysis. *Global*
876 *Change Biology* 18, 267–276.

877 Johnson, D.W., Murphy, J.D., Walker, R.F., Glass, D.W., Miller, W.W., 2007. Wildfire
878 effects on forest carbon and nutrient budgets. *Ecological Engineering* 31, 183–192.

879 Jomaa, I., Auda, Y., Saleh, B.A., Hamze, M., Safi, S., 2008. Landscape spatial dynamics
880 over 38 years under natural and anthropogenic pressures in Mount Lebanon. *Landscape*
881 *and Urban Planning* 87, 67–75.

882 Karnosky, D.F., Skelly, J.M., Percy, K.E., Chappelka, A.H., 2007. Perspectives regarding 50
883 years of research on effects of tropospheric ozone air pollution on US forests. Review.
884 *Environmental Pollution* 147, 489–506.

885 Kazakis, G., Ghosn, D., Vogiatzakis, I.N., Papanastasis V.P., 2007. Vascular plant diversity
886 and climate change in the alpine zone of the Lefka Ori, Crete. *Plant Conservation and*
887 *Biodiversity* 6, 29–41.

888 Keeley, J.E., Fotheringham, C.J., Baer-Keeley, M., 2005. Determinants of postfire recovery
889 and succession in Mediterranean-climate shrublands of California. *Ecological*
890 *Applications* 15, 1515–1534.

891 Kéfi, S., Rietkerk, M., Alados, C.L., Pueyo, Y., Papanastasis, V.P., Elaich, A., de Ruiter,
892 P.C., 2007. Spatial vegetation patterns and imminent desertification in Mediterranean arid
893 ecosystems. *Nature* 449, 213–217.

894 Kerdelhué, C., Zane, L., Simonato, M., Salvato, P., Rousset, J., Roques, A., Battistini, A.,
895 2009. Quaternary history and contemporary patterns in a currently expanding species.
896 *BMC Evolutionary Biology* 9, 220-233.

897 Kiss, L., Magnin, F., Torre, F., 2004. The role of landscape history and persistent
898 biogeographical patterns in shaping the responses of Mediterranean land snail
899 communities to recent fire disturbances. *Journal of Biogeography* 31, 145–157.

900 Klein, T., Di Matteo, G., Rotenberg, E., Cohen, S., Yakir, D., 2013. Differential
901 ecophysiological response of a major Mediterranean pine species across a climatic
902 gradient. *Tree Physiology* 33, 26–36.

903 Köchy, M., Mathaj, M., Jeltsch, F., 2008. Resilience of stocking capacity to changing
904 climate in arid to Mediterranean landscapes. *Regional Environmental Change* 8, 73–87.

905 Koulouri, M., Giourga, C., 2007. Land abandonment and slope gradient as key factors of soil
906 erosion in Mediterranean terraced lands. *Catena* 69, 274–281.

907 Kosmas, C., Danalatos, N.G., López-Bermúdez, F., Romero Díaz, M.A., 2002. The effect of
908 land use on soil erosion and land degradation under Mediterranean conditions. In:
909 Geeson, N.A. Brandt, C.J., Thornes, J.B. (Eds.), *Mediterranean Desertification: A Mosaic*
910 *of Processes and Responses*. John Wiley and Sons, Chichester, pp. 57–70.

911 Langley, J.A., Hungate, B.A., 2014. Plant community feedbacks and long-term ecosystem
912 responses to multi-factored global change. *AoB PLANTS* 6: plu035.

913 Latron, J., Llorens, P., Gallart, F., 2009. Hydrology of Mediterranean mountain areas. The
914 case of the Vallcebre research catchments (Eastern Pyrenees, Spain). *Geography*
915 *Compass* 3/6, 2045–2064.

916 Lavorel, S., Canadell, J., Rambal, S., Terradas, J., 1998. Mediterranean terrestrial
917 ecosystems: research priorities on global change effects. *Global Ecology and*
918 *Biogeography Letters* 7, 157–166.

919 Le Houérou, H.N., 1992. Vegetation and land-use in the Mediterranean Basin by the year
920 2050: a prospective study. In: Jeftić, L. Milliman, J.D., Sestini, G. (Eds.), *Climate Change*
921 *and the Mediterranean*. UNEP, Arnold, Boston, pp. 175–232.

922 Legakis, A., Adamopoulou, C., 2005. Temporal responses of soil invertebrate communities
923 to draught stress in two semiarid ecosystems of the Mediterranean. *Israel Journal of*
924 *Zoology* 51, 331–348.

925 Lehner, B., Döll, P., Alcamo, J., Henrichs, T., Kaspar, F., 2006. Estimating the impact of
926 global change on flood and drought risks in Europe: a continental, integrated analysis.
927 *Climate Change* 75, 273–299.

928 Leonardi, S., Gentilesca, T., Guerrieri, R., Ripullone, F., Magnani, F., Mencuccini, M.,
929 Noije, T.V., Borghetti, M., 2012. Assessing the effects of nitrogen deposition and climate
930 on carbon isotope discrimination and intrinsic water-use efficiency of angiosperm and
931 conifer trees under rising CO₂ conditions. *Global Change Biology* 18, 2925–2944.

932 Lespinas, F., Ludwig, W., Heussner, S., 2010. Impact of recent climate change on the
933 hydrology of coastal Mediterranean rivers in Southern France. *Climatic Change* 99, 425–
934 456.

935 Lesschen, J.P., Kok, K., Verburg, P.H., Cammeraat, L.H., 2007. Identification of vulnerable
936 areas for gully erosion under different scenarios of land abandonment in Southeast Spain.
937 *Catena* 71, 110–121.

938 Limousin, J.M., Rambal, S., Ourcival, J.M., Rocheteau, A., Joffre, R., Rodriguez-Cortina,
939 R., 2009. Long-term transpiration change with rainfall decline in a Mediterranean
940 *Quercus ilex* forest. *Global Change Biology* 15, 2163–2175.

941 Lindner, M., Maroschek, M., Netherer, S., Kremer, A., Barbati, A., Garcia-Gonzalo, J.,
942 Seidl, R., Delzon, S., Corona, P., Kolströma, M., Lexer, M.J., Marchetti, M., 2010.
943 Climate change impacts, adaptive capacity, and vulnerability of European forest
944 ecosystems. *Forest Ecology and Management* 259, 698–709.

945 Llorens, P., Domingo, F., 2007. Rainfall partitioning by vegetation under Mediterranean
946 conditions. A review of studies in Europe. *Journal of Hydrology* 335, 37–54.

947 Lloret, F., Estevan, H., Vayreda, J., Terradas, J., 2005. Fire regenerative syndromes of forest
948 woody species across fire and climatic gradients. *Oecologia* 146, 461–468.

949 Lloret, F., Piñol, J., Castellnou, M., 2009. Wildfires. In: Woodward, J. (Ed.), *The Physical*
950 *Geography of the Mediterranean*. Oxford University Press, New York, pp. 541–558.

951 Lloret, F., Escudero, A., Iriondo, J.M., Martínez-Vilalta, J., Valladares, F., 2012. Extreme
952 climatic events and vegetation: the role of stabilizing processes. *Global Change Biology*
953 18, 797–805.

954 Loepfe, L., Martinez-Vilalta, J., Oliveres, J., Piñol, J., Lloret, F., 2010. Feedbacks between
955 fuel reduction and landscape homogenisation determine fire regimes in three
956 Mediterranean areas. *Forest Ecology Management* 259, 2366–2374.

957 Loreto, F., Pinelli, P., Manes, F., Kollist, H., 2004. Impact of ozone on monoterpene
958 emission and evidence for an isoprene-like antioxidant action of monoterpenes emitted by
959 *Quercus ilex* leaves. *Tree Physiology* 24, 361–367.

960 MacDonalD, J.A., Dise, N.B., Matzner, E., Armbruster, M., Gundersen, P., Forsius, M.,
961 2002. Nitrogen input together with ecosystem nitrogen enrichment predict nitrate leaching
962 from European forests. *Global Change Biology* 8, 1028–1033.

963 Malavasi, M., Carboni, M., Cutini, M., Carranza, M.L., Acosta, A.T.R., 2014. Landscape
964 fragmentation, land-use legacy and propagule pressure promote plant invasion on coastal
965 dunes: a patch-based approach. *Landscape ecology* 29, 1541–1550.

966 Malcolm, J.R., Liu, C., Neilson, R.P., Hansen, L., Hannah, L., 2006. Global warming and
967 extinctions of endemic species from biodiversity hotspots. *Conservation Biology* 20, 538–
968 548.

969 Mairota, P., Leronni, V., Xi, W.M., Mladenoff, D.J., Nagendra, H., 2014. Using spatial
970 simulations of habitat modification for adaptive management of protected areas:
971 Mediterranean grassland modification by woody plant encroachment. *Environmental*
972 *Conservation* 41, 144–156.

973 Marlon, J.R., Bartlein, P.J., Carcaillet, C., Gavin, D.G., Harrison, S.P., Higuera, P.E., Joos,
974 F., Power, M.J., Prentice, I.C., 2009. Climate and human influences on global biomass
975 burning over the past two millennia. *Nature Geoscience* 1, 697–702.

976 Martin, P.H., Canham, C.D., Marks, P.L., 2009. Why forests appear resistant to exotic plant
977 invasions: intentional introductions, stand dynamics, and the role of shade tolerance.
978 *Frontiers in Ecology and the Environment* 7, 142–149.

979 Marzano, R., Lingua, E., Garbarino, M., 2012. Post-fire effects and short-term regeneration
980 dynamics following high-severity crown fires in a Mediterranean forest. *iForest* 5, 93–
981 100.

982 Matesanz, S., Escudero, A., Valladares, F., 2009. Impact of three global change drivers on a
983 Mediterranean shrub. *Ecology* 90, 2609–2621.

984 Matías, L., Zamora, R., Castro, J., 2012. Sporadic rainy events are more critical than
985 increasing of drought intensity for woody species recruitment in a Mediterranean
986 community. *Oecologia* 169, 833–44.

987 Matteucci, M., Gruening, C., Ballarin, I.G., Cescatti, A., 2014. Soil and ecosystem carbon
988 fluxes in a Mediterranean forest during and after drought. *Agrochimica* 58: 91–115.

989 McCarthy, M.A., Possingham, H.P., 2007. Active adaptive management for conservation.
990 *Conservation Biology* 21, 956–963.

991 McLaughlin, S.B., Nosal, M., Wullschleger, S.D., Sun, G., 2007. Interactive effects of ozone
992 and climate on tree growth and water use in a southern Appalachian forest in the USA.
993 *New Phytologist* 174, 109–124.

994 MEA, Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being:*
995 *Synthesis*. Island Press, Washington DC, 137 pp.

996 Milne, R., van Oijen, M., 2005. A comparison of two modelling studies of environmental
997 effects on forest carbon stocks across Europe. *Annals of Forest Science* 62, 911–923.

998 Misson, L., Rochetau, A., Rambal, S., Ourcival, J.-M., Limousin, J.-M., Rodriguez, R.,
999 2010. Functional changes in the control of carbon fluxes after 3 years of increased
1000 drought in a Mediterranean evergreen forest? *Global Change Biology* 16, 2461–2475.

1001 Mitsopoulos, I.D., Dimitrakopoulos, A.P., 2007. Canopy fuel characteristics and potential
1002 crown fire behavior in Aleppo pine (*Pinus halepensis* Mill.) forests. *Annals of forest*
1003 *science* 64, 287–299.

1004 Montès, N., Ballini, C., Bonin, G., Faures, J., 2004. A comparative study of aboveground
1005 biomass of three Mediterranean species in a post-fire succession. *Acta Oecologica* 25, 1–
1006 6.

1007 Mooney, H.A., Kalin Arroyo, M.T., Bond, W.J., Canadell, J., Hobbs, R.J., Lavorel, S.,
1008 Neilson, R.P., 2001. Mediterranean-climate ecosystems. In: Chapin III, F.S., Sala, O.E.,

1009 Huber-Sannwald, E. (Eds.), *Global Biodiversity in a Changing Environment: Scenarios*
1010 *for the 21st Century*. Springer-Verlag, New York, pp. 157–198.

1011 Moreira, F., Viedma, O., Arianoutsou, M., Curt, T., Koutsias, N., Rigolot, E., Barbati, A.,
1012 Corona, P., Vaz, P., Xanthopoulos, G., Mouillot, F., Bilgili, E., 2011. Landscape –
1013 wildfire interactions in southern Europe: Implications for landscape management. *Journal*
1014 *of Environment Management* 92, 2389–2402.

1015 Moreno, J.M., Fellous, J.L., 1997. Report of the Enrich/Start International Workshop on
1016 Global change and the Mediterranean Region. Informe Comité IGBP España, Madrid, 78
1017 pp.

1018 Morgan, J.A., Pataki, D.E., Körner, C., Clark, H., Del Grosso, S.J., Grünzweig, J.M., Knapp,
1019 A.K., Mosier, A.R., Newton, P.C., Niklaus, P.A., Nippert, J.B., Nowak, R.S., Parton,
1020 W.J., Polley, H.W., Shaw, M.R., 2004. Water relations in grassland and desert
1021 ecosystems exposed to elevated atmospheric CO₂. *Oecologia* 140, 11–25.

1022 Morin, X., Roy, J., Sonié, L., Chuine, I., 2010. Changes in leaf phenology of three European
1023 oak species in response to experimental climate change. *New Phytologist* 186, 900–910.

1024 Moriondo, M., Good, P., Durao, R., Bindi, M., Giannakopoulos, C., Corte-Real, J., 2006.
1025 Potential impact of climate change on fire risk in the Mediterranean area. *Climate*
1026 *Research* 31, 85–95.

1027 Mouillot, F., Rambal, S., Joffre, R., 2002. Simulating climate change impacts on fire
1028 frequency and vegetation dynamics in a Mediterranean-type ecosystem. *Global Change*
1029 *Biology* 8, 423–437.

1030 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J., 2000.
1031 Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.

1032 Nahm, M., Radoglou, K., Halyvopoulos, G., Geßler, A., Rennenberg, H., Fotelli, M.N.,
1033 2006. Physiological Performance of Beech (*Fagus sylvatica* L.) at its Southeastern

1034 Distribution Limit in Europe: Seasonal Changes in Nitrogen, Carbon and Water Balance.
1035 Plant Biology 8, 52–63.

1036 Naveh, Z., 2007. Conservation, restoration, and research priorities for Mediterranean
1037 uplands threatened by global climate change. In: Moreno, J., Oechel, W.E. (Eds.), Global
1038 Change and Mediterranean–Type Ecosystems. Ecological Studies 117, Springer, New
1039 York, pp. 482–508.

1040 Nicault, A., Alleaume, S., Brewer, S., Carrer, M., Nola, P., Guiot, J., 2008. Mediterranean
1041 drought fluctuation during the last 500 years based on tree-ring data. Climate Dynamics
1042 31, 227–245.

1043 Niedda, M., Pirastru, M., Castellini, M., Giadrossich, F., 2014. Simulating the hydrological
1044 response of a closed catchment-lake system to recent climate and land-use changes in
1045 semi-arid Mediterranean environment. Journal of Hydrology 517, 732–745.

1046 Niinemets, U., 2010. Mild versus severe stress and BVOCs: thresholds, priming and
1047 consequences. Trends in Plant Science 15, 145–153.

1048 Norby, R.J., DeLucia, E.H., Gielen, B., Calfapietra, C., Giardina, C.P., King, J.S., Ledford,
1049 J., McCarthy, H.R., Moore, D.J.P., Ceulemans, R., de Angelis, P., Finzi, A.C., Karnosky,
1050 D.F., Kubiske, M.E., Lukac, M., Pregitzer, K.S., Scarascia-Mugnozza, G.E., Schlesinger,
1051 W.H., Oren, R., 2005. Forest response to elevated CO₂ is conserved across a broad range
1052 of productivity. Proceedings of the National Academy of Sciences USA 102, 18052–
1053 18056.

1054 Noy-Meir, I., Gutman, M., Kaplan, Y., 1989. Responses of Mediterranean grassland plants
1055 to grazing and protection. Journal of Ecology 77, 290–310.

1056 Nunes, J.P., Nearing, M.A., 2011. Modelling impacts of climatic change. In: Morgan R.P.C.,
1057 Nearing, M.A. (Eds.), Handbook of Erosion Modelling. Wiley-Blackwell, Oxford, pp.
1058 289–312.

1059 Ochoa-Hueso, R., Allen, E.B., Branquinho, C., Cruz, C., Dias, T., Fenn, M.E., Manrique, E.,
1060 Pérez-Corona, M.E., Sheppard, L.J., Stock, W.D., 2011. Nitrogen deposition effects on
1061 Mediterranean-type ecosystems: an ecological assessment. *Environmental Pollution* 159,
1062 2265–2279.

1063 Oudin, L., Andréassian, V., Lerat, J., Michel, C., 2008. Has land cover a significant impact
1064 on mean annual streamflow? an international assessment using 1508 catchments. *Journal*
1065 *of Hydrology* 357, 303–316.

1066 Padgett, P.E., Allen, E.B., 1999. Differential responses to nitrogen fertilization in native
1067 shrubs and exotic annuals common to Mediterranean coastal sage scrub of California.
1068 *Plant Ecology* 144, 93–101.

1069 Paoletti, E., 2006. Impact of ozone on Mediterranean forests: a review. *Environmental*
1070 *Pollution* 144, 463–474.

1071 Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J.K., Thomas, C.D., Descimon, H.,
1072 Huntley, B., Kaila, L., Kullberg, J., Tammaru, T., Tennent, W.J., Thomas, J.A., Warren,
1073 M., 1999. Poleward shifts in geographical ranges of butterfly species associated with
1074 regional warming. *Nature* 399, 579–583.

1075 Paula, S., Arianoutsou, M., Kazanis, D., Tavsanoğlu, Ç., Lloret, F., Buhk, C., Ojeda, F.,
1076 Luna, B., Moreno, J.M., Rodrigo, A., Espelta, J.M., Palacio, S., Fernández-Santos, B.,
1077 Fernandes, P.M., Pausas, J. G., 2009. Fire-related traits for plant species of the
1078 Mediterranean Basin. *Ecology* 90, 1420–1420.

1079 Pausas, J.G., 1999. Mediterranean vegetation dynamics: modelling problems and functional
1080 types. *Plant Ecology* 140, 27–39.

1081 Pautasso, M., Dehnen-Schmutz, K., Holdenrieder, O., Pietravalle, S., Salama, N., Jeger,
1082 M.J., Lange, E., Hehl-Lange, S., 2010. Plant health and global change – some
1083 implications for landscape management. *Biological Reviews* 85, 729–755.

1084 Peñuelas, J., Estiarte, M., 1997 Trends in carbon composition and plant demand for N
1085 throughout this century. *Oecologia* 109, 69–73.

1086 Peñuelas, J., Llusà, J., 2001. The complexity of factors driving volatile organic compound
1087 emissions by plants. *Biologia Plantarum* 44, 481–487.

1088 Peñuelas, J., Staudt, M., 2010. BVOCs and global change. *Trends in Plant Science* 15, 133–
1089 144.

1090 Peñuelas, J., Canadell, J., Ogaya, R., 2011. Increased water-use efficiency during the 20th
1091 century did not translate into enhanced tree growth. *Global Ecology and Biogeography*
1092 20, 597–608.

1093 Peñuelas, J., Sardans, J., Rivas-Ubach, A., Janssens, I.A., 2012. The human-induced
1094 imbalance between C, N and P in Earth's life system. *Global Change Biology* 18, 3–6.

1095 Petit, S., Firbank, L., Wyatt and B., Howard, D., 2001. MIRABEL: Models for Integrated
1096 Review and Assessment of Biodiversity in European Landscapes. *A Journal of the*
1097 *Human Environment* 30, 81–88.

1098 Pimentel, C., Calvao, T., Santos, M., Ferreira, C., Neves, M., Nilsson, J., 2006.
1099 Establishment and expansion of a *Thaumetopoea pityocampa* (Den. and Schiff.) (Lep.
1100 Notodontidae) population with a shifted life cycle in a production pine forest, Central-
1101 Coastal Portugal. *Forest Ecology and Management* 233, 108–115.

1102 Pino, J., Arnan, X., Rodrigo, A., Retana, J., 2013. Post-fire invasion and subsequent
1103 extinction of *Conyza* spp. in Mediterranean forests is mostly explained by local factors.
1104 *Weed Research* 53, 470–478.

1105 Piñol, J., Beven, K., Viegas, D., 2005. Modelling the effect of fire-exclusion and prescribed
1106 fire on wildfire size in Mediterranean ecosystems. *Ecological Modelling* 183, 397–409.

1107 Poesen, J.W.A., Hooke, J.M., 1997. Erosion, flooding and channel management in
1108 Mediterranean environments of southern Europe. *Progress in Physical Geography* 21,
1109 157–199.

1110 Poirier, M., Durand, J.L., Volaire, F., 2012. Persistence and production of perennial grasses
1111 under water deficits and extreme temperatures: importance of intraspecific vs.
1112 interspecific variability. *Global Change Biology* 18, 3632–3646.

1113 Polce, C., Kunin, W.E., Biesmeijer, J.C., Dauber, J., Phillips, O.L., The ALARM Field Site
1114 Network, 2011. Alien and native plants show contrasting responses to climate and land
1115 use in Europe. *Global Change and Biogeography* 20, 367–379.

1116 Post, E., Pedersen, C., 2008. Opposing plant community responses to warming with and
1117 without herbivores. *Proceedings of the National Academy of Sciences* 105, 12353–12358.

1118 Preti, F., Forzieri, G., Chirico, G. B., 2011. Forest cover influence on regional flood
1119 frequency assessment in Mediterranean catchments. *Hydrology and Earth System
1120 Sciences* 15, 3077–3090.

1121 Pretto, F., Celesti-Grapow, L., Carli, E., Brundu, G., Blasi, C., 2012. Determinants of non-
1122 native plant species richness and composition across small Mediterranean islands.
1123 *Biological Invasions* 14, 2559–2572.

1124 Puerta-Piñero, C., Espelta, J.M., Sánchez-Humanes, B., Rodrigo, A., Coll, L., Brotons, L.,
1125 2012. History matters: Previous land use changes determine post-fire vegetation recovery
1126 in forested Mediterranean landscapes. *Forest Ecology and Management* 279, 121–127.

1127 Puigdefábregas, J., 1995. Desertification: stress beyond resilience, exploring a unifying
1128 process structure. *Ambio* 24, 311–313.

1129 Puigdefábregas, J., Mendizabal, T., 1998. Perspectives on desertification: western
1130 Mediterranean. *Journal of Arid Environments* 39, 209–224.

1131 Ramankutty, N., Foley, J.A., 1999. Estimating historical changes in global land cover:
1132 croplands from 1700 to 1992. *Global Biogeochemical Cycles* 13, 997–1027.

1133 Ramírez-Valiente, J.A., Sánchez-Gómez, D., Aranda, I., Valladares, F., 2010. Phenotypic
1134 plasticity versus local adaptation for leaf ecophysiological traits in thirteen contrasting
1135 cork oak populations under varying water availabilities. *Tree Physiology* 30, 618–627.

- 1136 Rao, L.E., Allen, E.B., 2010. Combined effects of precipitation and nitrogen deposition on
1137 native and invasive winter annual production in California deserts. *Oecologia* 162, 1035–
1138 1046.
- 1139 Reinhard, M., Rebetz, M., Schlaepfer, R., 2005. Recent climate change: rethinking drought
1140 in the context of Forest Fire Research in Ticino, South of Switzerland. *Theoretical and*
1141 *Applied Climatology* 82, 17–25.
- 1142 Resco de Dios, V., Fischer, C., Colinas, C., 2007. Climate change effects on Mediterranean
1143 forests and preventive measures. *New Forests* 33, 29–40.
- 1144 Retana, J., Espelta, J.M., Habrouk, A., Ordóñez, J.L., de Solà-Morales, F. 2002.
1145 Regeneration patterns of three Mediterranean pines and forest changes after a large
1146 wildfire in northeastern Spain. *Ecoscience* 9, 89–97.
- 1147 Ribas, A., Peñuelas, J., Elvira, S., Gimeno, B.S., 2005. Ozone exposure induces the
1148 activation of leaf senescence-related processes and morphological and growth changes in
1149 seedlings of Mediterranean tree species. *Environmental Pollution* 134, 291–300.
- 1150 Rosenblatt, A.E., Schmitz, O.J., 2014. Interactive effects of multiple climate change
1151 variables on trophic interactions: a meta-analysis. *Climate Change Responses* 1, 8.
- 1152 Ross, L.C., Lambdon, P.W., Hulme, P.E., 2008. Disentangling the roles of climate,
1153 propagule pressure and land use on the current and potential elevational distribution of the
1154 invasive weed *Oxalis pes-caprae* L. on Crete. *Perspectives in Plant Ecology, Evolution*
1155 *and Systematics* 10, 251–258.
- 1156 Roura-Pascual, N., Pons, P., Etienne, M., Lambert, B., 2005. Transformation of a rural
1157 landscape in the Eastern Pyrenees between 1953 and 2000. *Mountain Research and*
1158 *Development* 25, 252–261.
- 1159 Roura-Pascual, N., Bas, J.M., Thuiller, W., Hui, C., Krug, R.M., Brotons, L., 2009. From
1160 introduction to equilibrium: reconstructing the invasive pathways of the Argentine ant in
1161 a Mediterranean region. *Global Change Biology* 15, 2101–2115.

1162 Rutigliano, F.A., Castaldi, S., D'Ascoli, R., Papa, S., Carfora, A., Marzaioli, R., Fioretto, A.,
1163 2009. Soil activities related to nitrogen cycle under three plant cover types in
1164 Mediterranean environment. *Applied Soil Ecology* 43, 40–46.

1165 Sabaté, S., Gracia, C.A., Sánchez, A., 2002. Likely effects of climate change on growth of
1166 *Quercus ilex*, *Pinus halepensis*, *Pinus pinaster*, *Pinus sylvestris* and *Fagus sylvatica*
1167 forests in the Mediterranean region. *Forest Ecology and Management* 162, 23–37.

1168 Safieddine, S., Boynard, A., Coheur, P.-F., Hurtmans, D., Pfister, G., Quennehen, B.,
1169 Thomas, J. L., Raut, J.-C., Law, K. S., Klimont, Z., Hadji-Lazaro, J., George, M.,
1170 Clerbaux, C., 2014. Summertime tropospheric ozone assessment over the Mediterranean
1171 region using the thermal infrared IASI/MetOp sounder and the WRF-Chem model.
1172 *Atmospheric Chemistry and Physics* 14, 10119–10131.

1173 Sala, O.E., Chapin III, F.S., Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-
1174 Sanwald, E., Huenneke, L.F., Jackson, R.B., Kinzig, A., Leemans, R., Lodge, D.M.,
1175 Mooney, H.A., Oesterheld, M., Poff, N.L., Sykes, M.T., Walker, B.H., Walker, M., Wall,
1176 D.H., 2000. Global biodiversity scenarios for the year 2100. *Science* 287, 1770–1774.

1177 Salvati, L., Ranalli, F., Gitas, I., 2014. Landscape fragmentation and the agro-forest
1178 ecosystem along a rural-to-urban gradient: an exploratory study. *International Journal of*
1179 *Sustainable Development and World Ecology* 21, 160-167.

1180 Sánchez-Humanes, B., Espelta, J.M., 2011. Increased drought reduces acorn production in
1181 *Quercus ilex* coppices: thinning mitigates this effect but only in the short term. *Forestry*
1182 84, 73–82.

1183 Santos, H., Paiva, M.R., Tavares, C., Kerdelhué, C., Branco, M., 2011. Temperature niche
1184 shift observed in a Lepidoptera population under allochronic divergence. *Journal of*
1185 *Evolutionary Biology* 24, 1897–1905.

1186 Sarris, D., Christodoulakis, D., Körner, C., 2007. Recent decline in precipitation and tree
1187 growth in the eastern Mediterranean. *Global Change Biology* 13, 1187–1200.

1188 Scalerio, S. 2009. On top of a Mediterranean Massif: Climate change and conservation of
1189 orophilous moths at the southern boundary of their range (Lepidoptera: Macroheterocera).
1190 European Journal of Entomology 106, 231–239.

1191 Scarascia-Mugnozza, G., Oswald, H., Piussi, P., Radoglou, K., 2000. Forests of the
1192 Mediterranean region: gaps in knowledge and research needs. Forest Ecology and
1193 Management 132, 97–109.

1194 Scherber, C., 2015. Insect responses to interacting global change drivers in managed
1195 ecosystems. Current Opinion in Insect Science 11, 56–62.

1196 Schröter, D., Cramer, W., Leemans, R., Prentice, I.C., Araújo, M.B., Arnell, N.W., Bondeau,
1197 A., Bugmann, H., Carter, T.R., Gracia, C.A., de la Vega-Leinert, A.C., Erhard, M., Ewert,
1198 F., Glendining, M., House, J.I., Kankaanpää, S., Klein, R.J.T., Lavorel, S., Lindner, M.,
1199 Metzger, M.J., Meyer, J., Mitchell, T.D., Reginster, I., Rounsevell, M., Sabaté, S., Sitch,
1200 S., Smith, B., Smith, J., Smith, P., Sykes, M.T., Thonicke, K., Thuiller, W., Tuck, G.,
1201 Zaehle, S., Zierl, B., 2005. Ecosystem service supply and vulnerability to global change
1202 in Europe. Science 310, 1333–1337.

1203 Seixas Arnaldo, P., Oliveira, I., Santos, J., Leite, S., 2011. Climate change and forest
1204 plagues: the case of the pine processionary moth in Northeastern Portugal. Forest Systems
1205 20, 508–515.

1206 Seligman, N.G., Henkin, Z., 2000. Regeneration of a dominant Mediterranean dwarf-shrub
1207 after fire. Journal of Vegetation Science 11, 893–902.

1208 Shakesby, R., 2011. Post-wildfire soil erosion in the Mediterranean: Review and future
1209 research directions. Earth-Science Reviews 105, 71–100.

1210 Sherman, C., Sternberg, M., Steinberger, Y., 2012. Effects of climate change on soil
1211 respiration and carbon processing in Mediterranean and semi-arid regions: An
1212 experimental approach. European journal of Soil Biology 52, 48–58.

1213 Simoes, M.P., Madeira, M., Gazarini, L., 2008. The role of phenology, growth and nutrient
1214 retention during leaf fall in the competitive potential of two species of Mediterranean
1215 shrubs in the context of global climate changes. *Flora* 203, 578–589.

1216 Sirami, C., Brotons, L., Burfield, I., Fonderflick, J., Martin, J. 2008. Is land abandonment
1217 having an impact on biodiversity? A meta-analytical approach to bird distribution changes
1218 in the north-western Mediterranean. *Biological Conservation* 141, 450–459.

1219 Sitch, S., Cox, P.M., Collins, W.J., Huntingford, C., 2007. Indirect radiative forcing of
1220 climate change through ozone effects on the land-carbon sink. *Nature* 448, 791–794.

1221 Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M.,
1222 Miller, H.L., 2007. *Climate Change 2007: The Physical Science Basis*. Working Group I
1223 Contribution to the Fourth Assessment Report of the Intergovernmental Panel on Climate
1224 Change. Cambridge University Press, Cambridge, 996 pp.

1225 Speed, J.D.M., Austrheim, G., Hester, A.J., Mysterud, A., 2010. Experimental evidence for
1226 herbivore limitation of the treeline. *Ecology* 91, 3414–3420.

1227 Stavros, E.N., McKenzie, D., Larkin, N., 2014. The climate-wildfire-air quality system:
1228 interactions and feedbacks across spatial and temporal scales. *Wiley Interdisciplinary*
1229 *Reviews-Climate Change* 5, 719–733.

1230 Steffen, W., Sanderson, A., Tyson, P.D., Jäger, J., Matson, P.A., Moore III, B., Oldfield, F.,
1231 Richardson, K., Schellnhuber, H.J., Turner II, B.L., Wasson, R.J., 2004. *Global Change*
1232 *and the Earth System: A Planet Under Pressure*. Springer-Verlag, Berlin, Heidelberg,
1233 New York, 332 pp.

1234 Steinbrecher, R., Smiatek, G., Köble, R. Seufert, G., Theloke, J., Hauff, K., Ciccioli, P.,
1235 Vautard, R., Curci, G., 2009. Intra- and inter-annual variability of VOC emissions from
1236 natural and seminatural vegetation in Europe and neighbouring countries. *Atmospheric*
1237 *Environment* 43, 1380–1391.

- 1238 Sternberg, M., Yakir, D., 2015. Coordinated approaches for studying long-term ecosystem
1239 responses to global change. *Oecologia* 177, 921–924.
- 1240 Tessler, N., Wittenberg, L., Provizor, E., Greenbaum, N., 2014. The influence of short-
1241 interval recurrent forest fires on the abundance of Aleppo pine (*Pinus halepensis* Mill.) on
1242 Mount Carmel, Israel. *Forest ecology and management* 324, 109–116.
- 1243 Thornes, J.B., 2005. Coupling erosion, vegetation and grazing. *Land Degradation and*
1244 *Development* 16, 127–138.
- 1245 Thornes, J.B., 2009. Land degradation. In: Woodward, J. (Ed.), *The Physical Geography of*
1246 *the Mediterranean*. Oxford University Press, New York, pp. 563–581.
- 1247 Thornes, J.B., Brandt, C.J., 1994. Erosion-vegetation competition in a stochastic
1248 environment undergoing climatic change. In: Millington, A.C., Pye, K. (Eds.),
1249 *Environmental change in drylands: biogeographical and geomorphological perspectives*.
1250 John Wiley and Sons Ltd., Chichester, pp. 305–320.
- 1251 Tomaz, C., Alegria, C., Monteiro, J.M., Teixeira, M.C., 2013. Land cover change and
1252 afforestation of marginal and abandoned agricultural land: A 10 year analysis in a
1253 Mediterranean region. *Forest Ecology and Management* 308, 40–49.
- 1254 Tsiafouli, M.A., Kallimanis, A.S., Katana, E., Stamou, G.P., Sgardelis, S.P., 2005.
1255 Responses of soil microarthropods to experimental short-term manipulations of soil
1256 moisture. *Applied Soil Ecology* 29, 17–26.
- 1257 Tsitsoni, T. 1997. Conditions determining natural regeneration after wildfires in the *Pinus*
1258 *halepensis* (Miller, 1768) forests of Kassandra Peninsula (North Greece). *Forest Ecology*
1259 *and Management* 92, 199–208.
- 1260 Vacchiano, G., Stanchi, S., Marinari, G., Ascoli, D., Zanini, E., Motta, R., 2014. Fire
1261 severity, residuals and soil legacies affect regeneration of Scots pine in the Southern Alps.
1262 *Science of the Total Environment* 472, 778–788.

1263 Valentini, R., Matteucci, G., Dolman, A.J., Schulze, E.D., Rebmann, C., Moors, E.J.,
1264 Granier, A., Gross, P., Jensen, N.O., Pilegaard, K., Lindroth, A., Grelle, A., Bernhofer,
1265 C., Grünwald, T., Aubinet, M., Ceulemans, R., Kowalski, A.S., Vesala, T., Rannik, U.,
1266 Berbigier, P., Loustau, D., Gudmundsson, J., Thorgeirsson, H., Ibrom, A., Morgenstern,
1267 K., Clement, R., 2000. Respiration as the main determinant of carbon balance in
1268 European forests. *Nature* 404, 861–865

1269 Valladares, F., Zaragoza-Castells, J., Sánchez-Gómez, D., Matesanz, S., Alonso, B.,
1270 Portsmouth, A., Delgado, A., Atkin, O.K., 2008. Is shade beneficial for Mediterranean
1271 shrubs experiencing periods of extreme drought and late-winter frosts? *Annals of Botany*
1272 102, 923–933.

1273 Vega, J.M., Moneo, I., Armentia, A., Vega, J., de la Fuente, R., Fernandez, A., 2000. Pine
1274 processionary caterpillar as a new cause of immunologic contact urticaria. *Contact*
1275 *Dermatitis* 43, 129–132.

1276 Vennetier, M., Ripert, C., 2009. Forest flora turnover with climate change in the
1277 Mediterranean region: A case study in Southeastern France. *Forest Ecology and*
1278 *Management* 258, S56–S63.

1279 Vieira, J., Campelo, F., Nabais, C., 2010. Intra-annual density fluctuations of *Pinus pinaster*
1280 are a record of climatic changes in the western Mediterranean region. *Canadian Journal of*
1281 *Forest Research* 40, 1567–1575.

1282 Vilà, M., Lloret, F., Ogheri, E., Terradas, J., 2001. Positive firegrass feedback in
1283 Mediterranean basin shrublands. *Forest Ecology and Management* 147, 3–14.

1284 Vilà, M., Tessier, M., Suehs, C.M., Brundu, G., Carta, L., Galanidis, A., Lambdon, P.,
1285 Manca, M., Médail, F., Moragues, E., Traveset, A., Troumbis, A.Y., Hulm, P.E., 2006.
1286 Local and regional assessment of the impacts of plant invaders on vegetation structure
1287 and soil properties of Mediterranean islands. *Journal of Biogeography* 33, 853–861.

1288 Vilà-Cabrera, A., Rodrigo, A., Martínez-Vilalta, J., Retana, J., 2012. Lack of regeneration
1289 and climatic vulnerability to fire of Scots pine may induce vegetation shifts at the
1290 southern edge of its distribution. *Journal of Biogeography* 39, 488-496.

1291 Vitousek, P.M., 1994. Beyond global warming: ecology and global change. *Ecology* 75,
1292 1861–1876.

1293 Vonshak, M., Dayan, T., Ionescu-Hirsh, A., Freidberg, A., Hefetz, A., 2010. The little fire
1294 ant *Wasmannia auropunctata*: a new invasive species in the Middle East and its impact
1295 on the local arthropod fauna. *Biological Invasions* 12, 1825–1837.

1296 Waldboth, M., Oberhuber, W., 2009. Synergistic effect of drought and chestnut blight
1297 (*Cryphonectria parasitica*) on growth decline of European chestnut (*Castanea sativa*).
1298 *Forest Pathology* 39, 43–55.

1299 Walther, G.-R., Roques, A., Hulme, P.E., Sykes, M.T., Pysek, P., Kühn, I., Zobel, M.,
1300 Bacher, S., Botta-Dukát, Z., Bugmann, H., Czúcz, B., Dauber, J., Hickler, T., Jarosík, V.,
1301 Kenis, M., Klotz, S., Minchin, D., Moora, M., Nentwig, W., Ott, J., Panov, V.E.,
1302 Reineking, B., Robinet, C., Semchenko, V., Solarz, W., Thuiller, W., Vilà, M.,
1303 Vohland, K., Settele, J., 2009. Alien species in a warmer world: risks and opportunities.
1304 *Trends in Ecology and Evolution* 24, 686–693.

1305 Williams, D.W., Liebhold, A.M., 1995. Herbivorous insects and global change: potential
1306 changes in the spatial distribution forest defoliator of outbreaks. *Journal of Biogeography*
1307 22, 665–671.

1308 Wittenberg, L., Kutiel, H., Greenbaum, N., Inbar, M., 2007. Short-term changes in the
1309 magnitude, frequency and temporal distribution of floods in the Eastern Mediterranean
1310 region during the last 45 years — Nahal Oren, Mt. Carmel, Israel. *Geomorphology* 84,
1311 181–191.

1312 Wittig, V., Ainsworth, E.A., Naiduz, S.L., Karnosky, D., Long, S.P., 2009. Quantifying the
1313 impact of current and future tropospheric ozone on tree biomass, growth, physiology and
1314 biochemistry: a quantitative meta-analysis. *Global Change Biology* 15, 396–424.

1315 Woodward, J., 2009. *The Physical Geography of the Mediterranean*. Oxford University
1316 Press, New York, 700 pp.

1317 Zaimeche, S.E., 1994. The consequences of rapid deforestation: a North African example.
1318 *Ambio* 23, 136–140.

1319 Zedler, P.H., 1995. Fire frequency in southern California shrublands: biological effects and
1320 management options. In: Keeley, J.E., Scott, T. (Eds.), *Brushfires in California*
1321 *Wildlands: Ecology and Resource Management*. International Association of Wildland
1322 Fire, Fairfield, Wash, pp. 101–112.

1323 Zhao, M., Running, S.W., 2010. Drought-induced reduction in global terrestrial net primary
1324 production from 2000 through 2009. *Science* 329, 940–943.

1325

1326 Table 1. Principal effects derived from the combinations between global change factors in
 1327 the Mediterranean Basin region. Shaded cells correspond to repeated combinations and
 1328 combinations of the same factor (including land-use intensification and land abandonment as
 1329 the two opposite means of land-use change). As different pollutants could interact among
 1330 them, these same factor interactions are explained in the first section of the manuscript
 1331 together with other atmospheric chemical alterations. Numbered combinations are explained
 1332 in the second section of the manuscript.

1333

	Drought and other climatic events	Alteration of atmospheric composition	Land use intensification	Land abandonment	Wild fires
Alteration of atmospheric composition	Atmospheric alteration increase 1 Modification of plant ecophysiology	Interactions among pollutants			
Land use intensification	2 Alteration of water resources 3 Land degradation 4 Regeneration decline 5 Disease expansion 6 Increase of fire risk	Atmospheric alteration increase			
Land abandonment	2 Alteration of water resources 3 Land degradation	Atmospheric alteration increase			
Wild fires	3 Land degradation 4 Regeneration decline 6 Increase of fire risk	Atmospheric alteration increase	6 Increase of fire risk	6 Increase of fire risk	
Biological invasions	7 Increase of invasion risk	7 Increase of invasion risk	7 Increase of invasion risk	7 Increase of invasion risk	7 Increase of invasion risk

1334

1335

1336 **List of figure legends**

1337

1338 Figure 1. Results for the Mediterranean Basin from time-series analysis of Landsat 7 ETM+
1339 images in characterizing global forest extent and change from 2000 through 2012 (Hansen et
1340 al., 2013). Dark grey: forest cover in 2000; black: gain forest from 2000 to 2012; white:
1341 forest lost from 2000 to 2012. It is difficult to appreciate forest gain and losses due to the
1342 scattered nature of the process in the Region although lower scales could be accessed in the
1343 original webpage: <http://earthenginepartners.appspot.com/science-2013-global-forest>.

1344

1345 Figure 2. Types of combination among global change factors. Solid arrows represent
1346 positive effects while shaded arrows represent negative effects. Some interactions alter the
1347 effects of a single factor (a), as for example CO₂ increase affects drought effects on plant
1348 growth through stomatal closure. New possible impacts can be caused by the interaction (b),
1349 such as the expansion of forest pests caused by the alteration of forest structure and climate
1350 warming. Finally, other combinations cause an increase in the risk of one of the factors
1351 implied (c) such as fire, land-use change, N deposition and climate change effects on
1352 invasion.

1353

1354 Figure 3. Combined effects of land-use intensification and abandonment, fire and drought on
1355 soil erosion and water availability. Solid lines represent positive effects while dashed lines
1356 represent negative effects.

1357