

A Review of the Promotion of Fitness Measures and Health Outcomes in Youth

Abstract: *The relationship between physical fitness and health in adults is well established, yet until recently, empirical support for the impact of physical fitness levels on health markers in children has not been thoroughly documented. The aim of this review was to provide a summary of current literature that has examined the linkage between measures of health-related physical fitness (ie, cardiorespiratory fitness, musculoskeletal fitness, flexibility, and body composition) and health markers in youth. Specifically, this review focused on the findings from the recent 2012 Institutes of Medicine report on Fitness Measures in Youth as well as other subsequent review articles related to the topic. In addition, recommendations for health-related fitness assessments in youth populations are presented.*

Keywords: youth; health markers; health-related fitness; assessments

Introduction

Promoting physical fitness in youth in the United States has been a priority for decades. In 1955, President Dwight D. Eisenhower was prompted to create the

Council for Youth Fitness in reaction to the noted decline in children's physical fitness.¹ President John F. Kennedy's efforts were also vital to improving the health and well-being of America's youth by promoting the development of youth fitness standards. The first National Survey of Youth Fitness was conducted in 1958, with additional surveys conducted in 1965, 1975, and finally in

lifelong health. Interestingly, although the proposed link between fitness testing and health markers/outcomes in youth drove the changes in fitness testing over the decades, this link was never formally established based on research. It is also important to note that secular trends over the past few decades demonstrate decreased health-related fitness levels in youth across the world.³⁻⁵ This decline

 It is also important to note that secular trends over the past few decades demonstrate decreased health-related fitness levels in youth across the world. 

1985-1986. Thus, a 30-year span has passed since the last national survey of youth fitness was conducted. Initial youth fitness standards were based on performance in relation to standards of military preparedness.² Over time, there was a shift away from performance-related fitness and toward measures of health-related fitness testing because these assessments were thought to be better predictors and facilitators of

reiterates the importance of linking aspects of physical fitness to health in youth. In addition, demonstrating a link between fitness levels in youth and adult fitness is important to promote continued health across the lifespan.⁶⁻⁸

The noted decreases in fitness and physical activity levels and increased obesity in youth over the past few decades prompted the Institutes of Medicine (IOM) to convene a committee

DOI: 10.1177/1559827615619577. Manuscript received September 9, 2015; revised October 23, 2015; accepted November 6, 2015. From the Department of Physical Education and Athletic Training, University of South Carolina, Columbia, South Carolina. Address correspondence to: Danielle Nesbitt, MAT, Department of Physical Education and Athletic Training, University of South Carolina, 1300 Wheat Street, Columbia, SC 29201; e-mail: nesbitdr@email.sc.edu.

For reprints and permissions queries, please visit SAGE's Web site at <http://www.sagepub.com/journalsPermissions.nav>.

Copyright © 2015 The Author(s)

to examine Fitness Measures and Health Outcomes in Youth. The charge of this committee was to review the latest advancements in research related to fitness testing in youth and examine the extent to which health-related fitness is linked to health markers/outcomes in youth. In examining the scientific literature, the committee evaluated the relationships between individual fitness components and health markers/outcomes (eg, cardiovascular disease risk factors, musculoskeletal health, diabetes, obesity, and others).⁹ The committee structured their report into 4 classifications of fitness recommendations, which are addressed in the following subsections: (1) body composition, (2) flexibility, (3) cardiorespiratory endurance, (4) musculoskeletal fitness. In addition to the publication in 2012 of the IOM report on Fitness Measures in Youth,¹⁰ additional reviews related to this topic¹¹⁻¹³ have been published, and their findings also have been noted in this report.

Body Composition

Body composition is a well-established measure of health in youth and adults.¹⁴⁻¹⁶ It is defined as a physiological characteristic that affects an individual's ability to carry out daily tasks with vigor. For the purpose of the IOM review, the committee defined body composition as a marker of health but also noted it as a component of fitness and a modifier of fitness. After an extensive review of fitness test measures related to body composition in youth, the committee identified 10 health-related fitness test batteries (out of a total of 15 in the literature) that used body composition as a component of health-related fitness.

Methods for assessing body composition have generally been developed for adults and have consisted of separating body mass into several elements: fat-free mass, fat mass, total body water, fat-free dry mass, and bone mineral content. Over the past few decades, technology has allowed researchers to analyze body composition more comprehensively. Laboratory tests

such as hydrostatic weighing and dual-energy X-ray absorptiometry (DXA) are considered to be highly accurate forms of measuring whole body composition and reduce measurement variations. For the purpose of this review, the methodology used for each test will not be discussed in depth. However, these tests require specific training and expensive equipment. Thus, the use of field-based assessments (eg, skinfolds, body mass index [BMI], and waist circumference), which are noted as valid and reliable in youth, have been preferred for large-scale population-based studies.

The literature indicates that body composition status affects physiological and psychological health in youth in several different ways. Obesity in youth is associated with cardiovascular risk factors such as high blood pressure and abnormalities in left ventricular mass and/or function.¹⁷ Other health consequences such as diabetes and asthma have been linked to the rise in an individual's weight status.¹⁸ Children who are classified as obese also are more likely to exhibit more psychological or psychiatric problems than those children who are nonobese.¹⁸ Specifically, increased BMI is associated with lower self-perceptions of social acceptance and physical appearance.¹⁹⁻²¹ Children who demonstrate low perceptions of self-concept also are generally less willing to participate in activities with peers.²²⁻²⁴ Deficits in cognitive function also have been associated with obesity in children as well as adults.²⁵

Body composition status was also considered a modifying factor of fitness by the committee and independently affects an individual's current level of fitness and the continued development of fitness. Being obese promotes a negative cycle of global fitness as individuals with excess body mass have decreased cardiorespiratory fitness and decreased muscular endurance.^{26,27} Thus, these individuals are not able to persist in health-enhancing physical activities for as long as other individuals who are more fit. In addition, being obese limits an individual's functional capability to

successfully perform many activities that require the manipulation of one's entire body mass,²⁸⁻³³ which again, decreases time spent in physical activity. Overall, obesity in early childhood perpetuates negative developmental trajectories of fitness and physical activity across middle and late childhood^{26-28,30,32,33} and continued obesity into adolescence and adulthood.³⁴⁻³⁶

When assessing body composition across various ages in youth, various moderating factors such as age, sex, and maturation status influence body composition status and should be considered when interpreting the results. Variations in fat-free mass, fatness, and relative fat distribution in late childhood and adolescence also affect body composition status³⁷ and are important when interpreting results. In addition, the influences of these moderating factors vary by sex. During childhood, an individual has not reached maturity at the cellular level; thus, age differences that coincide with natural growth patterns between boys and girls makes it difficult to adequately assess a person's body composition across childhood.^{38,39} Fat mass and fat-free mass are factors that also are affected by age and sex. For girls, fat mass increases more rapidly than for boys during childhood, and this trend continues throughout adolescence. Thus, sex differences create a dispersion of body composition among adolescents.⁴⁰ Other important variations in body composition that are influenced by age and sex include weight-for-height, subcutaneous fat, and relative fat distribution.⁴¹⁻⁴³

Race and ethnicity also affect the interpretation of body composition assessments. Gender differences among ethnicities are similar; however, elements such as fat-free mass and bone mineral content differ from ethnicity to ethnicity.^{44,45} For example, American Indian/Native Alaskan children tend to demonstrate the highest prevalence of obesity, followed by Hispanic and non-Hispanic black children.⁴⁶ Overall, differences in body composition status may be affected by many modifying factors.

Selection of Test

The final task of the committee was to choose the most appropriate measures of body composition for youth populations. Taking into consideration issues with feasibility, cost, training, and validity, the IOM committee recommended 3 measures of body composition that should be used for testing youth on a population-based scale. These recommendations also were based on their relationship to health markers in youth. The tests include skinfold, BMI, and waist circumference assessments.

Sum of skinfolds are considered valid and reliable estimates of subcutaneous fat and predictors of percentage body fat in youth.^{47,48} Skinfolds have been measured using several different sites and have been standardized to ensure accuracy. When assessed correctly, the skinfold test is used to estimate the relative subcutaneous fat distribution of the trunk and extremities. The committee recommends the use of the triceps and the subscapular skinfold thickness body composition measurement in youth.^{10,49,50}

BMI is an indicator of mass for height and is the most common assessment for weight status used in youth. It is reasonably well correlated with fat mass and percentage body fat in heterogeneous samples of youth.⁵¹ However, this measure has several limitations. It is important to understand that individuals with the same BMI can differ in both fat mass and percentage body fat.¹⁶ This being said, BMI is suggested to be more appropriately used as an indicator of weight status and has the potential to be an indicator of fatness in the general population.⁵²⁻⁵⁴ Because of the sensitivity of the educational setting, the committee only recommends this measurement to be used in the school setting.

Waist circumference has been shown to be a valid measure of central adiposity and a good predictor of body fat distribution in youth.⁵⁵⁻⁵⁷ Studies have shown that waist circumference is strongly associated with abdominal adiposity in children.⁵⁸ Waist circumference also is noted to be a good indicator of abdominal adiposity, which

provides information about different dimensions of body composition that are linked to health risks⁵⁹; however, waist circumference may not be the best indicator for percentage body fat or fatness in youth.

Conclusions

The committee recommends that the interpretations of results of the above-noted body composition measures be approached in 2 ways. First, the committee recommends the use of the Centers for Disease Control's (CDC's) currently established cut-points for underweight, overweight, and obesity. The 2000 CDC BMI-for-age charts are widely used and recommended for children 2 to 20 years old in the United States.^{16,60} For the remaining 2 measurements, the cut-points should be set at levels that are analogous to those currently being applied by the CDC for BMI.

Flexibility

Flexibility is defined as the range of motion of muscle and connective tissues at a joint or group of joints. Previous literature examining the association between flexibility and any health outcome in adults has been unclear, and research addressing the association of flexibility and health-related markers in youth is limited. Similar to the literature dealing with adults, the links between flexibility and other outcomes in youth are difficult to explain. Secular trends have demonstrated that youth in the present day are less flexible than those in the 1980s.¹⁰ However, these data have not led to the establishment of a link to health markers.

The American College of Sports Medicine suggests that flexibility exercises may enhance postural stability, balance, and functional capacity⁶¹; however, research results have been inconclusive.¹⁰ Several studies with adults who have health issues have demonstrated improvements in function and pain, but very few have focused on changes in flexibility. Complicating the association between flexibility and health

outcomes, research examining flexibility and function in older populations are hard to generalize because of the individual's current physical status.

A literature search screened a total of 6016 studies that addressed some aspect of flexibility in youth. Because of limited resources and time, the CDC did not abstract those articles. Because of the lack of specific evidence, the IOM presented findings from studies conducted in youth regardless of the quality of evidence that supported a relationship between flexibility and health outcomes. Based on the 16 studies included, results demonstrated that programs designed for physical activity in youth improved flexibility even though it was not an intended outcome.⁶²⁻⁶⁴ In addition, research suggests that youth who are less active may have a higher risk for low-back pain, but a direct link to this potential health-related issue has not been established.⁶⁵

A total of 7 studies examined the relationship between flexibility and pain outcomes, but only 1 study directly linked health outcomes that are commonly associated with flexibility. Similar to body composition, differences in flexibility were noted between sex and ethnicities among youth. Women generally are more flexible than men,⁶⁶⁻⁶⁸ and longitudinal data demonstrate a decrease in flexibility scores in both sexes based on data collected in the 1980s. One recent longitudinal study indicated that girls actually improved their sit-and-reach scores over a 6-year period, whereas boys' performance decreased.⁶⁹

Selection of Test

Flexibility tests measure joint range of motion (ROM) and generally can be classified into laboratory test and field tests. Unlike other fitness components, flexibility is systemic in nature, yet assessments generally are specific to individual joints. The specificity of flexibility for individual joints of the body makes it difficult to isolate a single flexibility-related factor contributing to health outcomes. In lieu of these issues,

the most common fitness batteries have generally used 3 flexibility tests: shoulder stretch, trunk lift, and sit-and-reach tests. The sit-and-reach test, which is the most common test used in youth, has been used to assess low-back and hip flexibility; however, it has been shown to be more of an indicator of hip flexibility rather than low-back flexibility.¹⁰

Conclusions

Although validity and reliability of some of the flexibility tests in youth have been established, the degree to which a test, such as the sit-and-reach, is an indicator of overall flexibility is still unclear. Unlike other variables of health-related fitness, flexibility does not relate to other health outcomes in a linear fashion. Because of this lack of evidence, the committee felt that the flexibility component is not a priority when assessing fitness. However, it should be considered when conducting fitness assessments as a whole because of its potential link to health in adulthood. These tests can be used to educate youth as well as parents about their child's overall fitness level.

Cardiorespiratory Endurance

Cardiorespiratory endurance has been noted as a cornerstone of physical fitness throughout history and is defined as the ability to perform large-muscle, whole-body exercise at a moderate to high intensity for an extended period of time.⁷⁰ High levels of cardiorespiratory endurance are associated with a wide range of health benefits in adults, including a lower risk of cardiovascular disease,⁷¹⁻⁷³ type 2 diabetes,⁷⁴ hypertension,⁷⁵ certain cancers,⁷⁶ and premature mortality from all causes.^{7,8,72} Although health outcomes resulting from the development of cardiorespiratory endurance in adults are well known,¹² relationships between cardiorespiratory endurance and health outcomes in youth are not as well understood.

For the purposes of the IOM report, 4795 studies were initially identified, of whom 260 fit criteria for further

consideration. There were 56 experimental, 24 longitudinal prospective, and 29 quasi-experimental studies that were considered high-quality studies and were provided foundational data for the report. The following criteria were utilized to determine high-quality studies: the study must have (1) provided a link between a candidate measure of cardiorespiratory endurance to a positive health outcome; (2) provided a marker or risk factor in either adiposity, metabolic risk, cognitive, or "other"; and (3) presented direct or associational evidence to a health risk factor or outcome. Studies were excluded for the following reasons: poor design, inappropriate population and lack of power, inability to assess the independent effect of dietary intervention, or an insufficient change in the fitness measure of interest.

Overall, the committee concluded that sufficient evidence has established a link between cardiorespiratory endurance and health in youth. Specifically, studies showed linkages between high levels of cardiorespiratory endurance and a wide range of health benefits: (1) adiposity,⁷⁶⁻⁸⁰ (2) blood pressure,⁸¹⁻⁸³ (3) blood lipids,⁸³⁻⁸⁵ (4) glucose,⁸⁶ and (5) insulin sensitivity.^{85,87} Some studies also found a relationship with other pediatric health risk factors, including pulmonary function, depression, a positive self-concept, and bone health. However, the standards for noting risk in each category were not as well established.^{80,88,89}

Importantly, the following benefits of cardiorespiratory endurance are continued into adulthood: decreased risk of cardiovascular disease,⁷¹⁻⁷³ type 2 diabetes,^{74,70} certain cancers,⁷⁶ and premature all-cause mortality.^{7,8,72} It was noted that youth in the lowest 20th percentile of cardiorespiratory endurance demonstrate the highest risk for cardiovascular-related negative health outcomes. The recommendation to use the 20th percentile as a cut-point for determining health risk was derived from its association with morbidity and mortality throughout the lifespan.⁷² The 20th percentile is more conservative than the 30th percentile, as suggested by

Lobelo et al⁹⁰ and Welk et al,⁹¹ and is recommended for use based on the viewpoint that identifying a fit individual as low fit is a greater error than concluding that an individual who is low fit is fit.

Selection of Tests

Cardiorespiratory endurance capability is the greatest rate at which a person is able to consume oxygen during sustained, exhaustive exercise. The tests noted in the literature that most closely related to this definition and that have shown the greatest association with a positive change in health markers in youth were the shuttle run, treadmill, and cycle ergometer tests. It is important to note that although traditional distance-based fitness tests (eg, 1 or 2 mile runs) have a long history,^{86,88,92} there was limited evidence linking these types of tests to health markers in youth.^{49,86,88,92,93}

Shuttle run testing protocols noted in the literature included the 6-minute, 20-m shuttle run and various 20-m multistage incremental tests. The majority of associations to health were with adiposity (measured BMI). Other factors such as blood lipids were noted in some studies, but their correlations were limited because of the location of the studies, most of which took place in a school setting.⁹⁴⁻⁹⁷ Improvements in bone health, blood pressure, BMI, and blood lipids also were related to increased cardiorespiratory endurance over time increments of 1 to 5 years. The shuttle run has proven to be a safe test that can be effectively administered with an understanding of the procedures for testing and safety. For example, Ruiz et al⁹⁸ have reported testing more than 10 000 children, with no safety issues occurring during testing. The 20-m shuttle run (eg, PACER) demonstrates adequate reliability ($r = 0.78$ to 0.93)⁹⁹ and validity ($r = 0.87$ and 0.72)^{100,101} and is the protocol used most often in the literature when compared against VO_{2max} performance.

Treadmill maximal and submaximal endurance testing protocols demonstrate decreased adiposity (skinfold and DXA), BMI, and waist circumference as well as an increase in VO_{2max} with increased test

performance.⁷⁶⁻⁷⁹ Treadmill testing protocols also have proven themselves to be safe and reliable and require a minimal amount of space to perform. Most notably, the effectiveness of treadmill protocols was shown in a longitudinal study by Eisenmann et al,⁷⁶ spanning adolescence to adulthood (mean age = 26 years), where the time to reach exhaustion on the treadmill in adolescence was correlated with both adult body fatness and changes in body fatness from adolescence to adults. Treadmill testing also has demonstrated strong validity with other criterion tests ($r = 0.84$)¹⁰⁰ and is highly reliable, with test-retest correlation coefficients ranging from 0.89 to 0.98.^{102,103}

Strong evidence associated cardiorespiratory endurance with adiposity, cardiometabolic risk factors, and blood pressure.^{85,87,89,104-108} Specifically, an improvement in cardiorespiratory endurance as well as positive changes in total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, and adiposity measures were noted in youth tracked over a 6-year period, spanning from childhood to adolescence.^{106,108} Other studies also noted an improvement in depression and positive self-concept that were mitigated by increased fitness.^{89,105}

Limited data have addressed the role of several potential important modifiers that may have an impact on the effect of cardiorespiratory endurance on health. These factors include age, gender, race, body composition, and maturation status. Accounting for maturation stage,^{77,78} specific ethnicities and gender, and the impact of baseline fitness are important when considering cardiorespiratory fitness levels and their relationship to changes in health markers. Overall, the strongest associated evidence of the effect of independent moderating factors for cardiorespiratory endurance and health-related risk factors was adiposity. Further research is warranted to better understand the role of these modifiers on the relationship between cardiorespiratory endurance and health risk factors.

Conclusion

Based on the relationship to health markers and also feasibility for testing in youth, a timed or progressive shuttle run, such as the 20-m shuttle run or PACER test is most appropriate for measuring cardiorespiratory endurance in youth. The submaximal treadmill or cycle ergometer tests were noted as feasible alternative tests if testing limitations did not allow the PACER test.

Musculoskeletal Fitness

Musculoskeletal fitness, as defined by the IOM committee, is “a multidimensional construct comprising the integrated function of muscle strength, muscle endurance, and muscle power to enable the performance of work against one’s own body weight or an external resistance. (p. 155)”¹⁰ Each dimension of musculoskeletal fitness, although integrated, should be assessed independently against health markers and then interpreted collectively to provide an overall understanding of its impact on health.¹⁰⁹ Muscular strength is defined as the ability of a muscle group to develop maximal contractile force against a resistance in a single contraction type (concentric, isometric, or eccentric).¹¹⁰⁻¹¹² Muscular endurance is the ability of a muscle group to exert submaximal force for extended periods.¹¹¹ Finally, muscle power refers to the ability of the neuromuscular system to produce the greatest possible impulse (ie, force per unit time).¹¹² The following section discusses how each of these components to musculoskeletal fitness are tested and how each is related to health.

More than 11 different body-region-specific classes of fitness test items (eg, upper-body strength, lower-body strength, abdominal endurance) were identified in the literature dating back to the mid 1970s. In addition, multiple variations of body-site-specific tests (eg, pull-ups, modified pull-ups, and isometric bent arm hang) have been developed.¹¹³⁻¹¹⁶ Many field-based muscular strength and power tests are not considered to be as physiologically

valid as muscle endurance fitness tests in youth because they require one to control the speed of a specific load through a specific number of repetitions.

Of 2642 reports initially screened for muscular strength, 63 satisfied the CDC search criteria. Out of the 63 studies, 23 were classified as experimental, 22 as experimental with no control, 12 as quasi-experimental, and 6 as longitudinal. The muscle endurance search screened 6563 reports, 38 of which satisfied the CDC search criteria. Out of the 38 studies, 12 were classified as experimental, 15 as experimental with no control, 6 as quasi-experimental, and 5 as longitudinal. The committee also considered studies that used field-based power tests that incorporated a single maximal effort at a submaximal speed and a load (vertical and horizontal jump) as they are moderately to strongly related to criterion measures of muscular strength.^{117,118} The following criteria were utilized to determine high-quality studies: (1) each study was evaluated against a set of criteria (eg, study design, representativeness of the population, and freedom from bias), and only those of high quality were reviewed further; (2) the evidence was categorized as direct or associational based on the strength of the study design and the rigor of the statistical analysis; and (3) the strength of the evidence was categorized as sufficient or insufficient based on the number of studies with direct or indirect evidence, the study designs, and the statistical significance of the association.

As noted by the IOM committee, currently, there is a lack of high-quality experimental studies that demonstrate a link between performance on any single musculoskeletal fitness test and health outcomes or markers in youth. However, research documenting a variety of health benefits stemming from resistance training programs in youth has emerged in the past 15 years.¹¹⁹⁻¹²¹ Benefits from resistance training programs include improved (1) body composition,^{106,117,122,123} (2) blood glucose and insulin regulation,^{122,124} (3) systemic arterial blood pressure in prehypertensives,^{106,125} (4) blood lipid and lipoprotein profiles,^{106,126} and (5) bone

health and management of arthritic pain and disability.¹¹⁸ Resistance training also has resulted in (6) an increase in functional capacity, (7) improved balance, (8) a decrease in falls, (9) improved quality of life, (10) self-efficacy,^{119,127} and (11) decreased levels of depression and anxiety^{123,128} into adulthood.

Selection of Tests

Emerging evidence noted a relationship between handgrip and standing long jump tests with health markers (ie, bone health and body composition) in youth.^{106,117,126,129,130} These 2 tests demonstrate acceptable reliability and concurrent validity with lower- and upper-body criterion strength measures while maintaining applicability across a broad age range in both sexes and in both normal and special pediatric subpopulations. Therefore, these 2 tests were summarily recommended for use in youth fitness testing.

The handgrip strength test has shown moderate to strong construct validity ($r = 0.52-0.84$) with established upper-body and lower-body strength tests (ie, 1-RM bench press and leg press, isometric kinetic knee extensor torque, respectively)^{131,132} and strong reliability ($r = 0.71-0.90$) in children and adolescents.¹³³⁻¹³⁵ There is also minimal test-related learning fatigue effects.¹¹ Furthermore, this test requires a minimal level of experience to perform, which diminishes the amount of practice needed to achieve an accurate depiction of a person's muscular skeletal strength and power.

The standing long jump has been used as a test of lower-body muscular strength, power, and explosive strength. In youth, this test demonstrates a moderate correlation with 1-RM leg press/body weight (0.39)¹³² and isokinetic quadriceps torque ($r = 0.50$)¹³¹ and a strong correlation with total-body isometric strength (0.77).¹³⁶ Additionally, the SLJ correlates strongly ($r = 0.70-0.91$) with other lower- and upper-extremity field-based power tests such as vertical jump, countermovement vertical jump, and upper-body explosive throw in youth (ages = 6-17 years) while

controlling for gender, age, and BMI or weight.^{132,136} Controlling for height and body mass with both the standing long jump and grip strength provides a more valid assessment of lower-body strength and power across all ages.^{132,136}

Other musculoskeletal tests such as the push-up, modified pull-up, curl up, and more recently, the isometric knee extension also have been examined in youth studies; however, there was limited evidence linking these types of tests to health markers.

Limitations

The vast majority of studies examined in the IOM review were not designed to answer questions about the specific test used and their effects on health markers. Many studies also were underpowered in relation to their ability to detect significant relationships, and many studies examined narrow age ranges. Furthermore, a majority of the studies failed to consider potential study confounders (eg, a combination of endurance, strength, and power exercises without a focus on a particular dimension), leaving the researchers to make indirect inferences relating to the test's ability to affect health markers. Finally, many studies were limited to isolated regions of the body (eg, modified pull-up focusing on upper-body strength), removing the ability to accurately infer results related to total-body musculoskeletal fitness levels.

Conclusions

The committee's review concluded that there currently exists a sufficient amount of evidence asserting that musculoskeletal fitness is related to health in humans. This conclusion was based on the importance of musculoskeletal fitness, especially muscle strength and power, to health markers in adults. Handgrip and standing long jump have been shown to have a growing base of evidence supporting their association to musculoskeletal fitness in youth. Overall, additional high-quality research is needed to link any of the aforementioned musculoskeletal tests to independent health risk factors in youth.^{126,137-143}

Resistance training in youth is safe¹⁴⁴⁻¹⁵⁰; however, training programs must be implemented with caution because children generally do not have experience with various types of resistance training. Resistance training in a youth population requires a properly designed program that is professionally supervised to ensure safety and optimal benefits for youth. A properly designed and supervised resistance training program can (1) enhance the muscular strength and power, (2) decrease cardiovascular risk, (3) improve motor skill performance, (4) contribute to enhanced sports performance, (5) increase a young athlete's resistance to sports-related injuries, and (6) help promote and develop exercise habits during childhood and adolescence.¹⁴⁵ It is recommended that youth resistance training programs focus on effectively manipulating one's own body mass (eg, push-ups, pull-ups, bodyweight squats, body weight lunges) before adding external resistance, specifically with consideration for developmental status.¹⁵¹

Future Implications

A global understanding of the predictive utility of physical fitness testing on health markers/outcomes in youth has not yet been fully demonstrated by the existing literature (ie, flexibility and musculoskeletal endurance). The following recommendations were designed to help fill the eminent need for understanding in this area and to help guide the design of future studies:

1. Well-designed research studies aimed at advancing understanding of the associations between fitness components and health in youth are needed.
2. Longitudinal studies should be conducted to provide empirical evidence concerning how health markers related to fitness track across the lifespan.
3. Randomized controlled trials and longitudinal studies should be designed to address the relationship between body composition and fitness tests and

the potential effects of modifying factors (ie, age, gender, body composition) on the relationship between a change in a health-related fitness component and health markers in specific populations.

4. National surveys of health-related physical fitness in youth should include measures of cardiometabolic health, bone health, psychological health and neurocognitive function.
5. Once a health marker is confirmed, research should be conducted to establish and validate health-related cut-points for that test that can be used across all ages in youth.^{10,144}

Finally, effective communication of test results is vital when the goal is improved fitness of the test taker, so that a child or adolescent is not derailed from a path of achieving a positive goal. Improved fitness should be one goal of any fitness intervention or program; however, across a continuous background of interpretation, communication that is intended to be encouraging could be interpreted as belittling and could derail a child from a path of future fitness. Understanding that a positive attitude should be displayed at all times and how to communicate an interpretation of an individual's test results in a manner that is appropriate to attaining achievable fitness goals is paramount to the success of sustainable, lifelong fitness.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article. ^(AJLM)

References

1. Kraus H, Hirschland RP. Minimum muscular fitness tests in school children. *Res Q*. 1954;25:178-188.
2. Kennedy JF. The soft American. *Sports Illustrated*. 1960;13:15-17.
3. Rodrigues LP, Stodden DF, Lopes VP. Developmental pathways of change in fitness and motor competence are related to overweight and obesity status at the end of primary school. *J Sci Med Sport*. 2016;19:87-92.
4. Rodrigues LP, Leitao R, Lopes VP. Physical fitness predicts adiposity longitudinal changes over childhood and adolescence. *J Sci Med Sport*. 2012;38:394-402.
5. Lopes VP, Maia JAR, Rodrigues LP, et al. Motor coordination, physical activity and fitness as predictors of longitudinal change in adiposity during childhood. *Eur J Sport Sci*. 2012;12:284-391.
6. Blair SN, Goodyear NN, Gibbons LW, Cooper KH. Physical fitness and incidence of hypertension in healthy normotensive men and women. *JAMA*. 1984;252(4):487-490.
7. Blair SN, Kohl HW, Barlow CE. Physical-activity, physical-fitness, and all-cause mortality in women: do women need to be active? *J Am Coll Nutr*. 1993;12:368-371.
8. Blair SN, Kohl HW, Barlow CE, Paffenbarger RS, Gibbons LW, Macera CA. Changes in physical-fitness and all-cause mortality: a prospective-study of healthy and unhealthy men. *JAMA*. 1995;273:1093-1098.
9. Pate R, Oria M, Pillsbury L, eds. *Fitness Measures and Health Outcomes in Youth*. Washington, DC: National Academies Press; 2012.
10. Institute of Medicine. *Fitness Measures and Health Outcomes in Youth*. Washington, DC: National Academies Press; 2012.
11. Ortega FB, Artero EG, Ruiz JR, et al; HELENA Study Group. Reliability of health-related physical fitness tests in European adolescents. The HELANA Study. *Int J Obes*. 2008;32(suppl 5):S49-S57.
12. Ruiz JR, Castro-Piñero J, Artero EG, et al. Predictive validity of health-related fitness in youth: a systematic review. *Br J Sports Med*. 2009;43:909-923.
13. Smith JJ, Eather N, Morgan PJ, Plotnikoff RC, Faigenbaum AD, Lubans DR. The health benefits of muscular fitness for children and adolescents: a systematic review and meta-analysis. *Sports Med*. 2014;44:1209-1223.
14. Lohman TG. Applicability of body composition techniques and constants for children and youths. *Exerc Sport Sci Rev*. 1985;14:325-357.
15. Lohman TG. Assessment of body composition in children. *Pediatr Exerc Sci*. 1989;1:19-30.
16. Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness and risk among children. *Pediatrics*. 2009;124(suppl 1):S23-S34.
17. Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics*. 1999;103:1175-1182.
18. Reilly JJ, Methven E, McDowell ZC, et al. Health consequences of obesity. *Arch Dis Child*. 2003;88:748-752.
19. Brown KM, McMahon RP, Biro FM, et al. Changes in self-esteem in black and white girls between the ages of 9 and 14 years. The NHLBI Growth and Health Study. *J Adolesc Health*. 1998;23:7-19.
20. Hesketh K, Wake M, Wake E. Body mass index and parent reported self-esteem in elementary school children: evidence for a causal relationship. *Int J Obes Relat Metab Disord*. 2004;28:1233-1237.
21. Franklin J, Deyner G, Steinbeck KS, Caterson ID, Hill AJ. Obesity and risk of low self-esteem: a statewide survey of Australian children. *Pediatrics*. 2006;118:2481-2487.
22. Southall J, Okely AD, Steele JR. Actual and perceived physical competence in overweight and non-overweight children. *Pediatr Exerc Sci*. 2004;16:15-24.
23. Okely AD, Booth ML, Patterson JW. Relationship of physical activity to fundamental movement skills among adolescents. *Med Sci Sports Exerc*. 2001;33:1899-1904.
24. Ulrich BD. Perceptions of physical competence, motor competence and participation in organized sport: their interrelationships in young children. *Res Q Exerc Sport*. 1987;58:57-67.
25. Smith E, Hay P, Campbell L, Trolor JN. A review of the association between obesity and cognitive function across the lifespan: implications for novel approaches to prevention and treatment. *Obes Rev*. 2011;12:740-755.
26. Carnethon MR, Gulati M, Greenland P. Prevalence and cardiovascular disease correlates of low cardiorespiratory fitness in adolescents and adults. *JAMA*. 2005;294:2981-2988.
27. Ortega FB, Ruiz JR, Castillo MJ, Sjöström M. Physical fitness in childhood and adolescence: a powerful marker of health. *Int J Obes*. 2008;32:1-11.
28. D'Hondt E, Van Hoorne G, Van Hoye K, Deforche B, De Bourdeaudhuij I, Lenoir M. *Relationship Between Motor Skill and BMI in Flemish Primary School Children: Proceedings of 12de VK Symposium*. Ghent, Belgium: Vereniging voor Kinesiologie (VK); 2007:14.
29. D'Hondt E, Segers V, Deforche B, et al. The role of vision in obese and normal-weight children's gait control. *Gait Posture*. 2011;33:179-184.
30. D'hondt E, Deforche B, Vaeyens R, et al. Gross motor coordination in relation to weight status and age in 5-to 12-year-old

- boys and girls: a cross-sectional study. *Int J Pediatr Obes*. 2011;6(2, pt 2):e556-e564.
31. D'Hondt E, Deforche B, Gentier I, et al. A longitudinal study of gross motor coordination and weight status in children. *Obesity*. 2014;22:1505-1511.
 32. Gentier I, D'Hondt E, De Bourdeaudhuij I, Deforche B, Lenoir M. Gross motor coordination, physical activity and weight status: a longitudinal analysis. Paper presented at: International congress on Enhancement of Physical Activity and Motor Skills (ICEPA 2012); November 1-3, 2012, Jyväskylä, Finland.
 33. Strutzenberger G, Richter A, Schneider M, Mündermann A, Schwameder H. Effects of obesity on the biomechanics of stair-walking in children. *Gait Posture*. 2011;34:119-125.
 34. Carnethon MR, Gidding SS, Nehgme R, Sidney S, Jacobs DR Jr, Liu K. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *JAMA*. 2003;290:3092-3100.
 35. Guo SS, Chumlea WC. Tracking of body mass index in children in relation to overweight in adulthood. *Am J Clin Nutr*. 1999;0:145s-148s.
 36. Yang X, Telama R, Leskinen E, Mansikkaniemi K, Viikari J, Raitakari OT. Testing a model of physical activity and obesity tracking from youth to adulthood: the cardiovascular risk in young Finns study. *Int J Obes*. 2007;31:521-527.
 37. Eissa MA, Dai S, Mihalopoulos NL, Day RS, Harrist RB, Labarthe DR. Trajectories of fat mass index, fat free-mass index, and waist circumference in children: Project HeartBeat! *Am J Prev Med*. 2009;37:S34-S39.
 38. Hammer LD, Wilson DM, Litt IF, et al. Impact of pubertal development on body fat distribution among white, Hispanic and Asian female adolescents. *J Pediatr*. 1991;118:975-980.
 39. Blum WF, Englaro P, Hanitsch S, et al. Plasma leptin levels in healthy children and adolescents: dependence on body mass index, body fat mass, gender, pubertal stage, and testosterone. *J Clin Endocrinol Metab*. 1997;82:2904-2910.
 40. Wang Y. Is obesity associated with early sexual maturation? A comparison of the association in American boys versus girls. *Pediatrics*. 2002;109:903-910.
 41. Malina RM. Tracking of physical activity and physical fitness across the lifespan. *Res Q Exerc Sport*. 1996;67(suppl 3):S-48.
 42. Malina RM, Katzmarzyk PT. Validity of the body mass index as an indicator of the risk and presence of overweight in adolescents. *Am J Clin Nutr*. 1999;70:131s-136s.
 43. Malina RM, Bouchard C, Bar-Or O. *Growth, Maturation, and Physical Activity*. Champaign, IL: Human Kinetics; 2004.
 44. Katzmarzyk PT, Shen W, Baxter-Jones A, et al. Adiposity in children and adolescents: correlates and clinical consequences of fat stored in specific body depots. *Pediatr Obes*. 2012;7:e42-e61.
 45. Schutte JE, Townsend EJ, Hugg JO, Shoup RF, Malina RM, Blomqvist CG. Density of lean body mass is greater in blacks than in whites. *J Appl Physiol*. 1984;56:1647-1649.
 46. Anderson SE, Whitaker RC. Prevalence of obesity among US preschool children in different racial and ethnic groups. *Arch Pediatr Adolesc Med*. 2009;163:344-348.
 47. Slaughter MH, Lohman TG, Boileau R, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol*. 1988;60:709-723.
 48. Deurenberg P, Pieters JJ, Hautvast JG. The assessment of the body fat percentage by skinfold thickness measurements in childhood and young adolescence. *Br J Nutr*. 1990;63:293-303.
 49. Safrit MJ. The validity and reliability of fitness tests for children: a review. *Pediatr Exerc Sci*. 1990;2:9-28.
 50. Harrison GG, Buskirk ER, Lindsay Carter JE, et al. Skinfold thicknesses and measurement technique. In: Lohman TG, Roche AF, Martorell R, eds. *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics; 1988:55-80.
 51. Goran MI, Allison DB, Poehlman ET. Issues relating to normalization of body fat content in men and women. *Int J Obes Relat Metab Disord*. 1995;19:638-643.
 52. Dietz WH, Bellizzi MC. Introduction: the use of body mass index to assess obesity in children. *Am J Clin Nutr*. 1999;70:123s-125s.
 53. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240.
 54. Malone SK, Zemel BS. Measurement and interpretation of body mass index during childhood and adolescence. *J Sch Nurs*. 2015;31:261-271.
 55. Taylor RW, Jones IE, Williams SM, Goulding A. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3-19 y. *Am J Clin Nutr*. 2000;72:490-495.
 56. Brambilla P, Bedogni G, Moreno LA, et al. Crossvalidation of anthropometry against magnetic resonance imaging for the assessment of visceral and subcutaneous adipose tissue in children. *Int J Obes*. 2006;30:23-30.
 57. Després JP, Nadeau A, Tremblay A, et al. Role of deep abdominal fat in the association between regional adipose tissue distribution and glucose tolerance in obese women. *Diabetes*. 1989;38:304-309.
 58. Goran MI, Gower BA. Abdominal obesity and cardiovascular risk in children. *Coron Artery Dis*. 1998;9:483-488.
 59. Lee S, Kuk JL, Kim Y, Arslanian SA. Measurement site of visceral adipose tissue and prediction of metabolic syndrome in youth. *Pediatr Diabetes*. 2001;12(3, pt 2):250-257.
 60. National Center of Health Statistics. 2000 CDC growth charts: United States. http://www.cdc.gov/growthcharts/cdc_charts.htm. Accessed November 18, 2015.
 61. American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*. Philadelphia, PA: Lippincott Williams & Wilkins; 2013.
 62. Faude O, Kerper O, Mulhaupt M, et al. Football to tackle overweight in children. *Scand J Med Sci Sports*. 2010;20(suppl 1):103-110.
 63. Serbescu C, Flora D, Hantiu I, Greene D, Benhamou CL, Courteix D. Effect of a six-month training programme on the physical capacities of Romanian schoolchildren. *Acta Paediatr*. 2006;95:1258-1265.
 64. Jones M, Stratton G, Reilly T, Unnithan V. The efficacy of exercise as an intervention to treat recurrent nonspecific low back pain in adolescents. *Pediatr Exerc Sci*. 2007;19:349-359.
 65. Ahlqwist A, Hagman M, Kjellby-Wendt G, Beckung E. Physical therapy treatment of back complaints on children and adolescents. *Spine*. 2008;33:E721-E727.
 66. Alter M. *Science of Flexibility*. 3rd ed. Champaign, IL: Human Kinetics; 2004.
 67. Tremblay MS, Shields M, Laviolette M, Craig CL, Janssen I, Gorber SC. Fitness of Canadian children and youth: results from the 2007-2009 Canadian Health Measures Survey. *Health Rep*. 2010;21:7-20.
 68. Welk GJ, Meredith MD, Ihmels M, Seeger C. Distribution of health-related physical fitness in Texas youth: a demographic and geographic analysis. *Res Q Exerc Sport*. 2010;81(suppl 3):S6-S15.
 69. McMillan CS, Erdmann LD. Tracking adiposity and health-related physical fitness test performances from early childhood through elementary school. *Pediatr Exerc Sci*. 2010;22:231-244.
 70. Saltin B. Oxygen transport by the circulatory system during exercise in man.

- In: Keul J, ed. *Limiting Factors of Physical Performance*. Stuttgart, Germany: Thieme; 1973:235-252.
71. Arraiz GA, Wigle DT, Mao Y. Risk assessment of physical-activity and physical-fitness in the Canada Health Survey mortality follow-up study. *J Clin Epidemiol*. 1992;45:419-428.
 72. Blair SN, Kohl HW, Paffenbarger RS, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: a prospective study of healthy men and women. *JAMA*. 1989;262:2395-2401.
 73. Sandvik L, Erikssen J, Thaulow E, Erikssen G, Mundal R, Rodahl K. Physical-fitness as a predictor of mortality among healthy, middle-aged Norwegian men. *N Engl J Med*. 1993;328:533-537.
 74. Colberg SR, Sigal RJ, Fernhall B, et al. Exercise and type 2 diabetes. *Diabetes Care*. 2010;33:e147-e167.
 75. Oliveria SA, Kohl HW, Trichopoulos D, Blair SN. The association between cardiorespiratory fitness and prostate cancer. *Med Sci Sports Exerc*. 1996;28:97-104.
 76. Eisenmann JC, Wickel EE, Welk GJ, Blair SN. Relationship between adolescent fitness and fatness and cardiovascular disease risk factors in adulthood: The Aerobics Center Longitudinal Study (ACLS). *Am Heart J*. 2005;149:46-53.
 77. Johnson MS, Figueroa-Colon R, Herd SL, et al. Aerobic fitness, not energy expenditure, influences subsequent increase in adiposity in black and white children. *Pediatrics*. 2000;106:e50.
 78. Byrd-Williams CE, Shaibi GQ, Sun P, et al. Cardiorespiratory fitness predicts changes in adiposity in overweight Hispanic boys. *Obesity*. 2008;16:1072-1077.
 79. Twisk JWR, Kemper HC, van Mechelen W. Tracking of activity and fitness and the relationship with cardiovascular disease risk factors. *Med Sci Sports Exerc*. 2000;32:1455-1461.
 80. Nourry C, Deruelle F, Guinhouya C, et al. High-intensity intermittent running training improves pulmonary function and alters exercise breathing pattern in children. *Eur J Appl Physiol*. 2005;94:415-423.
 81. Martins C, Santos R, Gaya A, Twisk J, Ribeiro J, Mota J. Cardiorespiratory fitness predicts later body mass index, but not other cardiovascular risk factors from childhood to adolescence. *Am J Hum Biol*. 2009;21:121-123.
 82. Farpour-Lambert NJ, Aggoun Y, Marchand LM, Martin XE, Herrmann F, Beghetti M. Physical activity reduces systemic blood pressure and improves early markers of atherosclerosis in prepubertal obese children. *J Am Coll Cardiol*. 2009;54:2396-2406.
 83. Reed KE, Warburton DER, Macdonald HM, Naylor PJ, McKay HA. Action Schools! BC: a school-based physical activity intervention designed to decrease cardiovascular disease risk factors in children. *Prev Med*. 2008;46:525-531.
 84. Ben Ounis O, Elloumi M, Makni E, et al. Exercise improves the ApoB/ApoA-I ratio, a marker of the metabolic syndrome in obese children. *Acta Paediatr*. 2010;99:1679-1685.
 85. Kelly AS, Wetzsteon RJ, Kaiser DR, Steinberger J, Bank AJ, Dengel DR. Inflammation, insulin, and endothelial function in overweight children and adolescents: the role of exercise. *J Pediatr*. 2004;145:731-736.
 86. Chang C, Liu W, Zhao X, Li S, Yu C. Effect of supervised exercise intervention on metabolic risk factors and physical fitness in Chinese obese children in early puberty. *Obes Rev*. 2008;9(suppl 1):135-141.
 87. Lee K, Shin Y, Jun T, Song W. Aerobic exercise training-induced decrease in plasma visfatin and insulin resistance in obese female adolescents. *Int J Sport Nutr Exerc Metab*. 2010;20:275-281.
 88. Sidiropoulou MP, Fotiadou EG, Tsimaras VK, Zakas AP, Angelopoulou NA. The effect of interval training in children with exercise-induced asthma competing in soccer. *J Strength Cond Res*. 2007;21:446-450.
 89. Stella SG, Vilar AP, Lacroix C, et al. Effects of type of physical exercise and leisure activities on the depression scores of obese Brazilian adolescent girls. *Braz J Med Biol Res*. 2005;38:1683-1689.
 90. Lobelo F, Pate RR, Dowda M, Liese AD, Ruiz JR. Validity of cardiorespiratory fitness criterion-referenced standards for adolescents. *Med Sci Sports Exerc*. 2009;41:1222-1229.
 91. Welk GJ, Laurson KR, Eisenmann JC, Cureton KJ. Development of youth aerobic-capacity standards using receiver operating characteristic curves. *Am J Prev Med*. 2011;41(4, suppl 2):S111-S116.
 92. Mota J, Ribeiro JC, Carvalho J, Santos MP, Martins J. Cardiorespiratory fitness status and body mass index change over time: a 2-year longitudinal study in elementary school children. *Int J Pediatr Obes*. 2009;4:338-342.
 93. Freedson PS, Cureton KJ, Heath GW. Status of field-based fitness testing in children and youth. *Prev Med*. 2000;31:S77-S85.
 94. Aires L, Mendonça D, Silva C, et al. A 3-year longitudinal analysis of changes in body mass index. *Int J Sports Med*. 2010;31:133-137.
 95. Chen JL, Wall D, Kennedy C, Unnithan V, Yeh CH. Predictors of increased body mass index in Chinese children. *Prog Cardiovasc Nurs*. 2007;22:138-144.
 96. Kim J, Must A, Fitzmaurice GM, et al. Relationship of physical fitness to prevalence and incidence of overweight among schoolchildren. *Obes Res*. 2005;13:1246-1254.
 97. Martins C, Santos R, Gaya A, Twisk J, Ribeiro J, Mota J. Cardiorespiratory fitness predicts later body mass index, but not other cardiovascular risk factors from childhood to adolescence. *Am J Hum Biol*. 2009;21:121-123.
 98. Ruiz JR, Castro-Piñero J, España-Romero V, et al. Field-based fitness assessment in young people: the alpha health-related fitness test battery for children and adolescents. *Br J Sports Med*. 2011;45:518-524.
 99. Artero EG, España-Romero V, Castro-Piñero J, et al. Reliability of field-based fitness tests in youth. *Int J Sports Med*. 2011;32:159-169.
 100. Boreham CA, Paliczka VJ, Nichols AK. A comparison of the PWC170 and 20-MST tests of aerobic fitness in adolescent schoolchildren. *J Sports Med Phys Fitness*. 1990;30:19-23.
 101. Liu NYS, Plowman SA, Looney MA. The reliability and validity of the 20-meter shuttle test in American students 12 to 15 years old. *Res Q Exerc Sport*. 1992;63:360-365.
 102. Watkins J, Ewing BG. Physical working capacity and mile run performance in adolescent boys. *Br J Sports Med*. 1983;17:188-192.
 103. Watson AWS, Odonovan DJ. Physical working capacity of male adolescents in Ireland. *Ir J Med Sci*. 1976;145:383-391.
 104. Ben Ounis O, Elloumi M, Makni E, et al. Exercise improves the ApoB/ApoA-I ratio, a marker of the metabolic syndrome in obese children. *Acta Paediatr*. 2010;99:1679-1685.
 105. Dunton GF, Schneider M, Cooper DM. An investigation of psychosocial factors related to changes in physical activity and fitness among female adolescents. *Psychol Health*. 2007;22:929-944.
 106. Janz KF, Dawson JD, Mahoney LT. Increases in physical fitness during childhood improve cardiovascular health during adolescence: The Muscatine Study. *Int J Sports Med*. 2002;23(suppl 1):S15-S21.
 107. McMurray RG, Bangdiwala SI, Harrell JS, Amorim LD. Adolescents with metabolic syndrome have a history of low aerobic fitness and physical activity levels. *Dyn Med*. 2008;7:5.

108. Ortega FB, Labayen I, Ruiz JR, et al. Improvements in fitness reduce the risk of becoming overweight across puberty. *Med Sci Sports Exerc.* 2001;43:1891-1897.
109. Bouchard C, Blair SN, Haskell WL. Why study physical activity and health? In: Bouchard C, Blair SN, Haskell WL, eds. *Physical Activity and Health.* Champaign, IL: Human Kinetics; 2007:3-19.
110. Farpour-Lambert NJ, Blimkie CJR. Muscle strength. In: Armstrong N, van Mechelen W, eds. *Paediatric Exercise Science and Medicine.* 2nd ed. Oxford, UK: Oxford University Press; 2008:37-53.
111. Kell RT, Bell G, Quinney A. Musculoskeletal fitness, health outcomes and quality of life. *Sports Med.* 2001;31:863-873.
112. Sale DG, Norman RW. Testing strength and power. In: Macdougall JD, Wenger HA, Green HJ, eds. *Physiological Testing of the Elite Athlete.* Ottawa, Canada: Mutual Press; 1982:7-37.
113. AAHPERD (American Alliance for Health, Physical Education, Recreation and Dance). *Health Related Physical Fitness Test: Technical Manual.* Reston, VA: AAHPERD; 1984.
114. Jackson AS. The evolution and validity of health-related fitness. *Quest.* 2006;58:160-175.
115. Morrow JR, Zhu W, Franks BD, Meredith MD, Spain C. 1958-2008: 50 Years of youth fitness tests in the United States. *Res Q Exerc Sport.* 2009;80:1-11.
116. Plowman SA. Muscular strength, endurance, and flexibility assessments. In: Welk GJ, Meredith MD, eds. *Fitnessgram/Activitygram Reference Guide.* Dallas, TX: Cooper Institute; 2008:11.11-11.40.
117. Ingle L, Sleaf M, Tolfrey K. The effect of a complex training and detraining programme on selected strength and power variables in early pubertal boys. *J Sports Sci.* 2006;24:987-997.
118. Heinonen A, Sievanen H, Kannus P, Oja P, Pasanen M, Vuori I. High-impact exercise and bones of growing girls: a 9-month controlled trial. *Osteoporosis International.* 2000;11:1010-1017.
119. McCartney N, Phillips S. Physical activity, muscular fitness, and health. In: Bouchard C, Blair SN, Haskell WL, eds. *Physical Activity and Health.* Champaign, IL: Human Kinetics; 2007:231-246.
120. Reid KF, Fielding RA. Skeletal muscle power: a critical determinant of physical functioning in older adults. *Exerc Sport Sci Rev.* 2012;40:4-12.
121. Williams MA, Haskell WL, Ades PA, et al; American Heart Association Council on Clinical Cardiology, American Heart Association Council on Nutrition, and Physical Activity, and Metabolism. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: A scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation.* 2007;116:572-584.
122. Benson AC, Torode ME, Fiatarone Singh MA. The effect of high-intensity progressive resistance training on adiposity in children: a randomized controlled trial. *Int J Obes (Lond).* 2008;32:1016-1027.
123. Lubans DR, Aguiar EJ, Callister R. The effects of free weights and elastic tubing resistance training on physical self-perception in adolescents. *Psychol Sport Exerc.* 2010;11:497-504.
124. Shaibi GQ, Cruz ML, Ball GDC, et al. Effects of resistance training on insulin sensitivity in overweight Latino adolescent males. *Med Sci Sports Exerc.* 2006;38:1208-1215.
125. Naylor LH, Watts K, Sharpe JA, et al. Resistance training and diastolic myocardial tissue velocities in obese children. *Med Sci Sports Exerc.* 2008;40:2027-2032.
126. van der Heijden GJ, Wang ZYJ, Chu ZL, et al. Strength exercise improves muscle mass and hepatic insulin sensitivity in obese youth. *Med Sci Sports Exerc.* 2010;42:1973-1980.
127. Garber CE, Blissmer B, Deschenes MR, et al; American College of Sports Medicine. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43:1334-1359.
128. Velez A, Golem DL, Arent SM. The impact of a 12-week resistance training program on strength, body composition, and self-concept of Hispanic adolescents. *J Strength Cond Res.* 2010;24:1065-1073.
129. Minck MR, Ruitter LM, Van Mechelen W, Kemper HCG, Twisk JWR. Physical fitness, body fatness, and physical activity: The Amsterdam Growth and Health Study. *Am J Hum Biol.* 2000;2:593-599.
130. Morris FL, Naughton GA, Gibbs JL, Carlson JS, Wark JD. Prospective ten-month exercise intervention in premenarcheal girls: positive effects on bone and lean mass. *J Bone Miner Res.* 1997;2:1453-1462.
131. Holm I, Fredriksen P, Fosdahl M, Vollestad N. A normative sample of isotonic and isokinetic muscle strength measurements in children 7 to 12 years of age. *Acta Paediatr.* 2008;97:602-607.
132. Milliken LA, Faigenbaum AD, Loud RL, Westcott WL. Correlates of upper and lower body muscular strength in children. *J Strength Cond Res.* 2008;22:1339-1346.
133. Benefice E, Fouere T, Malina RM. Early nutritional history and motor performance of Senegalese children, 4-6 years of age. *Ann Hum Biol.* 1999;26:443-455.
134. Brunet M, Chaput JP, Tremblay A. The association between low physical fitness and high body mass index or waist circumference is increasing with age in children: The "Québec en Forme" project. *Int J Obes.* 2007;31:637-643.
135. Ruiz JR, Ortega FB, Gutierrez AM, Sjöström DM, Castillo MJ. Health-related fitness assessment in childhood and adolescence: a European approach based on the AVENA, EYHS and HELENA studies. *J Public Health.* 2006;14:269-277.
136. Castro-Piñero J, Artero EG, España-Romero V, et al. Criterion-related validity of field-based fitness tests in youth: a systematic review. *Br J Sports Med.* 2010;44:934-943.
137. Pate RR, Burgess ML, Woods JA, Ross JG, Baumgartner T. Validity of field-tests of upper-body muscular strength. *Res Q Exerc Sport.* 1993;64:17-24.
138. Engelman ME, Morrow JR Jr. Reliability and skinfold correlates for traditional and modified pull-ups in children grades 3-5. *Res Q Exerc Sport.* 1991;62:88-91.
139. Erbaugh SJ. Reliability of physical fitness tests administered to young children. *Percept Mot Skills.* 1990;71(3, pt 2):1123-1128.
140. Kollath JA, Safrit MJ, Zhu W, Gao IG. Measurement errors in modified pull-ups testing. *Res Q Exerc Sport.* 1991;62:432-435.
141. Saint Romain B, Mahar MT. Norm-referenced and criterion-referenced reliability of the push-up and modified pull-up. *Meas Phys Educ Exerc Sci.* 2001;5:67-80.
142. Farpour-Lambert NJ, Blimkie CJR. Muscle strength. In: Armstrong N, van Mechelen W eds. *Paediatric Exercise Science and Medicine.* 2nd ed. Oxford, UK: Oxford University Press; 2008:37-53.
143. Witzke KA, Snow CM. Effects of plyometric jump training on bone mass in adolescent girls. *Med Sci Sports Exerc.* 2000;32:1051-1057.
144. Faigenbaum AD, Kraemer WJ, Blimkie CJ, et al. Youth resistance training:

- updated position statement paper from the national strength and conditioning association. *J Strength Cond Res.* 2009;23:S60-S79.
145. Faigenbaum AD, Farrell A, Fabiano M, et al. Effects of integrative neuromuscular 12 training on fitness performance in children. *Pediatr Exerc Sci.* 2011;23:573-584.
146. Faigenbaum AD, Farrell AC, Fabiano M, et al. Effects of detraining on fitness 7 performance in 7-year-old children. *J Strength Cond Res.* 2013;27:323-330.
147. Department of Health and Human Services. *Physical Activity Guidelines for Americans.* Washington, DC: Department of Health and Human Services; 2008.
148. National Association for Sport and Physical Education. *Physical Education for Lifelong Fitness.* 2nd ed. Champaign, IL: Human Kinetics; 2005.
149. Myer GD, Faigenbaum AD, Chu DA, et al. Integrative training for children and adolescents: techniques and practices for reducing sports related injuries and enhancing athletic performance. *Phys Sports Med.* 2011;39:74Y84.
150. Myer GD, Faigenbaum AD, Ford KR, Best TM, Bergeron MF, Hewett TE. When to initiate integrative neuromuscular training to reduce sports-related injuries in youth? *Curr Sports Med Rep.* 2011;10:155.
151. Stodden DF, Brooks T. Promoting musculoskeletal fitness in youth: performance and health implication from a developmental perspective. *Strength Cond J.* 2013;35:54-62.