

A Series Expansion for the First Positive Zero of the Bessel Functions

By R. Piessens

Abstract. It is shown that the first positive zero $j_{\nu,1}$ of the Bessel function $J_\nu(x)$ is given by

$$j_{\nu,1} = 2(\nu + 1)^{1/2} \left[1 + \frac{(\nu + 1)}{4} - \frac{7(\nu + 1)^2}{96} + \frac{49(\nu + 1)^3}{1152} - \frac{8363(\nu + 1)^4}{276480} + \dots \right]$$

for $-1 < \nu < 0$.

1. It is well known that, when ν is real and $\nu > -1$, the Bessel function $J_\nu(x)$ has an infinite number of zeros and that all zeros are real (Watson [9]). We denote the s th positive zero of $J_\nu(x)$ by $j_{\nu,s}$.

Several approximations, asymptotic expansions or bounds for the zeros of Bessel functions exist (see [1], [2], [4], [6], [7], [9]). Especially McMahon's expansion for large zeros (see Abramowitz and Stegun [1]), Olver's asymptotic expansion for large orders and Olver's uniform asymptotic expansions (see Olver [6]) are interesting formulas, but, unfortunately, they are not applicable when s and ν are small. The purpose of this note is to give a series expansion for $j_{\nu,1}$ when $-1 < \nu < 0$.

2. Cayley [3] noticed that Graeffe's method for solving a polynomial equation can be applied for the efficient computation of

$$(1) \quad \sum_{s=1}^{\infty} j_{\nu,s}^{-2r} \equiv \sigma_\nu^{(r)}, \quad r = 1, 2, \dots$$

An upper bound for $j_{\nu,1}$ is given by Chambers [4]:

$$(2) \quad j_{\nu,1} < (\nu + 1)^{1/2} [(\nu + 2)^{1/2} + 1].$$

Further it is known that, when $k > 1$,

$$(3) \quad \lim_{\nu \rightarrow -1} j_{\nu,k} = j_{1,k-1} > 0.$$

Thus, the first term in the left side of (1) is dominant when $\nu \approx -1$, so that

$$(4) \quad j_{\nu,1} = 2(\nu + 1)^{1/2} \phi_r(\nu) + o((\nu + 1)^{r-1}), \quad \nu \rightarrow -1,$$

where

$$(5) \quad \phi_r(\nu) = \left[\frac{1}{2^{2r}(\nu + 1)^r \sigma_\nu^{(r)}} \right]^{1/2r}$$

is analytic at $\nu = -1$.

Received November 22, 1982.

1980 *Mathematics Subject Classification.* Primary 33A40, 49G10.

©1984 American Mathematical Society
 0025-5718/84 \$1.00 + \$.25 per page

By approximating $\phi_r(\nu)$ by a Taylor polynomial, we obtain

$$(6) \quad j_{\nu,1} = 2(\nu + 1)^{1/2} \sum_{k=0}^{r-1} C_k (\nu + 1)^k + o((\nu + 1)^{r-1}), \quad \nu \rightarrow -1,$$

where

$$(7) \quad C_k = \frac{1}{k!} \left. \frac{d^k \phi_r(\nu)}{d\nu^k} \right|_{\nu=-1}$$

is independent of r .

When $r \rightarrow \infty$, (6) becomes a series expansion for $j_{\nu,1}$, which, because of the presence of a branchpoint of $\phi_r(\nu)$ at $\nu = -2$, converges only in the interval $-1 < \nu < 0$.

Using REDUCE, which is a computer language for formula manipulation [5], we have computed C_k , $k = 0, 1, 2, 3, 4$, using (7), where $r = 8$ and

$$(8) \quad \phi_8(\nu) = \left[\frac{(\nu + 2)^4(\nu + 3)^2(\nu + 4)^2(\nu + 5)(\nu + 6)(\nu + 7)(\nu + 8)}{429\nu^5 + 7640\nu^4 + 53752\nu^3 + 185430\nu^2 + 311387\nu + 202738} \right]^{1/16}.$$

The result is

$$(9) \quad j_{\nu,1} = 2(\nu + 1)^{1/2} \left[1 + \frac{(\nu + 1)}{4} - \frac{7(\nu + 1)^2}{96} + \frac{49(\nu + 1)^3}{1152} - \frac{8363(\nu + 1)^4}{276480} + \dots \right].$$

In Table 1, we compare the exact values of $j_{\nu,1}$ with the approximate values given by (9), for $\nu = -3/4, -2/3, -1/2, -1/3, -1/4$ and also for $\nu = 0$ (although we were not able to prove the convergences of the expansion for $\nu = 0$).

TABLE 1
Values of the first zero $j_{\nu,1}$ of $J_\nu(x)$

ν	exact	approximation (9)
$-3/4$	1.058508	1.058489
$-2/3$	1.243046	1.242958
$-1/2$	1.570796	1.570056
$-1/3$	1.866351	1.863061
$-1/4$	2.006300	2.000273
0	2.404826	2.378740

3. An interesting application of (9) is the estimation of the smallest zero of Laguerre- and Gegenbauer-polynomials [8]. For example, the smallest zero ξ_n of the Laguerre polynomials $L_n^{(\alpha)}(x)$ is approximated by (see Tricomi [8])

$$(10) \quad \xi_n \approx \frac{j_{\alpha,1}}{4k_n} \left[1 + \frac{2(\alpha^2 - 1) + j_{\alpha,1}^2}{48k_n^2} \right],$$

where $k_n = n + (\alpha + 1)/2$. In Table 2, this approximation, where $j_{\alpha,1}$ is replaced by (9), is compared with the exact value of ξ_n .

TABLE 2
 Values of the smallest zero of $L_n^{(\alpha)}(x)$

α	n	exact	approximation (10)
- 3/4	3	0.089682	0.089679
	15	0.018520	0.018519
- 1/2	3	0.190163	0.189982
	15	0.040452	0.040415
- 1/4	3	0.299347	0.297530
	15	0.065463	0.065071
0	3	0.415775	0.406686
	15	0.093308	0.091294

Acknowledgement. This research is supported by the 'Nationaal Fonds voor Wetenschappelijk Onderzoek (Belgium)'.

Department of Computer Science
 University of Leuven
 Celestijnenlaan 200 A
 B-3030 Heverlee, Belgium

1. M. ABRAMOWITZ & I. STEGUN, *Handbook of Mathematical Functions*, Dover, New York, 1965.
2. M. BRANDERS, R. PIESSENS & M. DE MEUE, "Rational approximations for zeros of Bessel functions," *J. Comput. Phys.*, v. 42, 1981, pp. 403-405.
3. A. CAYLEY, "Addition to Lord Rayleigh's paper: On the numerical calculation of the roots of fluctuating functions," *Proc. London Math. Soc.*, v. 5, 1873-1874, pp. 123-124.
4. L. G. CHAMBERS, "An upper bound for the first zero of Bessel functions," *Math. Comp.*, v. 38, 1982, pp. 589-591.
5. A. C. HEARN, REDUCE 2—*A System and Language for Algebraic Manipulation*, Proc. Second Sympos. on Symbolic and Algebraic Manipulation, Los Angeles, 1971, pp. 128-133.
6. F. W. J. OLVER, *Royal Society Mathematical Tables*, Vol. 7, *Bessel Functions. Part III. Zeros and Associated Values*, The University Press, Cambridge, 1960.
7. N. M. TEMME, "An algorithm with ALGOL 60 program for the computation of the zeros of ordinary Bessel functions and those of their derivatives," *J. Comput. Phys.*, v. 32, 1979, pp. 270-279.
8. F. G. TRICOMI, *Vorlesungen über Orthogonalreihen*, Springer-Verlag, Berlin, 1955.
9. G. N. WATSON, *A Treatise on the Theory of Bessel Functions*, The University Press, Cambridge, 1944.